

1.1 A Brief Introduction to Differential Geometry

We can start by giving the most basic definitions that will be used throughout all the thesis. One of the most important one is the smooth manifold, but first we need to understand what a manifold is and why it is so important in physics.

Definition 1.1 (Topological manifold). *A topological manifold of dimension n is a topological space M that is Hausdorff, second countable, and locally Euclidean of dimension n : every point $p \in M$ has an open neighborhood homeomorphic to an open subset of $\mathbb{R}^n$. [1]*

Definition 1.2 (Chart, atlas). *A chart on M is a pair (U, φ) , with $U \subseteq M$ open and $\varphi : U \rightarrow \varphi(U) \subseteq \mathbb{R}^n$ a bijection onto an open subset. Two charts $(U, \varphi), (V, \psi)$ are C^∞ -compatible if the transition map*

$$\psi \circ \varphi^{-1} : \varphi(U \cap V) \rightarrow \psi(U \cap V)$$

is a C^∞ -diffeomorphism whenever $U \cap V \neq \emptyset$. An atlas is a collection of pairwise compatible charts covering M . The union of all the atlases on M is the maximal atlas.

Definition 1.3 (Smooth manifold). *A smooth manifold is a pair $(M, \mathcal{A})$, where M is a topological manifold and $\mathcal{A}$ is a maximal atlas on M (the smooth structure). We write $\dim M = n$.*

Definition 1.4 (Smooth map, diffeomorphism). *Let M^m, N^n be smooth manifolds. A map $f : M \rightarrow N$ is smooth at $p \in M$ if there exist charts (U, φ) around p and (V, ψ) around $f(p)$, with $f(U) \subseteq V$, such that*

$$\psi \circ f \circ \varphi^{-1} : \varphi(U) \rightarrow \psi(V)$$

is C^∞ in the usual sense on $\mathbb{R}^m$. f is smooth if smooth at every point. f is a diffeomorphism if it is a smooth bijection with smooth inverse. We write $C^\infty(M) := C^\infty(M, \mathbb{R})$.

The concept of manifold itself doesn't carry a vector space structure. Despite that we can always fix a point and define a tangent vector as an object with the same properties as a derivation. This creates a vector space at every point of M called *tangent space*.

Definition 1.5 (Tangent vector, tangent space). *Fix $p \in M$. A tangent vector at p is a linear map $v : C^\infty(M) \rightarrow \mathbb{R}$ satisfying the Leibniz rule*

$$v(fg) = f(p)v(g) + g(p)v(f), \quad f, g \in C^\infty(M).$$

The set of tangent vectors at p , with the natural vector space structure, is the tangent space $T_p M$.

Remark 1.6. Equivalently, a curve $\gamma : (-\varepsilon, \varepsilon) \rightarrow M$ with $\gamma(0) = p$ induces the derivation $v_\gamma(f) := \left. \frac{d}{dt} \right|_{t=0} f(\gamma(t))$, and every $v \in T_p M$ can be derived in this way. It can be proved that two curves induce the same tangent vector iff their local representatives agree to first order at p ; this equivalent identification $T_p M \cong \{\text{curves through } p\} / \sim$ is used later (e.g. for the flows ϕ_t^H).

Sometimes it is convenient to switch to a notation more explicit than the global description really implicit used most of the times, for that we can write our objects on manifolds in coordinates.

Definition 1.7 (Coordinate basis). *Given a chart $(U, \varphi = (x^1, \dots, x^n))$ around p , the vectors*

$$\left. \frac{\partial}{\partial x^i} \right|_p \in T_p M, \quad \left. \frac{\partial}{\partial x^i} \right|_p (f) := \left. \frac{\partial (f \circ \varphi^{-1})}{\partial x^i} \right|_{\varphi(p)}$$

form a basis of $T_p M$; hence $\dim T_p M = n$, and $v = v^i \partial_i|_p$ (using the Einstein summation notation).

Definition 1.8 (Tangent bundle). *The tangent bundle of M is*

$$TM := \bigsqcup_{p \in M} T_p M,$$

with projection $\pi : TM \rightarrow M$, $\pi(v) = p$ for $v \in T_p M$. A chart $(U, \varphi = (x^1, \dots, x^n))$ on M induces a chart on $\pi^{-1}(U)$ via $v = v^i \partial_i|_p \mapsto (x^1(p), \dots, x^n(p), v^1, \dots, v^n)$, giving TM a smooth structure of dimension $2n$ and making $\pi : TM \rightarrow M$ a rank- n vector bundle.

Definition 1.9 (Vector field). A vector field on M is called a smooth section of $\pi : TM \rightarrow M$, if it is a smooth map $X : M \rightarrow TM$ with $\pi \circ X = \text{id}_M$; equivalently, $p \mapsto X_p \in T_p M$ smooth in p . Equivalently again, X is a derivation of the $\mathbb{R}$ -algebra $C^\infty(M)$: linear, with $X(fg) = fX(g) + gX(f)$. We denote by $\mathfrak{X}(M)$ the $C^\infty(M)$ -module of vector fields on M .

Definition 1.10 (Cotangent space). The cotangent space at p is the dual vector space $T_p^* M := (T_p M)^*$. For $f \in C^\infty(M)$, its differential at p is the covector

$$df_p \in T_p^* M, \quad df_p(v) := v(f), \quad v \in T_p M.$$

In the coordinate basis, $\{dx^1|_p, \dots, dx^n|_p\}$ is dual to $\{\partial/\partial x^i|_p\}$, and $df_p = \frac{\partial f}{\partial x^i} \Big|_p dx^i|_p$.

Definition 1.11 (Cotangent bundle). The cotangent bundle is

$$T^* M := \bigsqcup_{p \in M} T_p^* M,$$

with projection $\pi : T^* M \rightarrow M$ and the induced smooth structure of dimension $2n$ (local coordinates (x^i, p_i)), making $T^* M \rightarrow M$ a rank- n vector bundle. A smooth section $\alpha : M \rightarrow T^* M$ is a differential 1-form; the space of 1-forms is $\Omega^1(M)$.

Definition 1.12 (Pushforward / Differential). Let $F : M \rightarrow N$ be a smooth map between smooth manifolds. For each point $p \in M$, the pushforward (or differential) of F at p is the linear map

$$F_* : T_p M \rightarrow T_{F(p)} N$$

defined as follows: for any tangent vector $X \in T_p M$, its image $F_* X \in T_{F(p)} N$ is the derivation that operates on any smooth function $f \in C^\infty(N)$ by

$$(F_* X)(f) = X(f \circ F).$$

Alternatively, using the geometric description of tangent vectors as equivalence classes of curves, let $\gamma : (-\epsilon, \epsilon) \rightarrow M$ be a smooth curve such that $\gamma(0) = p$ and $\gamma'(0) = X$. Then the pushforward $F_* X$ is given by the velocity vector of the composition curve $F \circ \gamma$ at $t = 0$:

$$F_* X = \frac{d}{dt} \Big|_{t=0} (F \circ \gamma)(t).$$

The pushforward map is also frequently denoted by dF_p .

Definition 1.13 (Pullback). Let $F : M \rightarrow N$ be a smooth map between smooth manifolds. The pullback operator F^* is defined on different classes of fields as follows:

1. **Smooth Functions:** For any smooth function $f \in C^\infty(N)$, the pullback $F^* f \in C^\infty(M)$ is defined by precomposition:

$$F^* f = f \circ F.$$

2. **Covectors:** For any point $p \in M$ and any cotangent vector (covector) $\omega \in T_{F(p)}^* N$, the pullback $F^* \omega \in T_p^* M$ is defined by its action on any tangent vector $X \in T_p M$:

$$(F^* \omega)(X) = \omega(F_* X).$$

3. **Differential Forms:** Let $\Omega \in \otimes^k(N)$ be a smooth differential k -form on N . The pullback $F^* \Omega$ is a smooth differential k -form on M defined pointwise for each $p \in M$ and all tangent vectors $X_1, \dots, X_k \in T_p M$ by

$$(F^* \Omega)_p(X_1, \dots, X_k) = \Omega_{F(p)}(F_* X_1, \dots, F_* X_k).$$

Definition 1.14 (Flow of a Vector Field). Let X be a smooth vector field on a smooth manifold M . An integral curve of X is a smooth curve $\gamma : I \rightarrow M$, where $I \subset \mathbb{R}$ is an open interval, such that $\gamma'(t) = X(\gamma(t))$ for all $t \in I$.

The local flow (or simply flow) of X is a smooth map $\theta : \mathcal{D} \rightarrow M$, where $\mathcal{D} \subset \mathbb{R} \times M$ is an open neighborhood of $\{0\} \times M$ known as the flow domain, satisfying the following properties:

1. For each $p \in M$, the slice $\mathcal{D}^{(p)} = \{t \in \mathbb{R} : (t, p) \in \mathcal{D}\}$ is an open interval containing 0, and the curve $\theta^{(p)} : \mathcal{D}^{(p)} \rightarrow M$ defined by $\theta^{(p)}(t) = \theta(t, p)$ is the unique maximal integral curve of X satisfying the initial condition $\theta(0, p) = p$. That is,

$$\frac{\partial \theta}{\partial t}(t, p) = X(\theta(t, p)).$$

2. **One-Parameter Group Property:** If we write $\theta_t(p) = \theta(t, p)$, then for any $s \in \mathcal{D}^{(p)}$ and $t \in \mathcal{D}^{(\theta_s(p))}$ such that $t + s \in \mathcal{D}^{(p)}$, we have

$$\theta_t(\theta_s(p)) = \theta_{t+s}(p).$$

If $\mathcal{D} = \mathbb{R} \times M$, the vector field X is called complete, and the maps $\theta_t : M \rightarrow M$ constitute a global one-parameter group of diffeomorphisms.

1.2 Lie Algebras and Representations

In this chapter, $\mathbb{K}$ denotes an arbitrary commutative field.

Definition 1.15 (Lie algebra). A Lie algebra over $\mathbb{K}$ is a $\mathbb{K}$ -vector space $\mathfrak{g}$ equipped with a bilinear map $[\cdot, \cdot] : \mathfrak{g} \times \mathfrak{g} \rightarrow \mathfrak{g}$ satisfying

$$[X, Y] = -[Y, X], \quad \text{antisymmetry} \quad (1.1)$$

$$[[X, Y], Z] + [[Y, Z], X] + [[Z, X], Y] = 0 \quad \text{Jacobi identity} \quad (1.2)$$

for all $X, Y, Z \in \mathfrak{g}$. [2]

Remark 1.16. Every associative algebra $(\mathcal{A}, \cdot)$ carries a canonical Lie algebra structure via the commutator $[a, b] := ab - ba$. It satisfy trivially the antisymmetry and Jacobi identity via:

$$[[a, b], c] = (abc - bac) - c(ab - ba) = abc - bac - cab + cba + bca - bca + acb - acb = -[[b, c], a] - [[c, a], b].$$

The Jacobi identity is used in a Lie algebra to mimic the chain property of a derivation. For example, this allows $\mathfrak{g}$ to perfectly describe infinitesimal calculus on manifolds.

Example 1.17. One of the most important examples of a Lie algebra is $\mathfrak{gl}(V)$, the space of all linear endomorphisms of a vector space V , equipped with the commutator $[A, B] = AB - BA$ as Lie bracket. When $V = \mathbb{C}^n$, $\mathfrak{gl}(V) \cong \mathfrak{gl}(n, \mathbb{C})$ can be identified with $M(n, \mathbb{C})$. A natural basis, denoted E_{ij} for $i, j = 1, \dots, n$, is given by the *matrix units*, i.e. the matrix with entry 1 in position (i, j) and 0 elsewhere, satisfying

$$[E_{ij}, E_{kl}] = \delta_{jk}E_{il} - \delta_{li}E_{kj}.$$

Example 1.18. Another example is $\mathfrak{su}(n)$. It consist of the $M(n, \mathbb{C})$ matrices which are traceless and anti-hermitian, this makes the dimension of the resulting Lie algebra $n^2 - 1$. In two dimensions, the basis of $\mathfrak{su}(2)$, consists of the Pauli operators σ_i which $i = x, y, z$ which are used for representing the spin of a fermion in quantum physics. In three dimensions, the basis is formed by the eight Gell-Mann matrices λ_a used in quantum chromodynamics.

Now, let us recall some basic concepts and fundamentals in the Lie theory. In general, an *homomorphism* is an application between the same mathematical structures that preserves the inner operations. When, in particular, the structure is a Lie algebra we can use the following definition.

Definition 1.19 (Homomorphism of Lie Algebras). Let $\mathfrak{g}$ and $\mathfrak{h}$ be Lie algebras over the same field $\mathbb{K}$. A linear map $\phi : \mathfrak{g} \rightarrow \mathfrak{h}$ is a Lie algebra homomorphism if it preserves the Lie bracket:

$$\phi([X, Y]) = [\phi(X), \phi(Y)] \quad \forall X, Y \in \mathfrak{g}.$$

Definition 1.20 (Isomorphism of Lie Algebras). An isomorphism is a Lie algebra homomorphism $\phi : \mathfrak{g} \rightarrow \mathfrak{h}$ that is a bijective linear map.

1.2.1 Lie Groups

Definition 1.21 (Lie Group). A Lie Group is a smooth manifold G together with a smooth map from $G \times G \rightarrow G$ that makes G into a group and such that the inverse map $g \rightarrow g^{-1}$ is a smooth map $G \rightarrow G$.

Also in this case we can specialize the definition of homomorphism in the case of Lie groups.

Definition 1.22 (Homomorphism of Lie Groups). A map $F : G \rightarrow H$ where G and H are Lie groups is called an homomorphism if it is a smooth map and an homomorphism between groups.

Let's look now at the tangent space of a Lie group. We can define an operation $L_g : G \rightarrow G$ by $L_g(h) = g \circ h$ called the "left multiplication" which is smooth. So the differential dL_g at the point h will be a linear map from $T_h(G)$ to $T_{gh}(G)$.

Now we can take a vector field X on G , and that is called *left-invariant* if

$$dL_g(X_h) = X_{gh} \quad \forall h \in G.$$

It is also true that for any given vector $v \in T_e(G)$ there is a unique left-invariant vector field X^v . It can be constructed starting from the vector tangent to the identity equal to v , so $X_e^v = v$ then we can use L_g to move it and construct the remaining vector space:

$$X_g^v = dL_g(X_e^v) = dL_g(v),$$

it can be shown easily that is left-invariant. Recall that if we think of vector fields as first-order differential operators, then the commutator of two vector fields is, again, a vector field. It is not difficult to show that the commutator of two left-invariant vector fields is, again, a left-invariant vector field.

Definition 1.23 (Lie algebra of a Lie group). The Lie algebra $\mathfrak{g}$ of a Lie group G is the tangent space at the identity with the bracket operation defined by

$$[v, w] := [X^v, X^w]_e.$$

We now show that every left-invariant vector field is complete, so that it integrates to a global flow.

Proposition 1.24 (Completeness of left-invariant vector fields). Let $X^v \in \mathfrak{X}(G)$ be the left-invariant vector field associated to $v \in \mathfrak{g} = T_e G$. Then X^v is complete: its maximal integral curve through e is defined for all $t \in \mathbb{R}$.

Proposition 1.25 (One-parameter subgroup). Let $\gamma_v : \mathbb{R} \rightarrow G$ be the integral curve of X^v through e . Then

$$\gamma_v(t+s) = \gamma_v(t)\gamma_v(s) \quad \forall t, s \in \mathbb{R},$$

γ_v is a Lie group homomorphism $\mathbb{R} \rightarrow G$ (a one-parameter subgroup).

Definition 1.26 (Exponential map). The exponential map of G is

$$\exp : \mathfrak{g} \rightarrow G, \quad \exp(v) := \gamma_v(1).$$

Proposition 1.27 (Basic properties of $\exp$). It can be shown that

1. $\exp(tv) = \gamma_v(t)$ for all $t \in \mathbb{R}$ (from $\gamma_{tv}(s) = \gamma_v(ts)$).
2. $\exp(0) = e$ and $d(\exp)_0 = \text{id}_{\mathfrak{g}}$; by the inverse function theorem $\exp$ restricts to a diffeomorphism from a neighborhood of $0 \in \mathfrak{g}$ onto a neighborhood of $e \in G$.
3. *Naturality*: if $F : G \rightarrow H$ is a homomorphism of Lie groups with differential $dF_e : \mathfrak{g} \rightarrow \mathfrak{h}$, then

$$F(\exp_G(v)) = \exp_H(dF_e(v)), \quad v \in \mathfrak{g}.$$

Remark 1.28. Global surjectivity of $\exp$ is not automatic in general (fails e.g. for $SL(2, \mathbb{R})$), but holds for G compact and connected — which covers every group used in this thesis ($SU(2)$, $U(n)$).

Remark 1.29 (Matrix Lie groups). For $G \subseteq GL(n, \mathbb{C})$ a closed subgroup $\mathfrak{g} \subset \mathfrak{gl}(n, \mathbb{C}) = M_n(\mathbb{C})$ is identified with a linear subspace of matrices via $T_e G \subset T_e GL(n, \mathbb{C}) \cong M_n(\mathbb{C})$, and left multiplication is literally matrix multiplication: $L_g(h) = gh$, so $dL_g(v) = gv$ for $v \in \mathfrak{g}$. The left-invariant vector field associated to v is therefore $X_g^v = gv$, and its integral curve through e solves the linear matrix ODE

$$\dot{\gamma}(t) = \gamma(t)v, \quad \gamma(0) = I,$$

whose unique solution is the matrix exponential series

$$\gamma(t) = e^{tv} := \sum_{k=0}^{\infty} \frac{(tv)^k}{k!}.$$

Hence $\exp(v) = e^v$: the abstract exponential map of Definition 1.26 coincides with the ordinary matrix exponential whenever G is realized as a matrix group. In particular

$$\mathfrak{g} = \{v \in M_n(\mathbb{C}) : e^{tv} \in G \ \forall t \in \mathbb{R}\},$$

which is often the fastest way to identify $\mathfrak{g}$ in practice, without passing through tangent vectors to curves at all.

Example 1.30 ($\mathfrak{su}(2)$ from $SU(2)$ via the matrix exponential). Let $G = SU(2) = \{g \in GL(2, \mathbb{C}) : g^\dagger g = I, \det g = 1\}$. Differentiating both defining conditions along $\gamma(t) = e^{tv}$, $\gamma(0) = I$, gives the linearized conditions on v :

$$\left. \frac{d}{dt} \right|_{t=0} (e^{tv})^\dagger e^{tv} = v^\dagger + v = 0, \quad \left. \frac{d}{dt} \right|_{t=0} \det e^{tv} = \text{tr}(v) = 0,$$

using $\det e^{tv} = e^{t \text{tr } v}$. So

$$\mathfrak{su}(2) = \{v \in M_2(\mathbb{C}) : v^\dagger = -v, \text{tr}(v) = 0\},$$

a 3-dimensional real vector space.

Take $v = -\frac{i\theta}{2}\sigma_3$, $\theta \in \mathbb{R}$, $\sigma_3 = \text{diag}(1, -1)$. Since $\sigma_3^2 = I$, the series splits into even and odd powers:

$$\exp(v) = e^{-i\theta\sigma_3/2} = \cos\left(\frac{\theta}{2}\right)I - i\sin\left(\frac{\theta}{2}\right)\sigma_3 = \begin{pmatrix} e^{-i\theta/2} & 0 \\ 0 & e^{i\theta/2} \end{pmatrix}.$$

This matrix is unitary with determinant 1 for every θ , confirming $\exp(v) \in SU(2)$; and $\theta \mapsto \exp(v)$ satisfies the one-parameter subgroup law, consistently with Proposition 1.25.

1.2.2 Representations of groups

The previously presented concepts can be used to state the definition of a representation (or *rep*). Given that we will use the following applied to various mathematical objects, we decided to first show the general definition on groups but it is clearly valid for Lie groups and Lie algebras with some modifies. [3]

Definition 1.31 (Representation of a group). Let G be a group and V a vector space over $\mathbb{K}$. A representation of G on V is an homomorphism between groups:

$$\rho : G \rightarrow GL(V),$$

where $GL(V)$ denotes the general linear group, the set of all the linear isomorphism $\psi : V \rightarrow V$, called automorphisms. A representation is often denoted with (V, ρ) .

In the case of Lie algebras this can be transported as the following.

Definition 1.32 (Representation of a Lie Algebra). A representation of a Lie Algebra $\mathfrak{g}$ is an homomorphism $\phi : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ where V is a $\mathbb{K}$ -vector space.

Remark 1.33. It is worth noting that this definition is slightly different from the previous one. In the case of a group, the range of the homomorphism ρ is the set of all the automorphisms. Instead in the definition 1.32, the applications inside $\mathfrak{gl}(V)$ are not all isomorphisms. The reason why is that $\mathfrak{gl}(V)$ arises as the *tangent space* to $GL(V)$ at the identity. Defining an exponential map $g(t) = e^{tX}$ one has $g'(0) = X$, which is generically singular even though every $g(t)$ is invertible. Algebraically, this reflects the fact that a Lie algebra homomorphism $\rho : \mathfrak{g} \rightarrow \mathfrak{gl}(V)$ need only preserve the bracket, $\rho([X, Y]) = [\rho(X), \rho(Y)]$, whereas a group homomorphism $\rho : G \rightarrow GL(V)$ must preserve inverses, which forces $\rho(g) \in GL(V)$ for every $g \in G$.

Example 1.34. Let us define the *adjoint representation* as the homomorphism $\text{ad} : \mathfrak{g} \rightarrow \mathfrak{gl}(\mathfrak{g})$ which it sends $x \mapsto \text{ad } x$ with $\text{ad } x$ being an endomorphism on $\mathfrak{g}$ defined as

$$\text{ad } x(y) := [x, y].$$

One of the most important proprieties of a representation is the faithfulness.

Definition 1.35 (Faithful Representation). A representation (V, ρ) is said to be faithful if ρ is injective that is

$$\ker \rho = \{g \in G \mid \rho(g) = \mathbb{1}_V\} = \{e\},$$

where e is the identity element of the group.

Remark 1.36. This definition is referring to the representations which don't lose the informations encoded in the group G . All the different elements of the group are being mapped into a different application.

Definition 1.37 (Equivalent representations). Let (V, ρ) and (W, π) be two representations of the group G , with V and W two $\mathbb{K}$ -vector spaces. The representations are said to be equivalent if there exists a vector space isomorphism $T : V \rightarrow W$ such that:

$$T\rho(g) = \pi(g)T \quad \forall g \in G.$$

Equivalently we can state that, in order to define a representation equivalent to (V, ρ) on a $\mathbb{K}$ -vector space W , we have to use the vector space isomorphism T and define the homomorphism of the equivalent representation as

$$\pi := T \circ \rho \circ T^{-1}$$

Now we are ready to define what it means to "decompose" representations into the so called *irreducible representations* or *irreps*.

Definition 1.38 (Irreducibility of a representation). Let (V, π) be a finite-dimensional representation of a group G , acting on a space V . We can define the following terms:

- A subspace W of V is called *invariant* if $(\pi(g))(w) \in W \quad \forall w \in W, \forall g \in G$.
- An invariant subspace W is called *nontrivial* if $W \neq \{\emptyset\}$ and $W \neq V$.
- A representation with no nontrivial invariant subspaces is called *irreducible*.

The terms *invariant*, *nontrivial*, and *irreducible* are defined analogously for representations of Lie algebras.

Thanks to that definition we can now state one of the most famous and important lemmas in representation theory. It

Lemma 1.39 (Schur's Lemma). Let $\pi : G \rightarrow GL(V)$ be a finite-dimensional irrep over $\mathbb{C}$.

1. If $T : V \rightarrow V$ is a linear map commuting with $\pi(g)$ for every g , then $T = \lambda \text{id}_V$ for some $\lambda \in \mathbb{C}$.
2. More generally, if $\pi_1 : G \rightarrow GL(V_1)$, $\pi_2 : G \rightarrow GL(V_2)$ are both irreducible and $T : V_1 \rightarrow V_2$ satisfies $T\pi_1(g) = \pi_2(g)T$ for all g , then either $T = 0$ or T is an isomorphism.

Example 1.40 (Schur's Lemma on $\mathfrak{su}(2)$). Let $\pi : \mathfrak{su}(2) \rightarrow \text{End}(V)$ be a finite-dimensional irreducible representation, and let

$$\hat{J}^2 := \hat{J}_x^2 + \hat{J}_y^2 + \hat{J}_z^2$$

be the total angular momentum squared operator. Using $[\hat{J}_i, \hat{J}_j] = i\epsilon_{ijk}\hat{J}_k$, a direct computation gives $[\hat{J}^2, \hat{J}_i] = 0$ for every i : $\hat{J}^2$ commutes with the entire representation.

By Schur's Lemma (part 1), any operator commuting with an irreducible representation is a scalar multiple of the identity:

$$\hat{J}^2 = j(j+1)\text{id}_V, \quad \text{for some } j \geq 0.$$

In this simple case we see how Schur's Lemma guarantees $\hat{J}^2$ takes one fixed value on the whole irrep, not just on individual states.

To conclude, let us state an important definition that will be needed in the next chapters.

Definition 1.41 (Strongly continuous unitary representation). Let $\mathcal{H}$ be a Hilbert space, and let $U(\mathcal{H})$ denote the group of unitary operators on $\mathcal{H}$. Then, a homomorphism $\Pi : G \rightarrow U(\mathcal{H})$, being G a topological group, is called a strongly continuous unitary representation of G if Π satisfies the following continuity condition called strong continuity:

$$A_n, A \in G \quad \text{and} \quad A_n \rightarrow A \quad \implies \quad \Pi(A_n)(v) \rightarrow \Pi(A)(v) \quad \forall v \in \mathcal{H}.$$

1.3 Elements of symplectic geometry

Definition 1.42 (Non-degenerate 2-form on a manifold). Let M be a smooth manifold and $\omega \in \Omega^2(M)$. We say ω is non-degenerate if, for every $p \in M$, the bilinear form ω_p is non-degenerate on $T_p M$: for every $X \in T_p M$,

$$(\omega_p(X, Y) = 0 \quad \forall Y \in T_p M) \implies X = 0.$$

Proposition 1.43. Let $\omega \in \Omega^2(M)$ be skew-symmetric, and for $p \in M$ set $r_p := \text{rank}(\omega_p)$. Then for every $p \in M$ there exists an integer n_p with $r_p = 2n_p$, and an ordered basis $\hat{e} = (e_i)$ of $T_p M$, with dual ordered basis (α^i) , such that

$$\omega_p = \sum_{i=1}^{n_p} \alpha^i \wedge \alpha^{i+n_p},$$

or, equivalently, the matrix of ω_p in this basis is

$$\begin{pmatrix} 0 & I & 0 \\ -I & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

where I is the $n_p \times n_p$ identity matrix and the bottom-right block has size $(d-2n_p) \times (d-2n_p)$, $d := \dim T_p M$.

Remark 1.44. This is pointwise linear algebra: (e_i) is a basis of $T_p M$ alone, chosen independently at each p , with no claim that $p \mapsto e_i(p)$ varies smoothly, or even continuously.

Corollary 1.45. Suppose ω is non-degenerate, then $r_p = \dim T_p M =: d$ for every $p \in M$, so the bottom-right block above has no size, and $n_p = d/2$ is the same integer at every point. In particular:

- d must be even: a skew-symmetric non-degenerate 2-form can exist only on an even-dimensional manifold (on an odd-dimensional one, every 2-form is degenerate at every point).
- $\omega^n := \omega \wedge \cdots \wedge \omega$ ($n = d/2$ factors) is nowhere-vanishing (or in the basis used above, $\omega_p^n = n! \alpha^1 \wedge \alpha^{n+1} \wedge \cdots \wedge \alpha^n \wedge \alpha^{2n} \neq 0$) so ω^n (not ω itself) is a volume form on M , that can be standardized as

$$\Omega_\omega := \frac{(-1)^{n/2}}{n!} \omega^n$$

Definition 1.46 (Musical isomorphisms). Let M be a manifold and $\omega \in \Omega^2(M)$ be non-degenerate. Then define the map $\flat : \mathfrak{X}(M) \rightarrow \Omega^1(M)$ so that a vector field $X \mapsto X^\flat = \iota_X \omega = \omega(X, \cdot)$ and the inverse map $\sharp : \Omega^1(M) \rightarrow \mathfrak{X}(M)$ so that the form $\alpha \mapsto \alpha^\sharp = \omega^\sharp(\alpha)$.

Remark 1.47. We can clearly see that $(X^\flat)^\sharp = X$ and $(\alpha^\sharp)^\flat = \alpha$ due to the non-degeneracy of ω .

Theorem 1.48 (Darboux Theorem). Suppose ω is a non-degenerate two-form on a $2n$ -manifold M . Then $d\omega = 0$ (where d denotes the exterior derivative) iff there is a chart (U, ϕ) at each $m \in M$ such that $\phi(m) = \mathbf{0}$ and $\phi(u) = (x^1(u), \dots, x^n(u), y^1(u), \dots, y^n(u))$ having

$$\omega|_U = \sum_{i=1}^n dx^i \wedge dy^i$$

Finally, we can state the next important definition that encodes all the structure to derive Hamiltonian mechanics in the clearest and natural way possible.

Definition 1.49 (Symplectic manifold). A symplectic form ω on a manifold M is a non-degenerate, closed ($d\omega = 0$), two-form on M . A symplectic manifold (M, ω) is a manifold M together with a symplectic form ω on M . The charts guaranteed by the Darboux theorem are called symplectic charts and the coordinate are called canonical coordinates.

Definition 1.50 (Canonical transformation). Let (M, ω) and (N, ρ) be symplectic manifolds, F a C^∞ -mapping $F : M \rightarrow N$ is called symplectic or canonical transformation if $F^* \rho = \omega$.

Also we have a particular property for canonical transformations.

Proposition 1.51. If (M, ω) and (N, ρ) be symplectic manifolds and $F : M \rightarrow N$ symplectic then F is volume preserving and $\det_{(\Omega_\omega, \Omega_\rho)} F = 1$ and F is a local diffeomorphism.

The next theorem can be really obscure but it is a key result that we cannot omit in this introduction to symplectic geometry. The problem we want to solve is the following: Imagine you need to construct a 1-form on a manifold M , so you would need a rule to give any tangent vector a real number. If M is already a cotangent bundle of another manifold Q (that's the case of the phase space in Hamiltonian mechanics), this is really helping because there is a natural way to accomplish this.

Every point in M so it's a 1-form, viewed in the perspective of Q , let's call it α_q . If we switch in the perspective of Q , now we can define the tangent vector at the point α_q , let's call it w_{α_q} . The natural way to create a 1-form now is to use α_q as a form defined on tangent vectors to Q . So we take w_{α_q} , use a pushforward to move it to the tangent bundle of Q and finally pairing it with α_q . This is what the canonical form θ_0 does.

Theorem 1.52 (Canonical forms). Let Q be an n -manifold and $M = T^*Q$. Consider now the maps $\tau_Q^* : T^*Q \rightarrow Q$ and $d\tau_Q^* : TM \rightarrow TQ$, which is the pushforward of τ_Q^* . Given a point $q \in Q$, we can define $\alpha_q \in T^*Q$ and $w_{\alpha_q} \in TM$ in the fiber above α_q . Define

$$\theta_{\alpha_q} : T_{\alpha_q} M \rightarrow \mathbb{R} \quad w_{\alpha_q} \mapsto \alpha_q(d\tau_Q^*(w_{\alpha_q}))$$

and the map $\theta_0 : \alpha_q \in \theta_{\alpha_q}$. In this case $\theta_0 \in \Omega^1(M)$ and $\omega_0 = -d\theta_0$ is a symplectic form on M . The forms θ_0 and ω_0 are called the canonical forms on M .

Remark 1.53. The superscript $*$ in τ_Q^* there simply labels it as the cotangent-bundle projection, as opposed to $\tau_Q : TQ \rightarrow Q$; it is not the pullback operator.

Corollary 1.54 (Canonical forms in coordinates). From the Theorem 1.52, in finite-dimensional manifolds written in local coordinates of $M(q^1, \dots, q^n, p_1, \dots, p_n)$, it follows that

$$\theta_0 = \sum_{i=1}^n p_i dq^i \quad \omega_0 = \sum_{i=1}^n dq^i \wedge dp_i$$

We can also prove that these canonical forms are not changed by any diffeomorphism on the manifold.

Theorem 1.55 (Invariance of the canonical forms). *Let Q be a manifold and $f : Q \rightarrow Q$ a diffeomorphism; define the lift of f as*

$$T^*f : T^*Q \rightarrow T^*Q \quad (T^*f(\alpha_q))(v) = \alpha_q(df(v))$$

*Then T^*f is symplectic and if I use the lift to pullback the canonical form $(T^*f)^*\theta_0 = \theta_0$ it comes right back.*

Remark 1.56. The lift of diffeomorphisms are defined to make the following diagram commute.

$$\begin{array}{ccc} T^*Q & \xrightarrow{T^*f} & T^*Q \\ \tau_Q^* \downarrow & & \downarrow \tau_Q^* \\ Q & \xleftarrow{f} & Q \end{array}$$

In coordinates, if $f(q^1, \dots, q^n) = (Q^1, \dots, Q^n)$, then the lift of f is a map that

$$(Q^1, \dots, Q^n, P_1, \dots, P_n) \mapsto (q^1, \dots, q^n, p_1, \dots, p_n)$$

where

$$p_j = \sum_{i=1}^n \frac{\partial Q^i}{\partial q^j} P_i$$

Notice also that

$$T^*(f \circ g) = T^*g \circ T^*f$$

that is a law of composition similar to $T(f \circ g) = Tf \circ Tg$.

1.4 Hamiltonian Vector Fields and Poisson Brackets

Definition 1.57 (Hamiltonian Vector Field). *Let (M, ω) be a symplectic manifold called Phase space and $H : M \rightarrow \mathbb{R}$ a given C^r -function. The vector field X_H determined by*

$$\omega(X_H, Y) = dH(Y) \quad \text{i.e.} \quad \iota_{X_H} \omega = dH, \quad (1.3)$$

is called the Hamiltonian vector field with H be the energy function. (M, ω, H) is an Hamiltonian system.

Proposition 1.58 (Hamilton Equations). *Let $(q^1, \dots, q^n, p_1, \dots, p_n)$ be canonical coordinates for ω , so $\omega = \sum dq^i \wedge dp_i$. Then in these coordinates (and dropping base points),*

$$X_H = \left(\frac{\partial H}{\partial p_i}, -\frac{\partial H}{\partial q^i} \right) = J \cdot dH \quad (1.4)$$

where $J = \begin{pmatrix} 0 & I \\ -I & 0 \end{pmatrix}$. Thus $(q(t), p(t))$ is an integral curve of X_H iff Hamilton's equations hold:

$$\boxed{\dot{q}^i = \frac{\partial H}{\partial p_i}, \quad \dot{p}_i = -\frac{\partial H}{\partial q^i}, \quad i = 1, \dots, n}$$

Proof. Let X_H be defined by the formula 1.4. We then have to verify 1.3. Now $i_{X_H} dq^i = \partial H / \partial p_i$, $i_{X_H} dp_i = -\partial H / \partial q^i$ by construction, so

$$i_{X_H} \omega = \sum i_{X_H} (dq^i \wedge dp_i) = \sum (i_{X_H} dq^i) \wedge dp_i - \sum dq^i \wedge i_{X_H} dp_i = \sum \frac{\partial H}{\partial p_i} dp_i + \frac{\partial H}{\partial q^i} dq^i = dH$$

□

Conservation of energy is easy to prove:

Proposition 1.59 (Conservation of energy). *Let (M, ω, X_H) be a Hamiltonian system and let $c(t)$ be an integral curve for X_H . Then $H(c(t))$ is constant in t .*

Proof. By the chain rule and 1.3,

$$\begin{aligned}
 \frac{d}{dt}H(c(t)) &= dH(c(t)) \cdot c'(t) \\
 &= dH(c(t)) \cdot X_H(c(t)) \\
 &= \omega(X_H(c(t)), X_H(c(t))) \\
 &= 0
 \end{aligned}$$

since ω is skew-symmetric. □

The next basic fact about Hamiltonian systems is that their flows consist of canonical transformations.

Proposition 1.60 (Liouville's Theorem). *Let (M, ω, X_H) be a Hamiltonian system, and F_t be the flow of X_H . Then for each t , $F_t^* \omega = \omega$, that is, F_t is symplectic (on its domain). Thus F_t also preserves the phase volume Ω_ω .*

Proof. We have

$$\begin{aligned}
 \frac{d}{dt}F_t^* \omega &= F_t^* L_{X_H} \omega \\
 &= F_t^* (i_{X_H} d\omega + di_{X_H} \omega) \\
 &= F_t^* (0 + ddH) \quad (\text{since } d\omega = 0 \text{ and } i_{X_H} \omega = dH) \\
 &= 0
 \end{aligned}$$

Thus $F_t^* \omega$ is constant in t . Since $F_0 = \mathbb{1}$, the equation $F_t^* \omega = \omega$ results. □

Bibliography

- [1] Ralph Abraham and Jerrold E. Marsden. *Foundations of Mechanics*. 2nd ed. Providence, R.I: AMS Chelsea Pub./American Mathematical Society. 826 pp. ISBN: 978-0-8218-4438-0.
- [2] James E. Humphreys. *Introduction to Lie Algebras and Representation Theory*. 11. print. Graduate Texts in Mathematics 9. New York, NY: Springer. 173 pp. ISBN: 978-0-387-90053-7.
- [3] Brian C. Hall. *Lie Groups, Lie Algebras, and Representations*. Vol. 222. Graduate Texts in Mathematics. New York, NY: Springer New York. ISBN: 978-1-4419-2313-4 978-0-387-21554-9. DOI: 10.1007/978-0-387-21554-9. URL: <http://link.springer.com/10.1007/978-0-387-21554-9>.