



Oral toxicity prediction results for input compound



Predicted LD50: 2000mg/kg

Predicted Toxicity Class: 4



Average similarity: 38.65%

Prediction accuracy: 23%



Name	S=C(N)N=N=C1C[C@H](S[C@@H]2[C@H](OC(C)=O)[C@H](OC(C)=O)[C@H](O[C@@H]3[C@H](OC(C)=O)[C@H](OC(C)=O)[C@H](OC(C)=O)[C@H](COC(C)=O)O3)[C@@H](COC(C)=O)O2)[C@@H]4O[C@H]/1OC4
Molweight	851.85
Number of hydrogen bond acceptors	24
Number of hydrogen bond donors	2
Number of atoms	57
Number of bonds	60
Number of rotatable bonds	22
Molecular refractivity	191.29
Topological Polar Surface Area	338.05
octanol/water partition coefficient(logP)	0.13

Toxicity Model Report

[Copy](#) [Excel](#) [CSV](#) [PDF](#)

Classification	Target	Shorthand	Prediction	Probability
Organ toxicity	Hepatotoxicity	dili	Inactive	0.57
Organ toxicity	Neurotoxicity	neuro	Inactive	0.59
Organ toxicity	Nephrotoxicity	nephro	Active	0.65
Organ toxicity	Respiratory toxicity	respi	Active	0.69
Organ toxicity	Cardiotoxicity	cardio	Inactive	0.52
Toxicity end points	Carcinogenicity	carcino	Active	0.51
Toxicity end points	Immunotoxicity	immuno	Inactive	0.91
Toxicity end points	Mutagenicity	mutagen	Active	0.51
Toxicity end points	Cytotoxicity	cyto	Inactive	0.70
Toxicity end points	BBB-barrier	bbb	Inactive	0.63
Toxicity end points	Ecotoxicity	eco	Inactive	0.56
Toxicity end points	Clinical toxicity	clinical	Inactive	0.55
Toxicity end points	Nutritional toxicity	nutri	Active	0.51
Tox21-Nuclear receptor signalling pathways	Aryl hydrocarbon Receptor (AhR)	nr_ahr	Inactive	0.88
Tox21-Nuclear receptor signalling pathways	Androgen Receptor (AR)	nr_ar	Inactive	0.94
Tox21-Nuclear receptor signalling pathways	Androgen Receptor Ligand Binding Domain (AR-LBD)	nr_ar_lbd	Inactive	0.96
Tox21-Nuclear receptor signalling pathways	Aromatase	nr_aromatase	Inactive	0.91
Tox21-Nuclear receptor signalling pathways	Estrogen Receptor Alpha (ER)	nr_er	Inactive	0.81
Tox21-Nuclear receptor signalling pathways	Estrogen Receptor Ligand Binding Domain (ER-LBD)	nr_er_lbd	Inactive	0.94
Tox21-Nuclear receptor signalling pathways	Peroxisome Proliferator Activated Receptor Gamma (PPAR-Gamma)	nr_ppar_gamma	Inactive	0.93
Tox21-Stress response pathways	Nuclear factor (erythroid-derived 2)-like 2/antioxidant responsive element (nrf2/ARE)	sr_are	Inactive	0.92
Tox21-Stress response pathways	Heat shock factor response element (HSE)	sr_hse	Inactive	0.92
Tox21-Stress response pathways	Mitochondrial Membrane Potential (MMP)	sr_mmp	Inactive	0.81
Tox21-Stress response pathways	Phosphoprotein (Tumor Suppressor) p53	sr_p53	Inactive	0.85
Tox21-Stress response pathways	ATPase family AAA domain-containing protein 5 (ATAD5)	sr_atad5	Inactive	0.93
Molecular Initiating Events	Thyroid hormone receptor alpha (THRα)	mie_thr_alpha	Inactive	0.79
Molecular Initiating Events	Thyroid hormone receptor beta (THRβ)	mie_thr_beta	Inactive	0.71
Molecular Initiating Events	Transthyretin (TTR)	mie_ttr	Inactive	0.69
Molecular Initiating Events	Ryanodine receptor (RYP)	mie_ryr	Inactive	0.68
Molecular Initiating Events	GABA receptor (GABAR)	mie_gabar	Inactive	0.79
Molecular Initiating Events	Glutamate N-methyl-D-aspartate receptor (NMDAR)	mie_nmdar	Inactive	0.93
Molecular Initiating Events	alpha-amino-3-hydroxy-5-methyl-4-isoxazolepropionate receptor (AMPA)	mie_ampar	Inactive	0.96
Molecular Initiating Events	Kainate receptor (KAR)	mie_kar	Inactive	0.98

