

# Deriving equations of motions of a double pendulum on a cart using the Lagrangian

Mirko van de Hoef

December 6, 2023

## 1 Introduction

This file is made to document the simulation of a double pendulum on a cart using lagrangian mechanics. In this file a double pendulum on a cart will be discussed.

## 2 The system

Where in figure 1:

- $M$ : Mass of the cart [kg]
- $m_1$ : Mass of the first pendulum [kg]
- $m_2$ : Mass of the second pendulum [kg]
- $\theta_1$ : Degrees first pendulum in reference to the cart [rad]
- $\theta_2$ : Degrees second pendulum in reference to the cart [rad]
- $\ell_1$ : Length of the first pendulum [m]
- $\ell_2$ : Length of the second pendulum [m]
- $x_c$ : Position of the cart [m]




Figure 1: The pendulum

### 3 Lagrangian

Let the position and velocity of the pendulum be

$$\begin{aligned} x_1 &= \ell_1 \sin(\theta_1) + x_c & \dot{x}_1 &= \ell_1 \dot{\theta}_1 \cos(\theta_1) + \dot{x}_c \\ y_1 &= -\ell_1 \cos(\theta_1) & \dot{y}_1 &= \ell_1 \dot{\theta}_1 \sin(\theta_1) \\ x_2 &= \ell_1 \sin(\theta_1) + \ell_2 \sin(\theta_2) + x_c & \dot{x}_2 &= \ell_1 \dot{\theta}_1 \cos(\theta_1) + \ell_2 \dot{\theta}_2 \cos(\theta_2) + \dot{x}_c \\ y_2 &= -\ell_1 \cos(\theta_1) - \ell_2 \cos(\theta_2) & \dot{y}_2 &= \ell_1 \dot{\theta}_1 \sin(\theta_1) + \ell_2 \dot{\theta}_2 \sin(\theta_2) \end{aligned}$$

To derive the equations of motion, the Lagrangian (1) will be used. First,  $T$ , the kinetic energy of the system will be calculated.

$$\mathcal{L} = T - V \quad (1)$$

$$\begin{aligned} T &= \frac{1}{2} M \dot{x}_c^2 + \frac{1}{2} m_1 (\dot{x}_1^2 + \dot{y}_1^2) + \frac{1}{2} m_2 (\dot{x}_2^2 + \dot{y}_2^2) \\ T &= \frac{1}{2} M \dot{x}_c^2 + \frac{1}{2} m_1 \left( (\ell_1 \dot{\theta}_1 \cos(\theta_1) + \dot{x}_c)^2 + (\dot{y}_1)^2 \right) \\ &\quad + \frac{1}{2} m_2 \left( (\ell_1 \dot{\theta}_1 \cos(\theta_1) + \dot{x}_c + \ell_2 \dot{\theta}_2 \cos(\theta_2))^2 + (\dot{y}_2)^2 \right) \\ T &= \frac{1}{2} M \dot{x}_c^2 + \frac{1}{2} m_1 (\ell_1^2 \dot{\theta}_1^2 \cos^2(\theta_1) + 2\ell_1 \dot{\theta}_1 \cos(\theta_1) \dot{x}_c + \dot{x}_c^2 + \dot{y}_1^2) + \frac{1}{2} m_2 (\dots) \\ T &= \frac{1}{2} M \dot{x}_c^2 + \frac{1}{2} m_1 (\dots) + \frac{1}{2} m_2 \left( (\ell_1 \dot{\theta}_1 \cos(\theta_1) + \dot{x}_c + \ell_2 \dot{\theta}_2 \cos(\theta_2))^2 + \dot{y}_2^2 \right) \\ T &= \frac{1}{2} M \dot{x}_c^2 + \frac{1}{2} m_1 (\ell_1^2 \dot{\theta}_1^2 \cos^2(\theta_1) + 2\ell_1 \dot{\theta}_1 \cos(\theta_1) \dot{x}_c + \dot{x}_c^2 + \dot{y}_1^2) \\ &\quad + \frac{1}{2} m_2 \left( (\ell_1^2 \dot{\theta}_1^2 \cos^2(\theta_1) + 2\ell_1 \dot{\theta}_1 \cos(\theta_1) \dot{x}_c + 2\ell_1 \ell_2 \dot{\theta}_1 \dot{\theta}_2 \cos(\theta_1) \cos(\theta_2) \dot{x}_c^2 + 2\dot{x}_c \ell_2 \dot{\theta}_2 \cos(\theta_2) + \ell_2^2 \dot{\theta}_2^2 \cos^2(\theta_2) + \dot{y}_2^2) \right) \\ T &= \frac{1}{2} M \dot{x}_c^2 + \frac{1}{2} (m_1 + m_2) (\ell_1^2 \dot{\theta}_1^2 \cos^2(\theta_1) + 2\ell_1 \dot{\theta}_1 \cos(\theta_1) \dot{x}_c) \\ &\quad + \frac{1}{2} m_1 (\dot{x}_c^2 + \dot{y}_1^2) + m_2 \left( (2\ell_1 \ell_2 \dot{\theta}_1 \dot{\theta}_2 \cos(\theta_1) \cos(\theta_2) \dot{x}_c^2 + 2\dot{x}_c \ell_2 \dot{\theta}_2 \cos(\theta_2) + \ell_2^2 \dot{\theta}_2^2 \cos^2(\theta_2) + \dot{y}_2^2) \right) \\ T &= \frac{1}{2} (M + m_1) \dot{x}_c^2 + \frac{1}{2} (m_1 + m_2) (\ell_1^2 \dot{\theta}_1^2 \cos^2(\theta_1) + 2\ell_1 \dot{\theta}_1 \cos(\theta_1) \dot{x}_c) \\ &\quad + \frac{1}{2} m_1 \dot{y}_1^2 + m_2 \left( (2\ell_1 \ell_2 \dot{\theta}_1 \dot{\theta}_2 \cos(\theta_1) \cos(\theta_2) \dot{x}_c^2 + 2\dot{x}_c \ell_2 \dot{\theta}_2 \cos(\theta_2) + \ell_2^2 \dot{\theta}_2^2 \cos^2(\theta_2) + \dot{y}_2^2) \right) \end{aligned} \quad (2)$$

Then, for  $V$ , the potential energy of the system

$$V = m_1 g(y_1) + m_2 g(y_2) \quad (3)$$

Lastly, using (2) and (3) the Lagrangian,  $\mathcal{L}$ , can be formulated

$$\begin{aligned} \mathcal{L} &= \frac{1}{2} (M + m_1) \dot{x}_c^2 + \frac{1}{2} (m_1 + m_2) (\ell_1^2 \dot{\theta}_1^2 \cos^2(\theta_1) + 2\ell_1 \dot{\theta}_1 \cos(\theta_1) \dot{x}_c) \\ &\quad + \frac{1}{2} m_1 \dot{y}_1^2 + m_2 \left( (2\ell_1 \ell_2 \dot{\theta}_1 \dot{\theta}_2 \cos(\theta_1) \cos(\theta_2) \dot{x}_c^2 + 2\dot{x}_c \ell_2 \dot{\theta}_2 \cos(\theta_2) + \ell_2^2 \dot{\theta}_2^2 \cos^2(\theta_2) + \dot{y}_2^2) \right) \\ &\quad - m_1 g y_1 - m_2 g y_2 \end{aligned} \quad (4)$$

## 4 Solve for $\ddot{\theta}_1$

Using the Lagrangian equation (19)

$$\frac{d}{dt} \left( \frac{\partial \mathcal{L}}{\partial \dot{q}} \right) = \frac{\partial \mathcal{L}}{\partial q} \quad (5)$$

Substituting (4) in equation (19) and solving for  $\theta_1$  gives:

$$\frac{\partial \mathcal{L}}{\partial \dot{\theta}_1} = (m_1 + m_2)\ell_1^2 \dot{\theta}_1 + m_2 \ell_1 \ell_2 \dot{\theta}_2 \cos(\theta_1 - \theta_2) + m_2 \dot{x}_c \ell_1 \cos(\theta_1) \quad (6)$$

$$\begin{aligned} \frac{d}{dt} \left( \frac{\partial \mathcal{L}}{\partial \dot{\theta}_1} \right) = & (m_1 + m_2)\ell_1^2 \ddot{\theta}_1 + m_2 \ell_1 \ell_2 \ddot{\theta}_2 \cos(\theta_1 - \theta_2) - m_2 \ell_1 \ell_2 \dot{\theta}_2 \sin(\theta_1 - \theta_2)(\dot{\theta}_1 - \dot{\theta}_2) \\ & + m_2 \ddot{x}_c \ell_1 \cos(\theta_1) - m_2 \dot{x}_c \ell_1 \sin(\theta_1) \dot{\theta}_1 \end{aligned} \quad (7)$$

$$\frac{\partial \mathcal{L}}{\partial \theta_1} = -m_2 \ell_1 \ell_2 \dot{\theta}_1 \dot{\theta}_2 \sin(\theta_1 - \theta_2) - (m_1 + m_2)g \ell_1 \sin(\theta_1) - m_2 \dot{x}_c \ell_1 \dot{\theta}_1 \sin(\theta_1) \quad (8)$$

Which give the following differential equation:

$$\begin{aligned} & (m_1 + m_2)\ell_1^2 \ddot{\theta}_1 + m_2 \ell_1 \ell_2 \ddot{\theta}_2 \cos(\theta_1 - \theta_2) - m_2 \ell_1 \ell_2 \dot{\theta}_2 \sin(\theta_1 - \theta_2)(\dot{\theta}_1 - \dot{\theta}_2) \\ & + m_2 \ddot{x}_c \ell_1 \cos(\theta_1) - m_2 \dot{x}_c \ell_1 \sin(\theta_1) \dot{\theta}_1 + m_2 \ell_1 \ell_2 \dot{\theta}_1 \dot{\theta}_2 \sin(\theta_1 - \theta_2) \\ & + (m_1 + m_2)g \ell_1 \sin(\theta_1) + m_2 \dot{x}_c \ell_1 \dot{\theta}_1 \sin(\theta_1) = 0 \end{aligned} \quad (9)$$

Which simplifies to:

$$\begin{aligned} & (m_1 + m_2)\ell_1 \ddot{\theta}_1 + m_2 \ell_2 \ddot{\theta}_2 \cos(\theta_1 - \theta_2) + m_2 \ell_2 \dot{\theta}_2^2 \sin(\theta_1 - \theta_2) \\ & + m_2 \ddot{x}_c \cos(\theta_1) + (m_1 + m_2)g \sin(\theta_1) = 0 \end{aligned} \quad (10)$$

With the differential equation solved for  $\ddot{\theta}_1$ :

$$\frac{m_2 \ell_2 \ddot{\theta}_2 \cos(\theta_1 - \theta_2) + m_2 \ell_2 \dot{\theta}_2^2 \sin(\theta_1 - \theta_2) + m_2 \ddot{x}_c \cos(\theta_1) + (m_1 + m_2)g \sin(\theta_1)}{-(m_1 + m_2)\ell_1} = \ddot{\theta}_1 \quad (11)$$

## 5 Solve for $\ddot{\theta}_2$

Using the Lagrangian equation (19)

$$\frac{d}{dt} \left( \frac{\partial \mathcal{L}}{\partial \dot{q}} \right) = \frac{\partial \mathcal{L}}{\partial q} \quad (12)$$

Substituting (4) in equation (19) and solving for  $\theta_1$  gives:

$$\frac{\partial \mathcal{L}}{\partial \dot{\theta}_1} = m_2 \ell_1 \ell_2 \dot{\theta}_1 \cos(\theta_1 - \theta_2) + m_2 \ell_2^2 \dot{\theta}_2 + m_2 \dot{x}_c \ell_2 \cos(\theta_2) \quad (13)$$

$$\begin{aligned} \frac{d}{dt} \left( \frac{\partial \mathcal{L}}{\partial \dot{\theta}_1} \right) &= m_2 \ell_1 \ell_2 \ddot{\theta}_1 \cos(\theta_1 - \theta_2) - m_2 \ell_1 \ell_2 \dot{\theta}_1 \sin(\theta_1 - \theta_2) (\dot{\theta}_1 - \dot{\theta}_2) \\ &\quad + m_2 \ell_2^2 \ddot{\theta}_2 + m_2 \ddot{x}_c \ell_2 \cos(\theta_2) - m_2 \dot{x}_c \ell_2 \sin(\theta_2) \dot{\theta}_2 \end{aligned} \quad (14)$$

$$\frac{\partial \mathcal{L}}{\partial \theta_1} = -m_2 \ell_1 \ell_2 \dot{\theta}_1 \dot{\theta}_2 \sin(\theta_1 - \theta_2) - m_2 \dot{x}_c \dot{\theta}_2 \ell_2 \sin(\theta_2) - m_2 g \ell_2 \sin(\theta_2) \quad (15)$$

Which give the following differential equation:

$$\begin{aligned} &m_2 \ell_1 \ell_2 \ddot{\theta}_1 \cos(\theta_1 - \theta_2) - m_2 \ell_1 \ell_2 \dot{\theta}_1 \sin(\theta_1 - \theta_2) (\dot{\theta}_1 - \dot{\theta}_2) \\ &+ m_2 \ell_2^2 \ddot{\theta}_2 + m_2 \ddot{x}_c \ell_2 \cos(\theta_2) - m_2 \dot{x}_c \ell_2 \sin(\theta_2) \dot{\theta}_2 + m_2 \ell_1 \ell_2 \dot{\theta}_1 \dot{\theta}_2 \sin(\theta_1 - \theta_2) \\ &+ m_2 \dot{x}_c \dot{\theta}_2 \ell_2 \sin(\theta_2) + m_2 g \ell_2 \sin(\theta_2) = 0 \end{aligned} \quad (16)$$

Which simplifies to:

$$m_2 \ell_2 \ddot{\theta}_2 + m_2 \ell_1 \ddot{\theta}_1 \cos(\theta_1 - \theta_2) - m_2 \ell_1 \dot{\theta}_1^2 \sin(\theta_1 - \theta_2) + m_2 \ddot{x}_c \cos(\theta_2) + m_2 g \sin(\theta_2) = 0 \quad (17)$$

With the differential equation solved for  $\ddot{\theta}_1$ :

$$\frac{m_2 \ell_1 \ddot{\theta}_1 \cos(\theta_1 - \theta_2) - m_2 \ell_1 \dot{\theta}_1^2 \sin(\theta_1 - \theta_2) + m_2 \ddot{x}_c \cos(\theta_2) + m_2 g \sin(\theta_2)}{-m_2 \ell_2} = \ddot{\theta}_2 \quad (18)$$

## 6 Solve for $\ddot{x}_c$

Using the Lagrangian equation (19)

$$\frac{d}{dt} \left( \frac{\partial \mathcal{L}}{\partial \dot{q}} \right) = \frac{\partial \mathcal{L}}{\partial q} \quad (19)$$

Substituting (4) in equation (19) and solving for  $\theta_1$  gives:

$$\frac{\partial \mathcal{L}}{\partial \dot{\theta}_1} = (M + m_1)\dot{x}_c + m_2\ell_1\dot{\theta}_1 \cos(\theta_1) + m_2\ell_2\dot{\theta}_2 \cos(\theta_2) \quad (20)$$

$$\frac{d}{dt} \left( \frac{\partial \mathcal{L}}{\partial \dot{\theta}_1} \right) = (M + m_1)\ddot{x}_c + m_2\ell_1\ddot{\theta}_1 \cos(\theta_1) - m_2\ell_1\dot{\theta}_1^2 \sin(\theta_1) + m_2\ell_2\ddot{\theta}_2 \cos(\theta_2) - m_2\ell_2\dot{\theta}_2^2 \sin(\theta_2) \quad (21)$$

$$\frac{\partial \mathcal{L}}{\partial \theta_1} = 0 \quad (22)$$

Which give the following differential equation:

$$(M + m_1)\ddot{x}_c + m_2\ell_1\ddot{\theta}_1 \cos(\theta_1) - m_2\ell_1\dot{\theta}_1^2 \sin(\theta_1) + m_2\ell_2\ddot{\theta}_2 \cos(\theta_2) - m_2\ell_2\dot{\theta}_2^2 \sin(\theta_2) = 0 \quad (23)$$

With the differential equation solved for  $\ddot{\theta}_1$ :

$$\frac{m_2\ell_1\ddot{\theta}_1 \cos(\theta_1) - m_2\ell_1\dot{\theta}_1^2 \sin(\theta_1) + m_2\ell_2\ddot{\theta}_2 \cos(\theta_2) - m_2\ell_2\dot{\theta}_2^2 \sin(\theta_2)}{m_2\ell_1 \cos(\theta_1)} - M + m_1 = \ddot{x}_c \quad (24)$$