

Ankle Injuries

SportVerse AI — Curated RAG Knowledge Base | Educational use only

Ankle injuries commonly occur after twisting, awkward landing, sudden direction changes, or contact. Sprains involve ligaments around the ankle and may cause pain, swelling, bruising, stiffness, and difficulty bearing weight.

Ask how the injury occurred, where pain is located, whether swelling or bruising is present, whether the athlete can bear weight, and whether there is deformity or numbness.

General recovery commonly progresses from protecting the injured area and controlling symptoms toward restoring movement, strength, balance, and sport-specific function.

Severe pain, significant swelling, visible deformity, inability to bear weight, numbness, or worsening symptoms warrant professional assessment. The AI should not label an injury as a definite sprain or fracture.

Safety requirements

- These documents are educational context for the SportVerse RAG prototype. They do not diagnose injuries or replace assessment by a qualified healthcare professional.
- The AI must use deterministic red-flag rules before generating general guidance. It must not downgrade an urgent/high-risk safety classification.
- Do not use this knowledge base to generate prescriptions, medication dosages, or definitive diagnoses.

Sources

NIAMS, Sports Injuries — <https://www.niams.nih.gov/health-topics/sports-injuries/diagnosis-treatment-and-steps-to-take>

Peer-reviewed ankle rehabilitation literature — <https://pmc.ncbi.nlm.nih.gov/articles/PMC164373/>

RAG implementation note: preserve this document's title and source metadata when indexing so the assistant can return source attribution with retrieved context.