

Muscle Strains

SportVerse AI — Curated RAG Knowledge Base | Educational use only

Muscle strains may occur during sprinting, jumping, sudden acceleration, overstretching, or high-force activity. Hamstring, calf, quadriceps, and adductor injuries are common in sport.

Possible symptoms include localized pain, tenderness, weakness, reduced movement, swelling, bruising, and pain during contraction or stretching. Symptoms alone do not establish the grade of a strain.

Rehabilitation generally progresses toward restoring range of motion, strength, coordination, and sport-specific function. Recovery time varies by injury severity and individual factors.

Significant weakness, severe pain, extensive bruising/swelling, inability to use the limb normally, or worsening symptoms should prompt professional evaluation.

Safety requirements

- These documents are educational context for the SportVerse RAG prototype. They do not diagnose injuries or replace assessment by a qualified healthcare professional.
- The AI must use deterministic red-flag rules before generating general guidance. It must not downgrade an urgent/high-risk safety classification.
- Do not use this knowledge base to generate prescriptions, medication dosages, or definitive diagnoses.

Sources

Evidence-Based Management and Factors Associated With Return to Play After Acute Hamstring Injury in Athletes — <https://pmc.ncbi.nlm.nih.gov/articles/PMC8649106/>

NIAMS, Sports Injuries — <https://www.niams.nih.gov/health-topics/sports-injuries/diagnosis-treatment-and-steps-to-take>

RAG implementation note: preserve this document's title and source metadata when indexing so the assistant can return source attribution with retrieved context.