

Dehydration in Athletes

SportVerse AI — Curated RAG Knowledge Base | Educational use only

Exercise increases fluid loss, especially in hot or humid conditions. Dehydration can impair performance and increase heat-related risk.

Possible signs include thirst, weakness, dizziness, fatigue, headache, reduced exercise tolerance, and muscle cramping. Severe or rapidly worsening symptoms require assessment.

Hydration needs vary by activity, environment, body size, and individual sweat rate. SportVerse should avoid prescribing a universal fluid volume for every athlete.

Fainting, confusion, severe weakness, or symptoms suggesting serious heat illness should trigger immediate cessation of activity and professional/emergency evaluation as appropriate.

Safety requirements

- These documents are educational context for the SportVerse RAG prototype. They do not diagnose injuries or replace assessment by a qualified healthcare professional.
- The AI must use deterministic red-flag rules before generating general guidance. It must not downgrade an urgent/high-risk safety classification.
- Do not use this knowledge base to generate prescriptions, medication dosages, or definitive diagnoses.

Sources

CDC, Heat and Athletes — <https://www.cdc.gov/heat-health/risk-factors/heat-and-athletes.html>

RAG implementation note: preserve this document's title and source metadata when indexing so the assistant can return source attribution with retrieved context.