

Return to Sport

SportVerse AI — Curated RAG Knowledge Base | Educational use only

Return to sport should be viewed as a progression rather than a fixed number of days. Recovery can include return to daily activity, training, sport participation, and eventually previous performance level.

Useful functional domains include range of motion, strength, movement/control, cardiovascular readiness, sport-specific ability, and psychological readiness where appropriate.

Return-to-sport criteria vary by injury and sport. Pain, swelling, instability, neurological symptoms, or worsening symptoms during progression should prompt stopping the activity and seeking professional guidance.

For concussion, return to sport should follow current healthcare guidance and a gradual progression. CDC materials state that athletes with suspected concussion should be removed from play and cleared by a healthcare provider before returning to play.

Safety requirements

- These documents are educational context for the SportVerse RAG prototype. They do not diagnose injuries or replace assessment by a qualified healthcare professional.
- The AI must use deterministic red-flag rules before generating general guidance. It must not downgrade an urgent/high-risk safety classification.
- Do not use this knowledge base to generate prescriptions, medication dosages, or definitive diagnoses.

Sources

Rehabilitation Using a Systematic and Holistic Approach for the Injured Athlete Returning to Sport — <https://pmc.ncbi.nlm.nih.gov/articles/PMC8811513/>

Defining Return to Sport: A Systematic Review — <https://pmc.ncbi.nlm.nih.gov/articles/PMC8320574/>

CDC HEADS UP — <https://www.cdc.gov/heads-up/>

RAG implementation note: preserve this document's title and source metadata when indexing so the assistant can return source attribution with retrieved context.