

MATH3871/MATH5960 Bayesian Inference and Computation, Lab 3 Exercises

1. Suppose that Y is an exponentially distributed random variable with mean $\lambda > 0$, and density $f(y) = \frac{1}{\lambda} \exp\left(-\frac{y}{\lambda}\right)$ for $y > 0$.

Derive the inverse-transform method to simulate from this density. Implement the sampling in R/Matlab and verify its correctness by plotting a histogram and the true pdf on the same figure.

2. Using R/Matlab approximate the following integrals via Monte Carlo integration:

(a)

$$\int_0^1 \exp[\exp(x)] dx.$$

(b)

$$\int_{-\infty}^{\infty} \exp(-x^2) dx.$$

(c)

$$\int_0^1 \int_0^1 \exp[(x+y)^2] dy dx.$$

(d)

$$\int_0^{\infty} \int_0^x \exp[-(x+y)] dy dx,$$

where you can use the indicator function

$$\mathbb{I}\{y < x\} = \begin{cases} 1 & \text{if } y < x \\ 0 & \text{otherwise} \end{cases}$$

to equate the integral to one in which the domains of integration both go from 0 to ∞ .

Solutions

1. Then the distribution function of an exponential random variable is

$$F(y) = \begin{cases} 1 - \exp\left(-\frac{y}{\lambda}\right) & y \geq 0 \\ 0 & \text{otherwise} \end{cases}$$

To implement the inverse transform method of simulation, we need to find $F^{-1}(p)$, $p \in (0, 1)$. For $p \in (0, 1)$, $F^{-1}(p)$ satisfies $F(F^{-1}(p)) = p$, or

$$\begin{aligned} 1 - \exp(-F^{-1}(p)/\lambda) &= p \\ \exp(-F^{-1}(p)/\lambda) &= 1 - p \\ -F^{-1}(p)/\lambda &= \ln(1 - p) \\ F^{-1}(p) &= -\lambda \ln(1 - p) \end{aligned}$$


So to simulate an exponential random variable with mean λ , we simulate a uniformly distributed random variable U on $[0, 1]$, and then compute $Y = -\lambda \ln(1 - U)$. Note that if U is uniformly distributed on $[0, 1]$, then $1 - U$ is also uniformly distributed on $[0, 1]$ so that we could simply compute $Y = -\lambda \ln U$ (simpler and faster).

Some R code to implement this is:

```

func=function(n,lambda) {
  return( -lambda*log(runif(n)) )
}
lambda=200
hist(func(5000,lambda),probability=T,xlab=expression(theta),ylab="Density",
main="Samples from f(x)",nclass=31,ylim=c(0,1/lambda))
xx=seq(0,qexp(0.999,1/lambda),length=100)
lines(xx,dexp(xx,1/lambda),col=2)

```



2. (a) Because the function $\exp(-x^2)$ is symmetric about the origin, $\int_{-\infty}^{\infty} \exp(-x^2) dx = 2 \int_0^{\infty} \exp(-x^2) dx$. Now transform the integral from $[0, \infty)$ to $[0, 1]$ by the transformation $u = 1/(1+x)$:

$$\int_0^{\infty} \exp(-x^2) dx = \int_0^1 \frac{1}{u^2} \exp\left(-\left(\frac{1}{u} - 1\right)^2\right) du$$

We can now approximate the integral as:

```

> u=runif(10000)
> 2*mean(1/u^2*exp(-(1/u-1)^2))
[1] 1.785992

```

(The true answer is $\sqrt{\pi} \approx 1.77$).

- (b) $\int_0^1 \int_0^1 \exp(-(x+y)^2) dy dx = \mathbb{E}[\exp(-(U_1 + U_2)^2)]$, where U_1, U_2 are independent uniformly distributed random variables on $[0, 1]$. Hence we can approximate the integral as:

```

> u1=runif(10000)
> u2=runif(10000)
> mean(exp((u1+u2)^2))
[1] 4.832706

```

- (c) We can write

$$\int_0^{\infty} \int_0^x \exp(-(x+y)) dy dx = \int_0^{\infty} \int_0^{\infty} \exp(-(x+y)) \mathbb{I}\{y < x\} dy dx.$$

Now make the transformations $u_1 = 1/(1+x)$ and $u_2 = 1/(1+y)$ so that the integrals on $[0, \infty)$ are transformed to $[0, 1]$. Hence we have

$$\int_0^{\infty} \int_0^x \exp(-(x+y)) \mathbb{I}\{y < x\} dy dx = \int_0^1 \int_0^1 \frac{1}{u_1^2 u_2^2} \exp\left(-\left(\frac{1}{u_1} + \frac{1}{u_2} - 2\right)\right) \mathbb{I}\{u_1 < u_2\} du_1 du_2.$$

So now we can approximate the integral by:

```

> u1=runif(10000)
> u2=runif(10000)
> z1=1/u1-1
> z2=1/u2-1
> mean(1/u1^2/u2^2*exp(-(z1+z2))*(u1<u2))
[1] 0.5084041

```

(The true answer is 0.5).