

# FINITE-QUARK-MASS EFFECTS ON THE HIGGS PRODUCTION CROSS SECTION IN THE GLUON-GLUON FUSION CHANNEL

Dissertation

zur Erlangung des Doktorgrades an der Fakultät für Mathematik, Informatik und  
Naturwissenschaften

Fachbereich Physik der Rheinisch-Westfälischen Technischen Hochschule

vorgelegt von

TOM CLAUS RUDOLF SCHELLENBERGER

Aachen

1. Juli 2025



*Finite-Quark-Mass Effects on the Higgs Production Cross Section in the Gluon-Gluon Fusion Channel*

© Tom Claus Rudolf Schellenberger 2025

**GUTACHTER DER DISSERTATION:**

Prof. Dr. Michał Czakon  
Prof. Dr. Robert Harlander

**ZUSAMMENSETZUNG DER PRÜFUNGSKOMMISSION:**

Prof. Dr. Michał Czakon  
Prof. Dr. Robert Harlander  
TBA  
TBA

**VORSITZENDER DER PRÜFUNGSKOMMISSION:**

TBA

**DATUM DER DISPUTATION:**

TBA

**DEKAN DER FAKULTÄT MIN:**

Prof. Dr. Carsten Honerkamp

## ABSTRACT

This is my Abstract

# ZUSAMMENFASSUNG

Dies ist meine Zusammenfassung



## ACKNOWLEDGMENTS

Here is where I thank god.





## PUBLICATIONS

During my PhD studies, I co-authored the following publications:

- [1] Michał Czakon, Felix Eschment, and Tom Schellenberger. “Revisiting the double-soft asymptotics of one-loop amplitudes in massless QCD.” In: *JHEP* 04 (2023), p. 065. DOI: [10.1007/JHEP04\(2023\)065](https://doi.org/10.1007/JHEP04(2023)065). arXiv: [2211.06465](https://arxiv.org/abs/2211.06465) [hep-ph]
- [2] Michał Czakon, Felix Eschment, and Tom Schellenberger. “Subleading effects in soft-gluon emission at one-loop in massless QCD.” In: *JHEP* 12 (2023), p. 126. DOI: [10.1007/JHEP12\(2023\)126](https://doi.org/10.1007/JHEP12(2023)126). arXiv: [2307.02286](https://arxiv.org/abs/2307.02286) [hep-ph]
- [3] Michał Czakon et al. “Top-Bottom Interference Contribution to Fully Inclusive Higgs Production.” In: *Phys. Rev. Lett.* 132.21 (2024), p. 211902. DOI: [10.1103/PhysRevLett.132.211902](https://doi.org/10.1103/PhysRevLett.132.211902). arXiv: [2312.09896](https://arxiv.org/abs/2312.09896) [hep-ph]
- [4] Michał Czakon et al. “Quark mass effects in Higgs production.” In: *JHEP* 10 (2024), p. 210. DOI: [10.1007/JHEP10\(2024\)210](https://doi.org/10.1007/JHEP10(2024)210). arXiv: [2407.12413](https://arxiv.org/abs/2407.12413) [hep-ph]

Among these, only the last two are directly relevant to this dissertation.



# CONTENTS

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>The Standard Model of Particle Physics</b>	<b>3</b>
2.1	Electroweak Symmetry breaking . . . . .	3
2.2	Cross Sections . . . . .	7
2.2.1	The Hard Scattering Amplitude . . . . .	8
2.2.2	The Phase-Space Integration . . . . .	8
2.2.3	The Parton Distribution Functions . . . . .	8
<b>3</b>	<b>Hadronic Higgs Production</b>	<b>11</b>
3.1	The Leading-Order Cross Section . . . . .	11
3.2	The Heavy-Top Limit . . . . .	11
3.3	Higher-Order Corrections . . . . .	11
3.4	Theory Status . . . . .	11
<b>4</b>	<b>Computational Details</b>	<b>13</b>
4.1	Computing the Amplitudes . . . . .	13
4.1.1	The Real-Real Corrections . . . . .	13
4.1.2	The Real-Virtual Corrections . . . . .	13
4.1.3	The Virtual-Virtual Corrections . . . . .	13
4.2	$\overline{\text{MS}}$ -scheme . . . . .	13
4.3	The 4-Flavour Scheme . . . . .	13
4.4	Performing the Phase-Space Integration . . . . .	13
<b>5</b>	<b>Results and Discussion</b>	<b>15</b>
5.1	Total Cross Section . . . . .	15
5.1.1	Effects of Finite Top-Quark Masses . . . . .	15
5.1.2	Effects of Finite Bottom-Quark Masses . . . . .	15
5.2	Differential Cross Section . . . . .	15
<b>6</b>	<b>Conclusions</b>	<b>17</b>
<b>A</b>	<b>Feynman Rules of the Standard Model</b>	<b>19</b>
	<b>Bibliography</b>	<b>23</b>

## ACRONYMS

SM	Standard model
VEV	Vacuum expectation value
SSB	Spontaneous symmetry breaking
PDF	Parton distribution function
QCD	Quantum chromodynamics
RGE	Renormalization group equation
LO	Leading order

## NOTATION AND CONVENTIONS

- In this thesis, I will be using the *Einstein summation convention*, by which any index—be it a Lorentz, color or flavor index—which appears twice is implicitly summed over.

- I will be using natural units

$$\hbar = c = 1. \quad (0.1)$$

- The electron charge is

$$-e, \quad e > 0. \quad (0.2)$$

- The metric is

$$g_{\mu\nu} = \begin{pmatrix} 1 & & & \\ & -1 & & \\ & & -1 & \\ & & & -1 \end{pmatrix}. \quad (0.3)$$

- The normalization of the Levi-Civita anti-symmetric tensor  $\epsilon_{\mu\nu\rho\sigma}$  is

$$\epsilon_{0123} = +1 \quad (0.4)$$

- The Pauli matrices are defined as

$$\sigma^1 \equiv \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 \equiv \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 \equiv \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \tau^i \equiv \sigma^i. \quad (0.5)$$



# 1 | INTRODUCTION

General introductions





# 2

## THE STANDARD MODEL OF PARTICLE PHYSICS

### 2.1 ELECTROWEAK SYMMETRY BREAKING

The standard model (SM) of particle physics is a theory describing all known matter and their fundamental interactions except for gravity. It unifies the electromagnetic, weak, and strong forces under a single theoretical framework. The matter content of the SM is classified into two primary groups: *fermions* and *bosons*. The fermions have spin 1/2, they are further subcategorized into *quarks* and *leptons*. Quarks participate in strong interactions, while leptons interact only via the electromagnetic and weak forces. In contrast, bosons have integer spin. There exists a single particle with spin 0, the *Higgs* boson, and four vector bosons, namely the gluon, the photon, and the *W* and *Z* boson. The vector bosons act as force carrier for the strong, the electromagnetic and the weak force respectively. Fermions are organized into three generations, with each generation containing two types of quarks (up-type and down-type) and two leptons (a charged lepton and its corresponding neutrino). These generations are shown in Fig. 2.1, along with their masses, charges, and spins.

The interactions between SM particles are described by a non-abelian gauge theory of the  $SU(3)_C \times SU(2)_L \times U(1)_Y$  group. Here  $SU(3)_C$  governs the strong interactions. It applies to all *colored* particles, i.e. quarks and gluons. The quarks transform under the fundamental representation of the  $SU(3)_C$  group.  $SU(2)_L \times U(1)_Y$  governs electroweak interactions. The  $SU(2)_L$  transformation acts non-trivially only on *left-handed* fermions which form doublets

$$L_{iL} \equiv \begin{pmatrix} \nu_{iL} \\ l_{iL} \end{pmatrix}, \quad Q_{iL} \equiv \begin{pmatrix} u_{iL} \\ d_{iL} \end{pmatrix}, \quad (2.1)$$


$$\nu_i = (\nu_e, \nu_\mu, \nu_\tau), \quad l_i = (e, \mu, \tau), \quad u_i = (u, c, t), \quad d_i = (d, s, b).$$

The phase transformation  $U(1)_Y$  acts on all particles except neutrinos according to their quantum number, the *hypercharge*  $Y$ . The symmetry is spontaneously broken to  $SU(3)_C \times U(1)_Q$ , where the  $U(1)_Q$  group corresponds to gauge transformation of the electromagnetic interaction, hence the subscript  $Q$  for the *electric charge*. To ensure that the particles have the correct charges, the hypercharge must satisfy the *Gell-Mann–Nishijima relation*:

$$\frac{Y}{2} = Q - I^3. \quad (2.2)$$

With the particle charges displayed in Fig. 2.1 we then get

	$L_{iL}$	$Q_{iL}$	$\nu_{iR}$	$l_{iR}$	$u_{iR}$	$d_{iR}$
$\frac{Y}{2}$	$-\frac{1}{2}$	$\frac{1}{6}$	0	-1	$\frac{2}{3}$	$-\frac{1}{3}$



**Figure 2.1:** Elementary particles of the SM. The was image generated with the help of Ref. [5].

The transformation properties of the gauge bosons is dictated by the covariance of the covariant derivative

$$D_\mu \equiv \partial_\mu - igA_\mu^a T_R^a - ig_2 W_\mu^a I^a + ig_Y \frac{Y}{2} B_\mu$$

$$T_R^a = \begin{cases} T^a & \text{for quarks,} \\ 0 & \text{for leptons,} \end{cases} \quad I^a = \begin{cases} \frac{\tau}{2} & \text{for left-handed fermions,} \\ 0 & \text{for right-handed fermions,} \end{cases} \quad (2.3)$$

where  $T^a$  and  $\tau^a$  are the *Gell-Mann* and *Pauli* matrices.

Before spontaneous symmetry breaking, the Lagrangian which governs the evolution of all matter fields must be invariant under the  $SU(3)_C \times SU(2)_L \times U(1)_Y$  gauge group. Up to a  $\mathbb{CP}$  violating term<sup>1</sup> the SM Lagrangian is the most general mass-dimension four Lagrangian for the described particle content

$$\mathcal{L}_{\text{SM}} = \mathcal{L}_G + \mathcal{L}_F + \mathcal{L}_Y + \mathcal{L}_H. \quad (2.4)$$

The gauge-field Lagrangian  $\mathcal{L}_G$  describes the free propagation and in the case of the non-abelian groups  $SU(3)_C$  and  $SU(2)_L$  also the self-interaction of the gauge bosons. It is given by

$$\mathcal{L}_G = -\frac{1}{4} G_{\mu\nu}^a G^{a\mu\nu} - \frac{1}{4} W_{\mu\nu}^a W^{a\mu\nu} - \frac{1}{4} B_{\mu\nu} B^{\mu\nu},$$

$$G_{\mu\nu}^a \equiv \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f^{abc} A_\mu^b A_\nu^c,$$

$$W_{\mu\nu}^a \equiv \partial_\mu W_\nu^a - \partial_\nu W_\mu^a + g_2 \epsilon^{abc} W_\mu^b W_\nu^c,$$

$$B_{\mu\nu} \equiv \partial_\mu B_\nu - \partial_\nu B_\mu. \quad (2.5)$$

<sup>1</sup> The absence of the  $\mathbb{CP}$  violating term  $\theta \frac{g^2}{64\pi^2} \epsilon^{\mu\nu\alpha\beta} F_{\mu\nu}^a F_{\alpha\beta}^a$  is an unsolved problem of particle physics, known as the strong CP problem.

The propagation of the fermions and their interaction with the gauge bosons is described by

$$\mathcal{L}_F = \bar{L}_{iL} i \not{D} L_{iL} + \bar{\nu}_{iR} i \not{D} \nu_{iR} + \bar{l}_{iR} i \not{D} l_{iR} + \bar{Q}_{iL} i \not{D} Q_{iL} + \bar{u}_{iR} i \not{D} u_{iR} + \bar{d}_{iR} i \not{D} d_{iR}. \quad (2.6)$$

The Higgs field is a doublet of the  $SU(2)_L$  group. We want the field to have a non-vanishing vacuum expectation value (VEV) to dynamically generate the fermion and boson masses. Of course, the vacuum cannot carry an electric charge, which means that the Higgs field must be electrically neutral along the direction of *spontaneous symmetry breaking* (SSB). We choose this to be the second component of the doublet. With the Gell-Mann–Nishijima relation we can then deduce that hypercharge of the doublet must be  $Y = +1$ . The Higgs doublet field thus takes the form

$$\Phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}, \quad (2.7)$$

where the superscript indicates the electric charge.

In order to generate a non-vanishing VEV, the Higgs field must be in a potential with a global minimum away from zero. Hence, the only gauge invariant mass-dimension four Lagrangian we can construct is

$$\begin{aligned} \mathcal{L}_H &= (D_\mu \Phi)^\dagger (D^\mu \Phi) - V(\Phi) \\ V(\Phi) &= \lambda(\Phi^\dagger \Phi)^2 - \mu^2 \Phi^\dagger \Phi, \quad \mu^2, \lambda > 0. \end{aligned} \quad (2.8)$$

The minimum of the Higgs potential  $V$  is at

$$\Phi_0^\dagger \Phi_0 = \frac{\mu^2}{2\lambda} \equiv \frac{v^2}{2} \neq 0. \quad (2.9)$$

After (SSB) we can expand the Higgs field around its minimum

$$\Phi = \begin{pmatrix} \phi^+ \\ \frac{1}{\sqrt{2}}(v + H + i\xi) \end{pmatrix} \quad (2.10)$$

The real scalar field  $H$  is the famous Higgs boson, whereas the fields  $\phi^\pm$  and  $\xi$  are unphysical since they can always be eliminated through a gauge transformation (*would-be Goldstone bosons*). After inserting the expansion in the Higgs Lagrangian, the mass of the Higgs can be read off from its square term

$$m_H = \sqrt{2}\mu. \quad (2.11)$$

SSB enables the generation of vector boson masses without breaking the gauge symmetry explicitly. If we insert the expanded Higgs field in the Higgs Lagrangian, we get quadratic terms of the gauge boson fields

$$\begin{aligned} \mathcal{L}_H &\supseteq \frac{v^2}{2} \left\{ g_2^2 \begin{pmatrix} 0 & 1 \end{pmatrix} I^a I^b \begin{pmatrix} 0 \\ 1 \end{pmatrix} W_\mu^a W^{b\mu} - g_2 g_Y \begin{pmatrix} 0 & 1 \end{pmatrix} I^a \begin{pmatrix} 0 \\ 1 \end{pmatrix} W_\mu^a B^\mu + \frac{g_Y^2}{4} B_\mu B^\mu \right\} \\ &= \frac{v^2}{2} \left\{ \frac{g_2^2}{4} [(W^1)^2 + (W^2)^2] + \frac{1}{4} \begin{pmatrix} B^\mu & W^{3\mu} \end{pmatrix} \begin{pmatrix} g_Y^2 & g_Y g_2 \\ g_Y g_2 & g_2^2 \end{pmatrix} \begin{pmatrix} B_\mu \\ W_\mu^3 \end{pmatrix} \right\}. \end{aligned} \quad (2.12)$$

By diagonalizing the mass matrix we obtain the physical states

$$\begin{pmatrix} A_\mu^\gamma \\ Z_\mu \end{pmatrix} = \begin{pmatrix} \cos \theta_W & -\sin \theta_W \\ \sin \theta_W & \cos \theta_W \end{pmatrix} \begin{pmatrix} B_\mu \\ W_\mu^3 \end{pmatrix}, \quad \cos \theta_W = \frac{g_2}{\sqrt{g_Y^2 + g_2^2}}, \quad \sin \theta_W = \frac{g_Y}{\sqrt{g_Y^2 + g_2^2}}. \quad (2.13)$$

In this new basis, we have one massless boson  $A_\mu^\gamma$ , which we identify as the photon and a charge neutral boson of mass

$$m_Z = \frac{v}{2} \sqrt{g_Y^2 + g_2^2}. \quad (2.14)$$

The vector bosons  $W^1$  and  $W^2$  are not eigenstates of the charge operator. We therefore define the new states

$$W_\mu^\pm = \frac{1}{\sqrt{2}} (W_\mu^1 \mp iW_\mu^2), \quad QW_\mu^\pm = \pm W_\mu^\pm, \quad (2.15)$$

which are eigenstates of  $Q$  and have mass

$$m_W = \frac{v}{2} g_2. \quad (2.16)$$

Last but not least, we discuss the Yukawa sector of the SM Lagrangian. Before SSB, fermions cannot generate masses because a mass term would mix the left- and right-handed components of the fields, thereby breaking the chiral gauge symmetry. Here, once again, the Higgs field comes to the rescue: by coupling the fermions with the Higgs field through a Yukawa interaction<sup>2</sup>

$$\mathcal{L}_Y = - \left( y_{ij}^\nu \bar{L}_{iL} \Phi^c \nu_{jR} + y_{ij}^l \bar{L}_{iL} \Phi l_{jR} + y_{ij}^d \bar{Q}_{iL} \Phi d_{jR} + y_{ij}^u \bar{Q}_{iL} \Phi^c u_{jR} \right) + \text{h.c.}, \quad (2.17)$$

where  $\Phi^c$  is the charge-conjugated field to  $\Phi$ , we do not explicitly break the symmetry. However, after SSB this Lagrangian will generate exactly the required mixing between left- and right-handed fields to generate the fermion masses. The Yukawa-interaction matrices  $y_{ij}^{\nu, l, d, u}$  can be shifted from the Yukawa sector to the fermion sector through a field redefinition. Indeed, if we apply the *singular value decomposition* of the Yukawa matrix

$$y = U_L^\dagger y_{\text{diag}} U_R, \quad \text{with} \quad (y_{\text{diag}})_{ij} = m_i \delta_{ij} \quad \text{and} \quad U_{L,R} \in \text{U}(3), \quad (2.18)$$

and redefine our fermion fields to be

$$f_{iR} \longrightarrow U_{Rij} f_{jR}, \quad f_{iL} \longrightarrow U_{Lij} f_{jL}, \quad f = \nu, l, u, d \quad (2.19)$$

the Yukawa Lagrangian becomes

$$\mathcal{L}_Y = - \sum_i \left( m_{\nu_i} \bar{\nu}_i \nu_i + m_{l_i} \bar{l}_i l_i + m_{u_i} \bar{u}_i u_i + m_{d_i} \bar{d}_i d_i \right) \left( 1 + \frac{H}{v} \right). \quad (2.20)$$

As an immediate consequence, we observe that the Yukawa coupling of the Higgs to the fermions is proportional to the mass of that fermion. The field redefinition is

<sup>2</sup> In the original formulation of the SM, there are no neutrino Yukawa interactions, since they were believed to be massless. Neutrino oscillation experiments have shown however, that neutrinos do in fact have finite masses.

a change from a flavor eigenbasis, which is diagonal in the couplings to the gauge bosons, to a mass eigenbasis. In the mass eigenbasis the part of fermion Lagrangian which contains the interaction to the electroweak gauge bosons after SSB is

$$\begin{aligned} \mathcal{L}_F \supseteq & \sum_f (-Q_f) e \bar{f}_i \not{A} f_i + \sum_f \frac{e}{\sin \theta_W \cos \theta_W} \bar{f}_i (I_f^3 \gamma^\mu P_L - \sin^2 \theta_W Q_f \gamma^\mu) f_i Z_\mu \\ & + \frac{e}{\sqrt{2} \sin \theta_W} \left( \bar{u}_i \gamma^\mu P_L (V_{\text{CKM}})_{ij} d_j W_\mu^+ + \bar{d}_i \gamma^\mu P_L (V_{\text{CKM}}^\dagger)_{ij} u_j W_\mu^- \right) \\ & + \frac{e}{\sqrt{2} \sin \theta_W} \left( \bar{\nu}_i \gamma^\mu P_L (V_{\text{PMNS}}^\dagger)_{ij} l_j W_\mu^+ + \bar{l}_i \gamma^\mu P_L (V_{\text{PMNS}})_{ij} \nu_j W_\mu^- \right). \end{aligned} \quad (2.21)$$

Here we identified the electromagnetic coupling constant

$$e = \frac{g_2 g_Y}{\sqrt{g_2^2 + g_Y^2}}, \quad (2.22)$$

as the factor in front of the photon interaction term. The operators  $P_{L,R}$  are just the projectors onto the left- and right-handed components

$$P_{L,R} = \frac{1 \mp \gamma^5}{2}. \quad (2.23)$$

The *CKM* and *PMNS matrices*<sup>3</sup> are the results of the field redefinitions

$$V_{\text{CKM}} \equiv U_L^{u\dagger} U_L^d, \quad V_{\text{PMNS}} \equiv U_L^{l\dagger} U_L^\nu. \quad (2.24)$$

Typically, one prefers to work in the mass eigenbasis of the quarks, while the neutrinos are kept in the flavor eigenbasis, in which case one encounters flavor changes (*neutrino oscillations*) through propagation. This is why the PMNS matrix is defined in terms of the complex conjugate of the CKM matrix equivalent in the lepton sector.

## 2.2 CROSS SECTIONS

Many of the great successes of the SM are its *cross section* predictions. The cross section is simply defined as the probability to create some final state from some initial state per unit of time per target particle divided by the incoming particle flux. This means that the cross section can be easily measured with a simple counting experiment. Experiments like **CMS** of **Atlas** do exactly that: they smash particles together and count how many times a certain final state was produced in some time interval. On the theory side, the cross section can be calculated with

$$d\hat{\sigma}_{ij \rightarrow n} = \frac{1}{F} d\Phi_n |M_{ij \rightarrow n}|^2, \quad (2.25)$$

where  $F$  is the *flux factor*<sup>4</sup>

$$F \equiv 4p_1 \cdot p_2, \quad (2.26)$$

$d\Phi_n$  is the *Lorentz invariant phase space measure*

$$d\Phi_n = \prod_{i=1}^n \frac{d^3 \mathbf{q}_i}{(2\pi)^3 2q_i^0} (2\pi)^4 \delta^{(4)} \left( p_1 + p_2 - \sum_{i=1}^n q_i \right), \quad (2.27)$$

<sup>3</sup> Named after Cabibbo, Kobayashi and Maskawa, and Pontecorvo, Maki, Nakagawa and Sakata.

<sup>4</sup> In the following we assume that the initial state particles are massless.

and  $M_{fi}$  is the *scattering amplitude* describing the hard interactions.

The computation of cross sections involves three basic steps:

1. the calculation of the hard scattering amplitude,
2. the phase-space integration,
3. and the convolution with *parton distribution functions* (PDFs).

In the following we will discuss them one-by-one.

### 2.2.1 The Hard Scattering Amplitude

The Hard Scattering Amplitude describes the transition probability from a certain initial state to a specific final state. Since the scattering is **hard**, the energy transfer between the particles during the scattering process is large compared to the QCD scale. This means we are in the perturbative regime of QCD, and we can perform an expansion in the coupling constant

$$M_{ij \rightarrow n} = \alpha_s^{n_{\text{Born}}} \left( M_{ij \rightarrow n}^{(0)} + \frac{\alpha_s}{\pi} M_{ij \rightarrow n}^{(1)} + \left( \frac{\alpha_s}{\pi} \right)^2 M_{ij \rightarrow n}^{(2)} + \mathcal{O}(\alpha_s^3) \right). \quad (2.28)$$

$n_{\text{Born}}$  is the power of the coupling constant at *leading order* (LO). The coefficients of the series are calculable graphically using *Feynman rules*.

### 2.2.2 The Phase-Space Integration


### 2.2.3 The Parton Distribution Functions

In hadron collisions, the initial state is not made up of elementary particles, but are bound states thereof. This means that during an inelastic scattering event, the partons which take part in the short-range interaction only carry a fraction of the original hadron momentum

$$p_1 = x_1 P_1, \quad p_2 = x_2 P_2. \quad (2.29)$$

Here  $p_1$  and  $p_2$  are the momenta of the partons and  $P_1$  and  $P_2$  are the momenta of the hadrons. Since the momentum of the parton can not be larger than that of the hadron,  $x_{1,2}$  is restricted to be less than one. Furthermore, since the energy of the parton must be positive the momentum fraction must also be positive. Otherwise, the momentum fraction is a priori unconstrained, we therefore integrate over all allowed values of  $x_1$  and  $x_2$

$$d\sigma_{H_1 H_2 \rightarrow n} = \sum_{i,j} \int_0^1 dx_1 dx_2 f_{H_1,i}(x_1, \mu_F) f_{H_2,j}(x_2, \mu_F) d\hat{\sigma}_{ij \rightarrow n}(x_1 P_1, x_2 P_2, \mu_F, \mu_R). \quad (2.30)$$



**Figure 2.2:** The various PDFs multiplied by  $x$  as a function of  $x$ . The plot was created using the LHAPDF6 [6] interface to the NNPDF31\_nnlo\_as\_0118 [7] PDF set at a scale of  $\mu_F = m_H/2$ .

$f_{H_k,i}(x_k, \mu_F)$  are the PDFs. They describe the probability of finding a parton  $i$  with momentum fraction  $x_k$  inside the hadron  $H_k$ . Momentum conservation then requires the normalization

$$\sum_i \int_0^1 dx x f_{H,i}(x, \mu_F) = 1. \quad (2.31)$$

The factorization theorem (2.30) is central in the SM as it tells us that the PDFs are universal quantities, i.e. they are not specific to any one process. It is a postulate of the parton model, in which hadrons are thought of as collection of the free elementary particles. In QCD however, the theorem requires proof [8]! The PDF for all light partons are displayed in Fig. 2.2. PDFs describe long range interactions, a regime in which QCD is non-perturbative. As such, PDFs are non-perturbative objects which have to be measured in experiments or be calculated non-perturbatively, e.g. on the lattice.

The scale  $\mu_F$  marks the boundary, over which we treat interactions perturbatively. This scale is unphysical in the sense that it is not a parameter in our theory, nor can it be measured in an experiment. This independence can be precisely formulated in terms of the *renormalization group equation* (RGE)

$$\frac{d}{d \ln \mu_F} f_{H,i}^B = 0, \quad (2.32)$$

which states that the *bare, unrenormalized* PDF can not depend on the unphysical scale  $\mu_F$ . If we now express the unrenormalized PDF in terms of its renormalized counterpart

$$f_{H,i}^B \equiv Z_{ij} \otimes f_{H,j}^R \equiv \int_0^1 dy dz Z_{ij}(y) f_{H,j}^R(z) \delta(x - yz), \quad (2.33)$$

then we can use the RGE to get the factorization scale dependents of the renormalized PDF

$$\begin{aligned} 0 &= \frac{d}{d \ln \mu_F} f_{H,i}^B = \frac{d\alpha_s}{d \ln \mu_F} \frac{dZ_{ij}}{d\alpha_s} \otimes f_{H,j}^R + Z_{ij} \otimes \frac{df_{H,j}}{d \ln \mu_F} \\ \Rightarrow \frac{df_{H,i}^R}{d \ln \mu_F} &= -Z_{ij}^{-1} \otimes (4\pi\beta - 2\epsilon\alpha_s) \frac{dZ_{jk}}{d\alpha_s} \otimes f_{H,k}^R = 2\alpha_s Z_{ij}^{-1} \otimes \frac{dZ_{ij}^{(1)}}{d\alpha_s} \otimes f_{H,k}^R. \end{aligned} \quad (2.34)$$

Here we used the definition of the  $d$ -dimensional  $\beta$ -function

$$\bar{\beta} \equiv \frac{1}{4\pi} \frac{d\alpha_s}{d \ln \mu_F} = \beta - \epsilon \frac{\alpha_s}{2\pi}, \quad \beta = \frac{\alpha_s}{2\pi} \frac{dZ^{(1)}}{d \ln \alpha_s}, \quad (2.35)$$

where  $Z_{\alpha_s}^{(1)}$  and  $Z_{ij}^{(1)}$  are the residues of the renormalization constants of the coupling constant and the PDFs respectively. At one loop, the PDF renormalization constant is designed to absorb the singularities from tree-level initial-state-collinear radiation, it therefore reads

$$Z_{ij}(z) = \delta(1-z)\delta_{ij} + \frac{\alpha_s}{2\pi\epsilon} P_{ij}^{(0)}(z), \quad (2.36)$$

with the *Altarelli-Parisi splitting kernels*

$$\begin{aligned} P_{qq}^{(0)}(z) &= P_{\bar{q}\bar{q}}^{(0)}(z) = C_F \left[ \frac{1+z^2}{(1-z)_+} + \frac{3}{2}\delta(1-z) \right], \\ P_{qg}^{(0)}(z) &= T_F [z^2 + (1-z)^2], \\ P_{gq}^{(0)}(z) &= C_F \left[ \frac{1+(1-z)^2}{z} \right], \\ P_{gg}^{(0)}(z) &= 2C_A \left[ \frac{z}{(1-z)_+} + \frac{1-z}{z} + z(1-z) \right] + \delta(1-z)\frac{\beta_0}{2}, \end{aligned} \quad (2.37)$$

and

$$\beta_0 = \frac{11}{3}C_A - \frac{4}{3}T_F n_l. \quad (2.38)$$

So even though the PDFs are non-perturbative, their dependence on the factorization scale is perturbative and calculable with the *Dokshitzer-Gribov-Lipatow-Altarelli-Parisi-evolution equation* [9, 10, 11]

$$\frac{df_{H,i}^R}{d \ln \mu_F} = \frac{\alpha_s}{\pi} P_{ij}^{(0)} \otimes f_{H,j}^R. \quad (2.39)$$



## 3 | HADRONIC HIGGS PRODUCTION

### 3.1 THE LEADING-ORDER CROSS SECTION

Here I compute the leading-order cross section for Higgs Production in the gluon-gluon fusion channel.

### 3.2 THE HEAVY-TOP LIMIT


Here I explain the heavy-top limit.

### 3.3 HIGHER-ORDER CORRECTIONS

Here I outline how to perform higher order corrections.

### 3.4 THEORY STATUS

Here I describe what is already known about the gluon-gluon fusion channel. I explain the theory uncertainties.



**Figure 3.1:** Illustration of the partonic luminosity of parton  $i$  and  $j$ . We choose exemplary parton combinations to represent gluon–gluon, gluon–valence-quark, gluon–sea-quark, valence-quark–sea-quark, valence-quark–valence-quark and sea-quark–sea-quark luminosities. For the luminosity of two different partons we include a factor 2 to account for the different flavor permutations. The luminosity was calculated using the LHAPDF6 [6] interface to the NNPDF31\_nnlo\_as\_0118 [7] PDF set at a scale of  $\mu_F = m_H/2$  with a fixed collider energy of  $\sqrt{S} = 13$  TeV.

## 4 | COMPUTATIONAL DETAILS

Description of this chapter.

### 4.1 COMPUTING THE AMPLITUDES

Here I

#### 4.1.1 The Real-Real Corrections

#### 4.1.2 The Real-Virtual Corrections

#### 4.1.3 The Virtual-Virtual Corrections

### 4.2 $\overline{\text{MS}}$ -SCHEME

### 4.3 THE 4-FLAVOUR SCHEME

### 4.4 PERFORMING THE PHASE-SPACE INTEGRATION



# 5 | RESULTS AND DISCUSSION

Description of this chapter.

## 5.1 TOTAL CROSS SECTION

### 5.1.1 Effects of Finite Top-Quark Masses

### 5.1.2 Effects of Finite Bottom-Quark Masses

## 5.2 DIFFERENTIAL CROSS SECTION



## 6 | CONCLUSIONS

Here are my conclusions.






In this chapter we list all Feynman rules of the SM. We choose to work in a unitary gauge, meaning that the all Goldstone-bosons decouple and there are no unphysical particles in the electroweak sector. In the QCD sector, we work in the  $R_\xi$  gauge, i.e. we have unphysical particles in the form of *Faddeev-Popov ghosts*. If not otherwise specified, the momenta on every line are defined as incoming.

### Propagators:


[illegible]

### Fermion–Gauge-Boson Vertices:




$$= ig\gamma^\mu T_{ij}^a$$

$a, \mu$




$$= -ie\gamma^\mu Q$$

$\mu$




$$= gf^{abc}p^\mu$$

$c, \mu$



$$= i\frac{e}{\sin\theta_W\cos\theta_W}\gamma^\mu \times (I^3P_L - Q\sin^2\theta_W)$$


$\mu$




$$= i \frac{e(V_{\text{CKM}})_{ij}}{\sqrt{2} \sin \theta_W} \gamma^\mu P_L$$

$$= i \frac{e(V_{\text{CKM}})_{ji}^*}{\sqrt{2} \sin \theta_W} \gamma^\mu P_L$$


### Gauge-Boson Self Interactions:




$$= g f^{abc} ((p_1^\rho - p_2^\rho) g^{\mu\nu} + (p_2^\mu - p_3^\mu) g^{\nu\rho} + (p_3^\nu - p_1^\nu) g^{\rho\mu})$$




$$= -i g^2 (f^{abe} f^{cde} (g^{\mu\rho} g^{\nu\sigma} - g^{\mu\sigma} g^{\nu\rho}) + f^{ace} f^{bde} (g^{\mu\nu} g^{\rho\sigma} - g^{\mu\sigma} g^{\nu\rho}) + f^{ade} f^{bce} (g^{\mu\nu} g^{\rho\sigma} - g^{\mu\rho} g^{\nu\sigma}))$$



$$= i \frac{e}{\sin \theta_W} ((p_1^\rho - p_2^\rho) g^{\mu\nu} + (p_2^\mu - p_3^\mu) g^{\nu\rho} + (p_3^\nu - p_1^\nu) g^{\rho\mu}) \times \begin{cases} -\sin \theta_W & \text{for } \gamma \\ \cos \theta_W & \text{for } Z \end{cases}$$



$$= i \frac{e^2}{\sin^2 \theta_W} (g^{\mu\sigma} g^{\nu\rho} + g^{\mu\rho} g^{\nu\sigma} - 2g^{\mu\nu} g^{\rho\sigma}) \times \prod_{i=1}^2 \begin{cases} -\sin \theta_W & \text{if } V_i = \gamma \\ \cos \theta_W & \text{if } V_i = Z \end{cases}$$




$$= i \frac{e^2}{\sin^2 \theta_W} (2g^{\mu\rho} g^{\nu\sigma} - g^{\mu\nu} g^{\rho\sigma} - g^{\mu\sigma} g^{\nu\rho})$$

### Higgs Interactions:




$$= -i \frac{m_q}{v} \delta_{ij}$$


$$= -i \frac{3m_H^2}{v}$$




$$= i \frac{e}{\sin \theta_W} m_W g^{\mu\nu}$$




$$= i \frac{e^2}{2 \sin^2 \theta_W} g^{\mu\nu}$$



$$= i \frac{e}{\sin \theta_W \cos \theta_W} m_Z g^{\mu\nu}$$



$$= i \frac{e^2}{2 \sin^2 \theta_W \cos^2 \theta_W} g^{\mu\nu}$$



$$= -i \frac{3m_H^2}{v^2}$$



## BIBLIOGRAPHY

- [1] Michał Czakon, Felix Eschment, and Tom Schellenberger. “Revisiting the double-soft asymptotics of one-loop amplitudes in massless QCD.” In: *JHEP* 04 (2023), p. 065. DOI: [10.1007/JHEP04\(2023\)065](https://doi.org/10.1007/JHEP04(2023)065). arXiv: [2211.06465](https://arxiv.org/abs/2211.06465) [hep-ph].
- [2] Michał Czakon, Felix Eschment, and Tom Schellenberger. “Subleading effects in soft-gluon emission at one-loop in massless QCD.” In: *JHEP* 12 (2023), p. 126. DOI: [10.1007/JHEP12\(2023\)126](https://doi.org/10.1007/JHEP12(2023)126). arXiv: [2307.02286](https://arxiv.org/abs/2307.02286) [hep-ph].
- [3] Michał Czakon et al. “Top-Bottom Interference Contribution to Fully Inclusive Higgs Production.” In: *Phys. Rev. Lett.* 132.21 (2024), p. 211902. DOI: [10.1103/PhysRevLett.132.211902](https://doi.org/10.1103/PhysRevLett.132.211902). arXiv: [2312.09896](https://arxiv.org/abs/2312.09896) [hep-ph].
- [4] Michał Czakon et al. “Quark mass effects in Higgs production.” In: *JHEP* 10 (2024), p. 210. DOI: [10.1007/JHEP10\(2024\)210](https://doi.org/10.1007/JHEP10(2024)210). arXiv: [2407.12413](https://arxiv.org/abs/2407.12413) [hep-ph].
- [5] Izaak Neutelings. *Izaak neutelings*. Mar. 2024. URL: [https://tikz.net/sm\\_particles/](https://tikz.net/sm_particles/).
- [6] Andy Buckley et al. “LHAPDF6: parton density access in the LHC precision era.” In: *Eur. Phys. J. C* 75 (2015), p. 132. DOI: [10.1140/epjc/s10052-015-3318-8](https://doi.org/10.1140/epjc/s10052-015-3318-8). arXiv: [1412.7420](https://arxiv.org/abs/1412.7420) [hep-ph].
- [7] Richard D. Ball et al. “Parton distributions from high-precision collider data.” In: *Eur. Phys. J. C* 77.10 (2017), p. 663. DOI: [10.1140/epjc/s10052-017-5199-5](https://doi.org/10.1140/epjc/s10052-017-5199-5). arXiv: [1706.00428](https://arxiv.org/abs/1706.00428) [hep-ph].
- [8] John C. Collins, Davison E. Soper, and George F. Sterman. “Factorization of Hard Processes in QCD.” In: *Adv. Ser. Direct. High Energy Phys.* 5 (1989), pp. 1–91. DOI: [10.1142/9789814503266\\_0001](https://doi.org/10.1142/9789814503266_0001). arXiv: [hep-ph/0409313](https://arxiv.org/abs/hep-ph/0409313).
- [9] Yuri L. Dokshitzer. “Calculation of the Structure Functions for Deep Inelastic Scattering and  $e^+e^-$  Annihilation by Perturbation Theory in Quantum Chromodynamics.” In: *Sov. Phys. JETP* 46 (1977), pp. 641–653.
- [10] V. N. Gribov and L. N. Lipatov. “Deep inelastic  $e p$  scattering in perturbation theory.” In: *Sov. J. Nucl. Phys.* 15 (1972), pp. 438–450.
- [11] Guido Altarelli and G. Parisi. “Asymptotic Freedom in Parton Language.” In: *Nucl. Phys. B* 126 (1977), pp. 298–318. DOI: [10.1016/0550-3213\(77\)90384-4](https://doi.org/10.1016/0550-3213(77)90384-4).