

Computational Identification of Non- β -Lactam PBP2a Inhibitors Against MRSA via Structure-Based Virtual Screening

July 30, 2026

Abstract

Methicillin-resistant *Staphylococcus aureus* (MRSA) remains a persistent clinical threat, with penicillin-binding protein 2a (PBP2a) as the primary resistance determinant. We performed a consensus rigid-docking virtual screen against the active site of MRSA PBP2a (PDB 3ZG0, 1VQQ, 4DKI) using AutoDock Vina 1.2.7. The compound library of 3,116 diverse compounds was assembled from multiple seed libraries and filtered through a cascade of β -lactam exclusion, similarity filtering, ADMET profiling, PAINS alerts, and Brenk alerts. Redocking validation of the native ligand ceftaroline yielded a core RMSD of 1.251 Å ("Validated" badge). Two lead compounds were identified with $SI \geq 2.0$ (Strong tier): BRICS_0022 ($SI = 2.13$, $E_{\text{PBP2a}} = -10.52 \text{ kcal mol}^{-1}$) and ALL_QU04 ($SI = 2.07$, $E_{\text{PBP2a}} = -10.07 \text{ kcal mol}^{-1}$), both forming hydrogen-bond contacts with the catalytic Ser403, Lys406, and Tyr446 triad. A third scaffold, ALL_QU05 ($E_{\text{PBP2a}} = -9.77 \text{ kcal mol}^{-1}$, $SI = 1.23$), is reported as a secondary candidate with favourable synthetic accessibility ($SA = 2.13$). Enrichment benchmarking against 21 known PBP2a binders yielded $AUC = 0.792$ and $EF_{1\%} = 8.14$. The pipeline provides a validated, reproducible framework for PBP2a-targeted virtual screening, prioritising BRICS_0022 and ALL_QU04 as drug-like, non- β -lactam scaffolds for experimental validation.

1 Introduction

β -Lactam antibiotics target penicillin-binding proteins (PBPs), the transpeptidases that cross-link the bacterial cell wall [1]. In MRSA, acquisition of the exogenous PBP2a confers resistance to virtually all β -lactams because PBP2a's active site has a markedly lower affinity for these substrates. Ceftaroline, a fifth-generation cephalosporin, overcomes this barrier and remains the only β -lactam approved for MRSA infections [2], though resistance has emerged.

Two complementary strategies exist for targeting PBP2a: (i) active-site inhibition, which competes with the native peptidoglycan substrate, and (ii) allosteric inhibition, which stabilises a closed conformation [3]. Several allosteric PBP2a inhibitors have been reported, including oxadiazole and quinazolinone series [4]. However, active-site-directed non- β -lactam

scaffolds remain underexplored due to the inherent challenge of achieving selective binding in a shallow, polar active site.

Here we present the AutoAntibiotic Discovery Pipeline v5.5.0, a fully automated, reproducible consensus docking pipeline that screens diverse non- β -lactam compounds against the PBP2a active site across three crystallographic conformers (3ZG0 holo, 1VQQ apo, 4DKI). The pipeline incorporates native-ligand redocking validation, enrichment benchmarking against known actives, MMFF94-based approximate rescoring, consensus scoring, selectivity profiling against human serine hydrolases (trypsin, CES1) and CYP3A4, and OpenMM Amber14 protein–ligand complex minimisation for the top candidates. The screening library of 3,116 compounds spanning 12 scaffold families was assembled from BRICS-recombined seed scaffolds, known PBP2a allosteric inhibitor series, and focused heterocyclic cores. This work prioritises BRICS_0022 and ALL_QU04 as drug-like, non- β -lactam candidates for experimental validation against MRSA PBP2a.

2 Methods

2.1 Target preparation

Five PDB structures were obtained from the RCSB Protein Data Bank: **3ZG0** (PBP2a holo, ceftaroline-bound, ligand AI8), **1VQQ** (PBP2a apo), **4DKI** (PBP2a alternative conformer), **1UTN** (human trypsin), and **1YAH** (human carboxylesterase 1). Each structure was cleaned by removing solvent molecules and non-target ligands. Polar hydrogens were added via RDKit (or OpenBabel fallback) and receptors were converted to PDBQT format using `obabel -xr`. The active-site grid centre for PBP2a was computed as the side-chain centroid of conserved catalytic residues Ser403, Lys406, and Tyr446. Allosteric-site centres used Tyr105, Gln199, and Glu237. Off-target grid centres used the catalytic triads: His57/Asp102/Ser195 for trypsin and Ser221/His468/Glu354 for CES1. Box sizes were auto-sized per-axis with a 20 Å maximum for PBP2a active site, 18 Å for the allosteric site, and 15 Å for off-targets, each with 2.0 Å padding.

2.2 Redocking validation

The docking protocol was validated by redocking the native co-crystallised ligand ceftaroline (PDB ID: 3ZG0, residue name AI8) into three prepared PBP2a receptor conformers. Vina rigid docking was performed at exhaustiveness 32 with three random seeds per conformer. Full-ligand RMSD was computed via Kabsch alignment over all 27 heavy atoms. Core RMSD used only the 12-atom fused ring scaffold. The best core RMSD across all conformers and seeds defined the protocol trust badge: ≤ 1.5 Å = “Validated”, ≤ 2.0 Å = “Validated (Marginal)”. The redocking box was auto-sized per axis from the native ligand’s spatial spread plus 5.0 Å padding.

2.3 Library generation and filtering

The screening library of 3,116 unique compounds was assembled from multiple seed libraries (including BRICS-recombined scaffolds, focused PBP2a allosteric libraries, and diverse het-

erocyclic cores) and filtered through a sequential cascade: (1) β -lactam SMARTS exclusion, (2) Tanimoto similarity to known antibiotics ($T_c < 0.3$), (3) ADMET filter (Lipinski rule-of-five + QED > 0.3), (4) PAINS alerts, and (5) Brenk alerts. The library spans 12 scaffold families: oxadiazoles, quinazolinones, penicillin-derived cores, benzothiazoles, pyrazolopyrimidines, triazolopyridines, benzothiazole-ureas, imidazopyridines, quinazoline-2,4-diones, pyrazole-sulfonamides, and triazolopyrimidines. Compounds were curated to ensure drug-likeness (MW 200–550, SA < 4.5) and a Bemis-Murcko framework cap (15%) prevented over-representation of any single scaffold class.

2.4 Consensus rigid docking

Docking was performed with AutoDock Vina 1.2.7 [5]. Each compound was docked independently against three PBP2a conformer PDBQTs at exhaustiveness 8 with num_modes 3. The consensus PBP2a binding energy was taken as the most negative (best) energy across all three conformers. Allosteric-site docking used the same protocol with the allosteric grid centre. The top 20 compounds by active-site energy were advanced to selectivity profiling.

2.5 Mechanism-restricted selectivity index

The selectivity index was defined as:

$$SI = \frac{|E_{PBP2a,active}|}{\frac{1}{|T|} \sum_{t \in T} |E_t|} \quad (1)$$

where T is the set of mechanism-relevant human serine hydrolases (trypsin, CES1) with valid negative binding energies. SI tiers: Strong (≥ 2.0), Promising (1.5–2.0), Below gate (< 1.5). Compounds with only one valid off-target energy received a provisional single-target ratio. An additional off-target, CYP3A4 (PDB 1TQN), was profiled for the top 10 candidates as a human liability target; CYP3A4 energies are reported separately and do not enter the SI denominator, consistent with the mechanism-restricted approach.

2.6 Interaction fingerprinting

Hydrogen-bond contacts between docked ligands and catalytic residues (Ser403, Lys406, Tyr446) were evaluated by measuring the minimum heavy-atom distance. A contact was classified as a confirmed hydrogen bond when the distance was $< 3.5 \text{ \AA}$ for Ser403 and Tyr446, and $< 3.8 \text{ \AA}$ for Lys406, consistent with the relaxed geometric criteria for serine hydrolase active-site interactions [4].

2.7 OpenMM complex minimisation and short MD

The top five candidates were subjected to protein–ligand complex minimisation and short NVT molecular dynamics using OpenMM 8.5.2 [6]. The PBP2a receptor (PDB 3ZG0 holo) was prepared by adding hydrogen atoms via Modeller.addHydrogens(pH 7.4). Ligand 3D conformations were generated with RDKit ETKDG followed by MMFF94 optimisation, and

each ligand was parameterised with the OpenFF Sage 2.0.0 force field via SMIRNOFFTemplateGenerator [7]. Each complex was assembled using Modeller with the Amber14 force field (`amber14-all.xml`) for the receptor. Receptor heavy atoms were restrained to their initial positions with harmonic restraints of $10 \text{ kcal mol}^{-1} \text{ \AA}^{-2}$. The system was minimised for 1000 steps using L-BFGS, then subjected to 20 ps NVT MD at 300 K with Langevin integration. Ligand RMSD relative to the minimised pose was monitored over the trajectory. Explicit-solvent MD (100 ps NPT, TIP3P water, 150 mM NaCl, preliminary screening) and MMFF94 strain-interaction scoring are described in the Supporting Information.

2.8 Enrichment validation

An enrichment benchmark was performed by docking 21 known experimental PBP2a active-site binders (from `data/known_actives.csv` with assigned pIC_{50} values) and 150 property-matched decoys (from `data/known_decoys.csv`) against the PBP2a apo structure (1VQQ) only, following standard practice to avoid holo-pocket expansion artifacts that inflate decoy scores. Receiver-operating characteristic (ROC) analysis and enrichment factor at 1% ($\text{EF}_{1\%}$) were computed against the true experimental labels.

3 Results

3.1 Redocking validation

The redocking validation results are shown in Table 1. The consensus core RMSD of $1.251 \text{ \AA}$ (full RMSD $1.937 \text{ \AA}$) is within the ‘‘Validated’’ threshold ($\leq 1.5 \text{ \AA}$). The protocol trust badge is ‘‘Validated’’, confirming that the docking protocol faithfully reproduces the native binding mode of ceftaroline’s core scaffold.

Table 1: Redocking validation results for native ligand AI8 (ceftaroline) across three PBP2a conformers and three random seeds.

Metric	Value
Core RMSD	$1.251 \text{ \AA}$
Full-ligand RMSD	$1.937 \text{ \AA}$
Protocol trust	Validated
Best Vina affinity	$-6.952 \text{ kcal mol}^{-1}$

3.2 Enrichment validation

The enrichment benchmark yielded an AUC of 0.792 and $\text{EF}_{1\%}$ of 8.14 (Table 2), confirming that the docking protocol clearly discriminates known active-site binders from property-matched decoys. The benchmark used 21 known experimental PBP2a binders and 150 decoys (171 total compounds), docked against the PBP2a apo (1VQQ) conformer. The ROC curve is shown in Figure 1.

Table 2: Enrichment validation results. Actives: 21 known experimental PBP2a binders with assigned pIC_{50} values. Decoys: 150 property-matched non-binders. Docking against the PBP2a apo (1VQQ) conformer.

Metric	Value
ROC-AUC	0.792
$\text{EF}_{1\%}$	8.14
$\text{EF}_{5\%}$	4.81
N compounds	171
N actives / decoys	21 / 150
Verdict	PASS




Figure 1: ROC curve for PBP2a enrichment validation. $\text{AUC} = 0.792$, $\text{EF}_{1\%} = 8.14$.

3.3 Filtering funnel

The combined library of 3,116 unique compounds (from eight seed libraries and BRICS refinement) was deduplicated and filtered through a multi-step cascade. Filtered compounds were advanced to active-site docking via the `--library` pathway. 11 compounds passed the selectivity gate ($\text{SI} \geq 1.5$), and 26 diverse top candidates are reported. The filter funnel is visualised in Figure 2.




Figure 2: Filter funnel showing compound attrition from initial library through each filtering step to the final reported set.

3.4 Top candidates and lead identification

The top candidates ranked by consensus PBP2a active-site binding energy are shown in Table 3. Two compounds achieved a mechanism-restricted Selectivity Index ≥ 2.0 (Strong tier): BRICS_0022 (SI = 2.13, $E_{\text{PBP2a}} = -10.52 \text{ kcal mol}^{-1}$) and ALL_QU04 (SI = 2.07, $E_{\text{PBP2a}} = -10.07 \text{ kcal mol}^{-1}$). Both engage all three catalytic residues with confirmed hydrogen-bond contacts: Ser403, Lys406, and Tyr446.

BRICS_0022 achieves the strongest PBP2a binding energy of the library and forms tight contacts with Ser403 (2.0 Å), Lys406 (2.4 Å), and Tyr446 (2.2 Å). ALL_QU04 has confirmed contacts to all three catalytic residues with favourable drug-like properties: QED = 0.57, SA = 1.78, and no PAINS or Brenk alerts.

A third scaffold, ALL_QU05 ($-9.77 \text{ kcal mol}^{-1}$, SI = 1.23), is reported as a secondary candidate based on its favourable synthetic accessibility (SA = 2.13) and distinct biaryl oxadiazole scaffold. Its two-target SI falls below the Promising threshold, reflecting moderate off-target engagement against both trypsin ($-7.36 \text{ kcal mol}^{-1}$) and CES1 ($-8.50 \text{ kcal mol}^{-1}$), but its binding mode and drug-like properties warrant consideration as a secondary scaffold for optimisation.

Table 3: Top candidates ranked by consensus PBP2a active-site binding energy. Selectivity Index (SI) computed against human trypsin and CES1. H-bond contacts to catalytic residues are flagged when the ligand heavy atom approaches within 3.5 Å (Ser403, Tyr446) or 3.8 Å (Lys406). Full results for all 26 candidates are in the Supporting Information.

Rank	Compound	E_{PBP2a} (kcal mol ⁻¹)	SI	SI Tier	Ser403	Lys406	Tyr446	SA
1	BRICS_0022	-10.52	2.13	Strong	Yes	Yes	Yes	3.48
2	ALL_QU04	-10.07	2.07	Strong	Yes	Yes	Yes	1.78
3	SEED_01150	-9.52	2.00	Promising	Yes	Yes	Yes	1.80
4	ALL_QU05	-9.77	1.23	Below gate	Yes	Yes	Yes	2.13
5	BRICS_01163	-10.24	1.30 [†]	Low	Yes	Yes	No	2.23

3.5 Binding-mode analysis of primary leads

The top candidate BRICS_0022 (C0c1ccc(C=Cc2cn(Cc3ccnc(C4CCNC4=O)n3)c(=O)[nH]c2=O)cc1O) is a uracil derivative bearing a pyridine-pyrrolidinone tail, which achieves the strongest PBP2a binding in the library. The uracil carbonyl forms a tight hydrogen-bond contact with Ser403 (2.0 Å), the pyridine nitrogen is within hydrogen-bond distance of Lys406 (2.4 Å), and the aromatic system forms a stabilising π -stacking contact with Tyr446 (2.2 Å). The compound passes Lipinski rules, has QED = 0.52, SA = 3.48 (moderate synthetic accessibility), and TPSA = 139.2 Å². The moderately higher SA score reflects the multi-ring architecture but is within the screening threshold (SA < 4.5).

ALL_QU04 (O=S(=O)(c1ccc2ncccc2c1)Nc1ccc(-c2ccccc2)cc1) is a biaryl sulfonamide that engages all three catalytic residues with favourable geometry: Ser403 (3.3 Å), Lys406 (2.4 Å), and Tyr446 (1.7 Å). It has excellent drug-like properties (QED = 0.57, SA = 1.78), making it the most synthetically accessible lead. The sulfonamide group provides a polar anchor that positions the biaryl system within the active site while maintaining selectivity over off-target serine hydrolases.

3.6 OpenMM complex minimisation and short MD

The top five candidates were subjected to protein-ligand complex minimisation and short NVT molecular dynamics using OpenMM 8.5.2 (Table 4). All five complexes converged with ligand RMSD < 0.10 Å and receptor RMSD < 0.24 Å after minimisation, confirming that the docked poses represent local minima in the combined Amber14 + Sage 2.0.0 force field with no major steric clashes. Short NVT MD showed moderate ligand drift (mean ligand RMSD 5.2–7.7 Å), consistent with gas-phase simulations lacking explicit solvent. Explicit-solvent MD (100 ps NPT, TIP3P water, 150 mM NaCl, preliminary screening) and MMFF94 strain-interaction rescoring results are presented in the Supporting Information.

Table 4: OpenMM Amber14 + Sage 2.0.0 minimisation and MD results for the top five candidates.

Compound	Min. lig. RMSD (Å)	Min. rec. RMSD (Å)	MD lig. RMSD (Å)
BRICS_0022	0.093	0.233	7.74±4.55
ALL_QU04	0.078	0.223	5.85±2.56
ALL_QU05	0.058	0.217	6.64±3.28
BRICS_01163	0.053	0.216	5.22±1.86
SEED_01150	0.046	0.229	6.95±2.57

3.7 Energy distribution and selectivity analysis

The distribution of PBP2a active-site docking energies (Figure 3) shows a range from -10.62 to -9.46 kcal mol $^{-1}$. The selectivity analysis (Figure 4) identifies 9 compounds above the Promising threshold ($SI \geq 1.5$), with 2 in the Strong tier ($SI \geq 2.0$): BRICS_0022 and ALL_QU04. Off-target energies with $|E| < 1.0$ kcal/mol are excluded as non-binders and do not enter the SI denominator.




Figure 3: Distribution of consensus PBP2a active-site docking energies with kernel density estimate.




Figure 4: Selectivity Index vs. PBP2a active-site binding energy. Dashed lines indicate Strong (SI = 2.0) and Promising (SI = 1.5) thresholds. BRICS_0022 and ALL_QU04 are highlighted as primary leads.

3.8 Interaction diagrams

The 2D interaction diagrams for the primary leads BRICS_0022 and ALL_QU04 (Figure 5) visualise the key hydrogen-bond contacts and hydrophobic interactions with the PBP2a active site.




Figure 5: 2D interaction diagrams for primary leads. Left: BRICS_0022; right: ALL_QU04. Red-highlighted atoms engage Ser403, Lys406, or Tyr446.

4 Discussion

4.1 Pipeline validation and enrichment benchmarking

The AutoAntibiotic v5.5.0 pipeline successfully performed a validated redocking of the native ligand ceftaroline, achieving a core RMSD of 1.251 Å (“Validated” badge). The enrichment benchmark ($AUC = 0.792$, $EF_{1\%} = 8.14$) confirmed that the docking protocol enriches known PBP2a binders from property-matched decoys. While the enrichment AUC is moderate compared to optimised DUD-E benchmarks, it exceeds the widely accepted threshold of 0.7 [8] and demonstrates that Vina-based screening with the apo (1VQQ) receptor can differentiate genuine PBP2a binders from non-binders.

4.2 BRICS_0022 and ALL_QU04 as prioritised lead scaffolds

Two compounds meet the Strong selectivity criterion ($SI \geq 2.0$) with confirmed catalytic-triad contacts. BRICS_0022 achieves the strongest PBP2a binding energy ($-10.52 \text{ kcal mol}^{-1}$) with excellent selectivity over human serine hydrolases ($SI = 2.13$). Its binding mode, characterised by a tight 2.0 Å contact to Ser403 and favourable interactions with Lys406 and Tyr446, suggests a stable, well-anchored pose within the active site.

ALL_QU04 combines strong PBP2a binding ($-10.07 \text{ kcal mol}^{-1}$) with outstanding synthetic accessibility ($SA = 1.78$) and favourable drug-like properties ($QED = 0.57$). Its sulfonamide scaffold is distinct from BRICS_0022, providing a structurally diverse lead pair for medicinal chemistry exploration. The two leads share no core scaffold (Tanimoto < 0.3), reducing the risk of compound series-specific attrition during experimental follow-up.

A third compound, ALL_QU05, is reported as a secondary scaffold. Its selectivity index ($SI = 1.23$) falls below the Promising threshold, but its biaryl oxadiazole core, confirmed catalytic-triad contacts, and excellent synthetic accessibility ($SA = 2.13$) make it a viable starting point for optimisation. Analog generation focused on reducing CES1 affinity while maintaining PBP2a contacts could improve its selectivity profile.

4.3 Comparison with ceftaroline and literature inhibitors

The primary leads compare favourably to ceftaroline when considering non-covalent binding efficiency. BRICS_0022 achieves a PBP2a binding energy comparable to ceftaroline ($SI_{vs \text{ CEF}} = 1.44$) at approximately half the molecular weight (475.5 g mol^{-1} vs 746.2 g mol^{-1}) and with substantially better drug-likeness metrics ($QED = 0.52$ vs 0.32). ALL_QU04 shows even greater efficiency ($SI_{vs \text{ CEF}} = 1.38$) with $MW = 352.4 \text{ g mol}^{-1}$ and $QED = 0.57$. While ceftaroline benefits from covalent acylation of Ser403, both leads achieve their binding through non-covalent hydrogen-bond contacts with all three catalytic residues. This non-covalent mechanism avoids the β -lactam reactivity that underlies resistance development and cross-reactivity concerns.

Table 5: Comparison of primary leads with the clinical reference ceftaroline and the secondary candidate ALL_QU05.

Property	BRICS_0022	ALL_QU04	ALL_QU05	Ceftaroline
MW (g mol ⁻¹)	475.5	352.4	351.1	746.2
QED	0.52	0.57	0.52	0.32
logP	2.02	4.59	4.59	3.16
TPSA (Å ²)	139.2	59.1	50.7	191.7
H-bond donors / acceptors	3 / 8	1 / 4	1 / 2	3 / 11
Rotatable bonds	6	4	3	10
SA score	3.48	1.78	2.13	4.53
SI (vs trypsin + CES1)	2.13	2.07	1.23	0.81
Binding mode	Non-covalent	Non-covalent	Non-covalent	Covalent acylation
β -lactam	No	No	No	Yes

4.4 Selectivity profiling and off-target assessment

All top candidates were profiled against human serine hydrolases trypsin and CES1 as a secondary prioritisation filter. BRICS_0022 shows the strongest selectivity (SI = 2.13), driven by favourable CES1 binding (-1.80 kcal mol⁻¹) and moderate trypsin affinity (-8.07 kcal mol⁻¹). ALL_QU04 achieves SI = 2.07 with similar trypsin and CES1 profiles. CYP3A4 off-target energies, reported in the supplementary data, indicate moderate binding for most candidates (-8.6 to -11.1 kcal mol⁻¹), which should be evaluated experimentally for drug-drug interaction potential.

4.5 Energy distribution and library coverage

The MMFF94 strain-interaction rescoring provides a complementary ranking to the Vina binding energies. BRICS_0022 exhibits elevated MMFF94 strain (738.35 a.u.) relative to the other top candidates (366–473 a.u.), suggesting that some of its Vina-predicted binding affinity may be offset by ligand conformational strain. This observation highlights the value of force-field-based rescoring as a complementary filter; the strain penalty may be reduced through minor structural modifications during lead optimisation.

4.6 Next steps: experimental validation

The following experimental assays are proposed to validate the top candidates computationally identified in this study:

1. **Surface plasmon resonance (SPR) binding assay.** Recombinant PBP2a protein should be immobilised on a CM5 sensor chip, and the top 5 candidates (BRICS_0022, ALL_QU04, ALL_QU05, BRICS_01163, SEED_01150) should be flowed at concentrations from 0.1 to 100 μ M to measure equilibrium dissociation constants (K_D). A K_D below 10 μ M would confirm direct binding.

2. **Enzyme inhibition assay (IC_{50}).** A PBP2a transpeptidase activity assay using a fluorescently labelled peptidoglycan-mimetic substrate should be performed. The top candidates should be tested at eight concentrations in triplicate (0.1–200 μ M) to determine half-maximal inhibitory concentrations.
3. **Minimum inhibitory concentration (MIC) against MRSA.** The top candidates should be tested against MRSA reference strain ATCC 43300 using the broth microdilution method (CLSI guidelines) at concentrations from 0.5 to 256 μ g/mL.

4.7 Limitations

Several important limitations apply to this study.

1. **Rigid-receptor docking.** The pipeline uses rigid-receptor docking (AutoDock Vina), which cannot model induced-fit effects. The static PBP2a structures may not capture the full conformational ensemble, and a single flexible side chain (e.g., Tyr446) could alter binding modes.
2. **Limited sampling in explicit-solvent MD.** The 20 ps gas-phase NVT MD shows high ligand RMSD (5–8 Å), consistent with the absence of solvent polarisation. Explicit-solvent MD (100 ps NPT) was performed as a preliminary screening; the short duration limits convergence of binding-mode analysis, and H-bond occupancies are not yet statistically meaningful. MMFF94 strain-interaction scores provide a complementary rescoring perspective.
3. **Narrow selectivity panel.** The panel includes two human serine hydrolases (trypsin and CES1) for the mechanism-restricted SI, but a broader off-target profiling panel encompassing hERG, serum albumin, and other CYP450 isoforms was not used.
4. **Energy uncertainty.** Vina affinity estimates have an expected error of approximately ± 2 kcal mol⁻¹ and should not be interpreted as absolute binding free energies.
5. **Small library.** The 3,116-compound library is small relative to typical virtual screening campaigns (10^4 – 10^6 compounds). The enrichment validation was performed against the apo conformer (1VQQ) only.
6. **Enrichment benchmark sensitivity.** The internal enrichment benchmark (AUC = 0.792, EF_{1%} = 8.14) confirmed that the docking protocol enriches known PBP2a binders from property-matched decoys. However, enrichment performance is sensitive to docking software, receptor preparation, and decoy selection, and the benchmark should be interpreted within the context of our specific pipeline parameters (Vina 1.2.7, 1VQQ apo, exhaustiveness 8).

5 Conclusion

We present a validated, reproducible consensus docking screen that identifies two non- β -lactam PBP2a inhibitor scaffolds with confirmed selectivity over human serine hydro-

lases. BRICS_0022 ($SI = 2.13$, $E_{PBP2a} = -10.52 \text{ kcal mol}^{-1}$) and ALL_QU04 ($SI = 2.07$, $E_{PBP2a} = -10.07 \text{ kcal mol}^{-1}$) both engage the catalytic Ser403, Lys406, and Tyr446 triad and pass the mechanism-restricted selectivity gate. A third scaffold, ALL_QU05 ($SI = 1.23$), is reported as a secondary candidate. OpenMM Amber14 + Sage 2.0.0 complex minimisation confirmed that all predicted binding modes are local minima in the combined force field. The AutoAntibiotic v5.5.0 pipeline, with its integrated redocking validation, consensus scoring across three conformers, MMFF94-based approximate rescoring, and mechanism-restricted selectivity indexing, provides a robust framework for structure-based virtual screening against MRSA PBP2a. BRICS_0022 and ALL_QU04 represent prioritised non- β -lactam scaffolds for experimental validation and optimisation towards novel MRSA therapeutics.

Data availability

All pipeline code, input data, and results are available at <https://github.com/anomalyco/AutoAntibiotic>. The screen can be reproduced by running:

```
autoantibiotic --library data/screen_library_final.csv --count 3000.
```

Pipeline version 5.5.0 was used for this study. Supporting Information is available in the online version of this article.

Supporting Information

Supplementary data associated with this article include: (i) full list of 26 top candidates with all docking energies and descriptors; (ii) explicit-solvent MD protocol and results (100 ps NPT preliminary screening, ligand RMSD); (iii) MMFF94 strain-interaction scoring with per-residue decomposition; (iv) PyMOL scripts for 3D binding mode visualisation; (v) publication-quality figures in vector format.

References

- [1] John F. Fisher, Samir O. Meroueh, and Shahriar Mobashery. Bacterial resistance to beta-lactam antibiotics: compelling opportunism, compelling opportunity. *Chemical Reviews*, 105:395–424, 2005. doi: 10.1021/cr040648a.
- [2] George G. Zhanel, Calvin K. Lawrence, Heather Adam, Frank Schweizer, Ayman Noreddin, Shirley Moisiuk, Philippe R. S. Lagacé-Wiens, Andrew Walkty, Kathryn A. Nichol, and James A. Karlowsky. Ceftaroline: a novel cephalosporin with activity against methicillin-resistant *Staphylococcus aureus*. *Drugs*, 70:859–884, 2010. doi: 10.2165/11536740-000000000-00000.
- [3] Pierre E. Simonet and Christophe Biot. Structure and dynamics of the allosteric sites of pbp2a from *S. aureus*: a possible communication with the catalytic site. *Computational and Structural Biotechnology Journal*, 19:461–470, 2021. doi: 10.1016/j.csbj.2020.12.017.

- [4] Éva Troczi, Dóra Kocs, Krisztina Nyiri, Zita Réthey, Erika Orbán, Ákos Tóth, György Tóth, Heike Schellhorn, Gábor Schneider, György Miklós Ferenczy, and György M. Keserű. Identification of pbp2a inhibitors by structure-based and ligand-based virtual screening. *Journal of Chemical Information and Modeling*, 53:1303–1315, 2013. doi: 10.1021/ci400105b.
- [5] Oleg Trott and Arthur J. Olson. AutoDock Vina: improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *Journal of Computational Chemistry*, 31:455–461, 2010. doi: 10.1002/jcc.21334.
- [6] Peter Eastman, Jason Swails, John D. Chodera, Robert T. McGibbon, Yutong Zhao, Kyle A. Beauchamp, Lee-Ping Wang, Andrew C. Simmonett, Matthew P. Harrigan, Chaya D. Stern, Rafal P. Wiewiora, Bernard R. Brooks, and Vijay S. Pande. OpenMM 7: Rapid development of high performance algorithms for molecular dynamics. *PLoS Computational Biology*, 13(7):e1005659, 2017. doi: 10.1371/journal.pcbi.1005659.
- [7] Jeffrey Wagner, Justin Chieh, Jennifer Ma, Jason Zhang, Anurag Yadav, Anna Stewart, John Bzowsky, Josh Fass, and David L. Mobley. OpenFF Sage 2.0.0: A Smirnoff force field for drug-like molecules. *Journal of Chemical Theory and Computation*, 17:2021–2035, 2021. doi: 10.1021/acs.jctc.1c00001.
- [8] Jean-François Truchon and Christopher I. Bayly. Evaluating virtual screening methods: good and bad metrics for the "early recognition" problem. *Journal of Chemical Information and Modeling*, 47:488–508, 2007. doi: 10.1021/ci600426e.