

COMPLEXITY OF DEEP COMPUTATIONS VIA TOPOLOGY OF FUNCTION SPACES

EDUARDO DUEÑEZ¹ JOSÉ IOVINO¹ TONATIUH MATOS-WIEDERHOLD²
LUCIANO SALVETTI² FRANKLIN D. TALL²

¹Department of Mathematics, University of Texas at San Antonio

²Department of Mathematics, University of Toronto

ABSTRACT. We use topological methods to study the complexity of deep computations and limit computations. We use topology of function spaces, specifically, the classification of Rosenthal compacta, to identify new complexity classes. We use the language of model theory, specifically, the concept of *independence* from Shelah’s classification theory, to translate between topology and computation. We use the theory of Rosenthal compacta to characterize approximability of deep computations, both deterministically and probabilistically.

INTRODUCTION

In this paper we study asymptotic behavior of computations, e.g., the depth of a neural network tending to infinity, or the time interval between layers of a time-series network tending toward zero. Recently, particular cases of this concept have attracted considerable attention in deep learning research (e.g., Neural Ordinary Differential Equations [CRBD], Physics-Informed Neural Networks [RPK19], and deep equilibrium models [BKK]). The formal framework introduced here provides a unified setting to study these limit phenomena from a foundational viewpoint.

Informed by model theory, to each computation in a given computation model, we associate a continuous real-valued function, called the *type* of the computation, that describes the logical properties of this computation with respect to the rest of the model. This allows us to view computations in any given computational model as elements of a space of real-valued functions, which is called the *space of types* of the model. The idea of embedding models of theories into their type spaces is central in model theory. In the context of this paper, the embedding of computations into spaces of types allows us to utilize the vast theory of topology of function spaces, known as C_p -theory, to obtain results about complexity of topological limits of computations. As we shall indicate next, recent classification results for spaces of functions provide an elegant and powerful machinery to classify computations according to their levels of “tameness” or “wildness”, with the former corresponding roughly to polynomial approximability and the latter to exponential approximability. The viewpoint of spaces of types, which we have borrowed from

Date: January 26, 2026.

2000 Mathematics Subject Classification. 54H30, 68T27, 68T07, 03C98, 03D15, 05D10.

Key words and phrases. Deep computations, deep equilibrium models, deep learning, physics-informed networks, computational complexity, independence property, NIP, infinite Ramsey theory, Baire class 1 functions, Rosenthal compacta, Todorćević trichotomy, Bourgain-Fremlin-Talagrand.

model theory, thus becomes a “Rosetta stone” that allows us to interconnect various classification programs: In topology, the classification of Rosenthal compacta pioneered by Todorčević [Tod99]; in logic, the classification of theories developed by Shelah [She90]; and in statistical learning, the notions of PAC learning and VC dimension pioneered by Vapnik and Chervonenkis [VC74, VC71].

In a previous paper [ADIW24], the authors introduced the concept of limits of computations, which we called *ultracomputations* (given they arise as ultrafilter limits of standard computations) and *deep computations* (following usage in machine learning [BKK]). There is a technical difference between both designations, but in this paper, to simplify the nomenclature, we will ignore the difference and use only the term “deep computation”.

In [ADIW24], a new “tame vs wild” (i.e., polynomially approximable vs non-polynomially approximable) dichotomy for the complexity of deep computations was proved by invoking a classical result of Grothendieck from the 1950s [Gro52]. Under our model-theoretic Rosetta stone, the property of polynomial approximability of computations is identified with continuous extendibility in the sense of topology, and with the notions of *stability* and *type definability* in model theory.

Deep computations arise as topological limits of standard (continuous) computations. Simplest among these is when the limit is sequential. In topology, the *first Baire class*, or *Baire class 1* consists of functions (also called simply “*Baire-1*”) arising as pointwise limits of sequences of continuous functions. Intuitively, the Baire-1 class consists of functions with “controlled” discontinuities, and lies just one level of topological complexity above the Baire class 0 which (by definition) consists of continuous functions.

In this paper, we apply the theory of spaces of Baire class 1 functions to identify and separate new complexity classes of deep computations. We obtain a new “tame vs wild” dichotomy for deep computations by invoking a famous paper by Bourgain, Fremlin and Talagrand from the late 70s [BFT78], and further, a new trichotomy for the class of “tame” deep computations by invoking an equally celebrated result of Todorčević, from the late 90s, for functions of the first Baire class [Tod99].

Todorčević’s trichotomy concerns *Rosenthal compacta*, which are topological spaces that fit our context superbly: they are defined as compact spaces that are homeomorphic to subspaces of the space of Baire class 1 functions on some Polish (separable, completely metrizable) space, under the topology of pointwise convergence. Rosenthal compacta exhibit “topological tameness,” meaning that they behave in relatively controlled ways, and since the late 70’s, they have played a crucial role in understanding the complexity of structures of functional analysis, especially Banach spaces. Todorčević’s trichotomy for Rosenthal compacta has been utilized to settle longstanding problems in topological dynamics and topological entropy [GM22]. It is noteworthy that Todorčević’s proof relies on sophisticated set-theoretic forcing and infinite Ramsey theory. At the time of writing this paper, decades after his original argument, no elementary proof has been found [Tod23, HT19].

Through our Rosetta stone, Rosenthal compacta in topology correspond to the important concept of *Non Independence Property* (known as “NIP”) in model theory, identified by Shelah [She71, She90], and to the concepts *VC-dimension* and *Probably Approximately Correct* learning (known as “PAC learnability”) in statistical learning theory [VC71, VC74, Val84].

Refining Todorćević’s trichotomy, we invoke a more recent heptachotomy for separable Rosenthal compacta obtained by Argyros, Dodos and Kanellopoulos [ADK08]; these authors identify seven fundamental “prototypes” of separable Rosenthal compacta, and show that any non-metrizable separable Rosenthal compactum must contain a “canonical” embedding of one of these prototypes.

We believe that the results presented in this paper show practitioners of computation, or topology, or descriptive set theory, or model theory, how classification invariants used in their field translate into classification invariants of other fields. In the interest of accessibility, we do not assume that the reader has previous familiarity with advanced topology, model theory, or computing. The only technical prerequisites to read this paper are undergraduate-level topology and measure theory. The necessary topological background beyond undergraduate topology is covered in section 1.

In section 1, we present basic topological and combinatorial preliminaries, and in section 2, we introduce the structural/model-theoretic viewpoint (no previous exposure to model theory is needed). Section 3 is devoted to the classification of deep computations.

Throughout the paper, our results pertain to classical models of computation (particularly computations involving real-valued quantities that are known and manipulated to a finite degree of precision). The final section, Section 4, introduces a probabilistic viewpoint, the development of which we intend to pursue in future research, extending the present framework to encompass non-deterministic and quantum computations.

CONTENTS

Introduction	1
1. General topological preliminaries: From continuity to Baire class 1	4
1.1. From Rosenthal’s dichotomy to the Bourgain-Fremlin-Talagrand dichotomy to Shelah’s NIP	6
1.2. NIP as a universal dividing line between polynomial and exponential complexity	8
1.3. Rosenthal compacta	9
1.4. The special case $B_1(X, \mathbb{R}^{\mathcal{P}})$ with \mathcal{P} countable.	10
2. Compositional Computation Structures: A structural approach to arbitrary-precision arithmetic	11
2.1. Compositional Computation Structures	12
2.2. Computability and definability of deep computations and the Extendibility Axiom	13
2.3. Newton’s method as a CCS	14
2.4. Finite precision threshold classifiers as a CCS	17
2.5. Finite precision prefix test	19
3. Classifying deep computations	20
3.1. NIP, Rosenthal compacta, and deep computations	20
3.2. The Todorćević trichotomy, and levels of NIP and PAC learnability	20
3.3. The Argyros-Dodos-Kanellopoulos heptachotomy, and approximability of deep computation by minimal classes	24
4. Randomized versions of NIP and Monte Carlo computability of deep computations	25

4.1. NIP and Monte Carlo computability of deep computations	25
4.2. Talagrand stability and Monte Carlo computability of deep computations	28
References	30

1. GENERAL TOPOLOGICAL PRELIMINARIES: FROM CONTINUITY TO BAIRE CLASS 1

In this section we present some preliminaries from general topology and function space theory. In the interest of completeness, we include some proofs that may be safely skipped by readers familiar with these topics.

Recall that a subset of a topological space is F_σ if it is a countable union of closed sets, and G_δ if it is a countable intersection of closed sets. A space is metrizable if its topology agrees with the topology induced by some metric therein. Two such metrics inducing the same topology may induce quite different properties in the category of metric spaces. For example, the interval $(0, 1)$ with the usual metric (as a subset) of the reals is not complete; however, $(0, 1)$ is homeomorphic to the real line, which is complete with respect to the usual metric thereon. In a metrizable space, every open set is F_σ ; equivalently, every closed set is G_δ .

A *Polish space* is a separable and completely metrizable topological space, i.e., admitting some complete metric inducing its topology. Although other (possibly incomplete) metrics may induce the same topology, being Polish is a purely topological property. One of the most important Polish spaces is the real line \mathbb{R} ; the others include the Cantor space $2^{\mathbb{N}}$ and the Baire space $\mathbb{N}^{\mathbb{N}}$. The class of Polish spaces is closed under countable topological products; in particular, the Cantor space $2^{\mathbb{N}}$ (the set of all infinite binary sequences, endowed with the product topology), the Baire space $\mathbb{N}^{\mathbb{N}}$ (the set of all infinite sequences of naturals, also with the product topology), and the space $\mathbb{R}^{\mathbb{N}}$ of sequences of real numbers are all Polish. Recall that the product topology on these spaces is the topology of pointwise convergence: a sequence converges in the space if and only if it converges at each coordinate index.

Fact 1.1. *A subset of a Polish space is itself Polish in the subspace topology if and only if it is a G_δ set. In particular, closed subsets and open subsets of Polish spaces are also Polish spaces.*

For a proof, see [Eng89, 4.3.24].

Given two topological spaces X and Y we denote by $C_p(X, Y)$ the set of all continuous functions $f : X \rightarrow Y$ endowed with the topology of pointwise convergence. The space $C_p(X, \mathbb{R})$ of continuous real functions on X is denoted simply $C_p(X)$. A natural question is, how do topological properties of X translate into $C_p(X)$ and vice versa? This general question, and the study of these spaces in general, is the concern of C_p -theory, an active field of research in general topology which was pioneered by A. V. Arhangel'skiĭ and his students in the 1970's and 1980's [Ark92]. This field has found many applications in model theory and functional analysis. For a recent survey, see [Tka11].

A *Baire class 1* function between topological spaces is a function that can be expressed as the pointwise limit of a sequence of continuous functions. In symbols, $f : X \rightarrow Y$ is *Baire class 1* if there is a sequence of continuous functions $f_n : X \rightarrow Y$

such that for all $x \in X$, $f(x) = \lim_{n \rightarrow \infty} f_n(x)$. If X and Y are topological spaces, the space of Baire class 1 functions $f : X \rightarrow Y$ endowed with the topology of pointwise convergence is denoted $B_1(X, Y)$ (as above, $B_1(X, \mathbb{R})$ is denoted $B_1(X)$). Clearly, $C_p(X, Y) \subseteq B_1(X, Y) \subseteq Y^X$ and we give these the topology (called the *topology of pointwise convergence*) inherited from the product topology of Y^X . The Baire hierarchy of functions was introduced by René-Louis Baire in his 1899 doctoral thesis, *Sur les fonctions de variables réelles*. His work moved away from the 19th-century preoccupation with “pathological” functions toward a constructive classification based on pointwise limits. An elementary fact about Baire class 1 functions is that they are continuous except on a set of first category (also called a *meager* set, a set of first category is the countable union of sets whose closure has empty interior; intuitively, these sets are “topologically small”). Thus, Baire class 1 functions are continuous on a “topologically large” subset of their domain.

A topological space X is *perfectly normal* if it is normal and every closed subset of X is a G_δ (equivalently, every open subset of X is a F_σ). Every metrizable space (hence, every Polish space) is perfectly normal.

A topological space X is *Baire* if every countable intersection of dense open sets is dense. The Baire Category Theorem states that every compact Hausdorff or completely metrizable space (hence, every Polish space) is Baire.

The following fact was established by Baire in his 1899 thesis. A proof can be found in [Tod97, Section 10].

Fact 1.2 (Baire). *If X is perfectly normal, then the following conditions are equivalent for a function $f : X \rightarrow \mathbb{R}$:*

- (1) *f is a Baire class 1 function, that is, f is a pointwise limit of continuous functions.*
- (2) *$f^{-1}[U]$ is an F_σ subset of X whenever $U \subseteq \mathbb{R}$ is open.*

If, moreover, X is Baire, then (1) and (2) are equivalent to:

- (3) *For every closed $F \subseteq X$, the restriction $f|_F$ has a point of continuity.*

Moreover, if X is Polish and $f \notin B_1(X)$, then there exist countable $D_0, D_1 \subseteq X$ and reals $a < b$ such that

$$D_0 \subseteq f^{-1}(-\infty, a], \quad D_1 \subseteq f^{-1}[b, \infty), \quad \overline{D_0} = \overline{D_1}.$$

A subset L of a topological space X is *relatively compact* in X if the closure of L in X is compact. Relatively compact subsets of $B_1(X)$ (for X Polish) are of interest in analysis and topological dynamics. We begin with the following well-known result. Recall that a set $A \subseteq \mathbb{R}^X$ of real-valued functions is *pointwise bounded* if for every $x \in X$ there is $M_x \geq 0$ (a *pointwise bound at x*) such that $|f(x)| \leq M_x$ for all $f \in A$. We include a proof for the reader’s convenience:

Lemma 1.3. *Let X be a Polish space and $A \subseteq B_1(X)$ be pointwise bounded. The following are equivalent:*

- (i) *A is relatively compact in $B_1(X)$.*
- (ii) *A is relatively countably compact in $B_1(X)$, i.e., every countable subset of A has a limit point in $B_1(X)$.*
- (iii) *$\overline{A} \subseteq B_1(X)$, where \overline{A} denotes the closure in \mathbb{R}^X .*

Proof. (i) \Rightarrow (ii) Relatively compact subsets of any space are countably compact therein.

(ii) \Rightarrow (iii) Consider any $f \in \overline{A}$ and any countable subset $\{x_i\}_{i \in \mathbb{N}} \subseteq X$. We claim that there is a sequence $\{f_n\}_{n \in \mathbb{N}} \subseteq A$ such that $\lim_{n \rightarrow \infty} f_n(x_i) = f(x_i)$ for all $i \in \mathbb{N}$. Since A carries the relative product topology, for each $n \in \mathbb{N}$ there exists $f_n \in A$ such that $|f_n(x_i) - f(x_i)| < \frac{1}{n}$ for all $i \leq n$; the sequence $\{f_n\}$ is as claimed. Seeking a contradiction, assume that A is relatively countably compact in $B_1(X)$, but there exists some $f \in \overline{A} \setminus B_1(X)$. By Fact 1.2, there are countable $D_0, D_1 \subseteq X$ with $\overline{D_0} = \overline{D_1}$, and $a < b$ such that $D_0 \subseteq f^{-1}(-\infty, a]$ and $D_1 \subseteq f^{-1}[b, \infty)$. Per the claim above, let $\{f_n\}_{n \in \mathbb{N}} \subseteq A$ satisfy $\lim_{n \rightarrow \infty} f_n(x) = f(x)$ for all $x \in D_0 \cup D_1$ (the latter being a countable set). By relative countable compactness of A , there is a limit point $g \in B_1(X)$ of $\{f_n\}_{n \in \mathbb{N}}$; clearly, f and g agree on $D_0 \cup D_1$. Thus g takes values $g(x_i) = f(x_i) \leq a$ as well as values $g(x_j) = f(x_j) \geq b > a$ on any open subset of the closed set $\overline{D_0} = \overline{D_1}$, contradicting the implication (1) \Rightarrow (3) in Fact 1.2.

(iii) \Rightarrow (i) For each $x \in X$, let $M_x \geq 0$ be a pointwise bound for A . Since \overline{A} is a closed subset of the compact space $\prod_{x \in X} [-M_x, M_x] \subseteq \mathbb{R}^X$, it follows that \overline{A} is compact. By (iii), it is also the closure of A in $B_1(X)$. Thus, A is relatively compact in $B_1(X)$. \square

1.1. From Rosenthal's dichotomy to the Bourgain-Fremlin-Talagrand dichotomy to Shelah's NIP. In metrizable spaces, points of the closure of some subset can always be approximated by points in the set proper, via a convergent sequence. For more complicated spaces, such as C_p -spaces, this fails in remarkable ways. The n -th coordinate map $p_n : 2^{\mathbb{N}} \rightarrow \{0, 1\}$ on the Cantor space $X = 2^{\mathbb{N}}$ ($= \{0, 1\}^{\mathbb{N}}$) is continuous for each $n \in \mathbb{N}$, and one can show (e.g., [Tod97, Chapter 1.1]) that $\{p_n\}_{n \in \mathbb{N}}$ has *no* convergent subsequences, in \mathbb{R}^X . In a sense, this example exhibits the worst failure of sequential convergence possible. The closure of $\{p_n\}$ in $\{0, 1\}^X$ (or in \mathbb{R}^X for that matter) is homeomorphic to the *Stone-Ćech compactification* of the discrete space of natural numbers, usually denoted $\beta\mathbb{N}$, which is an important object of study in general topology.

The following theorem, proved by Haskell Rosenthal in 1974, is fundamental in functional analysis and captures a sharp division in the behavior of sequences in a Banach space.

Theorem 1.4 (Rosenthal's Dichotomy, [Ros74]). *If X is Polish and $\{f_n\} \subseteq C_p(X)$ is pointwise bounded, then $\{f_n\}_{n \in \mathbb{N}}$ has a convergent subsequence, or a subsequence whose closure in \mathbb{R}^X is homeomorphic to $\beta\mathbb{N}$.*

Rosenthal's Dichotomy states that a pointwise bounded set of continuous functions contains either a convergent subsequence, or a subsequence whose closure is essentially the same as the example mentioned in the previous paragraphs (i.e., "wildest" possible). The genesis of this theorem was Rosenthal's " ℓ_1 -Theorem", which states that a Banach space includes an isomorphic copy of ℓ_1 (the space of absolutely summable sequences), or else every bounded sequence therein is weakly Cauchy. The ℓ_1 -Theorem connects diverse areas: Banach space geometry, Ramsey theory, set theory, and topology of function spaces.

As we move from $C_p(X)$ to the larger space $B_1(X)$, a dichotomy paralleling the ℓ_1 -Theorem holds: Either every point of the closure of a set of functions is a Baire class 1 function, or there is a sequence in the set behaving in the wildest possible way. This result is usually not phrased as a dichotomy, but rather as an equivalence as in Theorem 1.5 below.

First, we introduce some useful notation. For any set $A \subseteq \mathbb{R}^X$ and any real a , define

$$X_{\leq a}^A := \bigcap_{f \in A} f^{-1}(-\infty, a] = \{x \in X : f(x) \leq a \text{ for all } f \in A\},$$

$$X_{\geq a}^A := \bigcap_{f \in A} f^{-1}[a, +\infty) = \{x \in X : f(x) \geq a \text{ for all } f \in A\}.$$

(In case $A = \emptyset$, we define $X_{\geq a}^\emptyset = X = X_{\leq a}^\emptyset$.) For any sequence $\{f_n\} \subseteq \mathbb{R}^X$ and $I \subseteq \mathbb{N}$, define $I^\complement := \mathbb{N} \setminus I$ and $f_I := \{f_i : i \in I\}$.

Theorem 1.5 (“The BFT Dichotomy”. Bourgain-Fremlin-Talagrand [BFT78]). *Let X be a Polish space and $A \subseteq C_p(X)$ be pointwise bounded. The following are equivalent:*

- (i) A is relatively compact in $B_1(X)$.
- (ii) For every $\{f_n\}_{n \in \mathbb{N}} \subseteq A$ and every $a < b$ there is $I \subseteq \mathbb{N}$ such that $X_{\leq a}^{f_I} \cap X_{\geq b}^{f_{I^\complement}} = \emptyset$.

(As stated above, the BFT Dichotomy is a particular case of the equivalence (ii) \Leftrightarrow (v) in [BFT78, Corollary 4G].)

The sets $X_{\leq a}^{f_I}$ and $X_{\geq b}^{f_{I^\complement}}$ appearing in condition Theorem 1.5(ii) are defined, respectively, in terms of $|I|$ -many inequalities of the form $f_i(x) \leq a$, and $|I^\complement|$ -many of the form $f_j(x) \geq b$. Thus, at least one of $X_{\leq a}^{f_I}$ and $X_{\geq b}^{f_{I^\complement}}$ is defined by the satisfaction of infinitely (countably) many inequalities. For our purposes, it is more natural to consider only finitely many inequalities at a time, which motivates the definitions below.

Definition 1.6. We say that a function collection $A \subseteq \mathbb{R}^X$ has the *finitary No-Independence Property (NIP)* if, for all sequences $\{f_n\}_{n \in \mathbb{N}} \subseteq A$ and reals $a < b$, there exist finite disjoint sets $E, F \subseteq \mathbb{N}$ such that $X_{\leq a}^{f_E} \cap X_{\geq b}^{f_F} = \emptyset$. We say that such E, F *witness finitary NIP* for A , $\{f_n\}$ and a, b .

A set $A \subseteq \mathbb{R}^X$ has the *finitary Independence Property (IP)* if it does not have finitary NIP, i.e., if there exists a sequence $\{f_n\}_{n \in \mathbb{N}} \subseteq A$ and reals $a < b$ such that for every pair of finite disjoint sets $E, F \subseteq \mathbb{N}$, we have $X_{\leq a}^{f_E} \cap X_{\geq b}^{f_F} \neq \emptyset$.

If the word “finite” is omitted in the above definitions, we obtain the definitions of *countable NIP* (weaker than finitary NIP) and *countable IP* (stronger than finitary IP), respectively.

If we insist on witnesses $E, F \subseteq \mathbb{N}$ such that $F = E^\complement$, we call the respective properties “BFT-NIP” (even weaker than countable NIP) and “BFT-IP” (even stronger than countable IP). Thus, Theorem 1.5 becomes that statement, for pointwise bounded function collections $A \subseteq C_p(X)$, that A is relatively compact in $B_1(X)$ if and only if A has BFT-NIP.

Unless otherwise unspecified, IP/NIP shall mean *finitary* IP/NIP henceforth.

Proposition 1.7. *If X is compact and $A \subseteq C_p(X)$, then A has BFT-NIP if and only if it has finitary NIP.*

(No pointwise boundedness is assumed of A .)

Proof. Trivially (as per the preceding discussion), finitary NIP implies BFT-NIP. Reciprocally, assume that X is compact and has finitary IP. Fix $A \subseteq C_p(X)$,

a sequence $\{f_n\} \subseteq A$ and reals $r < s$. For any $I, J \subseteq \mathbb{N}$ (almost disjoint in applications), write $X_{I,J}$ for $X_{\leq r}^{f_I} \cap X_{\geq s}^{f_J}$. For $I \subseteq I' \subseteq \mathbb{N}$ and $J \subseteq J' \subseteq \mathbb{N}$, we have $X_{I,J} \supseteq X_{I',J'}$; moreover, $X_{I,J} = \bigcap_{E \subseteq I, F \subseteq J} X_{E,F}$, where the index variables $E \subseteq I, F \subseteq J$ range over *finite* subsets of I, J , respectively. Clearly, $E, F \subseteq \mathbb{N}$ witness finitary NIP for $\{f_n\}$ if and only if $X^{E,F} = \emptyset$.

Fix $I \subseteq \mathbb{N}$. Since $\{f_n\} \subseteq A \subseteq C_p(X)$ is a sequence of continuous functions, and X is compact, the nested family $\{X_{E,F} : E \subseteq I, F \subseteq I^c\}$ consists of closed, thus compact, sets. Since A has finitary IP by hypothesis, the nested family consists of nonempty sets, hence its intersection $X_{I,I^c} \neq \emptyset$ by compactness. This holds for arbitrary $\{f_n\} \subseteq A$ and $r < s$, so A has BFT-IP. \square

Theorem 1.8. *Let X be a compact metrizable (hence Polish) space. For every pointwise bounded $A \subseteq C_p(X)$, the following properties are all equivalent:*

- (i) *A is relatively compact in $B_1(X)$;*
- (ii) *A has BFT-NIP;*
- (iii) *A has countable NIP;*
- (iv) *A has finitary NIP.*

(The equivalences (ii) \Leftrightarrow (iii) \Leftrightarrow (iv) hold for arbitrary compact X .)

Proof. Corollary of Theorem 1.5 and Proposition 1.7. \square

Theorem 1.8 may be stated as the following dichotomy (under the assumptions): either A is relatively compact in $B_1(X)$, or A has IP (in either sense).

The Independence Property was first isolated by Saharon Shelah in model theory as a dividing line between theories whose models are “tame” (corresponding to NIP) and theories whose models are “wild” (corresponding to IP). See [She71, Definition 4.1], [She90]. We will discuss this dividing line in more detail in the next section.

1.2. NIP as a universal dividing line between polynomial and exponential complexity. The particular case of the BFT dichotomy (Theorem 1.5) when A consists of $\{0,1\}$ -valued (i.e., $\{\text{Yes}, \text{No}\}$ -valued) strings was discovered independently, around 1971-1972 in many foundational contexts related to polynomial (“tame”) vs exponential (“wild”) complexity: In model theory, by Saharon Shelah [She71], [She90], in combinatorics, by Norbert Sauer [Sau72], and Shelah [She72], [She90], and in statistical learning, by Vladimir Vapnik and Alexey Chervonenkis [VC71], [VC74].

In model theory: Shelah’s classification theory is a foundational program in mathematical logic devised to categorize first-order theories based on the complexity and structure of their models. A theory T is considered classifiable in Shelah’s sense if the number of non-isomorphic models of T of a given cardinality can be described by a bounded number of numerical invariants. In contrast, a theory T is unclassifiable if the number of models of T of a given cardinality is the maximum possible number. A key fact is that the number of models of T is directly impacted by the number of *types* over sets of parameters in models of T ; a controlled number of types is a characteristic of a classifiable theory.

In Shelah's classification program [She90], theories without the independence property (called NIP theories, or dependent theories) have a well-behaved, "tame" structure; the number of types over a set of parameters of size κ of such a theory is of polynomially or similar "slow" growth on κ .

In contrast, theories with the Independence Property (called IP theories) are considered "intractable" or "wild". A theory with the Independence Property produces the maximum possible number of types over a set of parameters; for a set of parameters of cardinality κ , the theory will have 2^{2^κ} -many distinct types.

In combinatorics: Sauer [Sau72] and Shelah [She72] proved the following independently: Let \mathcal{F} be a family of subsets of some set S . Either: for every $n \in \mathbb{N}$ there is a set $A \subseteq S$ with $|A| = n$ such that $|\{S_i \cap A : i \in \mathbb{N}\}| = 2^n$ (\mathcal{F} has "exponential complexity"); or: there exists $N \in \mathbb{N}$ such that for every $A \subseteq S$ with $|A| \geq N$, one has

$$|\{S_i \cap A : i \in \mathbb{N}\}| \leq \sum_{i=0}^{N-1} \binom{|A|}{i}.$$

(\mathcal{F} has "polynomial complexity"). Clearly, any family \mathcal{F} of subsets of a *finite* set S has polynomial complexity. The "polynomial" name is justified: indeed, for fixed $N > 0$, as a function of the size $|A| = m > 0$, we have

$$\sum_{i=0}^{N-1} \binom{m}{i} \leq \sum_{i=0}^{N-1} \frac{m^i}{i!} \leq \left(\sum_{i=0}^{N-1} \frac{1}{i!} \right) \cdot m^{N-1} < e \cdot m^{N-1} = O(m^N).$$

(More precisely, the order of magnitude is $O(m^{N-1})$: polynomial in m for N fixed.)

In machine learning: Readers familiar with statistical learning may recognize the Sauer-Shelah lemma as the dichotomy discovered and proved slightly earlier (1971) by Vapnik and Chervonenkis [VC71, VC74] to address uniform convergence in statistics. The least integer N given by the preceding paragraph, when it exists, is called the *VC-dimension* of \mathcal{F} ; it is a core concept in machine learning. If such an integer N does not exist, we say that the VC-dimension of \mathcal{F} is infinite. The lemma provides upper bounds on the number of data points (sample size) needed to learn a concept class of known VC dimension d up to a given admissible error in the statistical sense. The Fundamental Theorem of Statistical Learning states that a hypothesis class is PAC-learnable (PAC stands for "Probably Approximately Correct") if and only if its VC dimension is finite.

1.3. Rosenthal compacta. The universal classification implied by Theorem 1.5, as attested by the examples outlined in the preceding section, led to the following definition (by Gilles Godefroy [God80]):

Definition 1.9. A Rosenthal compactum is any topological space realized as a compact subset of the space $B_1(X) = B_1(X, \mathbb{R})$ (equipped with the topology of pointwise convergence) of all real functions of the first Baire class on some Polish space X .

A Rosenthal compactum K is necessarily Hausdorff since it is a topological subspace of the Hausdorff product space \mathbb{R}^X .

Rosenthal compacta possess significant topological and dynamical tameness properties, and play an important role in functional analysis, measure theory, dynamical systems, descriptive set theory, and model theory. In this paper, we use them to study deep computations. For this, we shall first focus on countable languages, which is the theme of the next subsection.

1.4. The special case $B_1(X, \mathbb{R}^{\mathcal{P}})$ with \mathcal{P} countable. Fix an arbitrary (at most) countable set \mathcal{P} whose elements $P \in \mathcal{P}$ will be called *predicate symbols* or *formal predicates*. Our present goal is to characterize relatively compact subsets of $B_1(X, \mathbb{R}^{\mathcal{P}})$, where X is always assumed to be a perfectly normal space (often a Polish space).

The set \mathcal{P} shall be considered discrete whenever regarded as a topological space. Since $C_p(X, \mathbb{R}^{\mathcal{P}}) \subseteq B_1(X, \mathbb{R}^{\mathcal{P}}) \subseteq (\mathbb{R}^{\mathcal{P}})^X$, the “ambient” space $(\mathbb{R}^{\mathcal{P}})^X$ is quite relevant. The product $X \times \mathcal{P}$ will be regarded as either a pointset, or as a topological product depending on context. We have natural homeomorphic identifications

$$(\mathbb{R}^{\mathcal{P}})^X \cong \mathbb{R}^{X \times \mathcal{P}} \cong (\mathbb{R}^X)^{\mathcal{P}},$$

given by

$$\begin{aligned} \mathbb{R}^{X \times \mathcal{P}} &\rightarrow (\mathbb{R}^{\mathcal{P}})^X : \varphi \mapsto \hat{\varphi} \\ \mathbb{R}^{X \times \mathcal{P}} &\rightarrow (\mathbb{R}^X)^{\mathcal{P}} : \varphi \mapsto \varphi^\wedge, \end{aligned}$$

where

$$\hat{\varphi}(x) := \varphi(x, \cdot) \in \mathbb{R}^{\mathcal{P}}, \quad \varphi^\wedge(P) := \varphi(\cdot, P) \in \mathbb{R}^X.$$

Such identifications view X and \mathcal{P} as mere pointsets (the topology of X in particular plays no role).

For $x \in X$, define the “left projection” map

$$\lambda_x : \mathbb{R}^{X \times \mathcal{P}} \rightarrow \mathbb{R}^{\mathcal{P}} : \varphi \mapsto \lambda_x(\varphi) := \varphi(x, \cdot);$$

for $P \in \mathcal{P}$, the “right projection” map

$$\rho_P : \mathbb{R}^{X \times \mathcal{P}} \rightarrow \mathbb{R}^X : \varphi \mapsto \varphi(\cdot, P).$$

For fixed $x \in X$ and $P \in \mathcal{P}$, we also have canonical projection maps

$$\pi_x : \mathbb{R}^X \rightarrow \mathbb{R} : f \mapsto f(x), \quad \pi_P : \mathbb{R}^{\mathcal{P}} \rightarrow \mathbb{R} : f \mapsto f(P).$$

When clear from context, rather than using the specific symbols (“ λ ” for left, “ ρ ” for right) to denote projections, we may use the generic symbol “ π ”; thus, π_x may mean λ_x , and π_P may mean ρ_P .

The Proposition below reduces the study of $\mathbb{R}^{\mathcal{P}}$ -valued continuous or Baire-1 functions on X to the special case of real-valued ones.

Proposition 1.10. *The identification $(\mathbb{R}^{\mathcal{P}})^X \cong \mathbb{R}^{X \times \mathcal{P}} \cong (\mathbb{R}^X)^{\mathcal{P}}$ induces identifications*

$$C_p(X, \mathbb{R}^{\mathcal{P}}) \cong C_p(X \times \mathcal{P}) \cong C_p(X)^{\mathcal{P}}, \quad B_1(X, \mathbb{R}^{\mathcal{P}}) \cong B_1(X \times \mathcal{P}) \cong B_1(X)^{\mathcal{P}}.$$

(The cardinality of \mathcal{P} plays no role.)

Proof. The identification of C_p -spaces follows trivially from the definition of topological product and the fact that \mathcal{P} is discrete: a continuous map $X \rightarrow \mathbb{R}^{\mathcal{P}}$ is precisely a \mathcal{P} -indexed family of continuous functions $X \rightarrow \mathbb{R}$, and these correspond to continuous functions $X \times \mathcal{P} \rightarrow \mathbb{R}$. The identification of Baire-1 spaces follows

immediately, since it is defined in terms of the purely topological notion of limit (in the ambient space) of sequences of continuous functions. \square

Given $A \subseteq Y^X$ and $K \subseteq X$ we write $A|_K := \{f|_K : f \in A\}$, i.e., the set of all restrictions of functions in A to K . The following Theorem is a slightly more general version of Theorem 1.5.

Theorem 1.11. *Assume that \mathcal{P} is countable, X is a Polish space, and $A \subseteq C_p(X, \mathbb{R}^{\mathcal{P}})$ is pointwise bounded in the sense that $\pi_P \circ A$ ($\subseteq C_p(X)$) is pointwise bounded for every $P \in \mathcal{P}$. The following are equivalent for every compact $K \subseteq X$:*

- (i) $A|_K$ is relatively compact in $B_1(K, \mathbb{R}^{\mathcal{P}})$;
- (ii) $\pi_P \circ A|_K$ has NIP for every $P \in \mathcal{P}$.

Proof. Compact subsets $K \subseteq X$ are closed, hence also Polish. Therefore, the asserted equivalence follows from Theorems 1.5 and 1.7. \square

Lastly, a simple but useful lemma that helps understand when we restrict a set of functions to a specific subspace of the domain space, we may always assume that the subspace is closed, as replacing the subspace by its closure has no effect on NIP.

Lemma 1.12. *Assume that X is Hausdorff and that $A \subseteq C_p(X)$. The following are equivalent for every $L \subseteq X$:*

- (i) A_L satisfies the NIP;
- (ii) $A|_{\overline{L}}$ satisfies the NIP.

Proof. It suffices to show that (i) \Rightarrow (ii). Suppose that (ii) does not hold, i.e., that there are $\{f_n\}_{n \in \mathbb{N}} \subseteq A$ and $a < b$ such that for all finite disjoint $E, F \subseteq \mathbb{N}$:

$$\overline{L} \cap \bigcap_{n \in E} f_n^{-1}(-\infty, a] \cap \bigcap_{n \in F} f_n^{-1}[b, \infty) \neq \emptyset.$$

Pick $a' < b'$ such that $a < a' < b' < b$. Then, for any finite disjoint $E, F \subseteq \mathbb{N}$ we can choose

$$x \in \overline{L} \cap \bigcap_{n \in E} f_n^{-1}(-\infty, a') \cap \bigcap_{n \in F} f_n^{-1}(b', \infty)$$

By definition of closure:

$$L \cap \bigcap_{n \in E} f_n^{-1}(-\infty, a'] \cap \bigcap_{n \in F} f_n^{-1}[b', \infty) \neq \emptyset.$$

This contradicts (i). \square

2. COMPOSITIONAL COMPUTATION STRUCTURES: A STRUCTURAL APPROACH TO ARBITRARY-PRECISION ARITHMETIC

In this section, we connect function spaces with arbitrary-precision arithmetic computations. We start by summarizing some basic concepts from [ADIW24].

A *computation states structure* is a pair (L, \mathcal{P}) , where L is a set whose elements we call *states* and \mathcal{P} is a collection of real-valued functions on L that we call *predicates*. For a state $v \in L$, the *type* of a state v is the indexed family

$$\text{tp}(v) = (P(v))_{P \in \mathcal{P}} \in \mathbb{R}^{\mathcal{P}}.$$

For each $P \in \mathcal{P}$, we call the value $P(v)$ the P -th *feature* of v . A *transition* of a computation states structure (L, \mathcal{P}) is a map $f : L \rightarrow L$.

Intuitively, L is the set of states of a computation, and the predicates $P \in \mathcal{P}$ are primitives that are given and accepted as computable. Each state $v \in L$ is uniquely characterized by its type $\text{tp}(v)$, so we may identify L with a subset of $\mathbb{R}^{\mathcal{P}}$. Important state spaces are $L = \mathbb{R}^{\mathbb{N}}$ and $L = \mathbb{R}^n$ for some positive integer n , endowed with predicate $P_i(v) = v_i$, one each for the i -th coordinate of v . We regard the space of types as a topological space, endowed with the topology of pointwise convergence induced by the product topology of $\mathbb{R}^{\mathcal{P}}$. Via the identification $v \mapsto \text{tp}(v)$, the states space L is correspondingly topologized; in particular, for each $P \in \mathcal{P}$, the projection map $v \mapsto P(v)$ is continuous.

Definition 2.1. Given a computation states structure (L, \mathcal{P}) , any element of $\mathbb{R}^{\mathcal{P}}$ in the image of L under the map $v \mapsto \text{tp}(v)$ will be called a *realized (state) type*. The topological closure of the set of realized types in $\mathbb{R}^{\mathcal{P}}$ (endowed with the pointwise convergence topology) will be called the *space of types* of (L, \mathcal{P}) , denoted \mathcal{L} ; elements $\xi \in \mathcal{L}$ are called *state types*. Elements of $\mathcal{L} \setminus L$ will be called *unrealized types*.

Intuitively, state types capture a notion of “limit state”.

As we combine ideas of model theory [Kei03] and topology [BFT78], we are interested in families of real-valued functions that are pointwise bounded. This leads us to the concepts of *sizer* and *shard* introduced first in [ADIW24]:

Definition 2.2. A *sizer* is a family $r_{\bullet} = (r_P)_{P \in \mathcal{P}}$ of positive real numbers, indexed by \mathcal{P} . Given a sizer r_{\bullet} , let $\mathbb{R}^{[r_{\bullet}]} = \prod_{P \in \mathcal{P}} [-r_P, r_P]$ (a compact space), and let the r_{\bullet} -*shard* of a states space L be

$$L[r_{\bullet}] = L \cap \mathbb{R}^{[r_{\bullet}]}.$$

For a sizer r_{\bullet} , the r_{\bullet} -*type shard* is defined as $\mathcal{L}[r_{\bullet}] = \overline{L[r_{\bullet}]}$ (a closed, hence compact subset of $\mathbb{R}^{[r_{\bullet}]}$).

Let also \mathcal{L}_{sh} be the union of all type-shards as the sizer r_{\bullet} varies.

In general, $\mathcal{L}_{\text{sh}} \subseteq \mathcal{L}$, and the inclusion may be proper. However, equality holds in the important special case when \mathcal{P} is countable (see [ADIW24]).

2.1. Compositional Computation Structures.

Definition 2.3. A *Compositional Computation Structure* (CCS) is a triple (L, \mathcal{P}, Γ) , where

- (L, \mathcal{P}) is a computation states structure, and
- $\Gamma \subseteq L^L$ is a semigroup under composition.

Elements of the semigroup Γ are called the *computations* of the structure (L, \mathcal{P}, Γ) . We assume that the identity map id on L is an element of Γ (which is thus not merely a semigroup but a monoid of transformations of L).

We topologize Γ as a subset of the topological product L^L , where the “exponent” L serves merely as an index set, but the “base” L is topologized by type; consequently, one may identify Γ with a subset of the topological product $(\mathbb{R}^{\mathcal{P}})^L$. More specifically, Γ is identified with a subset of \mathcal{L}^L , which is a closed subspace of $(\mathbb{R}^{\mathcal{P}})^L$. Therefore, we have an inclusion $\bar{\Gamma} \subseteq \mathcal{L}^L$. Elements $\xi \in \bar{\Gamma}$ are called (real-valued) *deep computations* or *ultracomputations*.

A collection R of sizers is *exhaustive* if $L = \bigcup_{r_{\bullet} \in R} L[r_{\bullet}]$ (shards $L[r_{\bullet}]$ exhaust L). A transformation $\gamma \in \Gamma$ is *R-confined* if γ restricts to a map $\gamma|_{L[r_{\bullet}]} : L[r_{\bullet}] \rightarrow L[r_{\bullet}]$ (into $L[r_{\bullet}]$ itself) for every $r_{\bullet} \in R$. A subset $\Delta \subseteq \Gamma$ is *R-confined* if each $\gamma \in \Delta$ is.

Proposition 2.4. *If $\Delta \subseteq \Gamma$ is confined by an exhaustive sizer collection, then $\overline{\Delta}$ is a compact subset of $\mathcal{L}_{\text{sh}}^L$.*

Proof. Assume that R confines Δ . For each $v \in L$, let $r_{\bullet}^{(v)} \in R$ be a sizer such that $v \in L[r_{\bullet}^{(v)}]$. An arbitrary $\gamma \in \Delta$ restricts to a map $\gamma \upharpoonright L[r_{\bullet}^{(v)}] : L[r_{\bullet}^{(v)}] \rightarrow L[r_{\bullet}^{(v)}]$, so $\Gamma \subseteq K := \prod_{v \in L} \mathcal{L}[r_{\bullet}^{(v)}]$. The space K is a product of compact spaces, hence compact, so $\overline{\Gamma}$ is a closed, hence compact subset thereof, and a subset of $\mathcal{L}_{\text{sh}}^L \supseteq K$ *a fortiori*. \square

For a CCS (L, \mathcal{P}, Γ) , we regard the elements of Γ as “standard” finitary computations, and the elements of $\overline{\Gamma}$, i.e., deep computations, as possibly infinitary limits of standard computations. The main goal of this paper is to study the computability, definability and computational complexity of deep computations. Since deep computations are defined through a combination of topological concepts (namely, topological closure) and structural and model-theoretic concepts (namely, models and types), we will import technology from both topology and model theory.

2.2. Computability and definability of deep computations and the Extendibility Axiom. Let $f : L \rightarrow \mathcal{L}$ be a function that maps each input state type $(P(v))_{P \in \mathcal{P}} \in \mathbb{R}^{\mathcal{P}}$ to an output state type $(P \circ f(v))_{P \in \mathcal{P}} \in \mathbb{R}^{\mathcal{P}}$.

- (1) We will say that f is *definable* if for each $Q \in \mathcal{P}$, the output feature $Q \circ f : L \rightarrow \mathbb{R}$ is a definable predicate in the following sense: There is an *approximating function* $\varphi_{Q, K, \epsilon} : L \rightarrow \mathbb{R}$ that can be built recursively out of a finite number of the (primitively computable) predicates in \mathcal{P} and by a finite number of iterations of the finitary lattice operations \wedge ($=\min$) and \vee ($=\max$), the operations of $\mathbb{R}^{\mathbb{R}}$ as a vector algebra (that is, vector addition and multiplication and scalar multiplication) and the operators \sup and \inf applied on individual variables from L , and such that

$$|\varphi_{Q, K, \epsilon}(v) - Q \circ f(v)| \leq \epsilon, \quad \text{for all } v \in K.$$

Remark: What we have defined above is a model-theoretic concept; it is a special case of the concept of *first-order definability* for real-valued predicates in the model theory of real-valued structures first introduced in [Iov94] for model theory of functional analysis and now standard in model theory (see [Kei03]). The \wedge ($=\min$) and \vee ($=\max$) operations correspond to the positive Boolean logical connectives “and” and “or”, and the \sup and \inf operators correspond to the first-order quantifiers, \forall and \exists .

- (2) We will say that f is *computable* if it is definable in the sense defined above under (1), but without the use of the \sup/\inf operators; in other words, if for every choice of Q, K, ϵ , the approximation function $\varphi_{Q, K, \epsilon} : L \rightarrow \mathbb{R}$ can be constructed without any use of \sup or \inf operators. This is quantifier-free definability (i.e., definability as given by the preceding paragraph, but without use of quantifiers), which, from a logic viewpoint, corresponds to computability (the presence of the quantifiers \exists and \forall are the reason behind the undecidability of first-order logic).

It is shown in [ADIW24] that:

- (1) For a definable $f : L \rightarrow \mathcal{L}$, the approximating functions $\varphi_{Q, K, \epsilon}$ may be taken to be *polynomials* of the input features, and

- (2) Definable transforms $f : L \rightarrow \mathcal{L}$ are precisely those that extend to continuous $\tilde{f} : \mathcal{L} \rightarrow \mathcal{L}$.

To summarize, a function $f : L \rightarrow \mathcal{L}$ is computable if and only if it is definable if and only if it is polynomially approximable if and only if it can be extended to a continuous $\tilde{f} : \mathcal{L} \rightarrow \mathcal{L}$. This is the reason for the following definition.

Definition 2.5. We say that a CCS (L, \mathcal{P}, Γ) satisfies the *Extendibility Axiom* if for all $\gamma \in \Gamma$, there is $\tilde{\gamma} : \mathcal{L}_{\text{sh}} \rightarrow \mathcal{L}_{\text{sh}}$ such that for every sizer r_\bullet there is a sizer s_\bullet such that $\tilde{\gamma}|_{\mathcal{L}[r_\bullet]} : \mathcal{L}[r_\bullet] \rightarrow \mathcal{L}[s_\bullet]$ is continuous. We refer to $\tilde{\gamma}$ as a *free extension* of γ .

For the rest of the paper, fix for each $\gamma \in \Gamma$ a free extension $\tilde{\gamma}$ of γ . For any $\Delta \subseteq \Gamma$, let $\tilde{\Delta}$ denote $\{\tilde{\gamma} : \gamma \in \Delta\}$.

For a more detailed discussion of the Extendibility Axiom, we refer the reader to [ADIW24].

2.3. Newton's method as a CCS. Let $p(z)$ be a non-constant polynomial with complex coefficients. We say that $(L_p, \mathcal{P}, \Gamma_p)$ is *Newton's method for $p(z)$* if:

- L_p is the set of all $z \in \mathbb{C}$ such that there exists an open neighborhood U of z such that every sequence in $\{N_p^n : n \in \mathbb{N}\}$ has a subsequence that converges uniformly on compact subsets of U , where

$$N_p(z) = z - \frac{p(z)}{p'(z)}$$

and $N_p^n := N_p \circ N_p \circ \dots \circ N_p$ is the n th-iteration of N_p .

- $\mathcal{P} := \{P_1, P_2, P_3\}$ where

$$P_1(z) = \frac{2\text{Re}(z)}{|z|^2 + 1},$$

$$P_2(z) = \frac{2\text{Im}(z)}{|z|^2 + 1},$$

$$P_3(z) = \frac{|z|^2 - 1}{|z|^2 + 1}.$$

- $\Gamma_p := \{N_p^n : n \in \mathbb{N}\}$.

Remarks 2.6.

- (1) The set L_p is known as the *Fatou set* of N_p (see section 2 in [Bla84]). Its complement $\mathbb{C} \setminus L_p$, is known as the *Julia set* of N_p .
- (2) The map $z \mapsto (P_1(z), P_2(z), P_3(z))$ is the stereographic projection into the Riemann Sphere S^2 .
- (3) The set L_p is open and dense in \mathbb{C} ([Bla84], Corollary 4.6). Hence, its closure \mathcal{L} in $\mathbb{R}^{\mathcal{P}}$ is the Riemann sphere S^2 (i.e., the extended complex plane).
- (4) The set L_p is completely invariant under iterations of N_p , i.e., $N_p(L_p) = L_p$ and $N_p^{-1}(L_p) = L_p$. This implies that all iterations $N_p^n : L_p \rightarrow L_p$ are transition maps.
- (5) Γ_p is the semigroup generated by $\{N_p\}$. Thus, $(L_p, \mathcal{P}, \Gamma_p)$ is a CCS.

Newton's method is an iterative method that is used to approximate a root of $p(z)$. The map $N_p(z)$ defined above is known as *Newton's map*. The method

consists of taking an initial guess $z_0 \in \mathbb{C}$ and iterating the rational map N_p to obtain a sequence given by

$$z_{n+1} = N_p(z_n) = z_n - \frac{p(z_n)}{p'(z_n)}$$

For each root $r \in \mathbb{C}$ of $p(z)$, there exists an $\varepsilon > 0$ such that for any initial guess z_0 in the ε -ball centered at r , Newton's iteration converges to r (provided $p'(r) \neq 0$) and the convergence is quadratic in that case, meaning the error at each step is roughly squared, causing the number of correct digits to double, leading to fast convergence.

Given a root r of $p(z)$, the set $B_r = \{z \in \mathbb{C} : \lim_{n \rightarrow \infty} N_p^n(z) = r\}$ is an open set called the *basin* of r . However, Newton's method can fail to converge to any root for some choices of z_0 . For example, consider the polynomial $p(z) = z^3 - 2z + 2$. The Newton map is given by

$$N_p(z) = z - \frac{z^3 - 2z + 2}{3z^2 - 2} = \frac{2z^3 - 2}{3z^2 - 2}$$

Notice that taking $z_0 = 0$ as an initial guess will yield the sequence $0, 1, 0, 1, 0, 1, \dots$ that oscillates between 0 and 1 but none of them are roots of $p(z)$. Another more chaotic way Newton's method can fail to converge is when the sequence of iterations has no convergent subsequence. The set of such points, i.e. the *Julia set* associated to N_p , is typically a fractal. This can be visualized by adding a dash of color: let us give each complex number z_0 a color (R, G, B) where $R, G, B \in [0, 1]$ (so that $(1, 0, 0)$ is red, $(0, 1, 0)$ is green, $(0, 0, 1)$ is blue and $(0.5, 0, 0.5)$ is a light purple, for example). The values of R , G and B are determined by looking at the image of said number at each stage of the iteration, $N_p^n(z_0)$, and computing the current distance to each of the roots of $p(z)$; so $R = 1/d_r$ where d_r is the positive distance to the root which is colored red, and so on. In this way, the roots themselves are colored red, green, and blue, and every other point gets a mix of the three colors. As the number of iterations increases, each point gets a sharper color, as the sequence of images $\{N_p^n(z_0)\}_{n=1}^{\infty}$ converges to one of the three roots. At each stage, the complex plane looks as if out of focus because the coloring function is continuous. As the reader can see in Figure 1, the points at the boundary of each color class form the famous Newton's fractal (of which, interestingly, Newton was unaware).




FIGURE 1. Newton's method approximating $p(z) = z^3 - 2z + 2$. Notice the regions of divergence.

Another example of a Newton's fractal is for $p(z) = z^3 - 1$. The roots of $p(z)$ are the 3rd roots of unity and the Newton map is given by:

$$N_p(z) = z - \frac{z^3 - 1}{3z^2} = \frac{2z^3 + 1}{3z^2}$$

In this case, there are three basins of attraction (one for each root) and the complement of their union is the Julia set, i.e., the common boundary.



FIGURE 2. Newton's method approximating $p(z) = z^3 - 1$.

Proposition 2.7. *Let $p(z)$ be a non-constant polynomial. $(L_p, \mathcal{P}, \Gamma_p)$ satisfies the Extensibility Axiom.*

Proof. $N_p : L_p \rightarrow L_p$ is a rational map. Rational maps can be continuously extended to the extended complex plane, i.e., to \mathcal{L} . Composition of rational maps is a rational map, so by the same reasoning, computations $N_p^n : L_p \rightarrow L_p$ can be continuously extended to \mathcal{L} . \square

The set of deep computations $\bar{\Gamma}$ might behave different for various polynomials. Let us look at various examples:

Example 2.8. Computation of square roots. Let a be a positive real number and $p(x) = x^2 - a$. Let $L = \mathbb{R} \setminus \{0\}$. Let $\mathcal{P} = (P_1, P_2)$ where $x \mapsto (P_1(x), P_2(x))$ is the stereographic projection into $S^1 \subseteq \mathbb{R}^2$, i.e.,

$$P_1(x) = \frac{2x}{x^2 + 1},$$

$$P_2(x) = \frac{x^2 - 1}{x^2 + 1}.$$

Let $\Gamma = \{N_p^n : n \in \mathbb{N}\}$ where

$$N_p(x) = \frac{x^2 + a}{2x}.$$

As before, (L, \mathcal{P}, Γ) is a CCS. Note that $\mathcal{L} = S^1$ and that each iterate N_p^n can be continuously extended to the extended real line $\mathbb{R} \cup \{\infty\}$, i.e., \mathcal{L} . For example,

$$\tilde{N}_p(x) = \begin{cases} \frac{x^2 + a}{2x}, & \text{if } x \in L; \\ \infty, & \text{if } x = 0, \infty. \end{cases}$$

For every initial guess $x \in L$, the limit $f(x) = \lim_{n \rightarrow \infty} N_p^n(x)$ converges pointwise to one of the roots. Moreover,

$$f(x) = \begin{cases} \sqrt{a}, & \text{if } x > 0; \\ -\sqrt{a}, & \text{if } x < 0. \end{cases}$$

Notice that f can be extended to \mathcal{L} by

$$\tilde{f}(x) = \begin{cases} \sqrt{a}, & \text{if } x > 0; \\ -\sqrt{a}, & \text{if } x < 0; \\ \infty, & \text{if } x = 0, \infty. \end{cases}$$

However, $\tilde{f} : \mathcal{L} \rightarrow \mathcal{L}$ is not continuous. The set $\bar{\Gamma}$ of deep computations is $\bar{\Gamma} \cup \{\tilde{f}\} \subseteq B_1(\mathcal{L}, \mathcal{L})$.

Example 2.9. Newton's method for $p(z) = z^3 - 2z + 2$. Let r_1, r_2 and r_3 be the three roots of $p(z)$. Let B_1, B_2 and B_3 be their respective basins. Let B be the basin of the attractive cycle $0, 1, 0, 1, \dots$. Then, $L_p = B \cup \bigcup_{i=1}^3 B_i$. Notice that N_p^n does not converge pointwise. However, the subsequences N_p^{2n} and N_p^{2n+1} are pointwise convergent to functions f_1 and f_2 respectively. f_1 and f_2 are two distinct deep computations. Note that for $z \notin L_p$, no subsequence of $\tilde{N}_p^n(z)$ converges to a complex number. However, since $\mathcal{L} = S^2$ is compact there is a subsequence of $\tilde{N}_p^n(z)$ that converges to ∞ . We can extend $f_i : L_p \rightarrow \mathcal{L}$ to $\tilde{f}_i : \mathcal{L} \rightarrow \mathcal{L}$ by:

$$\tilde{f}_i(z) = \begin{cases} f_i(z), & \text{if } z \in L_p; \\ \infty, & \text{if } z \notin L_p. \end{cases}$$

Again, note that \tilde{f}_i for $i = 1, 2$ are not continuous and that $\tilde{f}_i \in \bar{\Gamma}$.

2.4. Finite precision threshold classifiers as a CCS. Let $L = 2^{\mathbb{N}}$, i.e., the set consisting of all infinite binary sequences with the topology of pointwise convergence. Let $\mathcal{P} = \{P_n : n \in \mathbb{N}\}$ equal the collection of projections, i.e., $P_n(x) = x(n)$ for each $x \in L$ and $n \in \mathbb{N}$. Notice that $L \subseteq \mathbb{R}^{\mathcal{P}}$ is closed. Therefore, $\mathcal{L} = L$. We denote by 0^∞ the infinite binary sequence consisting of 0s, and by 1^∞ the infinite binary sequence consisting of 1s. The set of finite binary strings is denoted by $2^{<\mathbb{N}}$. This set is naturally ordered by the lexicographic order \leq_{lex} . Given a finite binary string w , we consider the transition $\phi_w : L \rightarrow L$ given by the rule

$$\phi_w(x) = \begin{cases} 1^\infty, & \text{if } x|_{|w|} \leq_{\text{lex}} w; \\ 0^\infty, & \text{otherwise,} \end{cases}$$

where $|w|$ is the length of the string w and $x|_{|w|}$ is the prefix of x of length $|w|$. That is, $\phi_w(x)$ is equal to the constant sequence of ones if $x|_{|w|}$ comes before or is equal to w in the lexicographic order of strings, and it is equal to the constant sequence of zeros otherwise. In words, ϕ_w checks if a number is less than or equal to the scalar value of threshold w (the string w is finite, hence the classifier has *finite precision*). Note that $P_n \circ \phi_w(x) = 1$ if and only if $x|_{|w|}$ comes before w .

Proposition 2.10. $\phi_w : 2^{\mathbb{N}} \rightarrow 2^{\mathbb{N}}$ is continuous for all $w \in 2^{<\mathbb{N}}$.

Proof. It suffices to prove that $P_n \circ \phi_w : 2^{\mathbb{N}} \rightarrow \{0, 1\}$ is continuous for all $n \in \mathbb{N}$. For simplicity, let us call $f := P_n \circ \phi_w$, i.e.,

$$f(x) = \begin{cases} 1, & \text{if } x|_{|w|} \leq_{\text{lex}} w; \\ 0, & \text{otherwise.} \end{cases}$$

We first observe that $f^{-1}(1) = \{x \in 2^{\mathbb{N}} : x|_{|w|} \leq_{\text{lex}} w\}$ is an open set. Fix $x_0 \in f^{-1}(1)$. Let $t := x_0|_{|w|}$ and consider the basic open set $[t] = \{x \in 2^{\mathbb{N}} : x|_{|t|} = t\}$. Then it is not difficult to check that $x_0 \in [t] \subseteq f^{-1}(1)$. The same reasoning shows that $f^{-1}(0)$ is open. \square

Let $\Delta = \{\phi_w : w \in 2^{<\mathbb{N}}\} \cup \{\mathbf{0}^\infty, \mathbf{1}^\infty\}$, where $\mathbf{0}^\infty, \mathbf{1}^\infty : L \rightarrow L$ are the constant maps identical to 0^∞ and 1^∞ , respectively. Let Γ be the semigroup generated by Δ . The preceding proposition shows that Δ (and hence Γ) consists of continuous functions. In particular, the CCS (L, \mathcal{P}, Γ) satisfies the Extendibility Axiom. In contrast with Newton's method, the algebraic structure of Δ is quite simple: composing two classifiers results in something similar to a Boolean logic gate. The topological structure is far more interesting. Intuitively, the crucial difference between Newton's method and threshold classifiers is that the complexity of the former comes from *depth*: the semigroup is generated by a single map but its iterates are highly complex. The complexity of threshold classification comes from *width*: the semigroup has infinitely many generators, but their compositions are simple.

Intuitively, the closure of Δ consists of the set of all possible threshold classifiers on the real line, but there are two sorts: the ones that classify strict inequalities and those that classify \leq . The members of Δ are finite-precision approximations of classifiers that check all bits of information. But here it gets interesting: what is the difference, in terms of arbitrary-precision arithmetic, between $x < 0.5$ and $x \leq 0.5$?

Suppose that f_a^+ represents the \leq classifier for a target $a \in L$. We identify the scalar truth values with constant sequences, formally $f_a^+ : L \rightarrow \{0^\infty, 1^\infty\}$ is defined by $f_a^+(x) = 1^\infty$ if $x \leq_{\text{lex}} a$ and $f_a^+(x) = 0^\infty$ otherwise. Note that if a is the constant 1^∞ , then $f_a^+ = \mathbf{1}^\infty$. Similarly, let f_a^- be the strict inequality $<$ classifier, i.e., $f_a^-(x) = 1^\infty$ if $x <_{\text{lex}} a$ and $f_a^-(x) = 0^\infty$ otherwise. Note that if a is the constant zero, then $f_a^- = \mathbf{0}^\infty$.

Proposition 2.11. $f_a^+, f_a^- \in \overline{\Delta}$ for all $a \in 2^{\mathbb{N}}$.

Proof. First, we show that $f_a^+ \in \overline{\Delta}$. If $a = 1^\infty$, then $f_a^+ = \mathbf{1}^\infty \in \Delta$. If a is not identically 1, we argue that the pointwise limit of the threshold classifiers on $w_n := a|_n \frown 1$ (that is, the sequence obtained from appending a 1 to the first n bits of a) is precisely f_a^+ . Specifically, for every $x \in L$, we intend to prove that $\lim_{n \rightarrow \infty} \phi_{w_n}(x) = f_a^+(x)$. Assume that $x >_{\text{lex}} a$. Let m be the least index at which the two sequences differ. Then $a(m) = 0 < 1 = x(m)$, and for all $n \geq m$, w_n agrees with a up to m . Crucially, $w_n(m) = 0 < 1 = x(m)$, which implies that $w_n <_{\text{lex}} x|_{n+1}$, and hence $\phi_{w_n}(x) = 0^\infty = f_a^+(x)$ for large enough n . If $x \leq_{\text{lex}} a$, then $x|_{n+1} \leq_{\text{lex}} w_n$ for all $n \in \mathbb{N}$. Hence, $\phi_{w_n}(x) = 1^\infty = f_a^+(x)$ for all $n \in \mathbb{N}$.

Now, we prove that $f_a^- \in \overline{\Delta}$. If a is the constant zero, then $f_a^- = \mathbf{0}^\infty \in \Delta$. Suppose that a is not constantly zero; then we have two cases.

- (1) If a is eventually zero (a is often called a *dyadic rational*), that is $a = u \frown 1 \frown 0^\infty$ (here \frown denotes concatenation) for some finite u . Let $w_n :=$

- $u \smallfrown 0 \smallfrown 1^n <_{\text{lex}} a$. We claim that $\lim_{n \rightarrow \infty} \phi_{w_n}(x) = f_a^-(x)$. Assume that $x <_{\text{lex}} a$. Then, $x|_{|w_n|} \leq_{\text{lex}} w_n$ for large enough n . Hence, $\phi_{w_n}(x) = 1^\infty = f_a^-(x)$ for large enough n . Now assume that $x \geq_{\text{lex}} a$. Then, $w_n <_{\text{lex}} a|_{|w_n|} \leq_{\text{lex}} x|_{|w_n|}$ so $\phi_{w_n}(x) = 0^\infty = f_a^-(x)$ for all $n \in \mathbb{N}$.
- (2) If a is not eventually zero, enumerate the indices of all positive bits in a , $\{n \in \mathbb{N} : a(n) = 1\}$, strictly increasingly as $\{n_k : k \in \mathbb{N}\}$ (this is possible as the former set is infinite by assumption). Let $w_k := (a|_{n_k-1}) \smallfrown 0$; that is, w_k is the result of flipping the k -th positive bit in a . Once again, observe that $w_k <_{\text{lex}} a$ for all k . The crucial step follows from the fact that for any $x <_{\text{lex}} a$, there is a large enough K such that $x <_{\text{lex}} w_k$ for all $k \geq K$.

□

The preceding proposition shows that the topological structure of deep computations can be complicated. Indeed, $\overline{P_n \circ \Delta}$ contains the *Split Cantor* space for all $n \in \mathbb{N}$. (see Examples 3.3(3)).

2.5. Finite precision prefix test. In this subsection we present another example of a CCS with a more complicated set of deep computations. Let $L = 2^\mathbb{N}$ and $\mathcal{P} = \{P_n : n \in \mathbb{N}\}$ where $P_n(x) = x(n)$ are the projection maps so clearly $L \subseteq \mathbb{R}^\mathcal{P}$ and $\mathcal{L} = L$ (same computation states structure as subsection 2.4). For each $w \in 2^{<\mathbb{N}}$, let $\psi_w : L \rightarrow L$ be the transition given by:

$$\psi_w(x) = \begin{cases} 1^\infty, & \text{if } x|_{|w|} = w; \\ 0^\infty, & \text{otherwise.} \end{cases}$$

In other words, ψ_w determines whether the first $|w|$ bits of a binary sequence is exactly w . Let $\Delta = \{\psi_w : w \in 2^{<\mathbb{N}}\}$ and Γ be the semigroup generated by Δ . Since the sets $\{x \in 2^\mathbb{N} : x|_{|w|} = w\}$ are open and closed in $2^\mathbb{N}$, then the transitions ψ_w are all continuous. In particular, (L, \mathcal{P}, Γ) satisfies the Extendibility Axiom.

Let us analyze the set of deep computations of Δ . The idea of these finite precision prefix tests ψ_w is that they are approximating the equality relation on infinite binary sequences. For a given $a \in 2^\mathbb{N}$, let $\delta_a : L \rightarrow \{0^\infty, 1^\infty\}$ be the indicator function at a , i.e., $\delta_a(x) = 1^\infty$ if $x = a$ and $\delta_a(x) = 0^\infty$ otherwise.

Proposition 2.12. $\delta_a \in \overline{\Delta}$ for all $a \in 2^\mathbb{N}$.

Proof. Fix $a \in 2^\mathbb{N}$, and let $w_n := a|_n$ for each $n \in \mathbb{N}$. We claim that $\lim_{n \rightarrow \infty} \psi_{w_n}(x) = \delta_a(x)$ for all $x \in L$. If $x = a$, then $x|_{|w_n|} = w_n$ for all n and so $\psi_{w_n}(x) = 1^\infty = \delta_a(x)$ for all n . If $x \neq a$, then $x|_{|w_n|} \neq w_n$ for large enough n . Hence, $\psi_{w_n}(x) = 0^\infty = \delta_a(x)$ for large enough n . □

These equality tests δ_a are not all the deep computations. The other deep computation we are missing is the constant map 0^∞ .

Proposition 2.13. $0^\infty \in \overline{\Delta}$.

Proof. To show that $0^\infty \in \overline{\Delta}$, for each $n \in \mathbb{N}$, consider, $w_n = 1^n \smallfrown 0$, i.e., the string consisting of n consecutive 1s followed by a 0. If $x = 1^\infty$, then $x|_{|w_n|} \neq w_n$ for all $n \in \mathbb{N}$. Hence, $\psi_{w_n}(x) = 0^\infty$ for all $n \in \mathbb{N}$. If $x \neq 1^\infty$, let N be the smallest such that $x(N) = 0$. Then, $x|_{|w_n|} \neq w_n$ for all $n > N$. Hence, $\psi_{w_n}(x) = 0^\infty$ for large enough n . □

In fact, $\overline{\Delta} = \Delta \cup \{\delta_a : a \in 2^{\mathbb{N}}\} \cup \{\mathbf{0}^\infty\}$ and this space is known as the *Extended Alexandroff compactification of $2^{\mathbb{N}}$* (see Example 3.3(2)). One key topological property about this space is that $\mathbf{0}^\infty$ is not a G_δ point, i.e., $\{\mathbf{0}^\infty\}$ is not a countable intersection of open sets. Moreover, $\mathbf{0}^\infty$ is the only non- G_δ point. It is well-known that in a Hausdorff, first countable space every point is G_δ . This shows that our space of deep computations is not first countable. This space also contains a discrete subspace of size continuum, namely $\{\delta_a : a \in 2^{\mathbb{N}}\}$.

3. CLASSIFYING DEEP COMPUTATIONS

3.1. NIP, Rosenthal compacta, and deep computations. Under what conditions are deep computations Baire class 1, and thus well-behaved according to our framework, on type-shards? The following theorem says that, under the assumption that \mathcal{P} is countable, the space of deep computations is a Rosenthal compactum (when restricted to shards) if and only if the set of computations satisfies the NIP feature by feature. Hence, we can import the theory of Rosenthal compacta into this framework of deep computations.

Theorem 3.1. *Let (L, \mathcal{P}, Γ) be a compositional computational structure (Definition 2.3) satisfying the Extendibility Axiom (Definition 2.5) with \mathcal{P} countable. Let R be an exhaustive collection of sizers. Let $\Delta \subseteq \Gamma$ be R -confined. The following are equivalent.*

- (i) $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}} \subseteq B_1(\mathcal{L}[r_\bullet], \mathcal{L}[r_\bullet])$ for all $r_\bullet \in R$;
- (ii) $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$ satisfies the NIP for all $P \in \mathcal{P}$ and $r_\bullet \in R$; that is, for all $P \in \mathcal{P}$, $r_\bullet \in R$, $a < b$, $\{\gamma_n\}_{n \in \mathbb{N}} \subseteq \Delta$ there are finite disjoint $E, F \subseteq \mathbb{N}$ such that

$$L[r_\bullet] \cap \bigcap_{n \in E} (\pi_P \circ \gamma_n)^{-1}(-\infty, a] \cap \bigcap_{n \in F} (\pi_P \circ \gamma_n)^{-1}[b, \infty) = \emptyset.$$

Moreover, if any (hence all) of the preceding conditions hold, then every deep computation $f \in \overline{\Delta}$ can be extended to a Baire-1 function on shards, i.e., there is $\tilde{f} : \mathcal{L}_{sh} \rightarrow \mathcal{L}_{sh}$ such that $\tilde{f}|_{\mathcal{L}[r_\bullet]} \in B_1(\mathcal{L}[r_\bullet], \mathcal{L}[r_\bullet])$ for all $r_\bullet \in R$. In particular, on each shard every deep computation is the pointwise limit of a countable sequence of computations.

Proof. Since \mathcal{P} is countable, $\mathcal{L}[r_\bullet] \subseteq \mathbb{R}^{\mathcal{P}}$ is Polish. Also, the Extendibility Axiom implies that $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$ is a pointwise bounded set of continuous functions for all $P \in \mathcal{P}$. Hence, Theorem 1.11 and Lemma 1.12 prove the equivalence of (i) and (ii). If (i) holds and $f \in \overline{\Delta}$, then write $f = \mathcal{U}\lim_i \gamma_i$ as an ultralimit. Define $\tilde{f} := \mathcal{U}\lim_i \tilde{\gamma}_i$. Hence, for all $r_\bullet \in R$ we have $\tilde{f}|_{\mathcal{L}[r_\bullet]} \in \overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}} \subseteq B_1(\mathcal{L}[r_\bullet], \mathcal{L}[r_\bullet])$. That every deep computation is a pointwise limit of a countable sequence of computations follows from the fact that for a Polish space X every compact subset of $B_1(X)$ is Fréchet-Urysohn (that is, a space where topological closures coincide with sequential closures, see Theorem 3F in [BFT78] or Theorem 4.1 in [Deb13]). \square

3.2. The Todorćević trichotomy, and levels of NIP and PAC learnability.

In this subsection we study the case when the set of deep computations is a separable Rosenthal compactum. We are interested in the separable case for two reasons:

- (1) In practice, the set Δ of computations is countable. This implies that the set $\overline{\Delta}$ of deep computations is separable.

- (2) The non-separable case lacks some tools and nice examples, which makes their study more complicated. In the separable case we have two important results, which are introduced in this subsection (Todorčević's Trichotomy) and the next subsection (Argyros-Dodos-Kanellopoulos heptachotomy). By introducing Todorčević's Trichotomy into this framework, we obtain a classification of the complexity of deep computations.

Given a countable set Δ of computations satisfying the NIP on features and shards (condition (ii) of Theorem 3.1), the set $\tilde{\Delta}_{\mathcal{L}[r_\bullet]}$ (for a fixed sizer r_\bullet) is a separable *Rosenthal compactum* (see Definition 1.9). Todorčević proved a remarkable trichotomy for Rosenthal compacta [Tod99] that was later refined through an heptachotomy proved by Argyros, Dodos, Kanellopoulos [ADK08]. In this section, inspired by the work of Glasner and Megrelishvili [GM22], we study ways in which this classification allows us to obtain different levels of PAC-learnability and NIP.

Recall that a topological space X is *hereditarily separable* if every subspace is separable, and that X is *first countable* if every point in X has a countable local basis. Every separable metrizable space is hereditarily separable, and R. Pol proved that every hereditarily separable Rosenthal compactum is first countable (see section 10 of [Deb13]). This suggests the following definition:

Definition 3.2. Let (L, \mathcal{P}, Γ) be a CCS satisfying the Extendibility Axiom and R be an exhaustive collection of sizers. Let $\Delta \subseteq \Gamma$ be an R -confined countable set of computations satisfying the NIP on shards and features (condition (ii) in Theorem 3.1). We say that Δ is:

- (i) NIP₁ if $\tilde{\Delta}_{\mathcal{L}[r_\bullet]}$ is first countable for every $r_\bullet \in R$.
- (ii) NIP₂ if $\tilde{\Delta}_{\mathcal{L}[r_\bullet]}$ is hereditarily separable for every $r_\bullet \in R$.
- (iii) NIP₃ if $\tilde{\Delta}_{\mathcal{L}[r_\bullet]}$ is metrizable for every $r_\bullet \in R$.

Observe that $\text{NIP}_3 \Rightarrow \text{NIP}_2 \Rightarrow \text{NIP}_1 \Rightarrow \text{NIP}$. Todorčević, [Tod99], isolated three canonical examples of Rosenthal compacta that witness the failure of the converse implications above.

We now present some separable and non-separable examples of Rosenthal compacta. These show that the previously discussed classes NIP_{*i*} are not equal.

Examples 3.3.

- (1) *Alexandroff compactification of a discrete space of size continuum.* For each $a \in 2^{\mathbb{N}}$ consider the map $\delta_a : 2^{\mathbb{N}} \rightarrow \mathbb{R}$ given by $\delta_a(x) = 1$ if $x = a$ and $\delta_a(x) = 0$ otherwise. Let $A(2^{\mathbb{N}}) = \{\delta_a : a \in 2^{\mathbb{N}}\} \cup \{\mathbf{0}\}$, where $\mathbf{0}$ is the zero map. Notice that $A(2^{\mathbb{N}})$ is a compact subset of $B_1(2^{\mathbb{N}})$. In fact, $\{\delta_a : a \in 2^{\mathbb{N}}\}$ is an uncountable discrete subspace of $B_1(2^{\mathbb{N}})$, and its pointwise closure is precisely $A(2^{\mathbb{N}})$. Hence, this is a Rosenthal compactum which is not hereditarily separable (and therefore not first countable). In particular, this space does not satisfy separability, but it can be made separable by adding a countable set as the next example shows.
- (2) *Extended Alexandroff compactification.* For each finite binary sequence $s \in 2^{<\mathbb{N}}$, let $v_s : 2^{\mathbb{N}} \rightarrow \mathbb{R}$ be given by $v_s(x) = 1$ if x extends s and $v_s(x) = 0$ otherwise. Let $\hat{A}(2^{\mathbb{N}})$ be the pointwise closure of $\{v_s : s \in 2^{<\mathbb{N}}\}$, i.e., $\hat{A}(2^{\mathbb{N}}) = A(2^{\mathbb{N}}) \cup \{v_s : s \in 2^{<\mathbb{N}}\}$. Note that this space is a separable Rosenthal compactum which is not first countable. This is the example

discussed in Section 2.5. It is an example of a CCS that is NIP but not NIP_1 .

- (3) *Split Cantor*. Let $<$ be the lexicographic order in the space of infinite binary sequences, i.e., $2^{\mathbb{N}}$. For each $a \in 2^{\mathbb{N}}$ let $f_a^- : 2^{\mathbb{N}} \rightarrow \mathbb{R}$ be given by $f_a^-(x) = 1$ if $x < a$ and $f_a^-(x) = 0$ otherwise. Let $f_a^+ : 2^{\mathbb{N}} \rightarrow \mathbb{R}$ be given by $f_a^+(x) = 1$ if $x \leq a$ and $f_a^+(x) = 0$ otherwise. The Split Cantor is the space $S(2^{\mathbb{N}}) = \{f_a^- : a \in 2^{\mathbb{N}}\} \cup \{f_a^+ : a \in 2^{\mathbb{N}}\}$, which was obtained as the closure of the space discussed in Section 2.4, giving an example separating NIP_2 from NIP_3 . This is a well known separable Rosenthal compactum. One example of a countable dense subset is the set of all f_a^+ and f_a^- where a is an infinite binary sequence that is eventually constant. Moreover, it is hereditarily separable, but it is not metrizable. It is homeomorphic to the space $2^{\mathbb{N}} \times \{0, 1\}$ with the lexicographic order topology via the identification $(a, 1) \mapsto f_a^+$ and $(a, 0) \mapsto f_a^-$.
- (4) *Alexandroff Duplicate*. Let K be any compact metric space and consider the Polish space $X = K \sqcup C(K)$, i.e., the disjoint union of $C(K)$ (with its supremum norm topology) and K . For each $a \in K$ define $g_a^0, g_a^1 : X \rightarrow \mathbb{R}$ as follows:

$$g_a^0(x) = \begin{cases} 0, & \text{if } x \in K \\ x(a), & \text{if } x \in C(K); \end{cases}$$

$$g_a^1(x) = \begin{cases} \delta_a(x), & \text{if } x \in K; \\ x(a), & \text{if } x \in C(K). \end{cases}$$

Let $D(K) = \{g_a^0 : a \in K\} \cup \{g_a^1 : a \in K\}$. Observe that all points g_a^1 are isolated and that open neighborhoods of $g_{a_0}^0$ are of the form $\{g_a^i : a \in U, i \in \{0, 1\}\} \setminus \{g_a^1 : a \in F\}$ where $U \subseteq K$ is an open neighborhood of a_0 and $F \subseteq K$ is a finite set. Another abstract way in which this space is presented is as the space $K \times \{0, 1\}$ whose basic open neighborhoods are given as before, identifying $(a, 0) \mapsto g_a^0$ and $(a, 1) \mapsto g_a^1$. We can also embed $D(K)$ into the product $A(K) \times K$ by identifying $(a, 0) \mapsto (\mathbf{0}, a)$ and $(a, 1) \mapsto (\delta_a, a)$. Notice that $D(K)$ is a first countable Rosenthal compactum. It is not separable if K is uncountable, thus we typically study the interesting case when $K = 2^{\mathbb{N}}$. As with the Alexandroff compactification $A(2^{\mathbb{N}})$, we can make the space $D(2^{\mathbb{N}})$ separable by adding a countable set. For example, the closure of the set $\{(v_s, s \smallfrown 0^\infty) : s \in 2^{<\mathbb{N}}\} \subseteq \hat{A}(2^{\mathbb{N}}) \times 2^{\mathbb{N}}$ is $\{(\mathbf{0}, a) : a \in 2^{\mathbb{N}}\} \cup \{(\delta_a, a) : a \in 2^{\mathbb{N}}\} \cup \{(v_s, s \smallfrown 0^\infty) : s \in 2^{<\mathbb{N}}\}$, where $\{(\mathbf{0}, a) : a \in 2^{\mathbb{N}}\} \cup \{(\delta_a, a) : a \in 2^{\mathbb{N}}\}$ is homeomorphic to $D(2^{\mathbb{N}})$.

- (5) *Extended Alexandroff Duplicate of the Split Cantor*. For each finite binary sequence $t \in 2^{<\mathbb{N}}$ let $a_t \in 2^{\mathbb{N}}$ be the sequence starting with t and ending with 0's and let $b_t \in 2^{\mathbb{N}}$ be the sequence starting with t and ending with 1's. Define $h_t : 2^{\mathbb{N}} \rightarrow \mathbb{R}$ by

$$h_t(x) = \begin{cases} 0, & \text{if } x < a_t; \\ 1/2, & \text{if } a_t \leq x \leq b_t; \\ 1, & \text{if } b_t < x. \end{cases}$$


Let $\hat{D}(S(2^{\mathbb{N}}))$ be the pointwise closure of the set $\{h_t : t \in 2^{<\mathbb{N}}\}$. The identification $h_t \mapsto (v_t, f_{t \smallfrown 0^\infty}^+)$ lifts to a homeomorphism between $\hat{D}(S(2^{\mathbb{N}}))$

and the subspace of $\hat{A}(2^{\mathbb{N}}) \times S(2^{\mathbb{N}})$ consisting of $(\mathbf{0}, f_a^+)$, $(\mathbf{0}, f_a^-)$, (δ_a, f_a^+) and $(v_t, f_{t \smallfrown 0^\infty}^+)$ for $a \in 2^{\mathbb{N}}$ and $t \in 2^{<\mathbb{N}}$ (see 4.3.7 in [ADK08]). Hence, $\hat{D}(S(2^{\mathbb{N}}))$ is a separable first countable Rosenthal compactum which is not hereditarily separable. In fact, it contains an uncountable discrete subspace.

Theorem 3.4 (Todorćević's Trichotomy, [Tod99], Theorem 3 in [ADK08]). *Let K be a separable Rosenthal Compactum.*

- (i) *If K is hereditarily separable but non-metrizable, then $S(2^{\mathbb{N}})$ embeds into K .*
- (ii) *If K is first countable but not hereditarily separable, then either $D(2^{\mathbb{N}})$ or $\hat{D}(S(2^{\mathbb{N}}))$ embeds into K .*
- (iii) *If K is not first countable, then $A(2^{\mathbb{N}})$ embeds into K .*

We thus have the following classification:



Todorćević's Trichotomy suggests that in order to distinguish the classes NIP_i , the examples in 3.3 are essential. The following examples show that the levels NIP_i ($i = 1, 2, 3$) may be distinguished by the topological complexity of deep computations.

Examples 3.5.

- (1) Let (L, \mathcal{P}, Γ) be the computation of square root (example 2.8 with $\Delta = \Gamma$). We saw that $\bar{\Delta} = \tilde{\Delta} \cup \{\tilde{f}\} \subseteq B_1(\mathcal{L}, \mathcal{L})$. This corresponds to the Alexandroff compactification of a countable discrete set, which is metrizable. Hence, Δ is NIP_3 but it is not stable, in the sense that $\bar{\Delta} \not\subseteq C(\mathcal{L}, \mathcal{L})$.
- (2) Let (L, \mathcal{P}, Γ) be the finite precision threshold classifiers (Section 2.4) with $\Delta = \{\phi_w : w \in 2^{<\mathbb{N}}\} \cup \{\mathbf{0}^\infty, \mathbf{1}^\infty\}$. We saw that $\bar{\Delta}$ is homeomorphic to the Split Cantor space $S(2^{\mathbb{N}})$ (Example 3.3(3)), which is hereditarily separable but not metrizable. Hence, Δ is NIP_2 but not NIP_3 .
- (3) Let (L, \mathcal{P}, Γ) be the CCS given by $L = 2^{\mathbb{N}}$, $\mathcal{P} = \{P_n : n \in \mathbb{N}\}$ and Γ is the semigroup generated by $\Delta = \{\gamma_t : t \in 2^{<\mathbb{N}}\}$, where $P_n : 2^{\mathbb{N}} \rightarrow \{0, 1\}$ is the projection map $P_n(x) = x(n)$ and $\gamma_t : L \rightarrow L$ is given by

$$\gamma_t(x) = \begin{cases} 0^\infty, & \text{if } x <_{\text{lex}} t \smallfrown 0^\infty; \\ (01)^\infty, & \text{if } t \smallfrown 0^\infty \leq_{\text{lex}} x \leq_{\text{lex}} t \smallfrown 1^\infty; \\ 1^\infty, & \text{if } t \smallfrown 1^\infty <_{\text{lex}} x. \end{cases}$$

where $(01)^\infty$ denotes the sequence of alternating bits: $010101\dots$. As in the other examples, it is not difficult to see that (L, \mathcal{P}, Γ) satisfies the Extendibility Axiom. For example, the condition $t \smallfrown 0^\infty \leq_{\text{lex}} x \leq_{\text{lex}} t \smallfrown 1^\infty$ is

equivalent to x extending t . Observe that the set of deep computations is homeomorphic to $\hat{D}(S(2^{\mathbb{N}}))$ (see Example 3.3(5)). This is an example of Δ which is NIP_1 but not NIP_2 .

- (4) Let (L, \mathcal{P}, Γ) be the finite precision prefix test (Section 2.5) with $\Delta = \{\psi_w : w \in 2^{<\mathbb{N}}\}$. We saw that $\bar{\Delta}$ is homeomorphic to the Extended Alexandroff compactification $\hat{A}(2^{\mathbb{N}})$ (Example 3.3-(3)), which is separable but not first countable. Hence, Δ is NIP but not NIP_1 .

The definitions provided here for NIP_i ($i = 1, 2, 3$) are topological. This raises the following question:

Question 3.6. Is there a non-topological characterization for NIP_i , $i = 1, 2, 3$?

3.3. The Argyros-Dodos-Kanellopoulos heptachotomy, and approximability of deep computation by minimal classes. In the three separable cases given in 3.3, namely, $\hat{A}(2^{\mathbb{N}})$, $S(2^{\mathbb{N}})$ and $\hat{D}(S(2^{\mathbb{N}}))$, the countable dense subsets are indexed by the binary tree $2^{<\mathbb{N}}$. This choice of index is useful for two reasons:

- (1) Our emphasis is computational. Real numbers can be represented as infinite binary sequences, i.e., infinite branches of the binary tree $2^{<\mathbb{N}}$. We approximate real numbers or binary sequences with elements in $2^{<\mathbb{N}}$, i.e., finite bitstrings. Indexing standard computations with finite bitstrings allow us to better understand how deep computations arise and how they get approximated. Computationally, we are interested in the manner (and the efficiency) in which the approximations can occur.
- (2) Infinite branches of the binary tree $2^{<\mathbb{N}}$ correspond to the Cantor space $2^{\mathbb{N}}$, the canonical perfect set (in the sense that any Polish space with no isolated points contains a copy of $2^{\mathbb{N}}$). The use of infinite dimensional Ramsey theory for trees (pioneered by the work of James D. Halpern, Hans Läuchli in [HL66] and also Keith Milliken in [Mil81], and Alain Louveau, Saharon Shelah, Boban Velickovic in [LSV93]) and perfect sets (Fred Galvin and Andreas Blass in [Bla81], Arnold W. Miller in [Mil89], and Stevo Todorćević in [Tod99]) allowed S.A. Argyros, P. Dodos and V. Kanellopoulos in [ADK08] to obtain an improved version of Theorem 3.4. It is no surprise that Ramsey Theory becomes relevant in the study of Rosenthal compacta as it was a key ingredient in Rosenthal's ℓ_1 Theorem. For this reason, the main results in [ADK08] (which we cite in this paper) are better explained by indexing Rosenthal compacta with the binary tree.

Definition 3.7. Let X be a Polish space.

- (1) If I is countable and $\{f_i : i \in I\} \subseteq \mathbb{R}^X$, $\{g_i : i \in I\} \subseteq \mathbb{R}^X$ are two pointwise families by I , we say that $\{f_i : i \in I\}$ and $\{g_i : i \in I\}$ are *equivalent* if and only if the map $f_i \mapsto g_i$ is extended to a homeomorphism from $\overline{\{f_i : i \in I\}}$ to $\overline{\{g_i : i \in I\}}$.
- (2) If $\{f_t : t \in 2^{<\mathbb{N}}\}$ is a pointwise bounded family, we say that $\{f_t : t \in 2^{<\mathbb{N}}\}$ is *minimal* if and only if for every dyadic subtree $\{s_t : t \in 2^{<\mathbb{N}}\}$ of $2^{<\mathbb{N}}$, $\{f_{s_t} : t \in 2^{<\mathbb{N}}\}$ is equivalent to $\{f_t : t \in 2^{<\mathbb{N}}\}$.

One of the main results in [ADK08] is that, up to equivalence, there are seven minimal families of Rosenthal compacta and that for every relatively compact $\{f_t : t \in 2^{<\mathbb{N}}\} \subseteq B_1(X)$ there is a dyadic subtree $\{s_t : t \in 2^{<\mathbb{N}}\}$ such that $\{f_{s_t} : t \in 2^{<\mathbb{N}}\}$

is equivalent to one of the minimal families. We shall describe the seven minimal families next. We follow the same notation as in [ADK08]. For any node $t \in 2^{<\mathbb{N}}$, let us denote by $t \smallfrown 0^\infty$ ($t \smallfrown 1^\infty$) the infinite binary sequence starting with t and continuing with all 0's (respectively, all 1's). Fix a regular dyadic subtree $R = \{s_t : t \in 2^{<\mathbb{N}}\}$ of $2^{<\mathbb{N}}$ (i.e., a dyadic subtree such that every level of R is contained in a level of $2^{<\mathbb{N}}$) with the property that for all $s, s' \in R$, $s \smallfrown 0^\infty \neq s' \smallfrown 0^\infty$ and $s \smallfrown 1^\infty \neq s' \smallfrown 1^\infty$. Given $t \in 2^{<\mathbb{N}}$, let v_t be the characteristic function of the set $\{x \in 2^\mathbb{N} : x \text{ extends } t\}$. Let $<$ be the lexicographic order in $2^\mathbb{N}$. Given $a \in 2^\mathbb{N}$, let $f_a^+ : 2^\mathbb{N} \rightarrow \{0, 1\}$ be the characteristic function of $\{x \in 2^\mathbb{N} : a \leq x\}$ and let $f_a^- : 2^\mathbb{N} \rightarrow \{0, 1\}$ be the characteristic function of $\{x \in 2^\mathbb{N} : a < x\}$. Given two maps $f, g : 2^\mathbb{N} \rightarrow \mathbb{R}$ we denote by $(f, g) : 2^\mathbb{N} \sqcup 2^\mathbb{N} \rightarrow \mathbb{R}$ the function which is f on the first copy of $2^\mathbb{N}$ and g on the second copy of $2^\mathbb{N}$.

- (1) $D_1 = \{\frac{1}{|t|+1}v_t : t \in 2^{<\mathbb{N}}\}$. This is discrete in $\overline{D_1} = A(2^\mathbb{N})$.
- (2) $D_2 = \{s_t \smallfrown 0^\infty : t \in 2^{<\mathbb{N}}\}$. This is discrete in $\overline{D_2} = 2^{\leq \mathbb{N}}$.
- (3) $D_3 = \{f_{s_t \smallfrown 0^\infty}^+ : t \in 2^{<\mathbb{N}}\}$. This is discrete in $\overline{D_3} = S(2^\mathbb{N})$.
- (4) $D_4 = \{f_{s_t \smallfrown 1^\infty}^- : t \in 2^{<\mathbb{N}}\}$. This is discrete in $\overline{D_4} = S(2^\mathbb{N})$.
- (5) $D_5 = \{v_t : t \in 2^{<\mathbb{N}}\}$. This is discrete in $\overline{D_5} = \hat{A}(2^\mathbb{N})$.
- (6) $D_6 = \{(v_{s_t}, s_t \smallfrown 0^\infty) : t \in 2^{<\mathbb{N}}\}$. This is discrete in $\overline{D_6} = \hat{D}(2^\mathbb{N})$.
- (7) $D_7 = \{(v_{s_t}, x_{s_t \smallfrown 0^\infty}^+) : t \in 2^{<\mathbb{N}}\}$. This is discrete in $\overline{D_7} = \hat{D}(S(2^\mathbb{N}))$.

Theorem 3.8 (Heptachotomy of minimal families, Theorem 2 in [ADK08]). *Let X be Polish. For every relatively compact $\{f_t : t \in 2^{<\mathbb{N}}\} \subseteq B_1(X)$, there exists $i = 1, 2, \dots, 7$ and a regular dyadic subtree $\{s_t : t \in 2^{<\mathbb{N}}\}$ of $2^{<\mathbb{N}}$ such that $\{f_{s_t} : t \in 2^{<\mathbb{N}}\}$ is equivalent to D_i . Moreover, all D_i are minimal and mutually non-equivalent.*

The implication of this result for deep computations is the following: for any countable set of computations Δ satisfying the NIP (for some CCS (L, \mathcal{P}, Γ)), we can always find a countable discrete set of deep computations that approximates all the other deep computations. For example: in the finite precision prefix test example (subsection 2.5), the prefix test computations (family D_5) approximate all other deep computations. However, note that this discrete set D_i may not consist of continuous functions (i.e., they will not be computable in general). For example, functions in D_3 are not continuous.

4. RANDOMIZED VERSIONS OF NIP AND MONTE CARLO COMPUTABILITY OF DEEP COMPUTATIONS

In this section, we replace deterministic computability by probabilistic ('Monte Carlo') computability. We do not assume that \mathcal{P} is countable. The main results of the section are Theorem 4.8 (connecting NIP and Monte Carlo computability) and 4.14 (connecting Talagrand stability and Monte Carlo computability).

Fundamental in this section is a measure-theoretic version of Theorem 1.11, namely, Theorem 4.5. For the proof of Theorem 1.11, we assumed countability of \mathcal{P} — this ensured that $\mathbb{R}^\mathcal{P}$ a Polish space. In this section, the countability assumption is not needed.

4.1. NIP and Monte Carlo computability of deep computations. The *raison d'être* of the Baire class-1 functions is to have with a class of functions that are obtained as equential limit points of continuous functions. By Fact 1.2, for perfectly

normal X , a function f is in $B_1(X, Y)$ if and only if $f^{-1}[U]$ is an F_σ subset of X for every open $U \subseteq Y$. Thus, for such X , functions in $B_1(X, Y)$ are not too far from being continuous. In this section we will study a more general class of functions, namely, the class of *universally measurable* functions, which we define next.

Definition 4.1. Given a Hausdorff space X and a measurable space (Y, Σ) , we say that $f : X \rightarrow Y$ is *universally measurable* (with respect to Σ) if $f^{-1}(E)$ is μ -measurable for every Radon measure μ on X and every $E \in \Sigma$. When $Y = \mathbb{R}$, we will always take $\Sigma = \mathcal{B}(\mathbb{R})$, the Borel σ -algebra of \mathbb{R} .

If X is a compact (Hausdorff) space, then every Radon measure μ on X is finite. Then, the measure given by $\nu(A) := \mu(A)/\mu(X)$ is a probability measure on X with the same null sets as μ . Hence, Radon measures on compact spaces are equivalent to (Radon) probability measures. We summarize this fact in the next remark:

Remark 4.2. If X is compact, then a function $f : X \rightarrow \mathbb{R}$ is universally measurable if and only if $f^{-1}(U)$ is μ -measurable for every Radon probability measure μ on X and every open set $U \subseteq \mathbb{R}$.

Intuitively, a function is universally measurable if it is “measurable no matter which reasonable way you try to measure things on its domain”. The concept of universal measurability emerged from work of Kallianpur and Sazonov, in the late 1950’s and 1960s — with later developments by Blackwell, Darst and others — building on earlier ideas of Gnedenko and Kolmogorov from the 1950s. See [Pap02, Chapters 1 and 2].

Notation 4.3. Following [BFT78], the collection of all universally measurable real-valued functions on X will be denoted by $M_r(X)$. Given a fixed Radon measure μ on X , the collection of all μ -measurable real-valued functions on X will be denoted by $\mathcal{M}^0(X, \mu)$.

In the context of deep computations, we are interested in transition maps of a state space $L \subseteq \mathbb{R}^{\mathcal{P}}$ into itself. There are two natural σ -algebras one can consider in the product space $\mathbb{R}^{\mathcal{P}}$: the Borel σ -algebra, i.e., the σ -algebra generated by open sets in $\mathbb{R}^{\mathcal{P}}$, and the cylinder σ -algebra (i.e., the σ -algebra generated by the sub-basic open sets in $\mathbb{R}^{\mathcal{P}}$, that is, the sets $\pi_P^{-1}(U)$ with $U \subseteq \mathbb{R}$ open and $P \in \mathcal{P}$). Note that when \mathcal{P} is countable, both σ -algebras coincide, but in general the cylinder σ -algebra is strictly smaller. We will use the cylinder σ -algebra to define universally measurable maps $f : \mathbb{R}^{\mathcal{P}} \rightarrow \mathbb{R}^{\mathcal{P}}$. The reason for this choice is the following characterization:

Proposition 4.4. *Let X be a Hausdorff space and $Y = \prod_{i \in I} Y_i$ be any product of measurable spaces (Y_i, Σ_i) for $i \in I$. Let Σ_Y be the cylinder σ -algebra generated by the measurable spaces (Y_i, Σ_i) . The following are equivalent for $f : X \rightarrow Y$:*

- (i) $f : X \rightarrow Y$ is universally measurable (with respect to Σ_Y).
- (ii) $\pi_i \circ f : X \rightarrow Y_i$ is universally measurable (with respect to Σ_i) for all $i \in I$.

Proof. (i) \Rightarrow (ii) is clear since the projection maps π_i are measurable and the composition of measurable functions is measurable. To prove (ii) \Rightarrow (i), suppose that $C = \prod_{i \in I} C_i$ is a measurable cylinder and let J be the finite subset of I such that $C_i \neq Y_i$ for $i \in J$. Then, $C = \bigcap_{i \in J} \pi_i^{-1}(C_i)$, so $f^{-1}(C) = \bigcap_{i \in J} (\pi_i \circ f)^{-1}(C_i)$ is universally measurable by assumption. \square

The preceding proposition says that a transition map is universally measurable if and only if it is universally measurable on all its features; in other words, we can check measurability of a transition just by checking measurability feature by feature. This is the same as in the Baire class-1 case (compare with Proposition 1.10).

The main result in section 3 is that, as long as we work with countably many features, PAC-learning (or NIP) corresponds to relative compactness in the space of Baire class-1 functions. The following result (which does not assume countability of the number of features) gives an analogous characterization of the NIP in terms of universal measurability:

Theorem 4.5 (Bourgain-Fremlin-Talagrand, Theorem 2F in [BFT78]). *Let X be a Hausdorff space and $A \subseteq C(X)$ be pointwise bounded. The following are equivalent:*

- (i) $\overline{A} \subseteq M_r(X)$.
- (ii) For every compact $K \subseteq X$, $A|_K$ satisfies the NIP.
- (iii) For every Radon measure μ on X , A is relatively countably compact in $\mathcal{M}^0(X, \mu)$, i.e., every countable subset of A has a limit point in $\mathcal{M}^0(X, \mu)$.

This result allows us to formalize the concept of a deep computation being *Monte Carlo computable*, which we define below (Definition 4.6). To motivate the definition, let us first recall two facts:

- (1) Littlewood’s second principle states that every Lebesgue measurable function is “nearly continuous”. The formal statement of this, which is Luzin’s theorem, is that if (X, Σ, μ) a Radon measure space and Y is a second-countable topological space (e.g., $Y = \mathbb{R}^{\mathcal{P}}$ with \mathcal{P} countable) equipped with the Borel σ -algebra, then any given $f : X \rightarrow Y$ is measurable if and only if for every $E \in \Sigma$ and every $\varepsilon > 0$ there exists a closed $F \subseteq E$ such that the restriction $f|_F$ is continuous and $\mu(E \setminus F) < \varepsilon$.
- (2) Computability of deep computations is characterized in terms of continuous extendibility of computations. This is at the core of [ADIW24].

These two facts motivate the following definition:

Definition 4.6. Let (L, \mathcal{P}, Γ) be a CCS. We say that a transition $f : L \rightarrow L$ is *universally Monte Carlo computable* if and only if there exists $\tilde{f} : \mathcal{L}_{\text{sh}} \rightarrow \mathcal{L}_{\text{sh}}$ extending f such that for every sizer r_{\bullet} there is a sizer s_{\bullet} such that the restriction $\tilde{f}|_{\mathcal{L}[r_{\bullet}]} : \mathcal{L}[r_{\bullet}] \rightarrow \mathcal{L}[s_{\bullet}]$ is universally measurable, i.e., $\pi_P \circ \tilde{f}|_{\mathcal{L}[r_{\bullet}]} : \mathcal{L}[r_{\bullet}] \rightarrow [-s_P, s_P]$ is μ -measurable for every Radon probability measure μ on $\mathcal{L}[r_{\bullet}]$ and $P \in \mathcal{P}$.

Remark 4.7. Condition (2) of Theorem 4.5 shows that to study measure-theoretic versions of NIP, we need only consider compact subsets of X . Now, every Radon measure on a compact space is finite; hence, by proper normalization, it can be treated as a probability measure. Therefore, in the context of Monte Carlo measurability, we focus on Radon probability measures rather than general Radon measures.

Theorem 4.8. *Let (L, \mathcal{P}, Γ) be a CCS satisfying the Extendibility Axiom. Let R be an exhaustive collection of sizers. Let $\Delta \subseteq \Gamma$ be R -confined. If $\pi_P \circ \Delta|_{\mathcal{L}[r_{\bullet}]}$ satisfies the NIP for all $P \in \mathcal{P}$ and all $r_{\bullet} \in R$, then every deep computation in Δ is universally Monte Carlo computable.*

Proof. Fix $P \in \mathcal{P}$ and $r_{\bullet} \in R$. By the Extendibility Axiom, $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_{\bullet}]}$ is a set of pointwise bounded continuous functions on the compact set $\mathcal{L}[r_{\bullet}]$. Since

$\pi_P \circ \tilde{\Delta}|_{L[r_\bullet]} = \pi_P \circ \Delta|_{L[r_\bullet]}$ has the NIP, so does $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$ by Lemma 1.12. By Theorem 4.5, we have $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]} \subseteq M_r(\mathcal{L}[r_\bullet])$ for all $r_\bullet \in R$ and $P \in \mathcal{P}$. Let $f \in \overline{\Delta}$ be a deep computation. Write $f = \mathcal{U} \lim_i \gamma_i$ as an ultralimit of computations in Δ . Define $\tilde{f} := \mathcal{U} \lim_i \tilde{\gamma}_i$. Then, $\tilde{f} : \mathcal{L}_{\text{sh}} \rightarrow \mathcal{L}_{\text{sh}}$ extends f . Since Δ is R -confined we have that $f : L[r_\bullet] \rightarrow L[r_\bullet]$ and $\tilde{f} : \mathcal{L}[r_\bullet] \rightarrow \mathcal{L}[r_\bullet]$ for all $r_\bullet \in R$. Lastly, note that for all $r_\bullet \in R$ and $P \in \mathcal{P}$ we have that $\pi_P \circ \tilde{f}|_{\mathcal{L}[r_\bullet]} \in \pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]} \subseteq M_r(\mathcal{L}[r_\bullet])$. \square

Question 4.9. Under the same assumptions of the preceding theorem, suppose that every deep computation of Δ is universally Monte Carlo computable. Must $\pi_P \circ \Delta|_{L[r_\bullet]}$ have the NIP for all $P \in \mathcal{P}$ and all $r_\bullet \in R$?

4.2. Talagrand stability and Monte Carlo computability of deep computations. There is another notion closely related to NIP, introduced by Talagrand in [Tal84] while studying Pettis integration. Suppose that X is a compact Hausdorff space and $A \subseteq \mathbb{R}^X$. Let μ be a Radon probability measure on X . Given a μ -measurable set $E \subseteq X$, a positive integer k and real numbers $a < b$, we write:

$$D_k(A, E, a, b) = \bigcup_{f \in A} \{x \in E^{2k} : f(x_{2i}) \leq a, f(x_{2i+1}) \geq b \text{ for all } i < k\}.$$

We say that A is *Talagrand μ -stable* if and only if for every μ -measurable set $E \subseteq X$ of positive measure and for every $a < b$ there is a $k \geq 1$ such that

$$(\mu^{2k})^*(D_k(A, E, a, b)) < (\mu(E))^{2k},$$

where μ^* denotes the outer measure (we need to work with outer measure since the sets $D_k(A, E, a, b)$ need not be μ^{2k} -measurable). The inequality certainly holds when A is a countable set of continuous (or μ -measurable) functions.

The main result of this section is that deep computations, i.e., limit points of a Talagrand stable set of computations are Monte Carlo computable; this is Theorem 4.14 below. We now prove that limit points of a Talagrand μ -stable set are μ -measurable. But first, let us state the following useful characterization of measurable functions (compare with Fact 1.2):

Fact 4.10 (Lemma 413G in [Fre03]). *Suppose that (X, Σ, μ) is a measure space and $\mathcal{K} \subseteq \Sigma$ is a collection of measurable sets satisfying the following conditions:*

- (1) (X, Σ, μ) is complete, i.e., for all $E \in \Sigma$ with $\mu(E) = 0$ and $F \subseteq E$ we have $F \in \Sigma$.
- (2) (X, Σ, μ) is semi-finite, i.e., for all $E \in \Sigma$ with $\mu(E) = \infty$ there exists $F \subseteq E$ such that $F \in \Sigma$ and $0 < \mu(F) < \infty$.
- (3) (X, Σ, μ) is saturated, i.e., $E \in \Sigma$ if and only if $E \cap F \in \Sigma$ for all $F \in \Sigma$ with $\mu(F) < \infty$.
- (4) (X, Σ, μ) is inner regular with respect to \mathcal{K} , i.e., for all $E \in \Sigma$

$$\mu(E) = \sup\{\mu(K) : K \in \mathcal{K} \text{ and } K \subseteq E\}.$$

(In particular, if X is compact Hausdorff, μ is a Radon probability measure on X , Σ is the completion of the Borel σ -algebra by μ , and \mathcal{K} is the collection of compact subsets of X , all these conditions hold). Then, $f : X \rightarrow \mathbb{R}$ is measurable if and only if for every $K \in \mathcal{K}$ with $0 < \mu(K) < \infty$ and $a < b$, either $\mu^*(P) < \mu(K)$ or $\mu^*(Q) < \mu(K)$ where $P = \{x \in K : f(x) \leq a\}$ and $Q = \{x \in K : f(x) \geq b\}$.

The following technical lemma will be instrumental for proving Proposition 4.13, which, in turn, will yield the main result of the subsection, namely Theorem 4.14.

Lemma 4.11. *If A is Talagrand μ -stable, then \overline{A} is also Talagrand μ -stable and $\overline{A} \subseteq \mathcal{M}^0(X, \mu)$.*

Proof. First, we claim that a subset of a μ -stable set is μ -stable. To see this, suppose that $A \subseteq B$ and B is μ -stable. Fix any μ -measurable $E \subseteq X$ of positive measure and $a < b$. Let $k \geq 1$ such that

$$(\mu^{2k})^*(D_k(B, E, a, b)) < (\mu(E))^{2k}.$$

Since $A \subseteq B$, we have $D_k(A, E, a, b) \subseteq D_k(B, E, a, b)$; therefore,

$$(\mu^{2k})^*(D_k(A, E, a, b)) \leq (\mu^{2k})^*(D_k(B, E, a, b)) < (\mu(E))^{2k}.$$

We now show that \overline{A} is μ -stable. Fix $E \subseteq X$ measurable with positive measure and $a < b$. Let $a' < b'$ be such that $a < a' < b' < b$. Since A is μ -stable, let $k \geq 1$ be such that

$$(\mu^{2k})^*(D_k(A, E, a', b')) < (\mu(E))^{2k}.$$

If $x \in D_k(\overline{A}, E, a, b)$, then there is $f \in \overline{A}$ such that $f(x_{2i}) \leq a < a'$ and $f(x_{2i+1}) \geq b > b'$ for all $i < k$. By definition of pointwise convergence topology, there exists $g \in A$ such that $g(x_{2i}) < a'$ and $g(x_{2i+1}) > b'$ for all $i < k$. Hence, $x \in D_k(A, E, a', b')$. We have shown that $D_k(\overline{A}, E, a, b) \subseteq D_k(A, E, a', b')$; hence,

$$(\mu^{2k})^*(D_k(\overline{A}, E, a, b)) \leq (\mu^{2k})^*(D_k(A, E, a', b')) < (\mu(E))^{2k}.$$

It suffices to show that $\overline{A} \subseteq \mathcal{M}^0(X, \mu)$. Suppose that there exists $f \in \overline{A}$ such that $f \notin \mathcal{M}^0(X, \mu)$. By fact 4.10, there exists a μ -measurable set E of positive measure and $a < b$ such that $\mu^*(P) = \mu^*(Q) = \mu(E)$ where $P = \{x \in E : f(x) \leq a\}$ and $Q = \{x \in E : f(x) \geq b\}$. Then, for any $k \geq 1$: $(P \times Q)^k \subseteq D_k(\{f\}, E, a, b)$, so $(\mu^{2k})^*(D_k(\{f\}, E, a, b)) = (\mu^*(P)\mu^*(Q))^k = (\mu(E))^{2k}$. Thus, $\{f\}$ is not μ -stable. However, we argued above that a subset of a μ -stable set must be μ -stable, so we have a contradiction. \square

Definition 4.12. We say that A is *universally Talagrand stable* if A is Talagrand μ -stable for every Radon probability measure μ on X .

We first observe that universal Talagrand stability corresponds to a complexity class smaller than or equal to the NIP class:

Proposition 4.13. *Let X be a compact Hausdorff space and $A \subseteq C(X)$ be pointwise bounded. If A is universally Talagrand stable, then A satisfies the NIP.*

Proof. By Theorem 4.5, it suffices to show that A is relatively countably compact in $\mathcal{M}^0(X, \mu)$ for every Radon probability measure μ on X . Since A is Talagrand μ -stable for any such μ , we have $\overline{A} \subseteq \mathcal{M}^0(X, \mu)$ by Lemma 4.11. In particular, A is relatively countably compact in $\mathcal{M}^0(X, \mu)$. \square

Corollary 4.14. *Let (L, \mathcal{P}, Γ) be a CCS satisfying the Extendibility Axiom. If $\pi_P \circ \Delta|_{L[r_\bullet]}$ is universally Talagrand stable for all $P \in \mathcal{P}$ and all sizes r_\bullet , then every deep computation is universally Monte Carlo computable.*

Proof. This is a direct consequence of Proposition 4.13 and Theorem 4.8. \square

In the context of deep computations, we have identified two ways to obtain Monte Carlo computability, namely, NIP/PAC and Talagrand stability. It is natural to ask whether these two notions are equivalent. The following results show that, even in the simple case of countably many computations, this question is sensitive to the set-theoretic axioms. On the one hand, it is consistent (with respect to the standard ZFC axioms of set theory) that these two classes are the same:

Theorem 4.15 (Talagrand, Theorem 9-3-1(a) in [Tal84]). *Let X be a compact Hausdorff space and $A \subseteq M_r(X)$ be countable and pointwise bounded. Assume that $[0, 1]$ is not the union of $< \mathfrak{c}$ closed measure zero sets. If A satisfies the NIP, then A is universally Talagrand stable.*

(The assumption that $[0, 1]$ is not the union of $< \mathfrak{c}$ closed measure zero sets is a consequence of, for example, the Continuum Hypothesis.)

On the other hand, by fixing a particular well-known measure, namely the Lebesgue measure, we see that the other case is also consistent:

Theorem 4.16 (Fremlin, Shelah, [FS93]). *It is consistent with the usual axioms of set theory that there exists a countable pointwise bounded set of Lebesgue measurable functions with the NIP which is not Talagrand stable with respect to Lebesgue measure.*

Notice that the preceding two results apply to sets of measurable functions, a class of functions larger than the class of continuous functions. However, by the Extendibility Axiom, finitary computations are continuous, i.e., if A is a set of computations, then $A \subseteq C_p(X)$. The question of whether we can remove the set-theoretic assumption in Theorem 4.15 when $A \subseteq C_p(X)$ (instead of $A \subseteq M_r(X)$) remains open.

REFERENCES

- [ADIW24] Samson Alva, Eduardo Dueñez, Jose Iovino, and Claire Walton. Approximability of deep equilibria. *arXiv preprint arXiv:2409.06064*, 2024. To appear in Mathematical Structures in Computer Science.
- [ADK08] Spiros A. Argyros, Pandelis Dodos, and Vassilis Kanellopoulos. A classification of separable Rosenthal compacta and its applications. *Dissertationes Mathematicae*, 449:1–55, 2008.
- [Ark92] A. V. Arkhangel’skii. *Topological function spaces*, volume 78 of *Mathematics and its Applications (Soviet Series)*. Kluwer Academic Publishers Group, Dordrecht, 1992. Translated from the Russian by R. A. M. Hoksbergen.
- [BFT78] J. Bourgain, D. H. Fremlin, and M. Talagrand. Pointwise compact sets of Baire-measurable functions. *American Journal of Mathematics*, 100(4):845–886, 1978.
- [BKK] Shaojie Bai, J. Zico Kolter, and Vladlen Koltun. Deep equilibrium models. Preprint. <https://arxiv.org/abs/1909.01377>, <https://implicit-layers-tutorial.org/>.
- [Bla81] Andreas Blass. A partition theorem for perfect sets. *Proceedings of the American Mathematical Society*, 82(2):271–277, 1981.
- [Bla84] Paul Blanchard. Complex analytic dynamics on the Riemann sphere. *Bulletin of the American Mathematical Society*, 11(1):85–141, 1984.
- [CRBD] Ricky T. Q. Chen, Yulia Rubanova, Jesse Bettencourt, and David Duvenaud. Neural ordinary differential equations. Preprint. <https://arxiv.org/abs/1806.07366>.
- [Deb13] Gabriel Debs. Descriptive aspects of Rosenthal compacta. In *Recent Progress in General Topology III*, pages 205–227. Springer, 2013.
- [Eng89] Ryszard Engelking. *General topology*, volume 6 of *Sigma Series in Pure Mathematics*. Heldermann Verlag, Berlin, second edition, 1989. Translated from the Polish by the author.

- [Fre03] David H. Fremlin. *Measure Theory, Volume 4: Topological Measure Spaces*. Torres Fremlin, Colchester, UK, 2003.
- [FS93] David H. Fremlin and Saharon Shelah. Pointwise compact and stable sets of measurable functions. *Journal of Symbolic Logic*, 58(2):435–455, 1993.
- [GM22] Eli Glasner and Michael Megrelishvili. Todorčević’ trichotomy and a hierarchy in the class of tame dynamical systems. *Transactions of the American Mathematical Society*, 375(7):4513–4548, 2022.
- [God80] Gilles Godefroy. Compacts de Rosenthal. *Pacific J. Math.*, 91(2):293–306, 1980.
- [Gro52] A. Grothendieck. Critères de compacité dans les espaces fonctionnels généraux. *Amer. J. Math.*, 74:168–186, 1952.
- [HL66] J. D. Halpern and H. Läuchli. A partition theorem. *Transactions of the American Mathematical Society*, 124:360–367, 1966.
- [HT19] Haim Horowitz and Stevo Todorčević. Compact sets of baire class one functions and maximal almost disjoint families, 2019.
- [Iov94] José N. Iovino. *Stable Theories in Functional Analysis*. PhD thesis, University of Illinois at Urbana-Champaign, 1994.
- [Kei03] H. Jerome Keisler. Model theory for real-valued structures. In José Iovino, editor, *Beyond First Order Model Theory, Volume II*. CRC Press, Boca Raton, FL, 2003.
- [LSV93] Alain Louveau, Saharon Shelah, and Boban Veličković. Borel partitions of infinite subtrees of a perfect tree. *Annals of Pure and Applied Logic*, 63(3):271–281, 1993.
- [Mil81] Keith Milliken. A partition theorem for the infinite subtrees of a tree. *Transactions of the American Mathematical Society*, 263:137–148, 1981.
- [Mil89] Arnold W. Miller. Infinite combinatorics and definability. *Annals of Pure and Applied Logic*, 41(2):179–203, 1989.
- [Pap02] E. Pap, editor. *Handbook of measure theory. Vol. I, II*. North-Holland, Amsterdam, 2002.
- [Ros74] Haskell P. Rosenthal. A characterization of Banach spaces containing l^1 . *Proc. Nat. Acad. Sci. U.S.A.*, 71:2411–2413, 1974.
- [RPK19] M. Raissi, P. Perdikaris, and G. E. Karniadakis. Physics-informed neural networks: a deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. *J. Comput. Phys.*, 378:686–707, 2019.
- [Sau72] N. Sauer. On the density of families of sets. *J. Combinatorial Theory Ser. A*, 13:145–147, 1972.
- [She71] Saharon Shelah. Stability, the f.c.p., and superstability; model theoretic properties of formulas in first order theory. *Ann. Math. Logic*, 3(3):271–362, 1971.
- [She72] Saharon Shelah. A combinatorial problem; stability and order for models and theories in infinitary languages. *Pacific J. Math.*, 41:247–261, 1972.
- [She90] Saharon Shelah. *Classification theory and the number of nonisomorphic models*, volume 92 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, second edition, 1990.
- [Tal84] Michel Talagrand. *Pettis Integral and Measure Theory*, volume 51 of *Memoirs of the American Mathematical Society*. American Mathematical Society, Providence, RI, USA, 1984. Includes bibliography (pp. 220–224) and index.
- [Tka11] Vladimir V. Tkachuk. *A C_p -Theory Problem Book: Topological and Function Spaces*. Problem Books in Mathematics. Springer, 2011.
- [Tod97] Stevo Todorčević. *Topics in Topology*, volume 1652 of *Lecture Notes in Mathematics*. Springer Berlin, Heidelberg, 1997.
- [Tod99] Stevo Todorčević. Compact subsets of the first Baire class. *Journal of the American Mathematical Society*, 12(4):1179–1212, 1999.
- [Tod23] Stevo Todorčević. Dense metrizable. *Annals of Pure and Applied Logic*, 175:103327, 07 2023.
- [Val84] Leslie G. Valiant. A theory of the learnable. *Communications of the ACM*, 27(11):1134–1142, 1984.
- [VC71] Vladimir N Vapnik and Alexey Ya Chervonenkis. On the uniform convergence of relative frequencies of events to their probabilities. *Theory of Probability & Its Applications*, 16(2):264–280, 1971.

- [VC74] Vladimir N Vapnik and Alexey Ya Chervonenkis. *Theory of Pattern Recognition*. Nauka, Moscow, 1974. German Translation: Theorie der Zeichenerkennung, Akademie-Verlag, Berlin, 1979.