

1  
2  
  
3  
4  
5  
6

COMPLEXITY OF DEEP COMPUTATIONS  
VIA TOPOLOGY OF FUNCTION SPACES

EDUARDO DUEÑEZ<sup>1</sup>      JOSÉ IOVINO<sup>1</sup>      TONATIUH MATOS-WIEDERHOLD<sup>2</sup>  
LUCIANO SALVETTI<sup>2</sup>      FRANKLIN D. TALL<sup>2</sup>

<sup>1</sup>Department of Mathematics, University of Texas at San Antonio  
<sup>2</sup>Department of Mathematics, University of Toronto

ABSTRACT. We study complexity of deep computations. We use topology of function spaces, specifically, the classification Rosenthal compacta, to identify new complexity classes. We use the language of model theory, specifically, the concept of the independence from Shelah’s classification theory, to translate between topology and computation.

7

0. INTRODUCTION

8      In this paper we study limit behavior of real-valued computations as the value  
9 of certain parameters of the computation model tend towards infinity, or towards  
10 zero, or towards some other fixed value, e.g., the depth of a neural network tending  
11 to infinity, or the time interval between layers of the network tending toward zero.  
12 Recently, particular cases of this situation have attracted considerable attention  
13 in deep learning research (e.g., Neural Ordinary Differential Equations [CRBD],  
14 Physics-Informed Neural Networks [RPK19], deep equilibrium models [BKK], etc).  
15 In this paper, we combine ideas of topology and model theory to study these limit  
16 phenomena from a unified viewpoint.  
17      Informed by model theory, to each computation in a given computation model,  
18 we associate a continuous real-valued function, called the *type* of the computation,  
19 that describes the logical properties of this computation with respect to the rest of  
20 the model. This allows us to view computations in any given computational model  
21 as elements of a space of real-valued functions, which is called the *space of types*  
22 of the model. The idea of embedding models of theories into their type spaces is  
23 central in model theory. The embedding of computations into spaces of types allows  
24 us to utilize the vast theory of topology of function spaces, known as  $C_p$ -theory,  
25 to obtain results about complexity of topological limits of computations. As we  
26 shall indicate next, recent classification results for spaces of functions provide an  
27 elegant and powerful machinery to classify computations according to their levels  
28 of “tameness” or “wildness”, with the former corresponding roughly to polyno-  
29 mial approximability and the latter to exponential approximability. The viewpoint  
30 of spaces of types, which we have borrowed from model theory, thus becomes a  
31 “Rosetta stone” that allows us to interconnect various classification programs: In  
32 topology, the classification of Rosenthal compacta pioneered by Todorćević [Tod99];  
33 in logic, the classification of theories developed by Shelah [She90]; and in statistical  
34 learning, the notion PAC learning and VC dimension pioneered by Vapkins and  
35 Chervonenkis [VC74, VC71].  
36      In a previous paper [ADIW24], we introduced the concept of limits of compu-  
37 tations, which we called *ultracomputations* (given they arise as ultrafilter limits of

standard computations) and *deep computations* (following usage in machine learning [BKK]). There is a technical difference between both designations, but in this paper, to simplify the nomenclature, we will ignore the difference and use only the term “deep computation”.

In [ADIW24], we proved a new “tame vs wild” (i.e., polynomial vs exponential) dichotomy for complexity of deep computations by invoking a classical result of Grothendieck from late 50s [Gro52]. Under our model-theoretic Rosetta stone, polynomial approximability in the sense of computation becomes identified with the notion of continuous extendability in the sense of topology, and with the notions of *stability* and *type definability* in the sense of model theory.

In this paper, we follow a more general approach, i.e., we view deep computations as pointwise limits of continuous functions. In topology, real-valued functions that arise as the pointwise limit of a sequence of continuous are called *functions of the first Baire class*, or *Baire class 1* functions, or *Baire-1* for short; Baire class 1 form a step above simple continuity in the hierarchy of functions studied in real analysis (Baire class 0 functions being continuous functions). Intuitively, Baire-1 functions represent functions with “controlled” discontinuities, so they are crucial in topology and set theory.

We prove a new “tame vs wild” Ramsey-theoretic dichotomy for complexity of general deep computations by invoking a famous paper by Bourgain, Fremlin and Talagrand from the late 70s [BFT78], and a new trichotomy for the class of “tame” deep computations by invoking an equally celebrated result of Todorćević, from the late 90s, for functions of the first Baire class [Tod99].

Todorćević’s trichotomy regards *Rosenthal compacta*; these are special classes of topological spaces, defined as compact spaces that can be embedded (homeomorphically identified as a subset) within the space of Baire class 1 functions on some Polish (separable, complete metric) space, under the pointwise convergence topology. Rosenthal compacta exhibit “topological tameness,” meaning that they behave in relatively controlled ways, and since the late 70’s, they have played a crucial role for understanding complexity of structures of functional analysis, especially, Banach spaces. Todorćević’s trichotomy has been utilized to settle longstanding problems in topological dynamics and topological entropy [GM22].

Through our Rosetta stone, Rosenthal compacta in topology correspond to the important concept of “No Independence Property” (known as “NIP”) in model theory, identified by Shelah [She71, She90], and to the concept of Probably Approximately Correct learning (known as “PAC learnability”) in statistical learning theory identified by Valiant [Val84].

Going beyond Todorćević’s trichotomy, we invoke a more recent heptachotomy for Rosenthal compacta obtained by Argyros, Dodos and Kanellopoulos [ADK08]. Argyros, Dodos and Kanellopoulos identified the fundamental “prototypes” of separable Rosenthal compacta, and proved that any non-metrizable separable Rosenthal compactum must contain a “canonical” embedding of one of these prototypes. They showed that if a separable Rosenthal compactum is not hereditarily separable, it must contain an uncountable discrete subspace of the size of the continuum.

We believe that the results presented in this paper show practitioners of computation, or topology, or descriptive set theory, or model theory, how classification invariants used in their field translate into classification invariants of other fields. However, in the interest of accessibility, we do not assume previous familiarity with

high-level topology or model theory, or computing. The only technical prerequisite of the paper is undergraduate-level topology. The necessary topological background beyond undergraduate topology is covered in section 1.

Throughout the paper, we focus on classical computation; however, by refining the model-theoretic tools, the results presented here can be extended to quantum computation and open quantum systems. This extension will be addressed in a forthcoming paper.

## CONTENTS

0. Introduction	1
1. General topological preliminaries: From continuity to Baire class 1	3
1.1. From Rosenthal's dichotomy to the Bourgain-Fremlin-Talagrand dichotomy to Shelah's NIP	5
1.2. NIP as universal dividing line between polynomial and exponential complexity	7
1.3. Rosenthal compacta	8
1.4. The special case $B_1(X, \mathbb{R}^{\mathcal{P}})$ with $\mathcal{P}$ countable.	8
2. Compositional computation structures.	10
3. Classifying deep computations	12
3.1. NIP, Rosenthal compacta, and deep computations	12
3.2. The Todorćević trichotomy and levels of PAC learnability	12
3.3. The Argyros-Dodos-Kanellopoulos heptachotomy, and approximability of deep computation by minimal classes	14
3.4. Bourgain-Fremlin-Talagrand, NIP, and essential computability of deep computations	16
3.5. Talagrand stability, NIP, and essential computability of deep computations	18
References	19

## 1. GENERAL TOPOLOGICAL PRELIMINARIES: FROM CONTINUITY TO BAIRE CLASS 1

In this section we give preliminaries from general topology and function space theory. We include some of the proofs for completeness, but the reader familiar with these topics may skip them.

Recall that a subset of a topological space is  $F_\sigma$  if it is a countable union of closed sets, and  $G_\delta$  if it is a countable intersection of closed sets. Note that in a metrizable space, every open set is  $F_\sigma$ ; equivalently, every closed set is  $G_\delta$ .

A *Polish space* is a separable and completely metrizable topological space. The most important examples are the reals  $\mathbb{R}$ , the Cantor space  $2^{\mathbb{N}}$  (the set of all infinite binary sequences, endowed with the product topology), and the Baire space  $\mathbb{N}^{\mathbb{N}}$  (the set of all infinite sequences of naturals, also with the product topology). Countable products of Polish spaces are Polish; this includes spaces like  $\mathbb{R}^{\mathbb{N}}$ , the space of sequences of real numbers.

In this paper, we shall discuss subspaces, and so there is a pertinent subtlety of the definitions worth mentioning: *completely metrizable space* is not the same as *complete metric space*; for an illustrative example, the interval  $(0, 1)$  with the metric

inherited from the reals not complete, but it is Polish since that is homeomorphic to the real line. Being Polish is a topological property.

The following result is a cornerstone of descriptive set theory, closely tied to the work of Waclaw Sierpiński and Kazimierz Kuratowski, with proofs often built upon their foundations and formalized later, notably, involving Stefan Mazurkiewicz's work on complete metric spaces.

**Fact 1.1.** *A subset  $A$  of a Polish space  $X$  is itself Polish in the subspace topology if and only if it is a  $G_\delta$  set. In particular, closed subsets and open subsets of Polish spaces are also Polish spaces.*

Given two topological spaces  $X$  and  $Y$  we denote by  $C_p(X, Y)$  the set of all continuous functions  $f : X \rightarrow Y$  endowed with the topology of pointwise convergence. When  $Y = \mathbb{R}$ , we denote this collection simply as  $C_p(X)$ . A natural question is, how do topological properties of  $X$  translate to  $C_p(X)$  and vice versa? These questions, and in general the study of these spaces, are the concern of  $C_p$ -theory, an active field of research in general topology which was pioneered by A. V. Arhangel'skiĭ and his students in the 1970's and 1980's. This field has found many applications in model theory and functional analysis. Recent surveys on the topics include [HT23] and [Tka11].

A *Baire class 1* function between topological spaces is a function that can be expressed as the pointwise limit of a sequence of continuous functions. If  $X$  and  $Y$  are topological spaces, the Baire class 1 functions  $f : X \rightarrow Y$  endowed with the topology of pointwise convergence is denoted  $B_1(X, Y)$ . As above, in the special case  $Y = \mathbb{R}$  we denote  $B_1(X, Y)$  as  $B_1(X)$ . Clearly,  $C_p(X, Y) \subseteq B_1(X, Y)$ . The Baire hierarchy of functions was introduced by French mathematician René-Louis Baire in his 1899 doctoral thesis, *Sur les fonctions de variables réelles*. His work moved away from the 19th-century preoccupation with "pathological" functions toward a constructive classification based on pointwise limits.

A topological space  $X$  is *perfectly normal* if it is normal and every closed subset of  $X$  is a  $G_\delta$  (equivalently, every open subset of  $X$  is a  $G_\delta$ ). Note that every metrizable space is perfectly normal.

The following fact was established by Baire in thesis. A proof can be found in Section 10 of [Tod97].

**Fact 1.2** (Baire). *If  $X$  is perfectly normal, then the following conditions are equivalent for a function  $f : X \rightarrow \mathbb{R}$ :*

- $f$  is a Baire class 1 function, that is,  $f \in B_1(X)$ .
- $f^{-1}[U]$  is an  $F_\sigma$  subset of  $X$  whenever  $U \subseteq \mathbb{R}$  is open.
- $f$  is a pointwise limit of continuous functions.
- For every closed  $F \subseteq X$ , the restriction  $f|_F$  has a point of continuity.

Moreover, if  $X$  is Polish and  $f \notin B_1(X)$ , then there exists countable  $D_0, D_1 \subseteq X$  and reals  $a < b$  such that

$$D_0 \subseteq f^{-1}(-\infty, a], \quad D_1 \subseteq f^{-1}[b, \infty), \quad \overline{D_0} = \overline{D_1}.$$

A subset  $L$  of a topological space  $X$  is *relatively compact* in  $X$  if the closure of  $L$  in  $X$  is compact. Relatively compact subsets of  $B_1(X)$  (for  $X$  Polish space) have been objects of interest for researchers in Analysis and Topological Dynamics.

We begin with the following well-known result. Recall that a set  $A \subseteq \mathbb{R}^X$  of real-valued functions is *pointwise bounded* if for every  $x \in X$  there is  $M_x > 0$  such that  $|f(x)| < M_x$  for all  $f \in A$ . We include a proof for the reader's convenience:

**Lemma 1.3.** *Let  $X$  be a Polish space and  $A \subseteq B_1(X)$  be pointwise bounded. The following are equivalent:*

- (i)  $A$  is relatively compact in  $B_1(X)$ .
- (ii)  $A$  is relatively countably compact in  $B_1(X)$ , i.e., every countable subset of  $A$  has an accumulation point in  $B_1(X)$ .
- (iii)  $\overline{A} \subseteq B_1(X)$ , where  $\overline{A}$  denotes the closure in  $\mathbb{R}^X$ .

*Proof.* Since  $A$  is pointwise bounded, for each  $x \in X$ , fix  $M_x > 0$  such that  $|f(x)| \leq M_x$  for every  $f \in A$ .

(i)  $\Rightarrow$  (ii) holds in general.

(ii)  $\Rightarrow$  (iii) Assume that  $A$  is relatively countably compact in  $B_1(X)$  and that  $f \in \overline{A} \setminus B_1(X)$ . By Fact 1.2, there are countable  $D_0, D_1 \subseteq X$  with  $\overline{D_0} = \overline{D_1}$ , and  $a < b$  such that  $D_0 \subseteq f^{-1}(-\infty, a]$  and  $D_1 \subseteq f^{-1}[b, \infty)$ . We claim that there is a sequence  $\{f_n\}_{n \in \mathbb{N}} \subseteq A$  such that  $\lim_{n \rightarrow \infty} f_n(x) = f(x)$  for all  $x \in D_0 \cup D_1$ . Indeed, use the countability to enumerate  $D_0 \cup D_1$  as  $\{x_n\}_{n \in \mathbb{N}}$ . Then for each positive  $n$  find  $f_n \in A$  with  $|f_n(x_i) - f(x_i)| < \frac{1}{n}$  for all  $i \leq n$ . The claim follows.

By relative countable compactness of  $A$ , there is an accumulation point  $g \in B_1(X)$  of  $\{f_n\}_{n \in \mathbb{N}}$ . It is straightforward to show that since  $f$  and  $g$  agree on  $D_0 \cup D_1$ ,  $g$  does not have a point of continuity on the closed set  $\overline{D_0} = \overline{D_1}$ , which contradicts Fact 1.2.

(iii)  $\Rightarrow$  (i) Suppose that  $\overline{A} \subseteq B_1(X)$ . Then  $\overline{A} \cap B_1(X) = \overline{A}$  is a closed subset of  $\prod_{x \in X} [-M_x, M_x]$ ; Tychonoff's theorem states that the product of compact spaces is always compact, and since closed subsets of compact spaces are compact,  $\overline{A}$  must be compact, as desired.  $\square$

**1.1. From Rosenthal's dichotomy to the Bourgain-Fremlin-Talagrand dichotomy to Shelah's NIP.** The fundamental idea that connects the rich theory here presented to real-valued computations is the concept of an *approximation*. In the reals, points of closure from some subset can always be approximated by points inside the set, via a convergent sequence. For more complicated spaces, such as  $C_p(X)$ , this fails in remarkable ways. To see an example, consider the Cantor space  $X = 2^{\mathbb{N}}$ , and for each  $n \in \mathbb{N}$  define  $p_n : X \rightarrow \{0, 1\}$  by  $p_n(x) = x(n)$  for each  $x \in X$ . Then  $p_n$  is continuous for each  $n$ , but one can show (see Chapter 1.1 of [Tod97] for details) that the only continuous functions in the closure of  $\{p_n\}_{n \in \mathbb{N}}$  are the functions  $p_n$  themselves; moreover, none of the subsequences of  $\{p_n\}_{n \in \mathbb{N}}$  converge. In some sense, this example is the worst possible scenario for convergence. The topological space obtained from this closure is well-known: it is the *Stone-Ćech compactification* of the discrete space of natural numbers, or  $\beta\mathbb{N}$  for short, and it is an important object of study in general topology.

The following theorem, established by Haskell Rosenthal in 1974, is fundamental in functional analysis, and describes a sharp division in the behavior of sequences within a Banach space:

**Theorem 1.4** (Rosenthal's Dichotomy, [Ros74]). *If  $X$  is Polish and  $\{f_n\} \subseteq C_p(X)$  is pointwise bounded, then either  $\{f_n\}_{n \in \mathbb{N}}$  contains a convergent subsequence or a subsequence whose closure (in  $\mathbb{R}^X$ ) is homeomorphic to  $\beta\mathbb{N}$ .*

In other words, a pointwise bounded set of continuous functions either contains a convergent subsequence, or a subsequence whose closure is essentially the same as the example mentioned in the previous paragraphs (the “wildest” possible scenario). Note that in the preceding example, the functions are trivially pointwise bounded in  $\mathbb{R}^X$  as the functions can only take values 0 and 1.

The genesis of Theorem 1.4 was Rosenthal’s  $\ell_1$  theorem, which states that the only reason why Banach space can fail to have an isomorphic copy of  $\ell_1$  (the space of absolutely summable sequences) is the presence of a bounded sequence with no weakly Cauchy subsequence. The theorem is famous for connecting diverse areas of mathematics: Banach space geometry, Ramsey theory, set theory, and topology of function spaces.

As we transition from  $C_p(X)$  to the larger space  $B_1(X)$ , we find a similar dichotomy. Either every point of closure of the set of functions will be a Baire class 1 function, or there is a sequence inside the set that behaves in the wildest possible way. The theorem is usually not phrased as a dichotomy but rather as an equivalence:

**Theorem 1.5** (The BFT Dichotomy. Bourgain-Fremlin-Talagrand, [BFT78, Theorem 4G]). *Let  $X$  be a Polish space and  $A \subseteq C_p(X)$  be pointwise bounded. The following are equivalent:*

- (i)  $A$  is relatively compact in  $B_1(X)$ , i.e.,  $\overline{A} \subseteq B_1(X)$ .
- (ii) For every  $\{f_n\}_{n \in \mathbb{N}} \subseteq A$  and every  $a < b$  there is  $I \subseteq \mathbb{N}$  such that

$$\bigcap_{n \in I} f_n^{-1}(-\infty, a] \cap \bigcap_{n \notin I} f_n^{-1}[b, \infty) = \emptyset.$$

**Definition 1.6.** We shall say that a set  $A \subseteq \mathbb{R}^X$  has the *Independence Property*, or IP for short, if it satisfies the following condition: There exists every  $\{f_n\}_{n \in \mathbb{N}} \subseteq A$  and  $a < b$  such that for every pair of disjoint sets  $E, F \subseteq \mathbb{N}$ , we have

$$\bigcap_{n \in E} f_n^{-1}(-\infty, a] \cap \bigcap_{n \in F} f_n^{-1}[b, \infty) \neq \emptyset.$$

If  $A$  satisfies the negation of this condition, we will say that  $A$  *satisfies NIP*, or that has the NIP.

*Remark 1.7.* Note that if  $X$  is compact and  $A \subseteq C_p(X)$ , then  $A$  has the NIP if and only if for every  $\{f_n\}_{n \in \mathbb{N}} \subseteq A$  and for every  $a < b$  there is  $I \subseteq \mathbb{N}$  such that

$$\bigcap_{n \in I} f_n^{-1}(-\infty, a] \cap \bigcap_{n \notin I} f_n^{-1}[b, \infty) = \emptyset.$$

To summarize, the particular case of Theorem 1.8 when for  $X$  compact can be stated in the following way:

**Theorem 1.8.** *Let  $X$  be a compact Polish space. Then, for every pointwise bounded  $A \subseteq C_p(X)$ , one and exactly one of the following two conditions must hold:*

- (i)  $\overline{A} \subseteq B_1(X)$ .
- (ii)  $A$  has NIP.

The Independence Property was first isolated by Saharon Shelah in model theory as a dividing line between theories whose models are “tame” (corresponding to NIP) theories of models are “wild” (corresponding to IP). See [She71, Definition 4.1], [She90].

254 **1.2. NIP as universal dividing line between polynomial and exponential**  
 255 **complexity.** The particular case of the BSF Dichotomy (Theorem 1.8) when  $A$   
 256 consists of  $\{0, 1\}$ -valued (i.e.,  $\{\text{Yes}, \text{No}\}$ -valued) strings was discovered indepen-  
 257 dently, around 1971-1972 in many foundational contexts related to polynomial  
 258 (“tame”) vs exponential (“wild”) complexity: In model theory, by Saharon She-  
 259 lah [She71],[She90], in combinatorics, by Norbert Sauer [Sau72], and Shelah [She72,  
 260 She90], and in statistical learning, by Vladimir Vapnik and Alexey Chervonenkis [VC71,  
 261 VC74].

262 **In model theory:** Shelah’s classification theory is a foundational program  
 263 in mathematical logic devised to categorize first-order theories based on  
 264 the complexity and structure of their models. A theory  $T$  is considered  
 265 classifiable in Shelah’s sense if the number of non-isomorphic models of  $T$   
 266 of a given cardinality can be described by a bounded number of numerical  
 267 invariants. In contrast, a theory  $T$  is unclassifiable if the number of models  
 268 of  $T$  of a given cardinality is the maximum possible number. This number  
 269 is directly impacted by the number of “types” over of parameters in models  
 270 of  $T$ ; a controlled number of types is a characteristic of a classifiable theory.

271 In Shelah’s classification program [She90], theories without the indepen-  
 272 dence property (called NIP theories, or dependent theories) have a well-  
 273 behaved, “tame” structure; the number of types over a set of parameters  
 274 of size  $\kappa$  of such a theory is of polynomially or similar “slow” growth on  $\kappa$ .  
 275 Theories with the Independence Property (called IP theories), in contrast,  
 276 are considered “intractable” or “wild”. A theory with the independence  
 277 property produces the maximum possible number of types over a set of  
 278 parameters; for a set of parameters of cardinality  $\kappa$ , the theory will have  
 279  $2^{2^\kappa}$ -many distinct types.

280 **In combinatorics:** Sauer [Sau72] and Shelah [She72] proved the following:  
 281 If  $\mathcal{F} = \{S_0, S_1, \dots\}$  is a family of subsets of some infinite set  $S$ , then  
 282 either for every  $n \in \mathbb{N}$ , there is a set  $A \subseteq S$  with  $|A| = n$  such that  
 283  $|\{S_i \cap A : i \in \mathbb{N}\}| = 2^n$  (yielding exponential complexity), or there exists  
 284  $N \in \mathbb{N}$  such that  $A \subseteq S$  with  $|A| \geq N$ , one has

$$|\{S_i \cap A : i \in \mathbb{N}\}| \leq \sum_{i=0}^{N-1} \binom{|A|}{i} \approx O(|A|^N)$$

285 for every  $A \subseteq S$  such that  $|A| \geq N$  (yielding polynomial complexity). This  
 286 answered a question of Erdős.

287 **In machine learning:** Readers familiar with statistical learning may recog-  
 288 nize the Sauer-Shelah lemma as the dichotomy discovered and proved  
 289 slightly earlier (1971) by Vapnik and Chervonenkis [VC71, VC74] to ad-  
 290 dress the problem of uniform convergence in statistics. The least integer  
 291  $N$  given by the preceding paragraph, when it exists, is called the *VC-*  
 292 *dimension* of  $\mathcal{F}$ . This is a core concept in machine learning. If such an  
 293 integer  $N$  does not exist, we say that the VC-dimension of  $\mathcal{F}$  is infinite. The  
 294 lemma provides upper bounds on the number of data points (sample size  $m$ )  
 295 needed to learn a concept class with VC dimension  $d \in \mathbb{N}$  by showing this  
 296 number grows polynomially with  $m$  and  $d$  (namely,  $\sum_{i=0}^d \binom{m}{i} \approx O(m^d)$ ),  
 297 not exponentially. The Fundamental Theorem of Statistical Learning states

that a hypothesis class is PAC-learnable (PAC stands for “Probably Approximately Correct”) if and only if its VC dimension is finite.

**1.3. Rosenthal compacta.** The comprehensiveness of Theorem 1.8, attested by the examples outlined in the preceding section, led to the following definition (isolated by Godefroy [God80]):

**Definition 1.9.** A Rosenthal compactum is a compact Hausdorff topological space  $K$  that can be topologically embedded as a compact subset into the space of all functions of the first Baire class on some Polish space  $X$ , equipped with the topology of pointwise convergence.

Rosenthal compacta are characterized by significant topological and dynamical tameness properties. They play a significant role in functional analysis, measure theory, dynamical systems, descriptive set theory, and model theory. In this paper, we introduce their applicability in deep computation. For this, we shall first focus on countable languages, which is the theme of the next section.

**1.4. The special case  $B_1(X, \mathbb{R}^{\mathcal{P}})$  with  $\mathcal{P}$  countable.** Our goal now is to characterize relatively compact subsets of  $B_1(X, Y)$  for the particular case when  $Y = \mathbb{R}^{\mathcal{P}}$  with  $\mathcal{P}$  countable. Given  $P \in \mathcal{P}$  we denote the projection map onto the  $P$ -coordinate by  $\pi_P : \mathbb{R}^{\mathcal{P}} \rightarrow \mathbb{R}$ . From a high-level topological interpretation, the subsequent lemma states that, in this context, the spaces  $\mathbb{R}$  and  $\mathbb{R}^{\mathcal{P}}$  are really not that different, and that if we understand the Baire class 1 functions of one space, then we also understand the functions of both.

**Lemma 1.10.** *Let  $X$  be a Polish space and  $\mathcal{P}$  be a countable set. Then,  $f \in B_1(X, \mathbb{R}^{\mathcal{P}})$  if and only if  $\pi_P \circ f \in B_1(X)$  for all  $P \in \mathcal{P}$ .*

*Proof.* Only one implication needs a proof. Suppose that  $\pi_P \circ f \in B_1(X)$  for all  $P \in \mathcal{P}$ . Let  $V$  be a basic open subset of  $\mathbb{R}^{\mathcal{P}}$ . That is, there exists a finite  $\mathcal{P}' \subseteq \mathcal{P}$  such that  $V = \bigcap_{P \in \mathcal{P}'} \pi_P^{-1}[U_P]$  where  $U_P$  is open in  $\mathbb{R}$ . Then,

$$f^{-1}[V] = \bigcap_{P \in \mathcal{P}'} (\pi_P \circ f)^{-1}[U_P]$$

is an  $F_\sigma$  set. Since  $\mathcal{P}$  is countable,  $\mathbb{R}^{\mathcal{P}}$  is second countable so every open set  $U$  in  $\mathbb{R}^{\mathcal{P}}$  is a countable union of basic open sets. Hence,  $f^{-1}[U]$  is  $F_\sigma$ .  $\square$

Below we consider  $\mathcal{P}$  with the discrete topology. For each  $f : X \rightarrow \mathbb{R}^{\mathcal{P}}$  denote  $\hat{f}(P, x) := \pi_P \circ f(x)$  for all  $(P, x) \in \mathcal{P} \times X$ . Similarly, for each  $g : \mathcal{P} \times X \rightarrow \mathbb{R}$  denote  $\check{g}(x)(P) := g(P, x)$ . Given  $A \subseteq (\mathbb{R}^{\mathcal{P}})^X$ , we denote  $\hat{A}$  as the set of all  $\hat{f}$  such that  $f \in A$ . Note that the map  $(\mathbb{R}^{\mathcal{P}})^X \rightarrow \mathbb{R}^{\mathcal{P} \times X}$  given by  $f \mapsto \hat{f}$  is a homeomorphism and its inverse is given by  $g \mapsto \check{g}$ .

**Lemma 1.11.** *Let  $X$  be a Polish space and  $\mathcal{P}$  be countable. Then,  $f \in B_1(X, \mathbb{R}^{\mathcal{P}})$  if and only if  $\hat{f} \in B_1(\mathcal{P} \times X)$ .*

*Proof.*  $(\Rightarrow)$  By Lemma 1.10, given an open set of reals  $U$ , we have  $f^{-1}[\pi_P^{-1}[U]]$  is  $F_\sigma$  for every  $P \in \mathcal{P}$ . Given that  $\mathcal{P}$  is a discrete countable space, we observe that

$$\hat{f}^{-1}[U] = \bigcup_{P \in \mathcal{P}} (\{P\} \times f^{-1}[\pi_P^{-1}[U]])$$

is an  $F_\sigma$  as well.



( $\Leftarrow$ ) By lemma 1.10 it suffices to show that  $\pi_P \circ f \in B_1(X)$  for all  $P \in \mathcal{P}$ . Fix an open  $U \subseteq \mathbb{R}$ . Write  $\hat{f}^{-1}[U] = \bigcup_{n \in \mathbb{N}} F_n$  where  $F_n$  is closed in  $\mathcal{P} \times X$ . Then,

$$(\pi_P \circ f)^{-1}[U] = \bigcup_{n \in \mathbb{N}} \{x \in X : (P, x) \in F_n\}$$

331 which is  $F_\sigma$ . □

332 Given  $A \subseteq Y^X$  and  $K \subseteq X$  we write  $A|_K := \{f|_K : f \in A\}$ , i.e., the set of  
 333 all restrictions of functions in  $A$  to  $K$ . The following Theorem is a slightly more  
 334 general version of Theorem 1.8.

335 **Theorem 1.12.** *Assume that  $\mathcal{P}$  is countable,  $X$  is a Polish space, and  $A \subseteq$   
 336  $C_p(X, \mathbb{R}^\mathcal{P})$  is such that  $\pi_P \circ A$  is pointwise bounded for all  $P \in \mathcal{P}$ . The follow-  
 337 ing are equivalent for every compact  $K \subseteq X$ :*

- 338 (1)  $\overline{A|_K} \subseteq B_1(K, \mathbb{R}^\mathcal{P})$ .  
 339 (2)  $\pi_P \circ A|_K$  has the NIP for every  $P \in \mathcal{P}$ .

*Proof.* (1) $\Rightarrow$ (2). Let  $P \in \mathcal{P}$ . Fix  $\{f_n\}_{n \in \mathbb{N}} \subseteq A$  and  $a < b$ . By (1), we have  $\overline{A|_K} \subseteq B_1(K, \mathbb{R}^\mathcal{P})$ . Applying the homeomorphism  $f \mapsto \hat{f}$  and using lemma 1.11 we get  $\hat{A}|_{\mathcal{P} \times K} \subseteq B_1(\mathcal{P} \times K)$ . By Theorem 1.8, there is  $I \subseteq \mathbb{N}$  such that

$$(\mathcal{P} \times K) \cap \bigcap_{n \in I} \hat{f}_n^{-1}(-\infty, a] \cap \bigcap_{n \notin I} \hat{f}_n^{-1}[b, \infty) = \emptyset$$

Hence,

$$K \cap \bigcap_{n \in I} (\pi_P \circ f_n)^{-1}(-\infty, a] \cap \bigcap_{n \notin I} (\pi_P \circ f_n)^{-1}[b, \infty) = \emptyset$$

By the compactness of  $K$ , there are finite  $E \subseteq I$  and  $F \subseteq \mathbb{N} \setminus I$  such that

$$K \cap \bigcap_{n \in E} (\pi_P \circ f_n)^{-1}(-\infty, a] \cap \bigcap_{n \in F} (\pi_P \circ f_n)^{-1}[b, \infty) = \emptyset$$

340 Thus,  $\pi_P \circ A|_L$  has the NIP.

341 (2) $\Rightarrow$ (1) Fix  $f \in \overline{A|_K}$ . By lemma 1.10 it suffices to show that  $\pi_P \circ f \in B_1(K)$   
 342 for all  $P \in \mathcal{P}$ . By (2),  $\pi_P \circ A|_K$  has the NIP. Hence, by Theorem 1.8 we have  
 343  $\overline{\pi_P \circ A|_K} \subseteq B_1(K)$ . But then  $\pi_P \circ f \in \overline{\pi_P \circ A|_K} \subseteq B_1(K)$ . □

344 Lastly, a simple but significant result that helps understand the operation of  
 345 restricting a set of functions to a specific subspace of the domain space  $X$ , of course  
 346 in the context of the NIP, is that we may always assume that said subspace is  
 347 closed. Concretely, whether we take its closure or not has no effect on the NIP:

348 **Lemma 1.13.** *Assume that  $X$  is Hausdorff and that  $A \subseteq C_p(X)$ . The following  
 349 are equivalent for every  $L \subseteq X$ :*

- 350 (i)  $A_L$  has the NIP.  
 351 (ii)  $A|_{\overline{L}}$  has the NIP.

*Proof.* It suffices to show that (i) $\Rightarrow$ (ii). Suppose that (ii) does not hold, i.e., that there are  $\{f_n\}_{n \in \mathbb{N}} \subseteq A$  and  $a < b$  such that for all finite disjoint  $E, F \subseteq \mathbb{N}$ :

$$\overline{L} \cap \bigcap_{n \in E} f_n^{-1}(-\infty, a] \cap \bigcap_{n \in F} f_n^{-1}[b, \infty) \neq \emptyset.$$

Pick  $a' < b'$  such that  $a < a' < b' < b$ . Then, for any finite disjoint  $E, F \subseteq \mathbb{N}$  we can choose

$$x \in \bar{L} \cap \bigcap_{n \in E} f_n^{-1}(-\infty, a') \cap \bigcap_{n \in F} f_n^{-1}(b', \infty)$$

By definition of closure:

$$L \cap \bigcap_{n \in E} f_n^{-1}(-\infty, a'] \cap \bigcap_{n \in F} f_n^{-1}[b', \infty) \neq \emptyset.$$

352 This contradicts (i). □

## 353 2. COMPOSITIONAL COMPUTATION STRUCTURES.

354 In this section, we connect function spaces with floating point computation. We  
355 start by summarizing some basic concepts from [ADIW24].

356 A *computation states structure* is a pair  $(L, \mathcal{P})$ , where  $L$  is a set whose elements we  
357 call *states* and  $\mathcal{P}$  is a collection of real-valued functions on  $L$  that we call *predicates*.  
358 For a state  $v \in L$ , *type* of  $v$  is the indexed family

$$\text{tp}(v) = (P(v))_{P \in \mathcal{P}} \in \mathbb{R}^{\mathcal{P}}.$$

359 For each  $P \in \mathcal{P}$ , we call real value  $P(v)$  the  $P$ -th *feature* of  $v$ . A *transition* of a  
360 computation states structure  $(L, \mathcal{P})$  is a map  $f : L \rightarrow L$ .

361 Intuitively,  $L$  is the set of states of a computation, and the predicates  $P \in \mathcal{P}$   
362 are primitives that are given and accepted as computational. We think of each  
363 state  $v \in L$  as being uniquely characterized by its type  $\text{tp}(v)$ ; thus, in practice, we  
364 identify  $L$  with a subset of  $\mathbb{R}^{\mathcal{P}}$ . A typical case will be when  $L = \mathbb{R}^{\mathbb{N}}$  or  $L = \mathbb{R}^n$   
365 for some positive integer  $n$  and there is a predicate  $P_i(v) = v_i$  for each of the  
366 coordinates  $v_i$  of  $v$ . We regard the space of types as a topological space, endowed  
367 with the topology of pointwise convergence inherited from  $\mathbb{R}^{\mathcal{P}}$ . In particular, for  
368 each  $P \in \mathcal{P}$ , the projection map  $v \mapsto P(v)$  is continuous.

369 **Definition 2.1.** Given a computation states structure  $(L, \mathcal{P})$ , any element of  $\mathbb{R}^{\mathcal{P}}$   
370 in the image of  $L$  under the map  $v \mapsto \text{tp}(v)$  will be called a *realized type*. The  
371 topological closure of the set of realized types in  $\mathbb{R}^{\mathcal{P}}$  (endowed with the point-  
372 wise convergence topology) will be called the *space of types* of  $(L, \mathcal{P})$ , denoted  $\mathcal{L}$ .  
373 Elements of  $\mathcal{L} \setminus L$  will be called *unrealized types*.

374 In traditional model theory, the space of types of a structure is viewed as a sort of  
375 compactification of the structure. However, the space  $\mathcal{L}$  is not necessarily compact.  
376 To bypass this obstacle, we follow the idea introduced in [ADIW24] of covering  $\mathcal{L}$   
377 by “thin” compact subspaces that we call *shards*. The formal definition of shard is  
378 next.

379 **Definition 2.2.** A *sizer* is a tuple  $r_{\bullet} = (r_P)_{P \in \mathcal{P}}$  of positive real numbers indexed  
380 by  $\mathcal{P}$ . Given a sizer  $r_{\bullet}$ , we define the  $r_{\bullet}$ -*shard* as:

$$L[r_{\bullet}] = L \cap \prod_{P \in \mathcal{P}} [-r_P, r_P].$$

381 For a sizer  $r_{\bullet}$ , the  $r_{\bullet}$ -*type shard* is defined as  $\mathcal{L}[r_{\bullet}] = \overline{L[r_{\bullet}]}$ . We define  $\mathcal{L}_{sh}$ , as  
382 the union of all type-shards.

383 **Definition 2.3.** A *Compositional Computation Structure* (CCS) is a triple  $(L, \mathcal{P}, \Gamma)$ ,  
384 where

- $(L, \mathcal{P})$  is a computation states structure
- $\Gamma \subseteq L^L$  is a semigroup under composition.

The elements of the  $\Gamma$  is called the *computations* of the structure  $(L, \mathcal{P}, \Gamma)$ .

If  $\Delta \subseteq \Gamma$ , we say that  $\Delta \subseteq \Gamma$  is *R-confined* if  $\gamma|_{L[r_\bullet]} : L[r_\bullet] \rightarrow L[r_\bullet]$  for every  $r_\bullet \in R$  and  $\gamma \in \Delta$ . Elements in  $\bar{\Delta} \subseteq \mathcal{L}_{sh}$  are called (real-valued) *deep computations* or *ultracomputations*.

A tenet of our approach is that a map  $f : L \rightarrow \mathcal{L}$  is to be considered “effectively computable” if, for each  $Q \in \mathcal{P}$ , the output feature  $Q \circ f : L \rightarrow \mathbb{R}$  is a *definable* predicate in the following sense:

Given any arbitrary  $\varepsilon > 0$  and any  $K \subseteq L$  wherein every input feature  $P(v)$  remains bounded in magnitude there is an  $\varepsilon$ -approximating continuous “algebraic” operator  $\varphi(P_1, \dots, P_n)$  of finitely many input features  $P_1, \dots, P_n$ , such that the following holds: for all  $v \in K$ , the output feature  $Q(f(v))$  is  $\varepsilon$ -approximated by  $\varphi(P_1(v), \dots, P_n(v))$ . By “algebraic”, we mean that the approximating operator  $\varphi(P_1, \dots, P_n)$  uses only, in addition to the primitives  $P_1, \dots, P_n$ , the algebra operations of  $\mathbb{R}^{\mathcal{P}}$ , i.e., vector addition, vector multiplication, and scalar addition.

It is shown in [ADIW24]) that:

- (1) For a definable  $f : L \rightarrow \mathcal{L}$ , the approximating operators  $\varphi$  may be taken to be *polynomials* of the input features, and
- (2) Definable transforms  $f : L \rightarrow \mathcal{L}$  are precisely those that extend to continuous  $\tilde{f} : \mathcal{L} \rightarrow \mathcal{L}$  (this is the property of *extendibility* mentioned above).

This motivates the following definition.

**Definition 2.4.** We say that a CCS  $(L, \mathcal{P}, \Gamma)$  satisfies the *Extendability Axiom* if for all  $\gamma \in \Gamma$ , there is  $\tilde{\gamma} : \mathcal{L}_{sh} \rightarrow \mathcal{L}_{sh}$  such that for every sizer  $r_\bullet$  there is a sizer  $s_\bullet$  such that  $\tilde{\gamma}|_{\mathcal{L}[r_\bullet]} : \mathcal{L}[r_\bullet] \rightarrow \mathcal{L}[s_\bullet]$  is continuous. We refer to  $\tilde{\gamma}$  as a *free* extension of  $\gamma$ .

By the preceding remarks, the Extendability Axiom says that the elements of the semigroup  $\Gamma$  are definable.

For an illustrative example, we can frame Newton’s polynomial root approximation method in the context of a CCS (see Example 5.6 of [ADIW24] for details) as follows. Begin by considering the extended complex numbers  $\hat{\mathbb{C}} := \mathbb{C} \cup \{\infty\}$  with the usual Riemann sphere topology that makes it into a compact space (where unbounded sequences converge to  $\infty$ ). In fact, not only is this space compact but it is covered by the shard given by the sizer  $(1, 1, 1)$  (the unit sphere is contained in the cube  $[-1, 1]^3$ ). The space  $\hat{\mathbb{C}}$  is homeomorphic to the usual unit sphere  $S^2 := \{(x, y, z) : x^2 + y^2 + z^2 = 1\}$  of  $\mathbb{R}^3$ , by means of the stereographic projection and its inverse  $\hat{\mathbb{C}} \rightarrow S^2$ . This function is regarded as a triple of predicates  $x, y, z : \hat{\mathbb{C}} \rightarrow [-1, 1]$  where each will map an extended complex number to its corresponding real coordinate on the cube  $[-1, 1]^3$ . Now fix the cubic complex polynomial  $p(s) := s^3 - 1$ , and consider the map which performs one step in Newton’s method at a particular (extended) complex number  $s$ , for finding a root of  $p$ ,  $\gamma_p : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$ . The explicit inner workings of  $\gamma_p$  are irrelevant for this example, except for the fact that it is a continuous mapping. It follows that  $(S^2, \{x, y, z\}, \{\gamma_p^k : k \in \mathbb{N}\})$  is a CCS. The idea is that repeated applications of  $\gamma_p(s), \gamma_p \circ \gamma_p(s), \gamma_p \circ \gamma_p \circ \gamma_p(s), \dots$  would approximate a root of  $p$  provided  $s$  was a good enough initial guess.

431 For a deeper discussion about this axiom, we refer the reader to [ADIW24].

### 432 3. CLASSIFYING DEEP COMPUTATIONS

433 **3.1. NIP, Rosenthal compacta, and deep computations.** Under what condi-  
 434 tions are deep computations Baire class 1, and thus well-behaved according to our  
 435 framework, on type-shards? The following Theorem says that, under the assump-  
 436 tion that  $\mathcal{P}$  is countable, the space of deep computations is a Rosenthal compactum  
 437 (when restricted to shards) if and only if the set of computations has the NIP,  
 438 feature by feature. Hence, we can import the theory of Rosenthal compacta into  
 439 this framework of deep computations.

440 **Theorem 3.1.** *Let  $(L, \mathcal{P}, \Gamma)$  be a compositional computational structure (Defini-*  
 441 *tion 2.3) satisfying the Extendability Axiom (Definition 2.4) with  $\mathcal{P}$  countable. Let*  
 442  *$R$  be an exhaustive collection of sizers. Let  $\Delta \subseteq \Gamma$  be  $R$ -confined. The following*  
 443 *are equivalent.*

- 444 (1)  $\tilde{\Delta}|_{\mathcal{L}[r_\bullet]} \subseteq B_1(\mathcal{L}[r_\bullet], \mathcal{L}[r_\bullet])$  for all  $r_\bullet \in R$ .  
 (2)  $\pi_P \circ \Delta|_{\mathcal{L}[r_\bullet]}$  has the NIP for all  $P \in \mathcal{P}$  and  $r_\bullet \in R$ ; that is, for all  $P \in \mathcal{P}$ ,  
 $r_\bullet \in R$ ,  $a < b$ ,  $\{\gamma_n\}_{n \in \mathbb{N}} \subseteq \Delta$  there are finite disjoint  $E, F \subseteq \mathbb{N}$  such that

$$L[r_\bullet] \cap \bigcap_{n \in E} (\pi_P \circ \gamma_n)^{-1}(-\infty, a] \cap \bigcap_{n \in F} (\pi_P \circ \gamma_n)^{-1}[b, \infty) = \emptyset.$$

445 Moreover, if any (hence all) of the preceding conditions hold, then every deep  
 446 computation  $f \in \tilde{\Delta}$  can be extended to a Baire-1 function on shards, i.e., there is  
 447  $\tilde{f} : \mathcal{L}_{sh} \rightarrow \mathcal{L}_{sh}$  such that  $\tilde{f}|_{\mathcal{L}[r_\bullet]} \in B_1(\mathcal{L}[r_\bullet], \mathcal{L}[r_\bullet])$  for all  $r_\bullet \in R$ . In particular, on  
 448 each shard every deep computation is the pointwise limit of a countable sequence of  
 449 computations.

450 *Proof.* Since  $\mathcal{P}$  is countable,  $\mathcal{L}[r_\bullet] \subseteq \mathbb{R}^{\mathcal{P}}$  is Polish. Also, the Extendability Axiom  
 451 implies that  $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$  is a pointwise bounded set of continuous functions for all  
 452  $P \in \mathcal{P}$ . Hence, Theorem 1.12 and Lemma 1.13 prove the equivalence of (1) and (2).  
 453 If (1) holds and  $f \in \tilde{\Delta}$ , then write  $f = \mathcal{U}\lim_i \gamma_i$  as an ultralimit. Define  $\tilde{f} := \mathcal{U}\lim_i \tilde{\gamma}_i$ .  
 454 Hence, for all  $r_\bullet \in R$  we have  $\tilde{f}|_{\mathcal{L}[r_\bullet]} \in \tilde{\Delta}|_{\mathcal{L}[r_\bullet]} \subseteq B_1(\mathcal{L}[r_\bullet], \mathcal{L}[r_\bullet])$ . That every  
 455 deep computation is a pointwise limit of a countable sequence of computations  
 456 follows from the fact that for a Polish space  $X$  every compact subset of  $B_1(X)$  is  
 457 Fréchet-Urysohn (that is, a space where topological closures coincide with sequential  
 458 closures, see Theorem 3F in [BFT78] or Theorem 4.1 in [Deb13]).  $\square$

459 **3.2. The Todorćević trichotomy and levels of PAC learnability.** Given a  
 460 countable set  $\Delta$  of computations satisfying the NIP on features and shards (con-  
 461 dition (2) of Theorem 3.1) we have that  $\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$  (for a fixed sizer  $r_\bullet$ ) is a separable  
 462 Rosenthal compactum (compact subset of  $B_1(P \times \mathcal{L}[r_\bullet])$ ). The work of Todorćević  
 463 ([Tod99]) and Argyros, Dodos, Kanellopoulos ([ADK08]) culminates in a trichotomy  
 464 theorem for separable Rosenthal Compacta. In this section, inspired by the work  
 465 of Glasner and Megrelishvili ([GM22]), we study ways in which this classification  
 466 allows us obtain different levels of PAC-learnability (NIP).

467 Recall that a topological space  $X$  is *hereditarily separable* (HS) if every subspace  
 468 is separable and that  $X$  is *first countable* if every point in  $X$  has a countable local  
 469 basis. Every separable metrizable space is hereditarily separable and it is a result

of R. Pol that every hereditarily separable Rosenthal compactum is first countable (see section 10 of [Deb13]). This suggests the following definition:

**Definition 3.2.** Let  $(L, \mathcal{P}, \Gamma)$  be a CCS satisfying the Extendability Axiom and  $R$  be an exhaustive collection of sizers. Let  $\Delta \subseteq \Gamma$  be an  $R$ -confined countable set of computations satisfying the NIP on shards and features (condition (2) in Theorem 3.1). We say that  $\Delta$  is:

- (i) NIP<sub>1</sub> if  $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$  is first countable for every  $r_\bullet \in R$ .
- (ii) NIP<sub>2</sub> if  $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$  is hereditarily separable for every  $r_\bullet \in R$ .
- (iii) NIP<sub>3</sub> if  $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$  is metrizable for every  $r_\bullet \in R$ .

Observe that  $\text{NIP}_3 \Rightarrow \text{NIP}_2 \Rightarrow \text{NIP}_1 \Rightarrow \text{NIP}$ . A natural question that would continue this work is to find examples of CCS that separate these levels of NIP. In [Tod99], Todorćević isolates three canonical examples of Rosenthal compacta that witness the failure of the converse implications above.

We now present some separable and non-separable examples of Rosenthal compacta:

### Examples 3.3.

- (1) *Alexandroff compactification of a discrete space of size continuum.* For each  $a \in 2^{\mathbb{N}}$  consider the map  $\delta_a : 2^{\mathbb{N}} \rightarrow \mathbb{R}$  given by  $\delta_a(x) = 1$  if  $x = a$  and  $\delta_a(x) = 0$  otherwise. Let  $A(2^{\mathbb{N}}) = \{\delta_a : a \in 2^{\mathbb{N}}\} \cup \{0\}$ , where 0 is the zero map. Notice that  $A(2^{\mathbb{N}})$  is a compact subset of  $B_1(2^{\mathbb{N}})$ , in fact  $\{\delta_a : a \in 2^{\mathbb{N}}\}$  is a discrete subspace of  $B_1(2^{\mathbb{N}})$  and its pointwise closure is precisely  $A(2^{\mathbb{N}})$ . Hence, this is a Rosenthal compactum which is not first countable. Notice that this space is also not separable.
- (2) *Extended Alexandroff compactification.* For each finite binary sequence  $s \in 2^{<\mathbb{N}}$ , let  $v_s : 2^{\mathbb{N}} \rightarrow \mathbb{R}$  be given by  $v_s(x) = 1$  if  $x$  extends  $s$  and  $v_s(x) = 0$  otherwise. Let  $\hat{A}(2^{\mathbb{N}})$  be the pointwise closure of  $\{v_s : s \in 2^{<\mathbb{N}}\}$ , i.e.,  $\hat{A}(2^{\mathbb{N}}) = A(2^{\mathbb{N}}) \cup \{v_s : s \in 2^{<\mathbb{N}}\}$ . Note that this space is a separable Rosenthal compactum which is not first countable.
- (3) *Split Cantor.* Let  $<$  be the lexicographic order in the space of infinite binary sequences, i.e.,  $2^{\mathbb{N}}$ . For each  $a \in 2^{\mathbb{N}}$  let  $f_a^- : 2^{\mathbb{N}} \rightarrow \mathbb{R}$  be given by  $f_a^-(x) = 1$  if  $x < a$  and  $f_a^-(x) = 0$  otherwise. Let  $f_a^+ : 2^{\mathbb{N}} \rightarrow \mathbb{R}$  be given by  $f_a^+(x) = 1$  if  $x \leq a$  and  $f_a^+(x) = 0$  otherwise. The split Cantor is the space  $S(2^{\mathbb{N}}) = \{f_a^- : a \in 2^{\mathbb{N}}\} \cup \{f_a^+ : a \in 2^{\mathbb{N}}\}$ . This is a separable Rosenthal compactum. One example of a countable dense subset is the set of all  $f_a^+$  and  $f_a^-$  where  $a$  is an infinite binary sequence that is eventually constant. Moreover, it is hereditarily separable but it is not metrizable.
- (4) *Alexandroff Duplicate.* Let  $K$  be any compact metric space and consider the Polish space  $X = C(K) \sqcup K$ , i.e., the disjoint union of  $C(K)$  (with its supremum norm topology) and  $K$ . For each  $a \in K$  define  $g_a^0, g_a^1 : X \rightarrow \mathbb{R}$  as follows:

$$g_a^0(x) = \begin{cases} x(a), & x \in C(K) \\ 0, & x \in K \end{cases}$$

$$g_a^1(x) = \begin{cases} x(a), & x \in C(K) \\ \delta_a(x), & x \in K \end{cases}$$

Let  $D(K) = \{g_a^0 : a \in K\} \cup \{g_a^1 : a \in K\}$ . Notice that  $D(K)$  is a first countable Rosenthal compactum. It is not separable if  $K$  is uncountable. The interesting case will be when  $K = 2^{\mathbb{N}}$ .

- (5) *Extended Alexandroff Duplicate of the split Cantor*. For each finite binary sequence  $t \in 2^{<\mathbb{N}}$  let  $a_t \in 2^{\mathbb{N}}$  be the sequence starting with  $t$  and ending with 0's and let  $b_t \in 2^{\mathbb{N}}$  be the sequence starting with  $t$  and ending with 1's. Define  $h_t : 2^{\mathbb{N}} \rightarrow \mathbb{R}$  by


$$h_t(x) = \begin{cases} 0, & x < a_t \\ 1/2, & a_t \leq x \leq b_t \\ 1, & b_t < x \end{cases}$$

Let  $\hat{D}(S(2^{\mathbb{N}}))$  be the pointwise closure of the set  $\{h_t : t \in 2^{<\mathbb{N}}\}$ . Hence,  $\hat{D}(S(2^{\mathbb{N}}))$  is a separable first countable Rosenthal compactum which is not hereditarily separable. In fact, it contains an uncountable discrete subspace (see Theorem 5 in [Tod99]).

**Theorem 3.4** (Todorčević's Trichotomy, [Tod99], Theorem 3 in [ADK08]). *Let  $K$  be a separable Rosenthal Compactum.*

- (i) *If  $K$  is hereditarily separable but non-metrizable, then  $S(2^{\mathbb{N}})$  embeds into  $K$ .*
- (ii) *If  $K$  is first countable but not hereditarily separable, then either  $D(2^{\mathbb{N}})$  or  $\hat{D}(S(2^{\mathbb{N}}))$  embeds into  $K$ .*
- (iii) *If  $K$  is not first countable, then  $A(2^{\mathbb{N}})$  embeds into  $K$ .*

We thus have the following classification:



The definitions provided here for  $NIP_i$  ( $i = 1, 2, 3$ ) are topological. This raises the following question:

**Question 3.5.** Is there a non-topological characterization for  $NIP_i$ ,  $i = 1, 2, 3$ ?

**3.3. The Argyros-Dodos-Kanellopoulos heptachotomy, and approximability of deep computation by minimal classes.** In the three separable three cases given in 3.3, namely,  $(\hat{A}(2^{\mathbb{N}}), S(2^{\mathbb{N}})$  and  $\hat{D}(S(2^{\mathbb{N}}))$ ), the countable dense subsets are indexed by the binary tree  $2^{<\mathbb{N}}$ . This choice of index is useful for two reasons:

- 529 (1) Our emphasis is computational. Elements of  $2^{<\mathbb{N}}$  represent finite bitstrings,  
 530 i.e., standard computations, while Rosenthal compacta represent deep com-  
 531 putations, i.e., limits of finite computations. Mathematically, deep compu-  
 532 tations are pointwise limits of standard computations; however, computa-  
 533 tionally, we are interested in the manner (and the efficiency) in which the  
 534 approximations can occur.
- 535 (2) The Ramsey theory of perfect subsets of the Cantor space  $2^{\mathbb{N}}$  can be im-  
 536 ported to analyze the behavior of the accumulation points. Since  $2^{<\mathbb{N}}$  is  
 537 countable, we can always choose this index for the countable dense subsets.  
 538 This is done in [ADK08].

539 **Definition 3.6.** Let  $X$  be a Polish space.

- 540 (1) If  $I$  is a countable and  $\{f_i : i \in I\} \subseteq \mathbb{R}^X$ ,  $\{g_i : i \in I\} \subseteq \mathbb{R}^X$  are two  
 541 pointwise families by  $I$ . We say that  $\{f_i : i \in I\}$  and  $\{g_i : i \in I\}$  are  
 542 *equivalent* if and only if the map  $f_i \mapsto g_i$  is extended to a homeomorphism  
 543 from  $\overline{\{f_i : i \in I\}}$  to  $\overline{\{g_i : i \in I\}}$ .
- 544 (2) If  $\{f_t : t \in 2^{<\mathbb{N}}\}$  is a pointwise bounded family, we say that  $\{f_t : t \in 2^{<\mathbb{N}}\}$   
 545 is *minimal* if and only if for every dyadic subtree  $\{s_t : t \in 2^{<\mathbb{N}}\}$  of  $2^{<\mathbb{N}}$ ,  
 546  $\{f_{s_t} : t \in 2^{<\mathbb{N}}\}$  is equivalent to  $\{f_t : t \in 2^{<\mathbb{N}}\}$ .

547 One of the main results in [ADK08] is that, up to equivalence, there are seven  
 548 minimal families of Rosenthal compacta and that for every relatively compact  $\{f_t : t \in 2^{<\mathbb{N}}\} \subseteq B_1(X)$  there is a dyadic subtree  $\{s_t : t \in 2^{<\mathbb{N}}\}$  such that  $\{f_{s_t} : t \in 2^{<\mathbb{N}}\}$  is equivalent to one of the minimal families. We shall describe the minimal families next. We follow the same notation as in [ADK08]. For any node  $t \in 2^{<\mathbb{N}}$ , let us denote by  $t \frown 0^\infty$  ( $t \frown 1^\infty$ ) the infinite binary sequence starting with  $t$  and continuing with all 0's (respectively, all 1's). Fix a regular dyadic subtree  $R = \{s_t : t \in 2^{<\mathbb{N}}\}$  of  $2^{<\mathbb{N}}$  (i.e., a dyadic subtree such that every level of  $R$  is contained in a level of  $2^{<\mathbb{N}}$ ) with the property that for all  $s, s' \in R$ ,  $s \frown 0^\infty \neq s' \frown 0^\infty$  and  $s \frown 1^\infty \neq s' \frown 1^\infty$ . Given  $t \in 2^{<\mathbb{N}}$ , let  $v_t$  be the characteristic function of the set  $\{x \in 2^{\mathbb{N}} : x \text{ extends } t\}$ . Let  $<$  be the lexicographic order in  $2^{\mathbb{N}}$ . Given  $a \in 2^{\mathbb{N}}$ , let  $f_a^+ : 2^{\mathbb{N}} \rightarrow \{0, 1\}$  be the characteristic function of  $\{x \in 2^{\mathbb{N}} : a \leq x\}$  and let  $f_a^- : 2^{\mathbb{N}} \rightarrow \{0, 1\}$  be the characteristic function of  $\{x \in 2^{\mathbb{N}} : a < x\}$ . Given two maps  $f, g : 2^{\mathbb{N}} \rightarrow \mathbb{R}$  we denote by  $(f, g) : 2^{\mathbb{N}} \sqcup 2^{\mathbb{N}} \rightarrow \mathbb{R}$  the function which is  $f$  on the first copy of  $2^{\mathbb{N}}$  and  $g$  on the second copy of  $2^{\mathbb{N}}$ .

- 562 (1)  $D_1 = \{\frac{1}{|t|+1}v_t : t \in 2^{<\mathbb{N}}\}$ . This is discrete in  $\overline{D_1} = A(2^{\mathbb{N}})$ .
- 563 (2)  $D_2 = \{s_t \frown 0^\infty : t \in 2^{<\mathbb{N}}\}$ . This is discrete in  $\overline{D_2} = 2^{\leq \mathbb{N}}$ .
- 564 (3)  $D_3 = \{f_{s_t \frown 0^\infty}^+ : t \in 2^{<\mathbb{N}}\}$ . This is discrete in  $\overline{D_3} = S(2^{\mathbb{N}})$ .
- 565 (4)  $D_4 = \{f_{s_t \frown 1^\infty}^- : t \in 2^{<\mathbb{N}}\}$ . This is discrete in  $\overline{D_4} = S(2^{\mathbb{N}})$ .
- 566 (5)  $D_5 = \{v_t : t \in 2^{<\mathbb{N}}\}$ . This is discrete in  $\overline{D_5} = \hat{A}(2^{\mathbb{N}})$ .
- 567 (6)  $D_6 = \{(v_{s_t}, s_t \frown 0^\infty) : t \in 2^{<\mathbb{N}}\}$ . This is discrete in  $\overline{D_6} = \hat{D}(2^{\mathbb{N}})$ .
- 568 (7)  $D_7 = \{(v_{s_t}, x_{s_t \frown 0^\infty}^+) : t \in 2^{<\mathbb{N}}\}$ . This is discrete in  $\overline{D_7} = \hat{D}(S(2^{\mathbb{N}}))$

569 **Theorem 3.7** (Heptacotomy of minimal families, Theorem 2 in [ADK08]). *Let*  
 570  *$X$  be Polish. For every relatively compact  $\{f_t : t \in 2^{<\mathbb{N}}\} \subseteq B_1(X)$ , there exists*  
 571  *$i = 1, 2, \dots, 7$  and a regular dyadic subtree  $\{s_t : t \in 2^{<\mathbb{N}}\}$  of  $2^{<\mathbb{N}}$  such that  $\{f_{s_t} :$   
 572  $t \in 2^{<\mathbb{N}}\}$  *is equivalent to  $D_i$ . Moreover, all  $D_i$  are minimal and mutually non-*  
 573 *equivalent.**

3.4. **Bourgain-Fremlin-Talagrand, NIP, and essential computability of deep computations.** We now turn to the question: what happens when  $\mathcal{P}$  is uncountable? Notice that the countability assumption is crucial in the proof of Theorem 1.12 essentially because it makes  $\mathbb{R}^{\mathcal{P}}$  a Polish space. For the uncountable case, we may lose Baire-1 definability so we shall replace  $B_1(X)$  by a larger class. Recall that the *raison d'être* of the class of Baire-1 functions is to have a class that contains the continuous functions but is closed under pointwise limits, and that (Fact 1.2) for perfectly normal  $X$ , a function  $f$  is in  $B_1(X, Y)$  if and only if  $f^{-1}[U]$  is an  $F_\sigma$  subset of  $X$  for every open  $U \subseteq Y$ . This motivates the following definition:

**Definition 3.8.** Given a Hausdorff space  $X$  and a measurable space  $(Y, \Sigma)$ , we say that  $f : X \rightarrow Y$  is *universally measurable* (with respect to  $\Sigma$ ) if  $f^{-1}(E)$  is Borel for every  $E \in \Sigma$ , i.e.,  $f^{-1}(E)$  is  $\mu$ -measurable for every Radon probability measure  $\mu$  on  $X$ . When  $Y = \mathbb{R}$  we will always take  $\Sigma = \mathcal{B}(\mathbb{R})$ , the Borel  $\sigma$ -algebra of  $\mathbb{R}$ . In this case, a function  $f : X \rightarrow \mathbb{R}$  is universally measurable if and only if  $f^{-1}(U)$  is  $\mu$ -measurable for every Radon probability measure  $\mu$  on  $X$  and every open set  $U \subseteq \mathbb{R}$ .

Intuitively, a function is universally measurable if it is “measurable no matter which reasonable way you try to measure things on its domain”. The concept of universal measurability emerged from work of Kallianpur and Sazonov, in the late 1950’s and 1960s, with later developments by Blackwell, Darst, and others, building on earlier ideas of Gnedenko and Kolmogorov from the 1950s. See [Pap02, Chapters 1 and 2].

Following [BFT78], the collection of all universally measurable real-valued functions will be denoted by  $M_r(X)$ . In the context of deep computations, we will be interested in transition maps from a state space  $L \subseteq \mathbb{R}^{\mathcal{P}}$  to itself. There are two natural  $\sigma$ -algebras one can consider in the product space  $\mathbb{R}^{\mathcal{P}}$ : the Borel  $\sigma$ -algebra, i.e., the  $\sigma$ -algebra generated by open sets in  $\mathbb{R}^{\mathcal{P}}$ , and the cylinder  $\sigma$ -algebra, i.e., the  $\sigma$ -algebra generated by the sub-basic open sets in  $\mathbb{R}^{\mathcal{P}}$ . Note that when  $\mathcal{P}$  is countable, both  $\sigma$ -algebras coincide but in general the cylinder  $\sigma$ -algebra is strictly smaller. We will use the cylinder  $\sigma$ -algebra to define universally measurable maps  $f : \mathbb{R}^{\mathcal{P}} \rightarrow \mathbb{R}^{\mathcal{P}}$ . The reason for this choice is the following characterization:

**Lemma 3.9.** *Let  $X$  be a Hausdorff space and  $Y = \prod_{i \in I} Y_i$  be any product of measurable spaces  $(Y_i, \Sigma_i)$  for  $i \in I$ . Let  $\Sigma_Y$  be the cylinder  $\sigma$ -algebra generated by the measurable spaces  $(Y_i, \Sigma_i)$ . Let  $f : X \rightarrow Y$ . The following are equivalent:*

- (i)  $f : X \rightarrow Y$  is universally measurable (with respect to  $\Sigma_Y$ ).
- (ii)  $\pi_i \circ f : X \rightarrow Y_i$  is universally measurable (with respect to  $\Sigma_i$ ) for all  $i \in I$ .

*Proof.* (i) $\Rightarrow$ (ii) is clear since the projection maps  $\pi_i$  are measurable and the composition of measurable functions is measurable. To prove (ii) $\Rightarrow$ (i), suppose that  $C = \prod_{i \in I} C_i$  is a measurable cylinder and let  $J$  be the finite set of  $i \in I$  such that  $C_i \neq Y_i$ . Then,  $C = \bigcap_{i \in J} \pi_i^{-1}(C_i)$ , so  $f^{-1}(C) = \bigcap_{i \in J} (\pi_i \circ f)^{-1}(C_i)$  is a universally measurable set by assumption.  $\square$

The preceding lemma says that a transition map is universally measurable if and only if it is universally measurable on all its features. In other words, we can check measurability of a transition just by checking measurability feature by feature. We will denote by  $M_r(X, \mathbb{R}^{\mathcal{P}})$  the collection of all universally measurable functions



619  $f : X \rightarrow \mathbb{R}^{\mathcal{P}}$  (with respect to the cylinder  $\sigma$ -algebra), endowed with the topology  
 620 of pointwise convergence.

621 We now wish to define the concept of a deep computation being computable  
 622 except a set of arbitrarily small measure “no matter which reasonable way you try  
 623 to measure things on its domain” (see the remarks following definition ). This is  
 624 definition below. To motivate the definition, we need to recall two facts:

- 625 (1) Littlewood’s second principle states that every Lebesgue measurable func-  
 626 tion is “nearly continuous”. The formal version of this, which is Luzin’s  
 627 theorem, states that if  $(X, \Sigma, \mu)$  a Radon measure space and  $Y$  be a second-  
 628 countable topological space (e.g.,  $Y = \mathbb{R}^{\mathcal{P}}$  with  $\mathcal{P}$  countable ) equipped with  
 629 a Borel algebra, then any given  $f : X \rightarrow Y$  is measurable if and only if for  
 630 every  $E \in \Sigma$  and every  $\varepsilon > 0$  there exists a closed  $F \subseteq E$  such that the  
 631 restriction of  $f$  to  $F$  is continuous.
- 632 (2) Computability of deep computations can be characterized in terms of con-  
 633 tinuous extendibility of computations. This is at the core of [ADIW24].

634 These facts motivate the following definition:

635 **Definition 3.10.** Let  $(L, \mathcal{P}, \Gamma)$  be a CCS. We say that a transition  $f : L \rightarrow L$   
 636 is *universally essentially computable* if and only if there exists  $\tilde{f} : \mathcal{L}_{sh} \rightarrow \mathcal{L}_{sh}$   
 637 extending  $f$  such that for every sizer  $r_{\bullet}$  there is a sizer  $s_{\bullet}$  such that the restriction  
 638  $\tilde{f}|_{\mathcal{L}[r_{\bullet}]} : \mathcal{L}[r_{\bullet}] \rightarrow \mathcal{L}[s_{\bullet}]$  is universally measurable, i.e.,  $\pi_P \circ \tilde{f}|_{\mathcal{L}[r_{\bullet}]} : \mathcal{L}[r_{\bullet}] \rightarrow [-s_P, s_P]$   
 639 is  $\mu$ -measurable for every Radon probability measure  $\mu$  on  $\mathcal{L}[r_{\bullet}]$ .

640 For a measure  $\mu$  on  $aX$ , the set of all  $\mu$ -measurable functions will be denoted by  
 641  $\mathcal{M}^0(X, \mu)$ .

642 We will need the following result about NIP and universally measurable func-  
 643 tions:

644 **Theorem 3.11** (Bourgain-Fremlin-Ta set lagrand, Theorem 2F in [BFT78]). *Let*  
 645  *$X$  be a Hausdorff space and  $A \subseteq C(X)$  be pointwise bounded. The following are*  
 646 *equivalent:*

- 647 (i)  $\overline{A} \subseteq M_r(X)$ .
- 648 (ii) For every compact  $K \subseteq X$ ,  $A|_K$  has the NIP.
- 649 (iii) For every Radon measure  $\mu$  on  $X$ ,  $A$  is relatively countably compact in  
 650  $\mathcal{M}^0(X, \mu)$ , i.e., every countable subset of  $A$  has an accumulation point in  
 651  $\mathcal{M}^0(X, \mu)$ .

652 Theorem 1.8 immediately yields the following.

653 **Theorem 3.12.** *Let  $(L, \mathcal{P}, \Gamma)$  be a CCS satisfying the Extendability Axiom. Let  $R$*   
 654 *be an exhaustive collection of sizers. Let  $\Delta \subseteq \Gamma$  be  $R$ -confined. If  $\pi_P \circ \Delta|_{\mathcal{L}[r_{\bullet}]}$  has*  
 655 *the NIP for all  $P \in \mathcal{P}$  and all  $r_{\bullet} \in R$ , then every deep computation is universally*  
 656 *essentially computable.*

657 *Proof.* By the Extendability Axiom, Theorem 1.8 and lemma 1.13 we have that  
 658  $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_{\bullet}]} \subseteq M_r(\mathcal{L}[r_{\bullet}])$  for all  $r_{\bullet} \in R$  and  $P \in \mathcal{P}$ . Let  $f \in \overline{\Delta}$  be a deep  
 659 computation. Write  $f = \mathcal{U} \lim_i \gamma_i$  as an ultralimit of computations in  $\Delta$ . De-  
 660 fine  $\tilde{f} := \mathcal{U} \lim_i \tilde{\gamma}_i$ . Then, for all  $r_{\bullet} \in R$  and  $P \in \mathcal{P}$   $\pi_P \circ \tilde{\gamma}_i|_{\mathcal{L}[r_{\bullet}]} \in M_r(\mathcal{L}[r_{\bullet}])$  for all  
 661  $i$ , so  $\pi_P \circ \tilde{f}|_{\mathcal{L}[r_{\bullet}]} \in \overline{\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_{\bullet}]}} \subseteq M_r(\mathcal{L}[r_{\bullet}])$ .  $\square$

**Question 3.13.** Under the same assumptions of the preceding theorem, suppose that every deep computation of  $\Delta$  is universally essentially computable. Must  $\pi_P \circ \Delta|_{L[r_\bullet]}$  have the NIP for all  $P \in \mathcal{P}$  and all  $r_\bullet \in R$ ?

**3.5. Talagrand stability, NIP, and essential computability of deep computations.** There is another notion closely related to NIP, introduced by Talagrand in [Tal84] while studying Pettis integration. Suppose that  $X$  is a compact Hausdorff space and  $A \subseteq \mathbb{R}^X$ . Let  $\mu$  be a Radon probability measure on  $X$ . Given a  $\mu$ -measurable set  $E \subseteq X$ , a positive integer  $k$  and real numbers  $a < b$ , we write:

$$D_k(A, E, a, b) = \bigcup_{f \in A} \{x \in E^{2k} : f(x_{2i}) \leq a, f(x_{2i+1}) \geq b \text{ for all } i < k\}$$

We say that  $A$  is *Talagrand  $\mu$ -stable* if and only if for every  $\mu$ -measurable set  $E \subseteq X$  of positive measure and for every  $a < b$  there is  $k \geq 1$  such that

$$(\mu^{2k})^*(D_k(A, E, a, b)) < (\mu(E))^{2k},$$

where  $\mu^*$  denotes the outer measure (we work with outer since the sets  $D_k(A, E, a, b)$  need not be  $\mu$ -measurable). This is certainly the case when  $A$  is a countable set of continuous (or  $\mu$ -measurable) functions.

The following lemma establishes that Talagrand stability is a way to ensure that deep computations are definable by measurable functions. We include a proof for the reader's convenience.

**Lemma 3.14.** *If  $A$  is Talagrand  $\mu$ -stable, then  $\overline{A}$  is also Talagrand  $\mu$ -stable and  $\overline{A} \subseteq \mathcal{M}^0(X, \mu)$ .*

*Proof.* First, observe that a subset of a  $\mu$ -stable set is  $\mu$ -stable. To show that  $\overline{A}$  is  $\mu$ -stable, observe that  $D_k(\overline{A}, E, a, b) \subseteq D_k(A, E, a', b')$  where  $a < a' < b' < b$  and  $E$  is a  $\mu$ -measurable set with positive measure. It suffices to show that  $\overline{A} \subseteq \mathcal{M}^0(X, \mu)$ . Suppose that there exists  $f \in \overline{A}$  such that  $f \notin \mathcal{M}^0(X, \mu)$ . By a characterization of measurable functions (see 413G in [Fre03]), there exists a  $\mu$ -measurable set  $E$  of positive measure and  $a < b$  such that  $\mu^*(P) = \mu^*(Q) = \mu(E)$  where  $P = \{x \in E : f(x) \leq a\}$  and  $Q = \{x \in E : f(x) \geq b\}$ . Then, for any  $k \geq 1$ :  $(P \times Q)^k \subseteq D_k(\{f\}, E, a, b)$  so  $(\mu^{2k})^*(D_k(\{f\}, E, a, b)) = (\mu^*(P)\mu^*(Q))^k = (\mu(E))^{2k}$ . Thus,  $\{f\}$  is not  $\mu$ -stable, but we argued before that a subset of a  $\mu$ -stable set must be  $\mu$ -stable.  $\square$

We say that  $A$  is *universally Talagrand stable* if  $A$  is Talagrand  $\mu$ -stable for every Radon probability measure  $\mu$  on  $X$ . An argument similar to the proof of 3.11, yields the following:

**Theorem 3.15.** *Let  $(L, \mathcal{P}, \Gamma)$  be a CCS satisfying the Extendability Axiom. If  $\pi_P \circ \Delta|_{L[r_\bullet]}$  is universally Talagrand stable for all  $P \in \mathcal{P}$  and all sizers  $r_\bullet$ , then every deep computation is universally essentially computable.*

It is then natural to ask: what is the relationship between Talagrand stability and the NIP? The following dichotomy will be useful.

**Lemma 3.16** (Fremlin's Dichotomy, 463K in [Fre03]). *If  $(X, \Sigma, \mu)$  is a perfect  $\sigma$ -finite measure space (in particular, for  $X$  compact and  $\mu$  a Radon probability measure on  $X$ ) and  $\{f_n : n \in \mathbb{N}\}$  be a sequence of real-valued measurable functions on  $X$ , then either:*

- 702 (i)  $\{f_n : n \in \mathbb{N}\}$  has a subsequence that converges  $\mu$ -almost everywhere, or  
 703 (ii)  $\{f_n : n \in \mathbb{N}\}$  has a subsequence with no  $\mu$ -measurable accumulation point  
 704 in  $\mathbb{R}^X$ .

705 The preceding lemma can be considered as a measure-theoretic version of Rosen-  
 706 thal's Dichotomy. Combining this dichotomy with the Theorem 3.11 we get the  
 707 following result:

708 **Theorem 3.17.** *Let  $X$  be a Hausdorff space and  $A \subseteq C(X)$  be pointwise bounded.*  
 709 *The following are equivalent:*

- 710 (i)  $\overline{A} \subseteq M_r(X)$ .  
 711 (ii) For every compact  $K \subseteq X$ ,  $A|_K$  has the NIP.  
 712 (iii) For every Radon measure  $\mu$  on  $X$ ,  $A$  is relatively countably compact in  
 713  $\mathcal{M}^0(X, \mu)$ , i.e., every countable subset of  $A$  has an accumulation point in  
 714  $\mathcal{M}^0(X, \mu)$ .  
 715 (iv) For every Radon measure  $\mu$  on  $X$  and every sequence  $\{f_n : n \in \mathbb{N}\} \subseteq A$ ,  
 716 there is a subsequence that converges  $\mu$ -almost everywhere.

717 *Proof.* Notice that the equivalence (i)-(iii) is Theorem 3.11. Notice that the equiv-  
 718 alence of (iii) and (iv) is Fremlin's Dichotomy (Theorem 3.16).  $\square$

719 Finally, it is natural to ask what the connection is between Talagrand stability  
 720 and NIP.

721 **Proposition 3.18.** *Let  $X$  be a compact Hausdorff space and  $A \subseteq C(X)$  be point-*  
 722 *wise bounded. If  $A$  is universally Talagrand stable, then  $A$  has the NIP.*

723 *Proof.* By Theorem 3.11, it suffices to show that  $A$  is relatively countably compact  
 724 in  $\mathcal{M}^0(X, \mu)$  for all Radon probability measure  $\mu$  on  $X$ . Since  $A$  is Talagrand  
 725  $\mu$ -stable for any such  $\mu$ , we have  $\overline{A} \subseteq \mathcal{M}^0(X, \mu)$ . In particular,  $A$  is relatively  
 726 countably compact in  $\mathcal{M}^0(X, \mu)$ .  $\square$

727 **Question 3.19.** Is the converse true?

728 The following two results suggest that the precise connection between Talagrand  
 729 stability and NIP may be sensitive to set-theoretic axioms (even assuming count-  
 730 ability of  $A$ ).

731 **Theorem 3.20** (Talagrand, Theorem 9-3-1(a) in [Tal84]). *Let  $X$  be a compact*  
 732 *Hausdorff space and  $A \subseteq M_r(X)$  be countable and pointwise bounded. Assume that*  
 733  *$[0, 1]$  is not the union of  $< \mathfrak{c}$  closed measure zero sets. If  $A$  has the NIP, then  $A$  is*  
 734 *universally Talagrand stable.*

735 **Theorem 3.21** (Fremlin, Shelah, [FS93]). *It is consistent that there exists a count-*  
 736 *able pointwise bounded set of Lebesgue measurable functions with the NIP which is*  
 737 *not Talagrand stable with respect to Lebesgue measure.*

## 738 REFERENCES

- 739 [ADIW24] Samson Alva, Eduardo Dueñez, Jose Iovino, and Claire Walton. Approximability of  
 740 deep equilibria. *arXiv preprint arXiv:2409.06064*, 2024. Last revised 20 May 2025,  
 741 version 3.  
 742 [ADK08] Spiros A. Argyros, Pandelis Dodos, and Vassilis Kanellopoulos. A classification of separ-  
 743 able Rosenthal compacta and its applications. *Dissertationes Mathematicae*, 449:1–55,  
 744 2008.

- 745 [Ark91] A. V. Arkhangel'skii. *Topological Function Spaces*. Springer, New York, 1st edition,  
746 1991.
- 747 [BD19] Shai Ben-David. Understanding machine learning through the lens of model theory.  
748 *The Bulletin of Symbolic Logic*, 25(3):319–340, 2019.
- 749 [BFT78] J. Bourgain, D. H. Fremlin, and M. Talagrand. Pointwise compact sets of Baire-  
750 measurable functions. *American Journal of Mathematics*, 100(4):845–886, 1978.
- 751 [BKK] Shaojie Bai, J. Zico Kolter, and Vladlen Koltun. Deep equilibrium models. Preprint.  
752 <https://arxiv.org/abs/1909.01377>, <https://implicit-layers-tutorial.org/>.
- 753 [CRBD] Ricky T. Q. Chen, Yulia Rubanova, Jesse Bettencourt, and David Duvenaud. Neural  
754 ordinary differential equations. Preprint. <https://arxiv.org/abs/1806.07366>.
- 755 [Deb13] Gabriel Debs. Descriptive aspects of Rosenthal compacta. In *Recent Progress in Gen-  
756 eral Topology III*, pages 205–227. Springer, 2013.
- 757 [Fre00] David H. Fremlin. *Measure Theory, Volume 1: The Irreducible Minimum*. Torres  
758 Fremlin, Colchester, UK, 2000. Second edition 2011.
- 759 [Fre01] David H. Fremlin. *Measure Theory, Volume 2: Broad Foundations*. Torres Fremlin,  
760 Colchester, UK, 2001.
- 761 [Fre03] David H. Fremlin. *Measure Theory, Volume 4: Topological Measure Spaces*. Torres  
762 Fremlin, Colchester, UK, 2003.
- 763 [FS93] David H. Fremlin and Saharon Shelah. Pointwise compact and stable sets of measurable  
764 functions. *Journal of Symbolic Logic*, 58(2):435–455, 1993.
- 765 [GM22] Eli Glasner and Michael Megrelishvili. Todorčević' trichotomy and a hierarchy in the  
766 class of tame dynamical systems. *Transactions of the American Mathematical Society*,  
767 375(7):4513–4548, 2022.
- 768 [God80] Gilles Godefroy. Compacts de Rosenthal. *Pacific J. Math.*, 91(2):293–306, 1980.
- 769 [Gro52] A. Grothendieck. Critères de compacité dans les espaces fonctionnels généraux. *Amer.  
770 J. Math.*, 74:168–186, 1952.
- 771 [HT23] Clovis Hamel and Franklin D. Tall.  $C_p$ -theory for model theorists. In Jose Iovino, edi-  
772 tor, *Beyond First Order Model Theory, Volume II*, chapter 5, pages 176–213. Chapman  
773 and Hall/CRC, 2023.
- 774 [Kec95] Alexander S. Kechris. *Classical Descriptive Set Theory*, volume 156 of *Graduate Texts  
775 in Mathematics*. Springer-Verlag, 1995.
- 776 [Kha20] Karim Khanaki. Stability, nip, and nsop; model theoretic properties of formulas via  
777 topological properties of function spaces. *Mathematical Logic Quarterly*, 66(2):136–  
778 149, 2020.
- 779 [Pap02] E. Pap, editor. *Handbook of measure theory. Vol. I, II*. North-Holland, Amsterdam,  
780 2002.
- 781 [Ros74] Haskell P. Rosenthal. A characterization of Banach spaces containing  $l^1$ . *Proc. Nat.  
782 Acad. Sci. U.S.A.*, 71:2411–2413, 1974.
- 783 [RPK19] M. Raissi, P. Perdikaris, and G. E. Karniadakis. Physics-informed neural networks: a  
784 deep learning framework for solving forward and inverse problems involving nonlinear  
785 partial differential equations. *J. Comput. Phys.*, 378:686–707, 2019.
- 786 [Sau72] N. Sauer. On the density of families of sets. *J. Combinatorial Theory Ser. A*, 13:145–  
787 147, 1972.
- 788 [She71] Saharon Shelah. Stability, the f.c.p., and superstability; model theoretic properties of  
789 formulas in first order theory. *Ann. Math. Logic*, 3(3):271–362, 1971.
- 790 [She72] Saharon Shelah. A combinatorial problem; stability and order for models and theories  
791 in infinitary languages. *Pacific J. Math.*, 41:247–261, 1972.
- 792 [She78] Saharon Shelah. *Unstable Theories*, volume 187 of *Studies in Logic and the Founda-  
793 tions of Mathematics*. North-Holland, 1978.
- 794 [She90] Saharon Shelah. *Classification theory and the number of nonisomorphic models*, vol-  
795 ume 92 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Pub-  
796 lishing Co., Amsterdam, second edition, 1990.
- 797 [Sim15a] Pierre Simon. *A Guide to NIP Theories*, volume 44 of *Lecture Notes in Logic*. Cam-  
798 bridge University Press, 2015.
- 799 [Sim15b] Pierre Simon. Rosenthal compacta and NIP formulas. *Fundamenta Mathematicae*,  
800 231(1):81–92, 2015.

- 801 [Tal84] Michel Talagrand. *Pettis Integral and Measure Theory*, volume 51 of *Memoirs of*  
802 *the American Mathematical Society*. American Mathematical Society, Providence, RI,  
803 USA, 1984. Includes bibliography (pp. 220–224) and index.
- 804 [Tal87] Michel Talagrand. The Glivenko-Cantelli problem. *The Annals of Probability*,  
805 15(3):837–870, 1987.
- 806 [Tka11] Vladimir V. Tkachuk. *A  $C_p$ -Theory Problem Book: Topological and Function Spaces*.  
807 Problem Books in Mathematics. Springer, 2011.
- 808 [Tod97] Stevo Todorćević. *Topics in Topology*, volume 1652 of *Lecture Notes in Mathematics*.  
809 Springer Berlin, Heidelberg, 1997.
- 810 [Tod99] Stevo Todorćević. Compact subsets of the first Baire class. *Journal of the American*  
811 *Mathematical Society*, 12(4):1179–1212, 1999.
- 812 [Val84] Leslie G. Valiant. A theory of the learnable. *Communications of the ACM*,  
813 27(11):1134–1142, 1984.
- 814 [VC71] Vladimir N Vapnik and Alexey Ya Chervonenkis. On the uniform convergence of rel-  
815 ative frequencies of events to their probabilities. *Theory of Probability & Its Applica-*  
816 *tions*, 16(2):264–280, 1971.
- 817 [VC74] Vladimir N Vapnik and Alexey Ya Chervonenkis. *Theory of Pattern Recognition*.  
818 Nauka, Moscow, 1974. German Translation: Theorie der Zeichenerkennung, Akademie-  
819 Verlag, Berlin, 1979.
- 820 [WYP22] Sifan Wang, Xinling Yu, and Paris Perdikaris. When and why PINNs fail to train: a  
821 neural tangent kernel perspective. *J. Comput. Phys.*, 449:Paper No. 110768, 28, 2022.