

COMPLEXITY OF DEEP COMPUTATIONS VIA TOPOLOGY OF FUNCTION SPACES

EDUARDO DUEÑEZ¹ JOSÉ IOVINO¹ TONATIUH MATOS-WIEDERHOLD²
LUCIANO SALVETTI² FRANKLIN D. TALL²

¹Department of Mathematics, University of Texas at San Antonio

²Department of Mathematics, University of Toronto

ABSTRACT. We use topological methods and a suitable framework to study the complexity of limit computations (such as the “deep computations” arising in recurrent neural networks). By using topology of function spaces —specifically, the classification of Rosenthal compacta— we identify new computational complexity classes. These classes form a strict hierarchy, each level of which is realized by classical computations such as the power method and Bresenham’s line algorithm, while Newton’s method transcends it. For iterative algorithms, the hierarchy coincides with a hierarchy of tame dynamical systems. The language of model theory, particularly, the concept of *independence* from Shelah’s classification theory, provides a medium to translate between topology and computation. Rosenthal compacta, and their measure-theoretic counterpart, give conditions for deep computations to be approximable, deterministically and probabilistically.

INTRODUCTION

In this paper we study asymptotic behavior of computations as parameters of the computation tend towards a limit, e.g., the depth of a neural network tending to infinity, or the time interval between layers of a time-series network tending toward zero. Recently, particular cases of this concept have attracted considerable attention in deep learning research (e.g., Neural Ordinary Differential Equations [CRBD], Physics-Informed Neural Networks [RPK19], and Deep Equilibrium Models [BKK]). The formal framework introduced here provides a unified setting to study these limit phenomena from a foundational viewpoint.

In a previous paper [ADIW26], the authors introduced limits of computations, which they called *ultracomputations*, since they arise as ultrafilter limits of standard computations, and *deep computations*, following usage in machine learning [BKK]. There is a technical difference between the two notions, but here we ignore it and speak of deep computations. Using a classical criterion of Grothendieck [Gro52], that paper proved a dichotomy between *stable* (polynomially approximable) and

Date: October 10, 2026.

2020 *Mathematics Subject Classification.* Primary 68Q15, 03C45, 54H30; Secondary 03C98, 03D15, 05D10, 26A21, 37B05, 54C35, 68T07.

Key words and phrases. Deep computations, limit computations, complexity classes, deep learning, recurrent neural networks, deep equilibrium models, compositional computation structures, independence property, NIP, model theory, Rosenthal compacta, Baire class 1 functions, Todorćević trichotomy, Argyros–Dodos–Kanellopoulos heptachotomy, Bourgain–Fremlin–Talagrand theorem, tame dynamical systems, infinite Ramsey theory.

unstable classes of deep computations. Polynomial approximability of computations corresponds to continuous extendibility in topology, and to *stability* and *type definability* in model theory.

In this paper we refine that dichotomy. An unstable class of deep computations may be *wild* (i.e., has Shelah’s *Independence Property* “IP”); otherwise, tame (when it is “NIP” —i.e., does *not* have IP). Within the tame unstable classes we identify three new levels giving a strict hierarchy of four tame classes beyond the stable ones, and seven minimal prototypes of tame families. The tools are more recent than those used in the first paper of this series [ADIW26]: on the topological side, the theory of *Rosenthal compacta* replaces Grothendieck’s criterion; on the model-theoretic side, *neostability* replaces stability.

The translation between topology and computation goes through model theory. To every intermediate state of a computation we associate its *type*: a “lookup table” of all observed (real) values of features of the computation at that instant. Those types form a space of real-valued functions —the *space of types*— on which computations act; the idea of embedding models into their type spaces is central in model theory. This embedding allows using the theory of function spaces (known as C_p -theory) to study topological limits of computations. The space of types thus becomes a “Rosetta stone” connecting three classification programs: in topology, the classification of Rosenthal compacta pioneered by Todorćević [Tod99]; in logic, the classification of theories developed by Shelah [She71, She90]; and in statistical learning, VC dimension and PAC learnability [VC71, VC74, Val84]. In such manner, Rosenthal compacta correspond to NIP —and also to an equivalent notion: finiteness of VC dimension (Section 1.5).

Deep computations arise as topological limits of standard (continuous) computations. Pointwise limits of sequences of continuous functions form the *first Baire class* —one topological rung above the class of continuous functions (intuitively, functions of Baire class 1 have “controlled” discontinuities). A *Rosenthal compactum* is a space of Baire class 1 functions, on a Polish (i.e., separable, completely metrizable) space, that is compact under the topology of *pointwise* convergence. Since the late 1970s, Rosenthal compacta have played a central role in functional analysis, especially in the theory of Banach spaces; the corresponding Todorćević’s trichotomy for such compacta [Tod99] has been used to organize tame dynamical systems into a hierarchy [GM22]. Todorćević’s proof relies on set-theoretic forcing and infinite Ramsey theory; decades later there is still no elementary proof, and his methods continue to be analyzed and exploited [Tod23, HT19]. Going beyond Todorćević’s trichotomy, Argyros, Dodos and Kanellopoulos [ADK08] have identified an heptachotomy consisting of seven minimal families of Baire class 1 functions, which we also use.

From the point of view of approximation, it is noteworthy that *compactness* and *sequential compactness* coincide in both the setting of Grothendieck [Gro52], and that of Bourgain, Fremlin and Talagrand [BFT78]. Consequently, when the computations are NIP, deep computations (which are, in principle, nonconstructive ultrafilter limits over a possibly uncountable set) can be approximated by *sequences* of standard computations.

We believe that the results presented in this paper show practitioners of computation, topology, descriptive set theory, or model theory how classification invariants

used in their field translate into classification invariants of the other fields. In the interest of accessibility, we do not assume familiarity with advanced topology, model theory, or computing. The only technical prerequisites are undergraduate-level topology and measure theory; the necessary background beyond them is covered in Section 1.

Main results. Theorem 3.1 shows that, when there are countably many predicates, a set of computations satisfies the NIP (one observable feature at a time, when restricted to “cylindrical state sets” that we call *shards*) if and only if its deep computations are of the first Baire class on each shard; equivalently, when the set of all such ultracomputations is a Rosenthal compactum; moreover, in such case, every deep computation is the limit of a *sequence* of computations. Using Todorćević’s trichotomy, we define three levels of tameness (Definition 3.2) and obtain the hierarchy

$$\text{stable} \Rightarrow \text{NIP}_3 \Rightarrow \text{NIP}_2 \Rightarrow \text{NIP}_1 \Rightarrow \text{NIP},$$

in which every implication is strict. Section 5 realizes each level by a classical algorithm:

- a rotation is stable;
- the power method is NIP_3 but not stable;
- Bresenham’s line algorithm is
 - NIP_2 but not NIP_3 in 2-dimensions (in the plane),
 - NIP_1 but not NIP_2 in 3-dimensions;
- tolerance tests, whose deep computations are exact equality tests, are NIP but not NIP_1 ;
- Newton’s method to find polynomial roots on the extended complex plane is not NIP .

For iterative algorithms, the hierarchy coincides with that of tame dynamical systems due to Glasner and Megrelishvili [GM22]. The heptachotomy of Argyros, Dodos and Kanellopoulos then shows that every tame family of computations indexed by binary tree contains, along a dyadic subtree, one of seven minimal families, which we read as primitives of binary search (Section 3.3). Finally, Theorem 6.7 and Corollary 6.14 relate NIP and Talagrand stability to Monte Carlo computability of deep computations.

Organization of the paper. In Section 1, we present the topological and combinatorial preliminaries, including five canonical Rosenthal compacta (Section 1.4) and the role of NIP in model theory, combinatorics and learning (Section 1.5). In Section 2, we introduce compositional computation structures, the structural and model-theoretic viewpoint (no previous exposure to model theory is needed), together with stability and the register construction; the section ends with two motivating examples, Newton’s method for square roots and Newton’s method on the Riemann sphere, whose different behaviors the classification is designed to separate (Section 2.5). Section 3 is devoted to the classification of deep computations; it contains the main theorems. Section 4 works out Newton’s method, threshold classifiers and prefix tests in detail, and Section 5 places examples at every level of the hierarchy, including classical iterative algorithms. Section 6 replaces topological limits by a probabilistic viewpoint, which we intend to develop further in future research. Section 7 collects open questions.

Throughout the paper, our results pertain to classical models of computation (particularly computations involving real-valued quantities that are known and manipulated to a finite degree of precision).

CONTENTS

Introduction	1
Main results	3
Organization of the paper	3
1. Topological preliminaries:	
From continuity to Baire class 1	5
1.1. From Rosenthal's dichotomy to the Bourgain-Fremlin-Talagrand dichotomy to Shelah's NIP	6
1.2. Rosenthal compacta	8
1.3. The special case $B_1(X, \mathbb{R}^{\mathcal{P}})$ with $\mathcal{P}$ countable	8
1.4. Five canonical Rosenthal compacta	10
1.5. NIP as a universal dividing line between polynomial and exponential complexity	11
Historical remarks	12
2. Compositional Computation Structures:	
A structural approach to arbitrary-precision computations	12
2.1. Computation States Structures	12
2.2. Compositional Computation Structures	14
2.3. Computability of deep computations and the Extendibility Axiom	15
2.4. Stability and register computations	16
2.5. Two motivating examples	17
3. Classifying deep computations	17
3.1. NIP, Rosenthal compacta, and deep computations	17
3.2. The Todorćević trichotomy and levels of NIP	18
3.3. The Argyros-Dodos-Kanellopoulos heptachotomy and binary search	20
4. Examples of Compositional Computation Structures	21
4.1. Example: Newton's CCS for the cubic $\rho(z) = z^3 - 2z + 2$	22
4.2. Example: Newton's CCS for 3-cyclotomic $\kappa(z) = z^3 - 1$	23
4.3. Example: Square roots	24
4.4. Example: Threshold classifiers on Cantor's set	24
4.5. Example: Prefix tests	25
5. Computations across the hierarchy	25
5.1. Iterative algorithms and tame dynamics	26
5.2. Stable and metrizable	26
5.3. Hereditarily separable but not metrizable	27
5.4. First countable but not hereditarily separable	28
5.5. NIP but not first countable	29
5.6. Beyond NIP: Newton's method on the Riemann sphere	30
5.7. Summary	31
6. Randomized versions of NIP and Monte Carlo computability of deep computations	31
6.1. NIP and Monte Carlo computability of deep computations	31
6.2. Talagrand stability and Monte Carlo computability of deep computations	33

7. Open questions	36
References	36

1. TOPOLOGICAL PRELIMINARIES: FROM CONTINUITY TO BAIRE CLASS 1

In this section we recall notions and results from general topology and function space theory. For simplicity, we assume that all spaces are Hausdorff unless explicitly stated otherwise.

A subset of a topological space is F_σ if it is a countable union of closed sets, and G_δ if it is a countable intersection of open sets. In a metric space, every open set is F_σ ; equivalently, every closed set is G_δ .

A *Polish space* is a separable and completely metrizable topological space, i.e., one admitting a complete metric that induces its topology. The class of Polish spaces is closed under countable topological products. Classical examples of Polish spaces (besides the real line) include the Cantor space $2^\mathbb{N}$ (the set of all infinite binary sequences, endowed with the product topology), the Baire space $\mathbb{N}^\mathbb{N}$ (the set of all infinite sequences of naturals, also with the product topology), and the space $\mathbb{R}^\mathbb{N}$ of sequences of real numbers.

Fact 1.1. [Eng89, 4.3.24] *A subset of a Polish space is itself Polish in the subspace topology if and only if it is a G_δ set. In particular, closed subsets and open subsets of Polish spaces are also Polish spaces.*

If X is a topological space and A is a subset of X , then A is said to be *nowhere dense* if its closure has an empty interior, and of the *first category* if A is a countable union of closed nowhere dense sets; otherwise, A is said to be of the *second category*. Sets $A \subseteq X$ of the first category are also called *meager*, and their complements $X \setminus A$ are called *comeager*. Thus, the set of meager subsets of X is the σ -ideal generated by the nowhere dense subsets of X . A topological space X is *Baire* if every comeager subset of X is dense. Equivalently, X is Baire if any countable intersection of open dense sets remains dense. The Baire Category Theorem states that compact Hausdorff spaces and all completely metrizable spaces (hence, all Polish spaces) are Baire.¹

Given two topological spaces X and Y we denote by $C_p(X, Y)$ the set of all continuous functions $f : X \rightarrow Y$ endowed with the topology of pointwise convergence, i.e., with the relative (subspace) product topology $C_p(X, Y) \subseteq Y^X$ —where Y carries its topology, but X serves merely as a (discrete) index set stripped of any topology. The space $C_p(X, \mathbb{R})$ of continuous real-valued functions on X is denoted simply $C_p(X)$.

A function $f \in Y^X$ between topological spaces X, Y is said to be of *Baire class 1*, or a *Baire-1 function*, if there is a sequence $(f_n)_{n \in \mathbb{N}} \subseteq C_p(X, Y)$ such that $f = \lim_{n \rightarrow \infty} f_n$ (i.e., $f(x) = \lim_{n \rightarrow \infty} f_n(x)$ for all $x \in X$). An elementary argument shows that Baire-1 functions are continuous everywhere except on a set of the first category. If X and Y are topological spaces, the space of Baire-1 functions $f : X \rightarrow Y$ endowed with the topology of pointwise convergence is denoted $B_1(X, Y)$; thus, $C_p(X, Y) \subseteq B_1(X, Y) \subseteq Y^X$ are topological inclusions. We abbreviate $B_1(X, \mathbb{R})$

¹More generally, any Čech-complete space is Baire [Eng89, Theorem 3.9.3].

as $B_1(X)$. Relatively compact subsets of $C_p(X)$ and of $B_1(X)$ (particularly for X Polish) are of interest in analysis and topological dynamics.

A topological space X is *perfectly normal* if it is normal and every closed subset of X is a G_δ set (equivalently, every open subset of X is an F_σ set). Every metrizable space (hence, every Polish space) is perfectly normal.

The following fact was proved by Baire in his groundbreaking thesis. For a proof, see [Tod97, Section 10].

Fact 1.2 (Baire). *If X is perfectly normal, then the following conditions are equivalent for a function $f : X \rightarrow \mathbb{R}$:*

- (1) *f is a Baire class 1 function, that is, f is a pointwise limit of continuous functions.*
- (2) *$f^{-1}[U]$ is an F_σ subset of X whenever $U \subseteq \mathbb{R}$ is open.*

If X is Baire, then (1) and (2) are equivalent to:

- (3) *For every closed $F \subseteq X$, the restriction $f|_F$ has a point of continuity.*

Moreover, if X is Polish and $f \notin B_1(X)$, then there exist countable $D_0, D_1 \subseteq X$ and reals $a < b$ such that

$$D_0 \subseteq f^{-1}(-\infty, a], \quad D_1 \subseteq f^{-1}[b, \infty), \quad \overline{D_0} = \overline{D_1}.$$

1.1. From Rosenthal's dichotomy to the Bourgain-Fremlin-Talagrand dichotomy to Shelah's NIP. In metrizable spaces, the closure of a subset consists of limits of convergent sequences therein. In general, such a characterization of the closure fails in remarkable ways.

The n -th coordinate map $p_n : 2^{\mathbb{N}} \rightarrow \{0, 1\}$ on the Cantor space $X = 2^{\mathbb{N}}$ ($= \{0, 1\}^{\mathbb{N}}$) is continuous for each $n \in \mathbb{N}$; however, $(p_n)_{n \in \mathbb{N}}$ has *no* convergent subsequences in $\mathbb{R}^X$ ([Tod97, Chapter 1.1]). In a sense, this example exhibits the worst failure of sequential convergence possible: The closure of $\{p_n\}$ in $\{0, 1\}^X$ (or in $\mathbb{R}^X$ for that matter) is homeomorphic to the Stone-Ćech compactification $\beta\mathbb{N}$ of the (discrete) space of natural numbers. The space $\beta\mathbb{N}$ has cardinality 2^{2^ω} (largest among all separable topological spaces) and, effectively, is the set of ultrafilters on $\mathbb{N}$.²

The following theorem, proved by Haskell Rosenthal in 1974, is fundamental in functional analysis and captures a sharp division in the behavior of sequences in a Banach space.

Theorem 1.3 (Rosenthal's Dichotomy, [Ros74]). *If X is Polish and $(f_n)_{n \in \mathbb{N}} \subseteq C_p(X)$ is pointwise bounded, then (f_n) has a convergent subsequence, or a subsequence whose closure in $\mathbb{R}^X$ is homeomorphic to $\beta\mathbb{N}$.*

The genesis of the dichotomy was *Rosenthal's ℓ_1 -Theorem*, which states that a Banach space includes an isomorphic copy of ℓ_1 (the space of absolutely summable sequences), or else every bounded sequence therein is weakly Cauchy. The ℓ_1 -Theorem connects diverse areas: Banach space geometry, Ramsey theory, set theory, and topology of function spaces.

As we move from $C_p(X)$ to the larger space $B_1(X)$, a dichotomy paralleling the ℓ_1 -Theorem holds. This result is usually not phrased as a dichotomy in the above manner, but rather as an equivalence (see Theorem 1.4 below).

²Our earlier proof [ADIW26] of the existence of deep equilibria in compositional computation structures relies on a classical result on the existence of idempotent elements of $\beta\mathbb{N}$.

First, we introduce some useful notation. For any $f \in \mathbb{R}^X$ and any real a , define

$$X_{\leq a}^f := f^{-1}(-\infty, a], \quad X_{\geq a}^f := f^{-1}[a, +\infty).$$

For any set $A \subseteq \mathbb{R}^X$ and any real a , define (by a slight abuse of notation)

$$X_{\leq a}^A := \bigcap_{f \in A} X_{\leq a}^f = \{x \in X : f(x) \leq a \text{ for all } f \in A\},$$

$$X_{\geq a}^A := \bigcap_{f \in A} X_{\geq a}^f = \{x \in X : f(x) \geq a \text{ for all } f \in A\}.$$

(In case $A = \emptyset$, we define $X_{\geq a}^\emptyset = X = X_{\leq a}^\emptyset$.) For any sequence $\{f_n\} \subseteq \mathbb{R}^X$ and $I \subseteq \mathbb{N}$, define $I^\complement := \mathbb{N} \setminus I$ and $f_I := \{f_i : i \in I\}$.

Theorem 1.4 (“The BFT Dichotomy”. Bourgain-Fremlin-Talagrand [BFT78]). *Let X be a Polish space and $A \subseteq C_p(X)$ be pointwise bounded. The following are equivalent:*

- (i) A is relatively compact in $B_1(X)$.
- (ii) For every $(f_n)_{n \in \mathbb{N}} \subseteq A$ and every $a < b$ there is $I \subseteq \mathbb{N}$ such that

$$X_{\leq a}^{f_I} \cap X_{\geq b}^{f_{I^\complement}} = \emptyset.$$

(As stated above, the BFT Dichotomy is a particular case of the equivalence (ii) $\Leftrightarrow$ (v) in [BFT78, Corollary 4G].)

The sets $X_{\leq a}^{f_I}$ and $X_{\geq b}^{f_{I^\complement}}$ appearing in condition Theorem 1.4(ii) are defined, respectively, in terms of $|I|$ -many inequalities of the form $f_i(x) \leq a$, and $|I^\complement|$ -many of the form $f_j(x) \geq b$. Thus, at least one of $X_{\leq a}^{f_I}$ and $X_{\geq b}^{f_{I^\complement}}$ is defined by the satisfaction of infinitely (countably) many inequalities. For our purposes, it is more natural to consider only finitely many inequalities at a time, which motivates the definitions below.

Definition 1.5. We say that a collection of functions $A \subseteq \mathbb{R}^X$ has the *finitary No-Independence Property (NIP)* if, for all sequences $(f_n)_{n \in \mathbb{N}} \subseteq A$ and reals $a < b$, there exist finite disjoint sets $E, F \subseteq \mathbb{N}$ such that $X_{\leq a}^{f_E} \cap X_{\geq b}^{f_F} = \emptyset$. We say that such E, F witness *finitary NIP* for (f_n) and a, b .

A set $A \subseteq \mathbb{R}^X$ has the *finitary Independence Property (IP)* if it does not have finitary NIP, i.e., if there exists a sequence $(f_n)_{n \in \mathbb{N}} \subseteq A$ and reals $a < b$ such that for every pair of finite disjoint sets $E, F \subseteq \mathbb{N}$, we have $X_{\leq a}^{f_E} \cap X_{\geq b}^{f_F} \neq \emptyset$.

If the word “finite” is omitted in the above definitions, we obtain the definitions of *countable NIP* (weaker than finitary NIP) and *countable IP* (stronger than finitary IP), respectively.

If we insist on witnesses $E, F \subseteq \mathbb{N}$ such that $F = E^\complement$, we call the respective properties “BFT-NIP” and “BFT-IP”. Thus, Theorem 1.4 becomes (for pointwise bounded function collections $A \subseteq C_p(X)$) that A is relatively compact in $B_1(X)$ if and only if A has BFT-NIP.

Unless otherwise specified, IP/NIP shall mean *finitary* IP/NIP henceforth.

Theorem 1.6. *Let X be a compact metrizable (hence Polish) space. For every pointwise bounded $A \subseteq C_p(X)$, the following properties are all equivalent:*

- (i) A is relatively compact in $B_1(X)$;

- (ii) A has BFT-NIP;
- (iii) A has countable NIP;
- (iv) A has finitary NIP.

(The equivalence (ii) $\Leftrightarrow$ (iv) holds for arbitrary compact X , without assuming pointwise boundedness. The equivalence (ii) $\Leftrightarrow$ (iii), for arbitrary X .)

Proof. (i) $\Leftrightarrow$ (ii) is Theorem 1.4. The implications (iv) $\Rightarrow$ (iii) and (ii) $\Rightarrow$ (iii) are trivial, and (iii) $\Rightarrow$ (ii) follows by taking $I = E$, since then $F \subseteq I^{\mathbb{C}}$. For (ii) $\Rightarrow$ (iv), suppose that (f_n) and $a < b$ witness finitary IP and fix $I \subseteq \mathbb{N}$. The closed sets $X_{\leq a}^{f_n}$ ($n \in I$) and $X_{\geq b}^{f_m}$ ($m \in I^{\mathbb{C}}$) have the finite intersection property, so by compactness $X_{\leq a}^{f_I} \cap X_{\geq b}^{f_{I^{\mathbb{C}}}} \neq \emptyset$. $\square$

Theorem 1.6 may be stated as the following dichotomy (under the assumptions) for any compact metrizable X : either A is relatively compact in $B_1(X)$, or A has IP (in either sense).

The Independence Property was first isolated by Saharon Shelah in model theory as a dividing line between theories whose models are “tame” (corresponding to NIP) and theories whose models are “wild” (corresponding to IP). See [She71, Definition 4.1], [She90]. We discuss this dividing line in Section 1.5.

1.2. Rosenthal compacta. The dichotomy of Theorem 1.4 led Godefroy [God80] to the following definition.

Definition 1.7. A Rosenthal compactum is any topological space realized as a compact subset of the space $B_1(X) = B_1(X, \mathbb{R})$ (equipped with the topology of pointwise convergence) of all real functions of the first Baire class on some Polish space X .

A Rosenthal compactum K is necessarily Hausdorff since it is a topological subspace of the Hausdorff product space $\mathbb{R}^X$.

Rosenthal compacta possess significant topological and dynamical tameness properties, and play an important role in functional analysis, measure theory, dynamical systems, descriptive set theory, and model theory. In this paper, we use them to study deep computations.

1.3. The special case $B_1(X, \mathbb{R}^{\mathcal{P}})$ with $\mathcal{P}$ countable. Fix an arbitrary (at most) countable set $\mathcal{P}$ whose elements $P \in \mathcal{P}$ will be called *predicate symbols* or *formal predicates*. Our present goal is to characterize relatively compact subsets of $B_1(X, \mathbb{R}^{\mathcal{P}})$, where X is always assumed to be a perfectly normal space (typically a Polish space).

The set $\mathcal{P}$ shall be considered discrete whenever regarded as a topological space. Since $C_p(X, \mathbb{R}^{\mathcal{P}}) \subseteq B_1(X, \mathbb{R}^{\mathcal{P}}) \subseteq (\mathbb{R}^{\mathcal{P}})^X$, the “ambient” space $(\mathbb{R}^{\mathcal{P}})^X$ is quite relevant. The product $X \times \mathcal{P}$ will be regarded as either a point set, or as a topological product depending on context. We have natural homeomorphic identifications

$$(\mathbb{R}^{\mathcal{P}})^X \cong \mathbb{R}^{X \times \mathcal{P}} \cong (\mathbb{R}^X)^{\mathcal{P}},$$

given by

$$\begin{aligned} \mathbb{R}^{X \times \mathcal{P}} &\rightarrow (\mathbb{R}^{\mathcal{P}})^X : \varphi \mapsto \hat{\varphi} \\ \mathbb{R}^{X \times \mathcal{P}} &\rightarrow (\mathbb{R}^X)^{\mathcal{P}} : \varphi \mapsto \varphi^\wedge, \end{aligned}$$

where

$$\hat{\varphi}(x) := \varphi(x, \cdot) \in \mathbb{R}^{\mathcal{P}}, \quad \varphi^\sim(P) := \varphi(\cdot, P) \in \mathbb{R}^X.$$

Such identifications view X and $\mathcal{P}$ as mere point sets (the topology of X in particular plays no role).

For $x \in X$, define the “left projection” map

$$\lambda_x : \mathbb{R}^{X \times \mathcal{P}} \rightarrow \mathbb{R}^{\mathcal{P}} : \varphi \mapsto \lambda_x(\varphi) := \hat{\varphi}(x);$$

for $P \in \mathcal{P}$, the “right projection” map

$$\rho_P : \mathbb{R}^{X \times \mathcal{P}} \rightarrow \mathbb{R}^X : \varphi \mapsto \rho_P(\varphi) := \varphi^\sim(P).$$

For fixed $x \in X$ and $P \in \mathcal{P}$, we also have canonical projection maps

$$\lambda_x : \mathbb{R}^X \rightarrow \mathbb{R} : f \mapsto f(x), \quad \rho_P : \mathbb{R}^{\mathcal{P}} \rightarrow \mathbb{R} : f \mapsto f(P).$$

When clear from context, rather than using the specific symbols (“ λ ” for left, “ ρ ” for right) to denote projections, we may use the generic symbol “ π ”; thus, π_x may mean λ_x , and π_P may mean ρ_P .

The Proposition below reduces the study of $\mathbb{R}^{\mathcal{P}}$ -valued continuous or Baire-1 functions on X to the special case of real-valued ones.

Proposition 1.8. *The identification $(\mathbb{R}^{\mathcal{P}})^X \cong \mathbb{R}^{X \times \mathcal{P}} \cong (\mathbb{R}^X)^{\mathcal{P}}$ induces identifications*

$$C_p(X, \mathbb{R}^{\mathcal{P}}) \cong C_p(X \times \mathcal{P}) \cong C_p(X)^{\mathcal{P}}, \quad B_1(X, \mathbb{R}^{\mathcal{P}}) \cong B_1(X \times \mathcal{P}) \cong B_1(X)^{\mathcal{P}}.$$

(The cardinality of $\mathcal{P}$ plays no role.)

Proof. The identification of C_p -spaces follows trivially from the definition of topological product and the fact that $\mathcal{P}$ is discrete: a continuous map $X \rightarrow \mathbb{R}^{\mathcal{P}}$ is precisely a $\mathcal{P}$ -indexed family of continuous functions $X \rightarrow \mathbb{R}$, and these correspond to continuous functions $X \times \mathcal{P} \rightarrow \mathbb{R}$. The identification of Baire-1 spaces follows immediately, since it is defined in terms of the purely topological notion of limit (in the ambient space) of sequences of continuous functions. $\square$

Given $A \subseteq Y^X$ and $K \subseteq X$ we write $A|_K := \{f|_K : f \in A\}$, i.e., the set of all restrictions of functions in A to K . The following Theorem is a slightly more general version of Theorem 1.4.

Theorem 1.9. *Assume that $\mathcal{P}$ is countable, X is a Polish space, and $A \subseteq C_p(X, \mathbb{R}^{\mathcal{P}})$ is pointwise bounded in the sense that $\pi_P \circ A (\subseteq C_p(X))$ is pointwise bounded for every $P \in \mathcal{P}$. The following are equivalent for every compact $K \subseteq X$:*

- (i) $A|_K$ is relatively compact in $B_1(K, \mathbb{R}^{\mathcal{P}})$;
- (ii) $\pi_P \circ A|_K$ has NIP for every $P \in \mathcal{P}$.

Proof. By Proposition 1.8 and Tychonoff’s theorem, (i) holds if and only if $\pi_P \circ A|_K$ is relatively compact in $B_1(K)$ for every $P \in \mathcal{P}$. Since K is compact metrizable, Theorem 1.6 shows that this is equivalent to (ii). $\square$

We conclude this section with an elementary but important observation.

Lemma 1.10. *Assume that X is Hausdorff and that $A \subseteq C_p(X)$. The following are equivalent for every $L \subseteq X$:*

- (i) $A|_L$ satisfies the NIP;
- (ii) $A|_{\overline{L}}$ satisfies the NIP.

Proof. (ii) $\Rightarrow$ (i) is trivial. Conversely, if $(f_n) \subseteq A$ and $a < b$ witness the IP on $\bar{L}$, fix $a < a' < b' < b$. For finite disjoint $E, F \subseteq \mathbb{N}$, the open set $\bigcap_{n \in E} f_n^{-1}(-\infty, a') \cap \bigcap_{n \in F} f_n^{-1}(b', \infty)$ meets $\bar{L}$, hence meets L ; so (f_n) and $a' < b'$ witness the IP on L . $\square$

1.4. Five canonical Rosenthal compacta. Five Rosenthal compacta, all built on the Cantor set, recur throughout the paper: they are the canonical members of the classes in Todorćević's trichotomy (Theorem 3.3), and the deep computations of our examples are copies of them. We fix their notation here. Recall that a topological space is *separable* if it has a countable dense subset, *hereditarily separable* if every subspace is separable, and *first countable* if every point has a countable base of neighborhoods.

Let $2^{\mathbb{N}}$ carry the product topology and the lexicographic order $<$ (whose order topology is the product topology). The length of a finite word $t \in 2^{<\mathbb{N}}$ is denoted $|t|$; for $a \in 2^{\mathbb{N}}$, we write $a \upharpoonright n$ for the word formed by the first n bits of a . Two words are *comparable* if one extends the other, and an *antichain* is a set of pairwise incomparable words. For a finite word $t \in 2^{<\mathbb{N}}$, let $t \smallfrown 0^\infty$ and $t \smallfrown 1^\infty$ be the sequences that begin with t and continue with all 0's, respectively all 1's; the set of sequences extending t is the clopen interval $[t \smallfrown 0^\infty, t \smallfrown 1^\infty]$.

For a statement S , we write $\llbracket S \rrbracket$ for its truth value: 1 if S is true and 0 if it is false. We introduce the following functions $2^{\mathbb{N}} \rightarrow \{0, 1\}$:

- the *prefix test* $v_t := \llbracket \cdot \upharpoonright |t| = t \rrbracket$, the characteristic function of the set of sequences extending a fixed $t \in 2^{<\mathbb{N}}$ (here, $x \upharpoonright n = x_{<n} = (x_i)_{i < n} \in 2^n \subseteq 2^{<\mathbb{N}}$ is the truncation of $x \in 2^{\mathbb{N}}$ to a string of finite length $n \in \mathbb{N}$);
- the *equality test* $\delta_a := \llbracket \cdot = a \rrbracket$, the characteristic function of the singleton $\{a\}$ for fixed $a \in 2^{\mathbb{N}}$;
- the *threshold tests* $\chi_{[a]} := \llbracket a \leq \cdot \rrbracket$ and $\chi_{(a)} := \llbracket a < \cdot \rrbracket$, the characteristic functions of the final intervals $\{x : a \leq x\}$ (closed) and $\{x : a < x\}$ (open), for fixed $a \in 2^{\mathbb{N}}$; the special case $\mathbf{0} := \chi_{(1^\infty)}$ is the identically zero function.

Prefix tests are continuous, and so are $\chi_{[t \smallfrown 0^\infty]}$ and $\chi_{(t \smallfrown 1^\infty)}$ —the all depend only on the first $|t|$ bits of its argument in $2^{\mathbb{N}}$; equality tests, and threshold tests at other points, are discontinuous Baire class 1 functions. All closures below are taken in the topology of pointwise convergence.

- (1) The *Alexandroff compactification* $A(2^{\mathbb{N}}) = \{\delta_a : a \in 2^{\mathbb{N}}\} \cup \{\mathbf{0}\}$. Each δ_a is isolated, and a neighborhood of $\mathbf{0}$ omits only finitely many δ_a . Thus, $A(2^{\mathbb{N}})$ is not separable, nor first countable at $\mathbf{0}$.
- (2) The *extended Alexandroff compactification* $\hat{A}(2^{\mathbb{N}}) = \overline{\{v_t : t \in 2^{<\mathbb{N}}\}} = \{v_t : t \in 2^{<\mathbb{N}}\} \cup A(2^{\mathbb{N}})$. As $n \rightarrow \infty$, the prefix tests $v_{a \upharpoonright n}$ converge to δ_a ; along an antichain they converge to $\mathbf{0}$. Thus, $\hat{A}(2^{\mathbb{N}})$ is separable but not first countable.
- (3) The *split Cantor set* $S(2^{\mathbb{N}}) = \{\chi_{[a]}, \chi_{(a)} : a \in 2^{\mathbb{N}}\}$. The continuous thresholds $\chi_{[t \smallfrown 0^\infty]}$, together with the isolated point $\mathbf{0} = \chi_{(1^\infty)}$, form a countable dense subset: thresholds approaching a from below converge to $\chi_{[a]}$, and from above to $\chi_{(a)}$. Thus, $S(2^{\mathbb{N}})$ is separable, first countable and hereditarily separable, but not metrizable.
- (4) The *Alexandroff duplicate* $D(2^{\mathbb{N}}) = \{(\mathbf{0}, a), (\delta_a, a) : a \in 2^{\mathbb{N}}\} \subseteq A(2^{\mathbb{N}}) \times 2^{\mathbb{N}}$. The points (δ_a, a) are isolated, and a neighborhood of $(\mathbf{0}, a)$ contains $(\mathbf{0}, b)$ and (δ_b, b) for all b near a with finitely many exceptions. Thus, $D(2^{\mathbb{N}})$

is first countable but not separable; its separable extension is $\hat{D}(2^{\mathbb{N}}) = \overline{\{(v_t, t \frown 0^\infty) : t \in 2^{<\mathbb{N}}\}} = D(2^{\mathbb{N}}) \cup \{(v_t, t \frown 0^\infty) : t \in 2^{<\mathbb{N}}\}$.

(5) The *extended duplicate of the split Cantor set*

$$\hat{D}(S(2^{\mathbb{N}})) = \overline{\{(v_t, \chi_{[t \frown 0^\infty]})\}_t} \subseteq \hat{A}(2^{\mathbb{N}}) \times S(2^{\mathbb{N}}),$$

which consists of the points $(v_t, \chi_{[t \frown 0^\infty]})$ and, for $a \in 2^{\mathbb{N}}$, the points $(\delta_a, \chi_{[a]})$, $(\mathbf{0}, \chi_{[a]})$ and $(\mathbf{0}, \chi_{(a)})$ [ADK08, 4.3.7]. Along the words $t = a \upharpoonright n$, the thresholds $t \frown 0^\infty$ increase to a , which gives the points $(\delta_a, \chi_{[a]})$. Thus, $\hat{D}(S(2^{\mathbb{N}}))$ is separable and first countable; since, the points $(\delta_a, \chi_{[a]})$ form an uncountable discrete subspace, it is not hereditarily separable.

The same constructions apply with $2^{\mathbb{N}}$ replaced by another compact metric space K ; we write $A(K)$, $D(K)$, and so on. A compactum of size continuum, such as $A([0, 1])$, is homeomorphic to $A(2^{\mathbb{N}})$.

1.5. NIP as a universal dividing line between polynomial and exponential complexity. The particular case of the BFT dichotomy (Theorem 1.4) in which A consists of $\{0, 1\}$ -valued (i.e., yes/no-valued) functions was discovered independently around 1971–1972 in three foundational contexts, each separating polynomial (“tame”) from exponential (“wild”) complexity: in model theory, by Saharon Shelah [She71, She90]; in combinatorics, by Norbert Sauer [Sau72] and by Shelah [She72, She90]; and in statistical learning, by Vladimir Vapnik and Alexey Chervonenkis [VC71, VC74].

In model theory: Shelah’s classification theory is a foundational program in mathematical logic devised to categorize first-order theories based on the complexity and structure of their models. A theory T is considered classifiable in Shelah’s sense if the number of non-isomorphic models of T of a given cardinality can be described by a bounded number of numerical invariants. In contrast, a theory T is unclassifiable if the number of models of T of a given cardinality is the maximum possible number. A key fact is that the number of models of T is directly impacted by the number of *types* over sets of parameters in models of T ; a controlled number of types is a characteristic of a classifiable theory.

In Shelah’s classification program [She90], theories *without* the independence property (called NIP theories, or *dependent* theories) have a well-behaved, “tame” structure: the number of types over a set of parameters of size κ grows slowly (polynomially, or nearly so) in κ .

In contrast, theories with the Independence Property (called IP theories) are considered “intractable” or “wild”. A theory with the Independence Property produces the maximum possible number of types over a set of parameters; for a set of parameters of cardinality κ , the theory will have 2^{2^κ} -many distinct types.

In combinatorics: Sauer [Sau72] and Shelah [She72] proved the following independently: Let $\mathcal{F} = \{S_i : i \in \mathbb{N}\}$ be a countable collection of subsets of a set S . Either: for every $n \in \mathbb{N}$ there is a set $A \subseteq S$ with $|A| = n$ such that $|\{S_i \cap A : i \in \mathbb{N}\}| = 2^n$ ($\mathcal{F}$ is a collection having “exponential complexity”);

or: there exists $N \in \mathbb{N}$ such that for every $A \subseteq S$ with $|A| \geq N$, one has

$$|\{S_i \cap A : i \in \mathbb{N}\}| \leq \sum_{i=0}^{N-1} \binom{|A|}{i}.$$

($\mathcal{F}$ is a collection having “polynomial complexity”). Clearly, any collection $\mathcal{F}$ of subsets of a *finite* set S has polynomial complexity. The “polynomial” name is justified: indeed, for fixed $N > 0$, as a function of the size $|A| = m > 0$, we have

$$\sum_{i=0}^{N-1} \binom{m}{i} \leq \sum_{i=0}^{N-1} \frac{m^i}{i!} \leq \left(\sum_{i=0}^{N-1} \frac{1}{i!} \right) \cdot m^{N-1} < e \cdot m^{N-1},$$

which is polynomial in m for fixed N .

In machine learning: Readers familiar with statistical learning may recognize the Sauer-Shelah lemma as the dichotomy discovered and proved slightly earlier (1971) by Vapnik and Chervonenkis [VC71, VC74] to address uniform convergence in statistics. The least integer N given by the preceding paragraph, when it exists, is called the *VC-dimension* of $\mathcal{F}$; it is a core concept in machine learning. If such an integer N does not exist, we say that the VC-dimension of $\mathcal{F}$ is infinite. The lemma provides upper bounds on the number of data points (sample size) needed to learn a concept class of known VC dimension d up to a given admissible error in the statistical sense. The Fundamental Theorem of Statistical Learning states that a hypothesis class is PAC-learnable (PAC stands for “Probably Approximately Correct”) if and only if its VC dimension is finite.

Historical remarks. The Baire hierarchy of functions was introduced by René-Louis Baire in his 1899 doctoral thesis, *Sur les fonctions de variables réelles*. The field of C_p -theory was pioneered by A. V. Arhangel’skiĭ and his students in the 1970s and 1980s [Ark92], and it has found many applications in model theory and functional analysis. For a recent survey, see [Tka11].

2. COMPOSITIONAL COMPUTATION STRUCTURES:

A STRUCTURAL APPROACH TO ARBITRARY-PRECISION COMPUTATIONS

In this section, we connect function spaces with arbitrary-precision computations. We start by summarizing some basic concepts from [ADIW26].

2.1. Computation States Structures. A *computation states structure* is a pair $(L, \mathcal{P})$, where L is a set whose elements we call *states* and $\mathcal{P}$ is a collection of real-valued functions on L that we call *predicates*, also called *features*. For each $P \in \mathcal{P}$, we call the value $P(v)$ the P -th (*observable*) *feature* of v .

A *transition* of a computation states structure $(L, \mathcal{P})$ is a map $f : L \rightarrow L$.

The *type* of a state $v \in L$ is the indexed family³

$$\text{tp}(v) = (P(v))_{P \in \mathcal{P}} \in \mathbb{R}^{\mathcal{P}}$$

of its features.

Intuitively, L is the set of states of a computation, and the predicates $P \in \mathcal{P}$ are primitives that are given and accepted as computable. (I.e., each symbol $P \in \mathcal{P}$ denotes a function $P(\cdot)$ taken as a computational primitive on L .)

We assume that each state $v \in L$ is uniquely characterized by its type $\text{tp}(v)$, so we may identify L with a topological subspace of $\mathbb{R}^{\mathcal{P}}$ and topologized as such (i.e., L is identified with its forward image $\text{tp}[L] \subseteq \mathbb{R}^{\mathcal{P}}$ under the type map). As a topological space, L is thus endowed with the topology of pointwise convergence induced by the product topology of $\mathbb{R}^{\mathcal{P}}$. In particular, for each $P \in \mathcal{P}$, the projection map $v \mapsto P(v)$ is continuous.

Important state spaces include $L = \mathbb{R}^{\mathbb{N}}$ (resp., $L = \mathbb{R}^n$ for some positive integer n), endowed with one predicate $P_i(v) = v_i$ (giving the i -th coordinate of v) for each $i \in \mathbb{N}$ (resp., for each $0 \leq i < n$). State subspaces $L \subseteq \mathbb{R}^{\mathbb{N}}$ and $L \subseteq \mathbb{R}^n$ are of primary interest in this paper. (Any computation states structure having a finite or countable family of predicates is effectively such a subspace.)

Definition 2.1. Given a computation states structure $(L, \mathcal{P})$, any type $\text{tp}(v) \in \mathbb{R}^{\mathcal{P}}$ of a state $v \in L$ will be called a *realized* (state) type. The topological closure of the set of realized types in $\mathbb{R}^{\mathcal{P}}$ (endowed with the pointwise convergence topology) will be called the *space of (state) types* of $(L, \mathcal{P})$, denoted $\mathcal{L}$; elements $\mathbf{v} \in \mathcal{L}$ are called *state types*. Elements $\mathbf{v} \in \mathcal{L} \setminus L$ are *unrealized* state types.

Intuitively, state types capture a notion of “limit state”. For each $P \in \mathcal{P}$, the P -th coordinate map (natural projection) $\mathbb{R}^{\mathcal{P}} \rightarrow \mathbb{R} : \mathbf{v} = (\mathbf{v}_Q)_{Q \in \mathcal{P}} \mapsto \mathbf{v}_P$, gives a continuous extension to $\mathcal{L}$ of the predicate $P(\cdot)$ on L ; abusing notation, the extension will still be denoted $P(\cdot)$ (or just P).

Remark 2.2. One may regard unrealized state types as “hyper-states”; when such a type is obtained as a sequential limit (or ultralimit) of realized state types, it may be called a “deep state”. A comment for readers versed in model theory: at least for the purpose of realizing state types (with parameters in the original states space L), the type space $(\mathcal{L}, \mathcal{P})$ serves as a “poor man’s” ultraproduct of $(L, \mathcal{P})$ wherein, by the very definition of $\mathcal{L}$, each type $\mathbf{v}$ is realized.

2.1.1. Bounds on precision and magnitude. Shards. In order for computation states structures to serve their purpose as models of real-world computing, we must keep in mind that computations involving real-valued quantities can only ever be carried out approximately, to a preset degree of precision (i.e., to within acceptable

³As in Proposition 1.8 above, we regard each predicate on L as named by a specific formal label P (a *predicate symbol*) which is given the semantic meaning of a *bona fide* real function $P(\cdot) : v \mapsto P(v)$ (the predicate proper) in the computation states structure. Such a formal symbol P becomes *de facto* the name of an observable feature having the real value $P(v)$ at any state $v \in L$. Whenever $\mathcal{P}$ is used as an index set (e.g., as in the topological product $\mathbb{R}^{\mathcal{P}}$ below), it is regarded as a set of formal symbols. This syntactic vs. semantic abuse of nomenclature is commonplace in mathematics, but the model-theoretic framework requires contextual awareness of the distinction. To disambiguate when necessary, we shall write $P(\cdot)$, implying the semantic meaning of a real function on L , in contrast to P , meant as a purely syntactic symbol. Nevertheless, by a standard abuse of nomenclature in appropriate contexts, we may also treat P as a function on states (i.e., take P to mean $P(\cdot)$ proper).

error bounds). More subtly, yet equally important in practice, one must also bound the magnitude of real quantities involved in all computations: one cannot possibly store—let alone compute with—arbitrarily large numbers! Thus, although the collection $\mathcal{P}$ of predicates may be infinite (possibly, uncountable even), actual computations involve choices, made at the outset, of bounds for errors and for absolute magnitudes of quantities involved. Heuristically, different parts of a computation may require suitably matching (different) error bounds, as well as dealing with quantities of very different sizes. Focusing for now on the second of these issues, our results shall generally apply to sets of computations for which arbitrary but fixed bounds for each predicate are set in advance. This leads to the concept of *shard* first introduced in [ADIW26], as per the following definition.

Definition 2.3. Fix a computation states structure $\underline{L} = (L, \mathcal{P})$, with $L \subseteq \mathbb{R}^{\mathcal{P}}$ implicitly identified with its type space. A *sizer* is a family $r_{\bullet} = (r_P)_{P \in \mathcal{P}}$ of positive real numbers, indexed by $\mathcal{P}$. Given a sizer $r_{\bullet}$, let $\mathbb{R}[r_{\bullet}] := \prod_{P \in \mathcal{P}} [-r_P, r_P]$ (a compact space), and let the $r_{\bullet}$ -*shard* of the states space L be

$$L[r_{\bullet}] = L \cap \mathbb{R}[r_{\bullet}].$$

For a sizer $r_{\bullet}$, the $r_{\bullet}$ -*type shard* is defined as $\mathcal{L}[r_{\bullet}] = \overline{L[r_{\bullet}]}$ (a closed, hence compact subset of $\mathbb{R}[r_{\bullet}]$).

Let also $\mathcal{L}_{\text{sh}}$ (the space of *shard-supported types*) be the union of all type-shards as the sizer $r_{\bullet}$ varies.

Evidently, $\mathcal{L}_{\text{sh}} \subseteq \mathcal{L}$, although the inclusion may be proper. However, the equality $\mathcal{L}_{\text{sh}} = \mathcal{L}$ holds in the important special case when $\mathcal{P}$ is (at most) countable (see [ADIW26]).

2.2. Compositional Computation Structures.

Definition 2.4. A *compositional computation structure* (CCS) is a triple $(L, \mathcal{P}, \Gamma)$, where

- $(L, \mathcal{P})$ is a computation states structure, and
- $\Gamma \subseteq L^L$ is a semigroup under composition.

Elements of the semigroup Γ are called the *computations* of the structure $(L, \mathcal{P}, \Gamma)$. The state space L is implicitly identified with a subset of the type space $\mathcal{L} \subseteq \mathbb{R}^{\mathcal{P}}$ of $(L, \mathcal{P})$. We also assume that the identity map id on L is an element of Γ (which is thus not merely a semigroup but a monoid of transformations of L).

We topologize Γ as a subset of the topological product L^L , where (as usual) the “exponent” L serves merely as an index set, but the “base” $L \subseteq \mathbb{R}^{\mathcal{P}}$ has the relative product topology (of pointwise convergence). In this manner, the semigroup Γ is identified with a subset of the topological product $\mathcal{L}^L \subseteq (\mathbb{R}^{\mathcal{P}})^L$, which leads to an inclusion $\bar{\Gamma} \subseteq \mathcal{L}^L$. Elements $\xi \in \bar{\Gamma}$ are called (real-valued) *deep computations* or *ultracomputations*.

A collection R of sizers is *exhaustive* if $L = \bigcup_{r_{\bullet} \in R} L[r_{\bullet}]$ (shards $L[r_{\bullet}]$ exhaust L). A transformation $\gamma \in \Gamma$ is *R-confined* if γ restricts to a map $\gamma|_{L[r_{\bullet}]} : L[r_{\bullet}] \rightarrow L[r_{\bullet}]$ (into $L[r_{\bullet}]$ itself) for every $r_{\bullet} \in R$. A subset $\Delta \subseteq \Gamma$ is *R-confined* if each $\gamma \in \Delta$ is.

Unlike its subspace L^L (a topological semigroup), the space $\mathcal{L}^L$ does not have a natural semigroup structure, so ultracomputations are not generally composable. Intuitively, ultracomputations are “final” or “ultimate”.

Proposition 2.5. *If $\Delta \subseteq \Gamma$ is confined by an exhaustive sizer collection, then $\overline{\Delta}$ is a compact subset of $\mathcal{L}_{\text{sh}}^L$.*

Proof. If R confines Δ , choose for each $v \in L$ some $r_{\bullet}^{(v)} \in R$ with $v \in L[r_{\bullet}^{(v)}]$. Then Δ lies in the compact space $\prod_{v \in L} \mathcal{L}[r_{\bullet}^{(v)}] \subseteq \mathcal{L}_{\text{sh}}^L$, which therefore contains $\overline{\Delta}$. $\square$

For a CCS $(L, \mathcal{P}, \Gamma)$, we regard the elements of Γ as “standard” finitary computations, and the elements of $\overline{\Gamma}$, i.e., deep computations, as possibly infinitary limits of standard computations.

Remarks 2.6. (For readers versed in model theory.)⁴

- (1) Compositional computation structures as defined above are not real-valued structures in a literal sense, because there is no intrinsic structural manner to ensure that Γ is realized as a literal subset of the product space L^L . However, in a semantically transparent manner, any CCS as above may alternatively be seen (per its original definition in [ADIW26]) as a real-valued structure $(L, \mathcal{P}, \Gamma, \circ, \text{ev})$, where
 - $(L, \mathcal{P})$ is a computation states structure,
 - $(\Gamma, \circ)$ is a semigroup,
 - $\text{ev} : \Gamma \times L \rightarrow L : (\gamma, v) \mapsto \text{ev}(\gamma, v)$ is a semigroup action of Γ on L (i.e., $\text{ev}(\gamma \circ \delta, v) = \text{ev}(\gamma, \text{ev}(\delta, v))$ for $\gamma, \delta \in \Gamma$ and $v \in L$).

Both definitions are reconciled upon identifying $\gamma \in \Gamma$ with the map $(\gamma, v) \mapsto \text{ev}(\gamma, v)$.

- (2) In the preceding sense of real-valued structures, the class of compositional computation structures (for a fixed predicate symbol collection $\mathcal{P}$) is first-order elementary. The limited notion of state type we have introduced (essentially, quantifier-free state types with parameters of sort L only) may be enriched to state types involving more formulas (e.g., with parameters in Γ , or including quantified formulas), and extended to “transform-types” (i.e., types of sort Γ). In particular, given a structure $\underline{C} = (L, \mathcal{P}, \Gamma, \circ, \text{ev})$, all types are realized in some ultrapower $\underline{C}^{(\mathbf{u})}$ of $\underline{C}$ (constructed in the usual manner detailed in the references above).⁵ This approach to spaces of types was introduced in Banach space theory by Krivine [Kri74, KM81, Kri72].

2.3. Computability of deep computations and the Extendibility Axiom.

Let $(L, \mathcal{P})$ be a computation states structure with *countable* predicate collection $\mathcal{P}$.

- (1) A *computable predicate* is any real function $\varphi : L \rightarrow \mathbb{R}$ whose restriction to an arbitrary shard is uniformly approximable by polynomials (with real coefficients) in the features $P(\cdot)$.
- (2) A transform $f : L \rightarrow \mathcal{L}$ is *computable* if, for each $Q \in \mathcal{P}$, the output feature $Q \circ f : L \rightarrow \mathbb{R}$ is a computable predicate.

It is shown in [ADIW26] that a transform $f : L \rightarrow \mathcal{L}$ is computable, that is, has uniformly polynomially approximable features on shards, if and only if it extends to a continuous map $\tilde{f} : \mathcal{L} \rightarrow \mathcal{L}$. This motivates the following definition.

⁴Refer to [Kei23] where a general framework for real-valued structures is introduced; see also [DnI17, section 6] for a basic tutorial.

⁵By contrast (cf., Remark 2.2), the ultrapower construction is not necessary to realize state types in a computation states structure.

A computation $\gamma \in \Gamma$ is *extendable* if it has some extension $\tilde{\gamma} : \mathcal{L}_{\text{sh}} \rightarrow \mathcal{L}_{\text{sh}}$ such that, for every sizer $r_\bullet$, there is a sizer $s_\bullet$ such that $\tilde{\gamma}|_{\mathcal{L}[r_\bullet]} : \mathcal{L}[r_\bullet] \rightarrow \mathcal{L}[s_\bullet]$ is continuous.

Such an extension is unique if it exists, since it is continuous on each type shard and $\mathcal{L}[r_\bullet]$ is dense in $\mathcal{L}[r_\bullet]$.

Abusing notation, for any $\Delta \subseteq \Gamma$ consisting of extendable computations, the set $\{\tilde{\gamma} : \gamma \in \Delta\}$ will be denoted $\tilde{\Delta}$.

Extendibility Axiom. *Every computation $\gamma \in \Gamma$ is extendable.*

For a more detailed discussion of the Extendibility Axiom, we refer the reader to [ADIW26].

For the rest of the paper, the Extendibility Axiom is imposed on all CCSs.

The following lemma extends deep computations to the type shards; it is used in Theorems 3.1 and 6.7.

Lemma 2.7. *Let R be an exhaustive collection of sizers and let $\Delta \subseteq \Gamma$ be R -confined. Every deep computation $f \in \tilde{\Delta}$ has an extension $\tilde{f} : \mathcal{L}_{\text{sh}} \rightarrow \mathcal{L}_{\text{sh}}$ such that, for every $r_\bullet \in R$, $\tilde{f}$ maps $\mathcal{L}[r_\bullet]$ into itself and $\tilde{f}|_{\mathcal{L}[r_\bullet]} \in \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$.*

Proof. Write $f = \mathcal{U} \lim_i \gamma_i$ with $\gamma_i \in \Delta$ and $\mathcal{U}$ an ultrafilter. Each $v \in \mathcal{L}_{\text{sh}}$ lies in some type shard $\mathcal{L}[r_\bullet]$, and by the Extendibility Axiom all the $\gamma_i(v)$ lie in a compact type shard $\mathcal{L}[s_\bullet]$, so $\tilde{f}(v) := \mathcal{U} \lim_i \gamma_i(v)$ exists, and $\tilde{f}$ extends f . If $r_\bullet \in R$, each γ_i maps $\mathcal{L}[r_\bullet]$ into itself, hence, by continuity, its closure $\mathcal{L}[r_\bullet]$ into itself; so does $\tilde{f}$, and $\tilde{f}|_{\mathcal{L}[r_\bullet]} = \mathcal{U} \lim_i \gamma_i|_{\mathcal{L}[r_\bullet]}$. $\square$

2.4. Stability and register computations. The computable maps are the continuous ones. At the top of the hierarchy of this paper sit the sets of computations all of whose deep computations are computable.

Definition 2.8. A set $\Delta \subseteq \Gamma$ of computations is *stable* if, for every sizer $r_\bullet$, every deep computation in $\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$ is continuous, that is, computable.

This is the class studied in [ADIW26], where it is characterized by Grothendieck's double limit criterion and identified with stability in the sense of model theory.

Many computations of interest are tests: they compute a value from the input and store it, without changing the input. The following construction turns any family of such tests into a CCS.

Definition 2.9. Let X be a compact metrizable space with a countable family $\mathcal{P}_X$ of continuous predicates that separates its points, and let F be a family of continuous functions $X \rightarrow [0, 1]$. The *register CCS* of F has states $L = X \times [0, 1]$, whose points (x, y) are an input x and a register y ; its predicates are those of $\mathcal{P}_X$, applied to x , together with the register predicate $(x, y) \mapsto y$; and its computations are

$$\Gamma_F = \{\gamma_f : f \in F\} \cup \{\text{id}\}, \quad \gamma_f(x, y) = (x, f(x)).$$

Lemma 2.10. *In the register CCS of F , the set Γ_F is a semigroup, the space of types is L itself, and the Extendibility Axiom holds. The deep computations of $\{\gamma_f : f \in F\}$ are the maps $(x, y) \mapsto (x, g(x))$ with g in the pointwise closure $\overline{F}$, and $\gamma_g \mapsto g$ is a homeomorphism onto $\overline{F}$. The set $\{\gamma_f : f \in F\}$ satisfies the NIP if and only if F does.*

Proof. We have $\gamma_f \circ \gamma_g = \gamma_f$, so Γ_F is closed under composition. The predicates are continuous and separate the points of the compact space L , so $v \mapsto \text{tp}(v)$ is a homeomorphism onto the space of types, which is a single shard, and each γ_f is its own continuous extension. A net γ_{f_i} converges pointwise exactly when f_i does, since the first coordinate is constant. For the last statement, composing γ_f with an input predicate gives a function that does not depend on f , and composing it with the register predicate gives f . $\square$

2.5. Two motivating examples. Before developing the classification, we describe two computations whose deep computations behave very differently. Both are treated in detail in Section 4.

Square roots. Newton's method for $p(x) = x^2 - a$, with $a > 0$, acts on the real projective line $\mathbb{RP}^1 = \mathbb{R} \cup \{\infty\}$ by $N_p(x) = (x^2 + a)/(2x)$. Its iterates are continuous, and they converge pointwise to the map $\mathfrak{f}$ that sends every positive number to $\sqrt{a}$, every negative number to $-\sqrt{a}$, and 0 and ∞ to ∞ (Figure 1). The deep computations are the iterates together with $\mathfrak{f}$. The map $\mathfrak{f}$ is discontinuous, so it is not computable and the iterates are not stable; but it is the limit of a *sequence* of computations, and the space of deep computations is countable.

Newton's method on the Riemann sphere. For $\rho(z) = z^3 - 2z + 2$, Newton's method on the Riemann sphere $\hat{\mathbb{C}}$ behaves in three ways at once (Figure 2 in Section 4.1). On the basins of the three roots the iterates converge. On the basin of the cycle $0 \mapsto 1 \mapsto 0$ they do not: the even and the odd iterates converge to two different deep computations. On the common boundary of these basins, a fractal set, the iterates are chaotic, and for the simpler polynomial $z^2 - 1$ we shall see that some deep computations are not even limits of sequences of computations (Proposition 5.6).

The classification of Section 3 is designed to separate such behaviors. Theorem 3.1 characterizes the sets of computations whose deep computations are limits of sequences, as for square roots, and Todorčević's trichotomy measures how complicated the space of such deep computations can be.

3. CLASSIFYING DEEP COMPUTATIONS

3.1. NIP, Rosenthal compacta, and deep computations. This is the main section of the paper. We use topology of function spaces to obtain a new hierarchy of complexity classes for deep computations. The main topological tool is Todorčević's trichotomy for Rosenthal compacta [Tod99]. Our inspiration is the work of Glasner and Megrelishvili on tame semigroups in topological dynamics [GM22].

Theorem 3.1, the cornerstone of the section, extends the results of [ADIW26] from continuous deep computations to deep computations of the first Baire class.

Theorem 3.1. *Let $(L, \mathcal{P}, \Gamma)$ be a compositional computation structure (Definition 2.4) satisfying the Extendibility Axiom, with $\mathcal{P}$ countable. Let R be an exhaustive collection of sizers, and suppose that $\Delta \subseteq \Gamma$ is R -confined. Then the following conditions are equivalent:*

- (i) $\overline{\Delta|_{\mathcal{L}[r_\bullet]}} \subseteq B_1(\mathcal{L}[r_\bullet], \mathcal{L}[r_\bullet])$ for all $r_\bullet \in R$;
- (ii) $\pi_P \circ \Delta|_{\mathcal{L}[r_\bullet]}$ satisfies the NIP for all $P \in \mathcal{P}$ and $r_\bullet \in R$; that is, for all $P \in \mathcal{P}$, $r_\bullet \in R$, $a < b$, $\{\gamma_n\}_{n \in \mathbb{N}} \subseteq \Delta$ there are finite disjoint $E, F \subseteq \mathbb{N}$ such that

$$L[r_\bullet] \cap \bigcap_{n \in E} (\pi_P \circ \gamma_n)^{-1}(-\infty, a] \cap \bigcap_{n \in F} (\pi_P \circ \gamma_n)^{-1}[b, \infty) = \emptyset.$$




FIGURE 1. Newton iterates N_p^n for $p(x) = x^2 - a$ on the real projective line, drawn on the circle with ∞ at both ends of each axis. The continuous maps N_p^n converge pointwise to the deep computation $\mathfrak{f}$, which sends the positive half line to $\sqrt{a}$, the negative half line to $-\sqrt{a}$, and 0 and ∞ to ∞ .

Furthermore, if any (hence all) of the preceding conditions hold, then every deep computation $f \in \overline{\Delta}$ can be extended to a Baire-1 function on shards: there is $\tilde{f} : \mathcal{L}_{sh} \rightarrow \mathcal{L}_{sh}$ such that $\tilde{f}|_{\mathcal{L}[r_\bullet]} \in B_1(\mathcal{L}[r_\bullet], \mathcal{L}[r_\bullet])$ for all $r_\bullet \in R$. In particular, on each shard, every deep computation is the pointwise limit of a countable sequence of computations.

Proof. Since $\mathcal{P}$ is countable, $\mathcal{L}[r_\bullet] \subseteq \mathbb{R}^{\mathcal{P}}$ is Polish. The Extendibility Axiom implies that $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$ is a pointwise bounded set of continuous functions for all $P \in \mathcal{P}$. Hence, Theorem 1.9 and Lemma 1.10 prove the equivalence of (i) and (ii). If (i) holds, the extension $\tilde{f}$ of Lemma 2.7 satisfies $\tilde{f}|_{\mathcal{L}[r_\bullet]} \in \overline{\tilde{\Delta}}|_{\mathcal{L}[r_\bullet]} \subseteq B_1(\mathcal{L}[r_\bullet], \mathcal{L}[r_\bullet])$ for all $r_\bullet \in R$. The assertion that every deep computation is a pointwise limit of a countable sequence of computations follows from the fact that for a Polish space X , every compact subset of $B_1(X)$ is Fréchet-Urysohn (that is, a space where any point in the closure of a set A is the limit of a sequence of points in A ; see Theorem 3F in [BFT78] or Theorem 4.1 in [Deb13]). $\square$

3.2. The Todorčević trichotomy and levels of NIP. Recall from Section 1.4 the notions of hereditarily separable and first countable spaces. Every separable metrizable space is hereditarily separable, and R. Pol proved that every hereditarily separable Rosenthal compactum is first countable (see section 10 of [Deb13]). This suggests the following definition:

Definition 3.2. Let $(L, \mathcal{P}, \Gamma)$ be a CCS satisfying the Extendibility Axiom and R be an exhaustive collection of sizers. Let $\Delta \subseteq \Gamma$ be an R -confined countable set of computations satisfying the NIP on shards and features (condition (ii) in Theorem 3.1). We say that Δ is:

- (i) NIP₁ if $\overline{\tilde{\Delta}}|_{\mathcal{L}[r_\bullet]}$ is first countable for every $r_\bullet \in R$;
- (ii) NIP₂ if $\overline{\tilde{\Delta}}|_{\mathcal{L}[r_\bullet]}$ is hereditarily separable for every $r_\bullet \in R$;


(iii) NIP_3 if $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$ is metrizable for every $r_\bullet \in R$.

Observe that $\text{NIP}_3 \Rightarrow \text{NIP}_2 \Rightarrow \text{NIP}_1 \Rightarrow \text{NIP}$. Moreover, a countable stable set of computations (Definition 2.8) is NIP_3 : on each shard its deep computations form a separable compact subset of a countable product of spaces $C_p(\mathcal{L}[r_\bullet])$, that is, a separable Eberlein compactum, and separable Eberlein compacta are metrizable [Ark92]. The five compacta of Section 1.4 show that the converse implications fail for Rosenthal compacta: $\hat{A}(2^\mathbb{N})$ is separable but not first countable; $\hat{D}(S(2^\mathbb{N}))$ and $\hat{D}(2^\mathbb{N})$ are separable and first countable but not hereditarily separable; and $S(2^\mathbb{N})$ is hereditarily separable but not metrizable. Todorćević proved that these examples are canonical [Tod99].

Theorem 3.3 (Todorćević’s Trichotomy, [Tod99], Theorem 3 in [ADK08]). *Let K be a separable Rosenthal compactum.*

- (i) *If K is hereditarily separable but non-metrizable, then $S(2^\mathbb{N})$ embeds into K .*
- (ii) *If K is first countable but not hereditarily separable, then either $D(2^\mathbb{N})$ or $\hat{D}(S(2^\mathbb{N}))$ embeds into K .*
- (iii) *If K is not first countable, then $A(2^\mathbb{N})$ embeds into K .*

We thus have the following classification for separable Rosenthal compacta:



The three levels can also be read without topology, as statements about how many inputs are needed to tell deep computations apart. Fix $r_\bullet \in R$ and write $K_{r_\bullet} = \overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$. Identify each $f \in K_{r_\bullet}$ with the real-valued function $(P, x) \mapsto P(f(x))$ on $X_{r_\bullet} = \mathcal{P} \times \mathcal{L}[r_\bullet]$, whose points we call *tests*: a test applies the feature P to the output on the input x . The topology of pointwise convergence on $K_{r_\bullet}$ is then the topology of pointwise convergence on tests. A set D of tests *separates* a set of deep computations if any two of them differ on some test in D .

Proposition 3.4. *Let Δ be as in Definition 3.2 and $r_\bullet \in R$.*

- (i) *$K_{r_\bullet}$ is metrizable if and only if one countable set of tests separates $K_{r_\bullet}$.*
- (ii) *$K_{r_\bullet}$ is first countable if and only if every $f \in K_{r_\bullet}$ is countably determined: some countable set D_f of tests separates f from every other member of $K_{r_\bullet}$.*
- (iii) *$K_{r_\bullet}$ is hereditarily separable if and only if there is no set $E \subseteq K_{r_\bullet}$ of cardinality $2^{\aleph_0}$ whose members are finitely separated: for each $f \in E$ there are finitely many tests and $\varepsilon_f > 0$ such that every $g \in E \setminus \{f\}$ differs from f by at least ε_f on one of them.*

Proof. (i) If a countable set D of tests separates $K_{r_\bullet}$, restriction to D is a continuous injection of the compact space $K_{r_\bullet}$ into the metrizable space $\mathbb{R}^D$, hence a

homeomorphism onto its image. Conversely, if $K_{r_\bullet}$ is metrizable, then $K_{r_\bullet} \times K_{r_\bullet}$ is second countable, so the open set $\{(f, g) : f \neq g\}$ is Lindelöf; it is covered by the open sets $\{(f, g) : f(t) \neq g(t)\}$, for tests t , and a countable subcover gives the required set D . (ii) If f is countably determined, $\{f\}$ is a countable intersection of open sets, and in a compact Hausdorff space such a point has a countable base of neighbourhoods. Conversely, if (U_k) is a countable base at f and each U_k contains a basic neighbourhood defined by a finite set F_k of tests, then $D_f = \bigcup_k F_k$ works. (iii) A set is finitely separated exactly when it is a discrete subspace, and, by Theorem 3.1, $K_{r_\bullet}$ is a separable Rosenthal compactum; for such compacta, hereditary separability is equivalent to the absence of discrete subspaces of cardinality $2^{\aleph_0}$, by a theorem of Todorćević [Tod99] (see also [GM22, Theorem 0.4]). $\square$

Thus Δ is NIP_3 when one countable set of tests decides every deep computation, NIP_1 when each deep computation is decided by countably many tests of its own, and NIP_2 when no continuum of deep computations can be told apart one by one with finitely many tests.

Section 5 gives computations at each level of the hierarchy, and shows that every implication between the classes stable, NIP_3 , NIP_2 , NIP_1 and NIP is strict.

The definitions of NIP_1 , NIP_2 and NIP_3 given here are topological; whether they admit a non-topological characterization is Question 7.1.

3.3. The Argyros–Dodos–Kanellopoulos heptachotomy and binary search.

In the separable compacta of Section 1.4, the countable dense subsets are indexed by the binary tree $2^{<\mathbb{N}}$. This indexing suits our computational emphasis: real numbers are represented by infinite binary sequences, i.e., branches of the binary tree, and approximated by finite bitstrings, i.e., its nodes.

Definition 3.5. Let X be a Polish space.

- (1) If I is countable and $\{f_i : i \in I\} \subseteq \mathbb{R}^X$, $\{g_i : i \in I\} \subseteq \mathbb{R}^X$ are two families indexed by I , we say that they are *equivalent* if the map $f_i \mapsto g_i$ extends to a homeomorphism from $\overline{\{f_i : i \in I\}}$ to $\overline{\{g_i : i \in I\}}$.
- (2) If $\{f_t : t \in 2^{<\mathbb{N}}\}$ is a pointwise bounded family, we say that it is *minimal* if for every dyadic subtree $\{s_t : t \in 2^{<\mathbb{N}}\}$ of $2^{<\mathbb{N}}$, $\{f_{s_t} : t \in 2^{<\mathbb{N}}\}$ is equivalent to $\{f_t : t \in 2^{<\mathbb{N}}\}$.

One of the main results in [ADK08] is that, up to equivalence, there are seven minimal families and that for every relatively compact $\{f_t : t \in 2^{<\mathbb{N}}\} \subseteq B_1(X)$ there is a dyadic subtree $\{s_t : t \in 2^{<\mathbb{N}}\}$ such that $\{f_{s_t} : t \in 2^{<\mathbb{N}}\}$ is equivalent to one of the minimal families. We shall describe the seven minimal families next. We follow the notation of [ADK08]. Fix a regular dyadic subtree $T = \{s_t : t \in 2^{<\mathbb{N}}\}$ of $2^{<\mathbb{N}}$ (i.e., a dyadic subtree such that every level of T is contained in a level of $2^{<\mathbb{N}}$) with the property that for all $s, s' \in T$, we have $s \frown 0^\infty \neq s' \frown 0^\infty$ and $s \frown 1^\infty \neq s' \frown 1^\infty$. We use the prefix tests v_t and the threshold tests $f_a^\pm$ of Section 1.4. A pair (f, g) of functions on $2^\mathbb{N}$ is regarded as a single function on the disjoint union $2^\mathbb{N} \sqcup 2^\mathbb{N}$.

- (1) $D_1 = \{\frac{1}{|t|+1} v_t : t \in 2^{<\mathbb{N}}\}$. Its closure is $D_1 \cup \{\mathbf{0}\}$, the one-point compactification of a countable discrete set.
- (2) $D_2 = \{s_t \frown 0^\infty : t \in 2^{<\mathbb{N}}\}$. This is discrete in $\overline{D_2} = 2^{\leq \mathbb{N}}$.
- (3) $D_3 = \{\chi_{[s_t \frown 0^\infty]} : t \in 2^{<\mathbb{N}}\}$. This is discrete in $\overline{D_3} = S(2^\mathbb{N})$.
- (4) $D_4 = \{\chi_{(s_t \frown 1^\infty)} : t \in 2^{<\mathbb{N}}\}$. This is discrete in $\overline{D_4} = S(2^\mathbb{N})$.
- (5) $D_5 = \{v_t : t \in 2^{<\mathbb{N}}\}$. This is discrete in $\overline{D_5} = \hat{A}(2^\mathbb{N})$.

- (6) $D_6 = \{(v_{s_t}, s_t \frown 0^\infty) : t \in 2^{<\mathbb{N}}\}$. This is discrete in $\overline{D_6} = \hat{D}(2^\mathbb{N})$.
 (7) $D_7 = \{(v_{s_t}, \chi_{[s_t \frown 0^\infty)} : t \in 2^{<\mathbb{N}}\}$. This is discrete in $\overline{D_7} = \hat{D}(S(2^\mathbb{N}))$.

Theorem 3.6 (Heptachotomy of minimal families, Theorem 2 in [ADK08]). *Let X be Polish. For every relatively compact $\{f_t : t \in 2^{<\mathbb{N}}\} \subseteq B_1(X)$, there exists $i = 1, 2, \dots, 7$ and a regular dyadic subtree $\{s_t : t \in 2^{<\mathbb{N}}\}$ of $2^{<\mathbb{N}}$ such that $\{f_{s_t} : t \in 2^{<\mathbb{N}}\}$ is equivalent to D_i . Moreover, the families D_i are minimal and mutually non-equivalent.*

The implication of this result for deep computations is the following: the building blocks of the families $D_1, \dots, D_7$ are primitives of binary search on $2^\mathbb{N}$, the space of binary expansions. A node $t \in 2^{<\mathbb{N}}$ is a dyadic interval $[t \frown 0^\infty, t \frown 1^\infty]$, the set of inputs that round to t at precision $|t|$. The prefix test v_t asks whether the input rounds to t ; the point $t \frown 0^\infty$ is the rounded value itself; the comparisons $\chi_{[t \frown 0^\infty}$ and $\chi_{(t \frown 1^\infty}$ ask whether the input lies to the right of the left endpoint, or strictly to the right of the right endpoint, of the interval. Table 1 reads the seven families accordingly.

Family	Computation at node t	Deep computations
D_1	prefix test weighted by $1/(t + 1)$	the test that never succeeds: confidence vanishes with depth
D_2	return the rounded value $t \frown 0^\infty$	the exact input: successive approximation
D_3	compare with the left endpoint	comparison with an arbitrary real, ties broken upward
D_4	compare with the right endpoint	comparison with an arbitrary real, ties broken downward
D_5	prefix test	exact equality, or no match
D_6	prefix test and rounded value	a dictionary lookup that returns its key
D_7	prefix test and comparison	a three-way comparison: below, inside or above

TABLE 1. The minimal families of Argyros, Dodos and Kanellopoulos as primitives of binary search.

In Section 5.4 we define the three-way comparison, one step of a binary search with three outcomes, and show that its closure is the closure of D_7 . By the theorem of Argyros, Dodos and Kanellopoulos, every NIP family of computations indexed by the binary tree contains, along a dyadic subtree, one of these seven primitives.

4. EXAMPLES OF COMPOSITIONAL COMPUTATION STRUCTURES

We now turn from the general theory to examples, and compute the deep computations of five compositional computation structures explicitly: Newton's method on the Riemann sphere for two cubic polynomials (Sections 4.1 and 4.2), Newton's method for square roots on the real line (Section 4.3), and threshold classifiers and prefix tests on the Cantor set (Sections 4.4 and 4.5). Section 5 then places these and other computations at each level of the hierarchy of Section 3.

We begin by formalizing the Newton-Raphson root-finding method as a CCS [Gal00], [BFB16, Section 2.3]. Let $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ be the extended complex plane (the Alexandroff —one-point— compactification of $\mathbb{C}$). We identify $\hat{\mathbb{C}}$ with the unit sphere $S^2 \subseteq \mathbb{R}^3$ (the Riemann sphere) through the inverse stereographic projection $z \mapsto (P_1(z), P_2(z), P_3(z))$, a homeomorphism $\hat{\mathbb{C}} \rightarrow S^2$, where $(P_1(\infty), P_2(\infty), P_3(\infty)) = (0, 0, 1)$ and, for $z \neq \infty$,

$$P_1(z) = \frac{2 \Re z}{|z|^2 + 1}, \quad P_2(z) = \frac{2 \Im z}{|z|^2 + 1}, \quad P_3(z) = \frac{|z|^2 - 1}{|z|^2 + 1}.$$

The extended complex plane $\hat{\mathbb{C}}$ is thus endowed with the real predicates P_1, P_2, P_3 with values in $[-1, 1]$; its topology is metrized by the pullback of the usual Euclidean metric on S^2 .

Newton CCSs: steps and fractals. Let $p = p(z)$ be a non-constant rational function with complex coefficients —i.e., an analytic (meromorphic) map $\hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$. The *Newton method step* (or *Newton map*) for p is the rational map

$$(4.1) \quad N_p : z \mapsto z - \frac{p(z)}{p'(z)}.$$

Successive iterations of the Newton-Raphson method (i.e., compositional powers $N_p^{(n)}$) may be regarded as computations of a CCS whose states space is $L = \hat{\mathbb{C}}$. The natural choice of predicates is $\mathcal{P} = (P_1, P_2, P_3)$, the inverse stereographic projection coordinates. (The real- and imaginary part maps $z \mapsto \Re z$, $z \mapsto \Im z$ are not defined at $z = \infty$.)

Let $\Gamma_p := \{N_p^{(n)} : n \in \mathbb{N}\}$ be the semigroup of iterates of the Newton map (4.1). The CCS $(\hat{\mathbb{C}}, (P_1, P_2, P_3), \Gamma_p)$ is *Newton's method for $p(z)$* (or the *Newton CCS for $p(z)$*). Note that the states space $\hat{\mathbb{C}}$ is compact and the Newton map N_p is continuous thereon, so the Extendibility Axiom is satisfied by Newton CCSs.

Given an initial estimate z_0 , one hopes for the sequence of Newton steps applied to z_0 (namely, $N_p^{(\bullet)}(z_0) := (z_n)_{n \in \mathbb{N}}$, where $z_n := N_p^{(n)}(z_0)$) to converge to a root r of p .

Fatou and Julia sets. Recall that the *Fatou set* $\mathcal{F}_p$ consists of all points that have a neighborhood on which every sequence of iterates of p has a uniformly convergent subsequence; the *Julia set* $\mathcal{J}_p := \hat{\mathbb{C}} \setminus \mathcal{F}_p$ is its complement: it is (necessarily closed and) typically a fractal. The Julia set of the Newton map N_p is called the *Newton fractal* of p .

4.1. Example: Newton's CCS for the cubic $\rho(z) = z^3 - 2z + 2$. The polynomial $\rho(z) = z^3 - 2z + 2$ has Newton map

$$N_\rho : z \mapsto z - \frac{z^3 - 2z + 2}{3z^2 - 2} = \frac{2z^3 - 2}{3z^2 - 2}.$$

Denote the real root of ρ by r_0 , and the roots in the upper and the lower half plane by r_+ and r_- . Starting from $z_0 = 0$, the Newton steps alternate $0, 1, 0, 1, \dots$, and neither 0 nor 1 is a root of ρ .

Let B_0, B_+, B_- be the basins of the roots r_0, r_+, r_- , and let B_{01} be the basin of the attracting cycle $0, 1, 0, 1, \dots$. The Fatou set of N_ρ is $B_{01} \cup B_0 \cup B_+ \cup B_-$, and each of these four sets is nonempty and open. Since B_{01} is nonempty, the iterates $N_\rho^{(n)}$ do not converge pointwise; however, the even and the odd iterates converge

pointwise on B_{01} , to two distinct deep computations f_0 (constant 0 on B_{01}) and f_1 (constant 1 on B_{01}).⁶ Figure 2 shows f_0 and f_1 . On the Newton fractal of ρ , the behavior of the iterates is chaotic.




FIGURE 2. The deep computations of Newton's method for $\rho(z) = z^3 - 2z + 2$: the limits of the even iterates $N_\rho^{(2k)}$ and of the odd iterates $N_\rho^{(2k+1)}$, on the whole plane and near the point 1. Points in the basins of the real root and of the roots in the upper and lower half planes (white circles) are purple, orchid and gold, paler where convergence is slower. The remaining open region is the basin of the superattracting cycle $0 \mapsto 1 \mapsto 0$ (white squares). A point there is lilac if the iterate in question is near 0 and plum if it is near 1, so the two colors exchange between even and odd iterates: the two deep computations agree on the basins of the roots and differ on the basin of the cycle. The boundaries of the colored regions form the Newton fractal of ρ .

4.2. Example: Newton's CCS for 3-cyclotomic $\kappa(z) = z^3 - 1$. To visualize the iterates themselves, rather than their limits, color each z_0 by blending the colors of the roots with relative weights $|z_n - r_i|^{-3}$, where $z_n := N_\rho^{(n)}(z_0)$. This coloring is a continuous function of z_0 , so at first it is blurred; as n increases it sharpens into large regions of nearly constant color, separated by narrow regions where the color varies rapidly, that is, where the coloring is nearly discontinuous. Figure 3 shows this for $\kappa(z) = z^3 - 1$, whose roots are the cube roots of unity $r_0 = 1$ and $r_\pm = \exp(\pm 2\pi i/3)$, and whose Newton map is

$$N_\kappa : z \mapsto z - \frac{z^3 - 1}{3z^2} = \frac{2z^3 + 1}{3z^2}.$$

Unlike N_ρ , the map N_κ has no attracting cycle: the Fatou set is the union of the three basins, and on each of them the iterates converge to the corresponding root.

On the Riemann sphere, Newton's method is chaotic on its Julia set, and this chaos is visible to the predicates P_1, P_2, P_3 : Newton CCSs on the Riemann sphere can fail to be NIP (Proposition 5.6).

⁶The point ∞ is a repelling fixed point of N_ρ : if $|z|$ is large, then $N_\rho(z) = \frac{2}{3}z + O(1)$. Hence ∞ lies on the Newton fractal of ρ , and $f_0(\infty) = \infty = f_1(\infty)$.




FIGURE 3. Newton's method for $\kappa(z) = z^3 - 1$ after one and three iterations, and its limit. The coloring blends the colors of the three cube roots of unity (white circles) according to the distance of $N_\kappa^{(n)}(z)$ to each of them; the limit is constant on each basin and discontinuous across the Newton fractal; in the right panel, points are paler where convergence is slower.

4.3. Example: Square roots. When $p : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$ has real coefficients, it restricts to a map $\mathbb{RP}^1 \rightarrow \mathbb{RP}^1$ on the real-projective line $\mathbb{RP}^1 = \mathbb{R} \cup \{\infty\}$. In this case, the Newton CCS for p may be regarded as having states space $\mathbb{RP}^1$. We shall explain the details in the simplest case of a polynomial $p(x) = x^2 - a$ (real $a > 0$) to illustrate iterative square-root approximations.

Let $\mathbb{R} \rightarrow S^1 : x \mapsto (P_1(x), P_2(x))$ be the (inverse) stereographic projection of the real line onto $S^1 \subseteq \mathbb{R}^2$, i.e.,

$$P_1(x) = \frac{2x}{x^2 + 1}, \quad P_2(x) = \frac{x^2 - 1}{x^2 + 1},$$

extended to a projective transformation of the real projective line $\mathbb{RP}^1 = \mathbb{R} \cup \{\infty\}$ by $\infty \mapsto (P_1(\infty), P_2(\infty)) = (0, 1)$; this gives a compact computation states structure $(\mathbb{RP}^1, (P_1, P_2))$. The Newton map N_p is

$$x \mapsto x - \frac{x^2 - a}{2x} = \begin{cases} \frac{x^2 + a}{2x}, & x \neq 0, \infty; \\ \infty, & x = 0, \infty. \end{cases}$$

These Newton steps converge to one of the square roots $\pm\sqrt{a}$ starting from any initial point $x_0 \neq 0, \infty$. More explicitly, the unique deep equilibrium $\mathfrak{f} : \mathbb{RP}^1 \rightarrow \mathbb{RP}^1$ of (the sequence of iterates of) N_p is given by

$$\mathfrak{f}(x) = \lim_{n \rightarrow \infty} N_p^n(x) = \begin{cases} \sqrt{a}, & x \in (0, \infty); \\ -\sqrt{a}, & x \in (\infty, 0); \\ \infty, & x = 0, \infty. \end{cases}$$

Thus, $\mathfrak{f}$ is discontinuous at $x = 0, \infty$ (Figure 1).

4.4. Example: Threshold classifiers on Cantor's set. The Cantor space $2^{\mathbb{N}}$, with the coordinate predicates $P_n(x) = x_n$, is a computation states structure. Since $2^{\mathbb{N}}$ is closed in $\mathbb{R}^{\mathbb{N}}$, it is its own space of types. We use the notation of Section 1.4.

A *finite-precision threshold classifier* decides whether the input lies at or above a dyadic threshold: for $t \in 2^{<\mathbb{N}}$, it is the threshold test $\chi_{[t \smallfrown 0^\infty]}$. It depends only on the first $|t|$ bits of the input, so it is continuous, and it is a Boolean combination of the coordinates $x_0, \dots, x_{|t|-1}$. We also count the constant $\mathbf{0} = \chi_{(1^\infty)}$, the classifier that never fires, as a finite-precision classifier.

The *threshold ultra-classifiers* are the pointwise limits of finite-precision classifiers. By item (3) of Section 1.4, they are exactly the threshold tests $\chi_{[a]}$ and $\chi_{(a]}$ for $a \in 2^\mathbb{N}$: thresholds increasing to a give $\chi_{[a]}$, and thresholds decreasing to a give $\chi_{(a]}$. Thus the space of ultra-classifiers is the split Cantor set $S(2^\mathbb{N})$. Apart from the thresholds of the form $\chi_{[t \smallfrown 0^\infty]}$ and $\chi_{(t \smallfrown 1^\infty]}$, an ultra-classifier is discontinuous at its threshold a , and, unlike a finite-precision classifier, it is not a Boolean combination of finitely many coordinates. A countable family of simple Boolean tests thus produces, in the limit, a compactum that is not metrizable.

4.5. Example: Prefix tests. On the same computation states structure on the Cantor space, the *prefix test* v_t , for $t \in 2^{<\mathbb{N}}$, asks whether the input begins with t . It is continuous, and it is a Boolean combination of the coordinates $x_0, \dots, x_{|t|-1}$. The *ultra-prefix tests* are the pointwise limits of prefix tests.

Proposition 4.2. *The ultra-prefix tests are the prefix tests v_t , the constant $\mathbf{0}$, and the equality tests δ_a for $a \in 2^\mathbb{N}$. Each of them is a sequential limit of prefix tests, hence Baire-1 on $2^\mathbb{N}$, and the space of ultra-prefix tests is the extended Alexandroff compactification $\hat{A}(2^\mathbb{N})$.*

Proof. For $a \in 2^\mathbb{N}$ we have $v_{a \smallfrown n}(y) = \llbracket y \smallfrown n = a \smallfrown n \rrbracket \rightarrow \llbracket y = a \rrbracket$, so $v_{a \smallfrown n} \rightarrow \delta_a$. The words $1^n 0$ ($n \in \mathbb{N}$) are pairwise incomparable, so at most one of the tests $v_{1^n 0}$ takes the value 1 at any given y , and $v_{1^n 0} \rightarrow \mathbf{0}$.

Conversely, let $\varphi = \lim_i v_{t_i}$ for a net (t_i) of words. Regard finite and infinite words as elements of the compact space $\{0, 1, *\}^\mathbb{N}$, padding finite words with blanks $*$; passing to a subnet, $t_i \rightarrow a$. If a is a finite word, then $t_i = a$ eventually, and $\varphi = v_a$. If $a \in 2^\mathbb{N}$ and $y \neq a$, let k be the first position where y and a differ; eventually $t_i \smallfrown (k+1) = a \smallfrown (k+1)$, so t_i is not a prefix of y and $v_{t_i}(y) = 0$. Hence φ vanishes off a , that is, $\varphi = \delta_a$ or $\varphi = \mathbf{0}$. $\square$

Unlike prefix tests, equality tests are discontinuous. As noted in Section 1.4, the space $\hat{A}(2^\mathbb{N})$ is not first countable: every neighborhood of $\mathbf{0}$ contains all but finitely many of the uncountably many tests δ_a , so no countable family of neighborhoods of $\mathbf{0}$ has intersection $\{\mathbf{0}\}$.

5. COMPUTATIONS ACROSS THE HIERARCHY

In this section we place examples of computations at every level of the hierarchy

$$\text{stable} \Rightarrow \text{NIP}_3 \Rightarrow \text{NIP}_2 \Rightarrow \text{NIP}_1 \Rightarrow \text{NIP},$$

and show that each implication is strict. For most levels we give two kinds of example. The first is a static example from Section 4, turned into a CCS by the register construction of Definition 2.9; its deep computations are one of the five compacta of Section 1.4. The second is a classical iterative algorithm, a single step map applied over and over. For iterative algorithms, the hierarchy coincides with a hierarchy studied in topological dynamics by Glasner and Megrelishvili [GM22], and we begin with this observation. Table 2 at the end of the section summarizes the examples.

5.1. Iterative algorithms and tame dynamics. A *cascade CCS* is a compositional computation structure $(L, \mathcal{P}, \Gamma)$ in which L is a compact metrizable space, $\mathcal{P}$ is a countable family of bounded continuous predicates that separates the points of L , and $\Gamma = \{\gamma^n : n \in \mathbb{N}\}$ is the semigroup of iterates of a single continuous map $\gamma : L \rightarrow L$. Since $\mathcal{P}$ separates points and L is compact, the map $v \mapsto \text{tp}(v)$ is a homeomorphism of L onto the space of types, so L is covered by a single shard, every type is realized, and the Extendibility Axiom holds with $\tilde{\gamma} = \gamma$. The deep computations of a cascade CCS form the pointwise closure

$$E(L, \gamma) = \overline{\{\gamma^n : n \in \mathbb{N}\}} \subseteq L^L,$$

which is the *enveloping semigroup* of the dynamical system (L, γ) introduced by Ellis. A dynamical system is called *tame* if its enveloping semigroup consists of Baire class 1 maps [Köh95, Gla06]. Theorem 3.1 therefore reads as follows.

Proposition 5.1. *A cascade CCS is NIP if and only if the dynamical system (L, γ) is tame. In that case it is NIP_1 , NIP_2 or NIP_3 if and only if $E(L, \gamma)$ is, respectively, first countable, hereditarily separable or metrizable.*

For metrizable systems, Glasner and Megrelishvili call these three classes *tame*₁, *tame*₂ and *hereditarily non-sensitive*, and prove that they are strictly nested [GM22, Proposition 0.6]. The dynamical point of view supplies a criterion for wildness: by a theorem of Kerr and Li [KL07], a tame system given by a homeomorphism has topological entropy zero. An algorithm that produces information at a positive exponential rate, such as a pseudorandom number generator, is wild by design; a numerical method that settles down is tame.

5.2. Stable and metrizable. A *stable example: rotation*. Let $L = \mathbb{T} = \mathbb{R}/\mathbb{Z}$, with the predicates $\cos 2\pi x$ and $\sin 2\pi x$, and let $\gamma(x) = x + \alpha$ with α irrational. A pointwise limit of rotations is a rotation, and the rotations $n\alpha$ are dense, so the deep computations are all the rotations $x \mapsto x + c$, $c \in \mathbb{T}$. They are continuous, so the rotation CCS is stable, and its space of deep computations is the circle group, a metrizable compactum.

NIP₃ but not stable: square roots. The square-root CCS of Section 4.3 has the deep computations $\Gamma \cup \{\mathfrak{f}\}$, a countable compactum, hence metrizable; and $\mathfrak{f}$ is discontinuous at 0 and ∞ . So it is NIP_3 but not stable.

NIP₃ but not stable: the power method. The power method computes a dominant eigenvector of a matrix by repeated multiplication and normalization. Let $A = \text{diag}(\lambda_1, \dots, \lambda_d)$ with $\lambda_1 > \lambda_2 > \dots > \lambda_d > 0$, let $L = \mathbb{RP}^{d-1}$, and let $\mathcal{P}$ consist of the predicates $p_{ij}([v]) = v_i v_j / \|v\|^2$, the entries of the orthogonal projection onto the line $[v]$. The step of the algorithm is $\gamma([v]) = [Av]$. For $[v] \in \mathbb{RP}^{d-1}$, let $i(v)$ be the least index with $v_i \neq 0$, and put $\pi([v]) = [e_{i(v)}]$.

Proposition 5.2. *The deep computations of the power method are $E(L, \gamma) = \{\gamma^n : n \in \mathbb{N}\} \cup \{\pi\}$. Hence the power method is NIP_3 . Its only nontrivial deep computation π is of Baire class 1 and discontinuous, so the power method is not stable.*

Proof. For $i = i(v)$, we have $\lambda_i^{-n} A^n v = v_i e_i + \sum_{j>i} (\lambda_j/\lambda_i)^n v_j e_j \rightarrow v_i e_i$, so $\gamma^n \rightarrow \pi$ pointwise. Every cluster point of $\{\gamma^n\}$ is therefore either some γ^m or π , and $E(L, \gamma)$ is a countable compact Hausdorff space, hence metrizable. The map π is a pointwise limit of continuous maps, and it is discontinuous at every $[v]$ with $i(v) > 1$, since $[v + te_1] \rightarrow [v]$ as $t \rightarrow 0$ while $\pi([v + te_1]) = [e_1] \neq \pi([v])$. $\square$

The deep computation π , which returns *the dominant eigenvector of the input*, is not computable in the sense of Section 2, since it does not extend continuously to the space of types; yet it is the limit of a sequence of computations, and the closure in which it lives is as simple as possible.

5.3. Hereditarily separable but not metrizable. Threshold classifiers. By Lemma 2.10, the finite-precision threshold classifiers of Section 4.4 form a register CCS whose deep computations are the ultra-classifiers. Their space is a copy of $S(2^{\mathbb{N}})$ (Section 4.4), so this CCS is NIP_2 but not NIP_3 . The classifiers are NIP because the sets they define are final segments of the lexicographic order, and no infinite family of final segments is independent.

Drawing a line. Let $0 < \alpha < 1$ be irrational. To draw the line $y = \alpha x + \rho$ on a screen, Bresenham's algorithm [Bre65] moves one pixel to the right at each step and decides whether to move one pixel up. The n -th decision is $b_n(\rho) = \lfloor (n+1)\alpha + \rho \rfloor - \lfloor n\alpha + \rho \rfloor = \llbracket \{n\alpha + \rho\} \in [1 - \alpha, 1) \rrbracket$, where $\{\cdot\}$ denotes the fractional part, and the state of the algorithm is the error term ρ modulo 1. The step $\rho \mapsto \rho + \alpha$ is a rotation of the circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$, but the decision bit is not a continuous function of ρ . The standard remedy [MH40, Lot02] is to take as states the decision sequences themselves.

Let $D = \{-k\alpha : k \in \mathbb{Z}\} \subseteq \mathbb{T}$, a countable dense set containing both endpoints 0 and $1 - \alpha$ of the decision interval. The *split circle* $\mathbb{T}_D$ is obtained from $\mathbb{T}$ by replacing each $t \in D$ by two points $t^- < t^+$, with the circular order topology; it is a compact metrizable space, homeomorphic to the Cantor set. The rotation by α preserves D and lifts to a homeomorphism R of $\mathbb{T}_D$ that preserves the labels $\pm$. The decision interval becomes clopen in $\mathbb{T}_D$, and the coding map $\mathbb{T}_D \rightarrow 2^{\mathbb{Z}}$, $y \mapsto (\llbracket R^n y \in [1 - \alpha, 1) \rrbracket)_{n \in \mathbb{Z}}$, is a homeomorphism onto the *Sturmian shift* X_α that conjugates R to the shift σ . The *Bresenham CCS* has states $L = X_\alpha$, the coordinate predicates, and the semigroup $\Gamma = \{\sigma^n : n \in \mathbb{N}\}$.

For $c \in \mathbb{T}$, let R_c^+ (respectively R_c^-) be the map of $\mathbb{T}_D$ that sends a point over y_0 to $(y_0 + c)^+$ (respectively $(y_0 + c)^-$) when $y_0 + c \in D$, and to $y_0 + c$ otherwise. The *double arrow circle* $\mathbb{D}$ is $\mathbb{T} \times \{-, +\}$ with the lexicographic circular order and its order topology; it is compact, first countable, hereditarily separable and not metrizable [Eng89, 3.10.C].

Theorem 5.3. *The Bresenham CCS is NIP. Its deep computations are*

$$E(X_\alpha, \sigma) = \{\sigma^n : n \in \mathbb{N}\} \cup \{R_c^+, R_c^- : c \in \mathbb{T}\},$$

where each σ^n is isolated and $c^\pm \mapsto R_c^\pm$ is a homeomorphism of $\mathbb{D}$ onto the remaining part. In particular, the Bresenham CCS is NIP_2 but not NIP_3 .

Proof. For NIP , note that the predicates $P_j \circ \sigma^n$ are $\{0, 1\}$ -valued, and that the j -th coordinate of $\sigma^n y$ is 1 exactly when $R^{n+j} y$ lies in the clopen arc $[1 - \alpha, 1)$ of $\mathbb{T}_D$, that is, when y lies in an arc of $\mathbb{T}_D$. A family of arcs of a circularly ordered set has Vapnik–Chervonenkis dimension at most 3, so it has no infinite independent subfamily.

For the deep computations, let γ be the limit of σ^{n_i} along an ultrafilter, and let $c = \lim n_i \alpha$ in $\mathbb{T}$. If n_i is eventually constant, then γ is a power of σ . Otherwise, since the points $n\alpha$ are distinct, the ultrafilter decides whether $n_i \alpha$ approaches c from the right or from the left. In the first case, for every point y over y_0 the points $R^{n_i} y$ over $y_0 + n_i \alpha$ approach $y_0 + c$ strictly from the right, and they converge in

$\mathbb{T}_D$ to $(y_0 + c)^+$ if $y_0 + c \in D$ and to $y_0 + c$ otherwise; so $\gamma = R_c^+$. The second case gives R_c^- . Conversely, every c is approached from both sides by the points $n\alpha$ with $n \geq N$, for every N , so every $R_c^\pm$ is a deep computation, and $R_c^+ \neq R_c^-$, since they differ at the points over $d - c$ with $d \in D$. The same argument shows that $c_i^{\varepsilon_i} \rightarrow c^+$ in $\mathbb{D}$ implies $R_{c_i}^{\varepsilon_i} \rightarrow R_c^+$ pointwise, and symmetrically for c^- ; so $c^\pm \mapsto R_c^\pm$ is a continuous bijection from the compact space $\mathbb{D}$ onto a Hausdorff space, hence a homeomorphism. Finally, fix $d \in D$ and consider the two points $(d - n\alpha)^\pm$ of $\mathbb{T}_D$, which σ^n sends to d^- and d^+ . Every map $R_c^\pm$ sends them to the same point, and so does every pointwise limit of σ^{m_i} with $m_i \rightarrow \infty$, by the argument above. Hence no net of such maps, and no net of maps σ^m with $m \neq n$ and m bounded, converges to σ^n , so σ^n is isolated. Thus $E(X_\alpha, \sigma)$ is the union of a countable set and a copy of $\mathbb{D}$; it is hereditarily separable and not metrizable. $\square$

The two deep computations R_c^+ and R_c^- are the two ways of drawing the line infinitely far away, where the accumulated error has reached c : they agree except at the pixels where the line passes exactly through a corner, and there they apply the two opposite rounding conventions. Figure 4 shows the two conventions.




FIGURE 4. Left: the pixels drawn by Bresenham's algorithm for $\alpha = (\sqrt{5} - 1)/2$, with the decision bits b_n . Right: where the line passes exactly through a pixel corner, the deep computations R_c^+ and R_c^- break the tie in opposite directions; they agree on every other pixel.

The split Cantor set of the threshold classifiers of Section 4.4 is the same phenomenon in a static form.

5.4. First countable but not hereditarily separable. Three-way comparison. For $t \in 2^{<\mathbb{N}}$, let $h_t: 2^{\mathbb{N}} \rightarrow \{0, \frac{1}{2}, 1\}$ take the value 0 below the interval $[t \frown 0^\infty, t \frown 1^\infty]$, the value $\frac{1}{2}$ on it, and 1 above it. This is one step of a binary search with three outcomes, as in a search tree: the input lies below, inside or above the dyadic interval of t . Each h_t is continuous, and its superlevel sets are final segments of the lexicographic order, so the family $\{h_t\}$ is NIP. The map $h_t \mapsto (v_t, f_{t \frown 0^\infty}^+)$ extends to a homeomorphism of the closure of $\{h_t\}$ onto $\hat{D}(S(2^{\mathbb{N}}))$ [ADK08, 4.3.7]. By Lemma 2.10, the register CCS of the three-way comparisons is NIP_1 but not NIP_2 .

Drawing a line in space. Let α and β be irrational numbers in $(0, 1)$ such that $1, \alpha, \beta$ are linearly independent over $\mathbb{Q}$. To draw a line in $\mathbb{Z}^3$ with direction $(1, \alpha, \beta)$,

the three-dimensional Bresenham algorithm runs the planar algorithm in two coordinates at once. Its CCS has states $L = X_\alpha \times X_\beta$, the coordinate predicates of both factors, and the semigroup generated by $\sigma \times \sigma$.

Theorem 5.4. *The three-dimensional Bresenham CCS is NIP_1 but not NIP_2 .*

Proof. Each predicate is a coordinate of one factor, so the CCS is NIP by Theorem 5.3. The deep computations form a closed subspace of $E(X_\alpha, \sigma) \times E(X_\beta, \sigma)$, which is first countable as a product of two first countable spaces; so the CCS is NIP_1 . By Kronecker's theorem, the points $(n\alpha, n\beta)$ with $n \geq N$ are dense in $\mathbb{T}^2$ for every N , so every point (c, c') is approached by them from inside each of the four open quadrants at (c, c') ; arguing as in the proof of Theorem 5.3, every pair $(R_c^\varepsilon, R_{c'}^\varepsilon)$ is a deep computation. Consider the pairs (R_c^+, R_{-c}^-) , for $c \in \mathbb{T}$. In $\mathbb{D} \times \mathbb{D}$, a neighborhood of $(c^+, (-c)^+)$ is $[c^+, (c + \delta)^-] \times [(-c)^+, (-c + \delta)^-]$, and it contains $(s^+, (-s)^+)$ only if $c \leq s < c + \delta$ and $-c \leq -s < -c + \delta$, that is, only if $s = c$. So these pairs form an uncountable discrete subspace, and the space of deep computations is not hereditarily separable. $\square$

The uncountable discrete set consists of the deep computations that break ties upward in both coordinates at points where the two accumulated errors are opposite. Each of the two planar algorithms is NIP_2 , and drawing in two directions at once leaves the class: like the Sorgenfrey plane, a product of two hereditarily separable spaces need not be hereditarily separable. Compare [GM22, Proposition 4.1], where a product of two tame₂ systems is shown not to be tame₂.

5.5. NIP but not first countable. *Prefix tests.* By Lemma 2.10, the prefix tests of Section 4.5 form a register CCS whose deep computations are the ultra-prefix tests, a copy of $\hat{A}(2^\mathbb{N})$ (Proposition 4.2). Any two prefix tests are either nested or disjoint, so no infinite family of them is independent, and the CCS is NIP but not NIP_1 .

Tolerance tests. Floating-point code does not test real numbers for equality; it tests whether $|x - a|$ is below a tolerance. For a rational $q \in [0, 1]$ and $n \geq 1$, let $h_{q,n}(x) = \max(0, 1 - n|x - q|)$, the degree to which x equals q at precision $1/n$. The *tolerance CCS* is the register CCS (Definition 2.9) of the family $\{h_{q,n}\}$ on $X = [0, 1]$: the *tolerance test* $\gamma_{q,n}$ writes $h_{q,n}(x)$ into the register.

Proposition 5.5. *The tolerance CCS is NIP but not NIP_1 . Its deep computations include the exact equality tests $(x, y) \mapsto (x, \llbracket x = a \rrbracket)$ for every real $a \in [0, 1]$, and the test $(x, y) \mapsto (x, 0)$ that never succeeds.*

Proof. By Lemma 2.10, it suffices to consider the functions $h_{q,n}$. For NIP, the sets $\{h_{q,n} \geq b\}$ are intervals, and an independent sequence of the $h_{q,n}$ would give an independent sequence of intervals, which does not exist. Given a , choose rationals $q_k \rightarrow a$ and integers $n_k \rightarrow \infty$ with $n_k|q_k - a| \rightarrow 0$; then $h_{q_k, n_k}(a) \rightarrow 1$ and $h_{q_k, n_k}(x) \rightarrow 0$ for $x \neq a$, so $\llbracket x = a \rrbracket$ is a deep computation. Exact equality tests at distinct points a_k converge pointwise to 0. The pointwise topology on $\{\llbracket \cdot = a \rrbracket : a \in [0, 1]\} \cup \{0\}$ is that of the Alexandroff compactification of a discrete set of size continuum, which is not first countable at 0; hence neither is the space of deep computations. $\square$

Exact equality is thus the deep computation of tolerance testing (Figure 5).




FIGURE 5. Tolerance tests $h_{q,n}$ with $q \rightarrow a$ and $n \rightarrow \infty$, chosen so that $n|q - a| \rightarrow 0$, converge pointwise to the exact equality test $\llbracket x = a \rrbracket$, which is 1 at a and 0 elsewhere.

The failure of first countability has a concrete meaning: a neighborhood of the test that never succeeds is specified by finitely many inputs at which the test must fail, and no countable family of such specifications suffices, because exact tests can succeed at uncountably many inputs.

5.6. Beyond NIP: Newton's method on the Riemann sphere. Newton's method on the Riemann sphere is chaotic on its Julia set, and the chaos is visible to the predicates of the CCS. For a polynomial p of degree $m \geq 2$ with simple roots, the Newton map N_p is a rational map of degree m , whose topological entropy is $\log m > 0$ [Lyu83], and positive entropy is incompatible with tameness for invertible systems [KL07]. The Newton map is not invertible, so we do not claim the general statement here; for square roots the independence can be exhibited by hand.

Proposition 5.6. *Let $p(z) = z^2 - 1$, and let $N(z) = \frac{1}{2}(z + \frac{1}{z})$ be its Newton map on $\hat{\mathbb{C}}$. The functions $P_2 \circ N^{3k}$, $k \in \mathbb{N}$, form an independent sequence: for all finite disjoint sets $E, F \subseteq \mathbb{N}$ there is $z \in \hat{\mathbb{C}}$ with $P_2(N^{3k}z) \geq \frac{1}{2}$ for $k \in E$ and $P_2(N^{3k}z) \leq -\frac{1}{2}$ for $k \in F$. In particular, the Newton CCS of p is not NIP.*

Proof. The Möbius transformation $\phi(z) = (z - 1)/(z + 1)$ conjugates N to the squaring map: $\phi(N(z)) = \phi(z)^2$. It maps the imaginary axis, together with ∞ , onto the unit circle, with $\phi^{-1}(e^{i\theta}) = i \cot(\theta/2)$. Hence, for $z = i \cot(\theta/2)$, we have $N^n(z) = i \cot(2^n\theta/2)$ and

$$P_2(N^n(z)) = \frac{2 \cot(2^{n-1}\theta)}{\cot^2(2^{n-1}\theta) + 1} = \sin(2^n\theta).$$

Write $\theta = 2\pi \sum_{j \geq 1} x_j 2^{-j}$ with bits x_j . The fractional part of $2^{3k}\theta/2\pi$ has binary expansion $0.x_{3k+1}x_{3k+2}x_{3k+3} \dots$. If $(x_{3k+1}, x_{3k+2}, x_{3k+3}) = (0, 1, 0)$, this fractional part lies in $[\frac{1}{4}, \frac{3}{8}]$, so $\sin(2^{3k}\theta) \geq \sin \frac{3\pi}{4} > \frac{1}{2}$; if the block is $(1, 1, 0)$, it lies in $[\frac{3}{4}, \frac{7}{8}]$, so $\sin(2^{3k}\theta) < -\frac{1}{2}$. Choosing the block $(0, 1, 0)$ for $k \in E$ and $(1, 1, 0)$ for $k \in F$ gives the required z . $\square$

The same algorithm is tame on the real line. The Julia set of N is the imaginary axis together with ∞ , which meets $\mathbb{RP}^1$ only at ∞ , and on $\mathbb{RP}^1$ the iterates converge

to the deep computation of Section 4.3. Whether Newton’s method is tame depends on the space of states, not on the formula for the step.

5.7. **Summary.** Table 2 collects the examples of this section.

Level	Static example	Iterative algorithm	Deep computations contain
stable	—	rotation	the circle group
NIP ₃ , not stable	square roots	power method	a countable compactum
NIP ₂ , not NIP ₃	threshold classifiers	Bresenham, plane	$S(2^{\mathbb{N}})$; a double arrow circle
NIP ₁ , not NIP ₂	three-way comparison	Bresenham, space	$\hat{D}(S(2^{\mathbb{N}}))$; $\mathbb{D} \times \mathbb{D}$
NIP, not NIP ₁	prefix tests	—	$\hat{A}(2^{\mathbb{N}})$
	tolerance tests		$A(2^{\mathbb{N}})$
not NIP	—	Newton on $\hat{\mathbb{C}}$	—

TABLE 2. Computations at each level of the hierarchy.

6. RANDOMIZED VERSIONS OF NIP AND MONTE CARLO COMPUTABILITY OF DEEP COMPUTATIONS

Sections 3–5 studied deep computations as *topological* limits: pointwise limits of computations, of the first Baire class on the type shards, and approximated by sequences of computations. That theory rests on the type shards being Polish, and hence on the assumption that the set $\mathcal{P}$ of predicates is countable (it is used in the proof of Theorem 1.9). In this section we drop that assumption and replace topology by measure: instead of asking that a deep computation be of the first Baire class, we ask that it be *universally measurable*, that is, measurable with respect to every Radon probability measure on the type shards, so that it can be evaluated almost surely by random sampling. We call such deep computations Monte Carlo computable. The role of the Bourgain–Fremlin–Talagrand dichotomy is taken by its measure-theoretic form (Theorem 6.5), and that of NIP is shared with Talagrand stability.

The main results of the section are Theorem 6.7, which connects NIP and Monte Carlo computability, and Corollary 6.14, which connects Talagrand stability and Monte Carlo computability.

6.1. NIP and Monte Carlo computability of deep computations. By Fact 1.2, for perfectly normal X , a function f is in $B_1(X, Y)$ if and only if $f^{-1}[U]$ is an F_σ subset of X for every open $U \subseteq Y$. This motivates the following definition.

Definition 6.1. Given a Hausdorff space X and a measurable space (Y, Σ) , we say that $f : X \rightarrow Y$ is *universally measurable* (with respect to Σ) if $f^{-1}(E)$ is μ -measurable for every Radon measure μ on X and every $E \in \Sigma$. When $Y = \mathbb{R}$, we will always take $\Sigma = \mathcal{B}(\mathbb{R})$, the Borel σ -algebra of $\mathbb{R}$.

If X is a compact (Hausdorff) space, then Radon measures on X are finite, and hence, through renormalization, we can regard them as probability measures. We thus have the following remark:

Remark 6.2. If X is compact, then a function $f : X \rightarrow \mathbb{R}$ is universally measurable if and only if $f^{-1}(U)$ is μ -measurable for every Radon probability measure μ on X and every open set $U \subseteq \mathbb{R}$.

Intuitively, a function is universally measurable if it is “measurable no matter which reasonable way you try to measure things on its domain”. The concept of universal measurability emerged from work of Kallianpur and Sazonov, in the late 1950’s and 1960s — with later developments by Blackwell, Darst and others — building on earlier ideas of Gnedenko and Kolmogorov from the 1950s. See [Pap02, Chapters 1 and 2].

Notation 6.3. Following [BFT78], the collection of all universally measurable real-valued functions on X will be denoted by $M_r(X)$. Given a fixed Radon measure μ on X , the collection of all μ -measurable real-valued functions on X will be denoted by $\mathcal{M}^0(X, \mu)$.

In the context of deep computations, we are interested in transition maps of a state space $L \subseteq \mathbb{R}^{\mathcal{P}}$ into itself. In the product space $\mathbb{R}^{\mathcal{P}}$, we have two natural σ -algebras: the Borel σ -algebra, i.e., the σ -algebra generated by open sets in $\mathbb{R}^{\mathcal{P}}$, and the cylinder σ -algebra (i.e., the σ -algebra generated by the sub-basic open sets in $\mathbb{R}^{\mathcal{P}}$, that is, the set $\pi_i^{-1}(U)$ with $U \subseteq \mathbb{R}$ open). In general, the cylinder σ -algebra is strictly smaller; however, when $\mathcal{P}$ is countable, both σ -algebras coincide. We will use the cylinder σ -algebra to define universally measurable maps $f : \mathbb{R}^{\mathcal{P}} \rightarrow \mathbb{R}^{\mathcal{P}}$. The reason for this choice is the following characterization:

Proposition 6.4. *Let X be a Hausdorff space and $Y = \prod_{i \in I} Y_i$ be any product of measurable spaces (Y_i, Σ_i) for $i \in I$. Let Σ_Y be the cylinder σ -algebra generated by the measurable spaces (Y_i, Σ_i) . The following are equivalent for $f : X \rightarrow Y$:*

- (i) $f : X \rightarrow Y$ is universally measurable (with respect to Σ_Y).
- (ii) $\pi_i \circ f : X \rightarrow Y_i$ is universally measurable (with respect to Σ_i) for all $i \in I$.

Proof. (i) $\Rightarrow$ (ii) is clear since the projection maps π_i are measurable and the composition of measurable functions is measurable. To prove (ii) $\Rightarrow$ (i), note that if $C = \prod_{i \in I} C_i$ is a measurable cylinder and J is the finite subset of I such that $C_i \neq Y_i$ for $i \in J$, then, $f^{-1}(C) = \bigcap_{i \in J} (\pi_i \circ f)^{-1}(C_i)$ is universally measurable by assumption. $\square$

The preceding proposition says that we can check measurability of a transition just by checking measurability feature by feature. This is analogous to the Baire class 1 setting (see Proposition 1.8).

We saw in section 1 that, when the set $\mathcal{P}$ of predicates is countable, PAC-learning (or NIP) corresponds to relative compactness in the space of Baire class 1 functions (see Theorems 1.4, 1.6, 1.9). The following result (which does not assume countability of $\mathcal{P}$) gives an analogous characterization of the NIP in terms of universal measurability:

Theorem 6.5 (Bourgain-Fremlin-Talagrand, Theorem 2F in [BFT78]). *Let X be a Hausdorff space and $A \subseteq C(X)$ be pointwise bounded. The following are equivalent:*

- (i) $\overline{A} \subseteq M_r(X)$.
- (ii) For every compact $K \subseteq X$, $A|_K$ satisfies the NIP.
- (iii) For every Radon measure μ on X , A is relatively countably compact in $\mathcal{M}^0(X, \mu)$, i.e., every countable subset of A has a limit point in $\mathcal{M}^0(X, \mu)$.

This result allows us to formalize the concept of a deep computation being *Monte Carlo computable*, which we define below (Definition 6.6). To motivate the definition, let us first recall two facts:

- (1) Littlewood's second principle states that every Lebesgue measurable function is "nearly continuous". The formal statement of this, which is Luzin's theorem, is that if (X, Σ, μ) a Radon measure space and Y is a second-countable topological space (e.g., $Y = \mathbb{R}^{\mathcal{P}}$ with $\mathcal{P}$ countable) equipped with the Borel σ -algebra, then any given $f : X \rightarrow Y$ is measurable if and only if for every $E \in \Sigma$ and every $\varepsilon > 0$ there exists a closed $F \subseteq E$ such that the restriction $f|_F$ is continuous and $\mu(E \setminus F) < \varepsilon$.
- (2) Computability of deep computations is characterized in terms of continuous extendibility of computations. This is at the core of [ADIW26].

These two facts motivate the following definition:

Definition 6.6. Let $(L, \mathcal{P}, \Gamma)$ be a CCS. We say that a transition $f : L \rightarrow L$ is *universally Monte Carlo computable* if and only if there exists an extension $\tilde{f} : \mathcal{L}_{\text{sh}} \rightarrow \mathcal{L}_{\text{sh}}$ of f satisfying the following property: for every sizer $r_{\bullet}$ there is a sizer $s_{\bullet}$ such that the restriction $\tilde{f}|_{\mathcal{L}[r_{\bullet}]} : \mathcal{L}[r_{\bullet}] \rightarrow \mathcal{L}[s_{\bullet}]$ is universally measurable, i.e., $\pi_P \circ \tilde{f}|_{\mathcal{L}[r_{\bullet}]} : \mathcal{L}[r_{\bullet}] \rightarrow [-s_P, s_P]$ is μ -measurable for every Radon probability measure μ on $\mathcal{L}[r_{\bullet}]$ and $P \in \mathcal{P}$.

Note that, since each type-shard $\mathcal{L}[r_{\bullet}]$ is compact, by Remark 6.2, in the context of Monte Carlo computability, we restrict our attention to Radon probability measures instead of general Radon measures.

Theorem 6.7. Let $(L, \mathcal{P}, \Gamma)$ be a CCS satisfying the Extendibility Axiom. Let R be an exhaustive collection of sizers. Let $\Delta \subseteq \Gamma$ be R -confined. If $\pi_P \circ \Delta|_{\mathcal{L}[r_{\bullet}]}$ satisfies the NIP for all $P \in \mathcal{P}$ and all $r_{\bullet} \in R$, then every deep computation in Δ is universally Monte Carlo computable.

Proof. Fix $P \in \mathcal{P}$ and $r_{\bullet} \in R$. By the Extendibility Axiom, $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_{\bullet}]}$ is a set of pointwise bounded continuous functions on the compact set $\mathcal{L}[r_{\bullet}]$. By Lemma 1.10, $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_{\bullet}]}$ has the NIP, hence so does its restriction to every compact $K \subseteq \mathcal{L}[r_{\bullet}]$, and Theorem 6.5 gives $\overline{\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_{\bullet}]}} \subseteq M_r(\mathcal{L}[r_{\bullet}])$. If $f \in \overline{\Delta}$ and $\tilde{f}$ is the extension of Lemma 2.7, then $\pi_P \circ \tilde{f}|_{\mathcal{L}[r_{\bullet}]}$ belongs to this closure. $\square$

Whether the converse of Theorem 6.7 holds is Question 7.2.

6.2. Talagrand stability and Monte Carlo computability of deep computations. There is another notion closely related to NIP, introduced by Talagrand in [Tal84] while studying Pettis integration. Suppose that X is a compact Hausdorff space and $A \subseteq \mathbb{R}^X$. Let μ be a Radon probability measure on X . Given a μ -measurable set $E \subseteq X$, a positive integer k and real numbers $a < b$, let

$$D_k(A, E, a, b) = \bigcup_{f \in A} \{x \in E^{2k} : f(x_{2i}) \leq a, f(x_{2i+1}) \geq b \text{ for all } i < k\}.$$

Definition 6.8. We say that A is *Talagrand μ -stable* if and only if for every μ -measurable set $E \subseteq X$ of positive measure and for every $a < b$ there is $k \geq 1$ such that

$$(\mu^{2k})^*(D_k(A, E, a, b)) < (\mu(E))^{2k},$$

where μ^* denotes the outer measure.

Note that the reference to outer measure in Definition 6.8 is necessary since $D_k(A, E, a, b)$ need not be measurable. The inequality certainly holds when A is a countable set of continuous (or μ -measurable) functions.

Remarks 6.9. Let $X, \mu, A \subseteq \mathbb{R}^X$, and $E \subseteq X$ be as above. Then,

- (1) If $A \subseteq B \subseteq \mathbb{R}^X$, then

$$D_k(A, E, a, b) \subseteq D_k(B, E, a, b).$$

- (2) If $a < a' < b' < b$, then

$$D_k(\bar{A}, E, a, b) \subseteq D_k(A, E, a', b').$$

[*Proof:* If $x \in D_k(\bar{A}, E, a, b)$, then there is $f \in \bar{A}$ such that $f(x_{2i}) \leq a < a'$ and $f(x_{2i+1}) \geq b > b'$ for all $i < k$. By definition of pointwise convergence topology, there exists $g \in A$ such that $g(x_{2i}) < a'$ and $g(x_{2i+1}) > b'$ for all $i < k$. Hence, $x \in D_k(A, E, a', b')$.]

- (3) Any subset of a Talagrand μ -stable set is Talagrand μ -stable. (This is immediate from (1).)
- (4) If A is Talagrand μ -stable, then so is $\bar{A}$. (This is immediate from (2).)

The main result of this section is that deep computations that are limit points of a Talagrand stable set of computations are Monte Carlo computable; this is Corollary 6.14 below. To prove it, we first show that the closure of a Talagrand μ -stable set is μ -measurable (lemma 6.11); but first, we need the following useful characterization of measurable functions (compare with Fact 1.2):

Fact 6.10 (Lemma 413G in [Fre03]). *Suppose that (X, Σ, μ) is a measure space and let $\mathcal{K} \subseteq \Sigma$ be a collection of measurable sets satisfying the following conditions:*

- *(X, Σ, μ) is complete, i.e., for all $E \in \Sigma$ with $\mu(E) = 0$ and $F \subseteq E$ we have $F \in \Sigma$.*
- *(X, Σ, μ) is semi-finite, i.e., for all $E \in \Sigma$ with $\mu(E) = \infty$ there exists $F \subseteq E$ such that $F \in \Sigma$ and $0 < \mu(F) < \infty$.*
- *(X, Σ, μ) is saturated, i.e., $E \in \Sigma$ if and only if $E \cap F \in \Sigma$ for all $F \in \Sigma$ with $\mu(F) < \infty$.*
- *(X, Σ, μ) is inner regular with respect to $\mathcal{K}$, i.e., for all $E \in \Sigma$*

$$\mu(E) = \sup\{\mu(K) : K \in \mathcal{K} \text{ and } K \subseteq E\}.$$

(In particular, if X is compact Hausdorff, $\mathcal{K}$ is the collection of compact subsets of X , μ is a Radon probability measure on X , Σ is the completion of the Borel σ -algebra by μ , and all these conditions hold). Then, $f : X \rightarrow \mathbb{R}$ is measurable if and only if for every $K \in \mathcal{K}$ with $0 < \mu(K) < \infty$ and $a < b$, either $\mu^*(K_{\leq a}) < \mu(K)$ or $\mu^*(K_{\geq b}) < \mu(K)$, where $K_{\leq a} = \{x \in K : f(x) \leq a\}$ and $K_{\geq b} = \{x \in K : f(x) \geq b\}$.

The following technical lemma will be instrumental for proving Proposition 6.13, which, in turn, will yield the main result of the subsection, namely Corollary 6.14

Lemma 6.11. *If A is Talagrand μ -stable, then $\overline{A}$ is also Talagrand μ -stable and $\overline{A} \subseteq \mathcal{M}^0(X, \mu)$.*

Proof. By Remarks 6.9, $\overline{A}$ is Talagrand μ -stable, and hence so is $\{f\}$ for every $f \in \overline{A}$. It therefore suffices to show that if $\{f\}$ is Talagrand μ -stable, then f is μ -measurable. If not, Fact 6.10 gives a set E of positive measure and $a < b$ with $\mu^*(E_{\leq a}) = \mu^*(E_{\geq b}) = \mu(E)$, where $E_{\leq a} = \{x \in E : f(x) \leq a\}$ and $E_{\geq b} = \{x \in E : f(x) \geq b\}$. Then $(E_{\leq a} \times E_{\geq b})^k \subseteq D_k(\{f\}, E, a, b)$ for every $k \geq 1$, and this product has outer measure $(\mu(E))^{2k}$ (see [Tal84]), so $\{f\}$ is not Talagrand μ -stable. $\square$

Definition 6.12. We say that A is *universally Talagrand stable* if A is Talagrand μ -stable for every Radon probability measure μ on X .

We first observe that universal Talagrand stability implies NIP:

Proposition 6.13. *Let X be a compact Hausdorff space and $A \subseteq C(X)$ be pointwise bounded. If A is universally Talagrand stable, then A satisfies the NIP.*

Proof. By Lemma 6.11, $\overline{A} \subseteq \mathcal{M}^0(X, \mu)$ for every Radon probability measure μ on X , that is, $\overline{A} \subseteq M_r(X)$. By Theorem 6.5 (i) $\Rightarrow$ (ii) with $K = X$, A satisfies the NIP. $\square$

Corollary 6.14. *Let $(L, \mathcal{P}, \Gamma)$ be a CCS satisfying the Extendibility Axiom, let R be an exhaustive collection of sizers, and let $\Delta \subseteq \Gamma$ be R -confined. If $\pi_P \circ \Delta|_{L[r_\bullet]}$ is universally Talagrand stable for all $P \in \mathcal{P}$ and all $r_\bullet \in R$, then every deep computation in $\overline{\Delta}$ is universally Monte Carlo computable.*

Proof. This follows directly from Proposition 6.13 and Theorem 6.7. $\square$

In the context of deep computations, we have identified two ways to obtain Monte Carlo computability, namely, NIP/PAC and Talagrand stability. It is natural to ask whether these two notions are equivalent. The following results show that, even in the simple case of countably many computations, this question is sensitive to the set-theoretic axioms. On the one hand, it is consistent (with respect to the standard ZFC axioms of set theory) that these two classes are the same:

Theorem 6.15 (Talagrand, Theorem 9-3-1(a) in [Tal84]). *Let X be a compact Hausdorff space and $A \subseteq M_r(X)$ be countable and pointwise bounded. Assume that $[0, 1]$ is not the union of $< \mathfrak{c}$ closed measure zero sets. If A satisfies the NIP, then A is universally Talagrand stable.*

(The assumption that $[0, 1]$ is not the union of $< \mathfrak{c}$ closed measure zero sets is a consequence of, for example, the Continuum Hypothesis.)

On the other hand, the opposite is also consistent for the particular case of the Lebesgue measure:

Theorem 6.16 (Fremlin, Shelah, [FS93]). *It is consistent with the usual axioms of set theory that there exists a countable pointwise bounded set of Lebesgue measurable functions with the NIP which is not Talagrand stable with respect to Lebesgue measure.*

Notice that the preceding two results apply to sets of measurable functions, a class of functions larger than the class of continuous functions. However, by the Extendibility Axiom, finitary computations are continuous, i.e., if A is a set of computations, then $A \subseteq C_p(X)$. The example of Fremlin and Shelah therefore does not settle the question for computations, and we leave it open (Question 7.3).

7. OPEN QUESTIONS

We collect the questions left open by this paper. The first concerns the hierarchy of Section 3.2, whose levels were defined through the topology of the space of deep computations.

Proposition 3.4 describes the three levels by the number of tests needed to tell deep computations apart, but these descriptions still involve infinitely many tests.

Question 7.1. Is there a finitary, combinatorial characterization of NIP_1 , NIP_2 and NIP_3 , in the spirit of the characterization of NIP by finite patterns in Theorem 3.1?

Theorem 6.7 shows that NIP suffices for Monte Carlo computability; we do not know whether it is necessary.

Question 7.2. Under the assumptions of Theorem 6.7, suppose that every deep computation of Δ is universally Monte Carlo computable. Must $\pi_P \circ \Delta|_{L[r_\bullet]}$ have the NIP for all $P \in \mathcal{P}$ and all $r_\bullet \in R$?

Finally, Section 6 gives two routes to Monte Carlo computability, NIP and universal Talagrand stability, and the results of Talagrand and of Fremlin and Shelah leave open whether they can differ for computations.

Question 7.3. Is it consistent with the usual axioms of set theory that some countable, pointwise bounded set of continuous functions on a compact Hausdorff space satisfies the NIP but is not universally Talagrand stable? Equivalently, can the two routes to Monte Carlo computability of deep computations differ for countable sets of computations?

REFERENCES

- [ADIW26] Samson Alva, Eduardo Dueñez, José Iovino, and Claire Walton. Approximability of deep computations. *Mathematical Structures in Computer Science*, 2026. To appear; arXiv:2409.06064.
- [ADK08] Spiros A. Argyros, Pandelis Dodos, and Vassilis Kanellopoulos. A classification of separable Rosenthal compacta and its applications. *Dissertationes Mathematicae*, 449:1–55, 2008.
- [Ark92] A. V. Arkhangel’skii. *Topological function spaces*, volume 78 of *Mathematics and its Applications (Soviet Series)*. Kluwer Academic Publishers Group, Dordrecht, 1992. Translated from the Russian by R. A. M. Hoksbergen.
- [BFB16] Richard L. Burden, J. Douglas Faires, and Annette M. Burden. *Numerical Analysis*. Cengage Learning, Boston, MA, 10th edition, 2016.
- [BFT78] J. Bourgain, D. H. Fremlin, and M. Talagrand. Pointwise compact sets of Baire-measurable functions. *American Journal of Mathematics*, 100(4):845–886, 1978.
- [BKK] Shaojie Bai, J. Zico Kolter, and Vladlen Koltun. Deep equilibrium models. Preprint. <https://arxiv.org/abs/1909.01377>, <https://implicit-layers-tutorial.org/>.
- [Bre65] Jack E. Bresenham. Algorithm for computer control of a digital plotter. *IBM Systems Journal*, 4(1):25–30, 1965.
- [CRBD] Ricky T. Q. Chen, Yulia Rubanova, Jesse Bettencourt, and David Duvenaud. Neural ordinary differential equations. Preprint. <https://arxiv.org/abs/1806.07366>.
- [Deb13] Gabriel Debs. Descriptive aspects of Rosenthal compacta. In *Recent Progress in General Topology III*, pages 205–227. Springer, 2013.
- [DnI17] Eduardo Dueñez and José N. Iovino. Model theory and metric convergence I: Metastability and dominated convergence. In *Beyond First Order Model Theory, Volume I*, pages 131–187. CRC Press, Boca Raton, FL, 2017.
- [Eng89] Ryszard Engelking. *General topology*, volume 6 of *Sigma Series in Pure Mathematics*. Heldermann Verlag, Berlin, second edition, 1989. Translated from the Polish by the author.

- [Fre03] David H. Fremlin. *Measure Theory, Volume 4: Topological Measure Spaces*. Torres Fremlin, Colchester, UK, 2003.
- [FS93] David H. Fremlin and Saharon Shelah. Pointwise compact and stable sets of measurable functions. *Journal of Symbolic Logic*, 58(2):435–455, 1993.
- [Gal00] A. Galántai. The theory of Newton’s method. *Journal of Computational and Applied Mathematics*, 124(1):25–44, 2000. Numerical Analysis 2000. Vol. IV: Optimization and Nonlinear Equations.
- [Gla06] Eli Glasner. On tame dynamical systems. *Colloquium Mathematicum*, 105(2):283–295, 2006.
- [GM22] Eli Glasner and Michael Megrelishvili. Todorčević’ trichotomy and a hierarchy in the class of tame dynamical systems. *Transactions of the American Mathematical Society*, 375(7):4513–4548, 2022.
- [God80] Gilles Godefroy. Compacts de Rosenthal. *Pacific J. Math.*, 91(2):293–306, 1980.
- [Gro52] A. Grothendieck. Critères de compacité dans les espaces fonctionnels généraux. *Amer. J. Math.*, 74:168–186, 1952.
- [HT19] Haim Horowitz and Stevo Todorčević. Compact sets of baire class one functions and maximal almost disjoint families, 2019.
- [Kei23] H. Jerome Keisler. Model theory for real-valued structures. In José Iovino, editor, *Beyond First Order Model Theory, Volume II*. CRC Press, Boca Raton, FL, 2023.
- [KL07] David Kerr and Hanfeng Li. Independence in topological and C^* -dynamics. *Mathematische Annalen*, 338(4):869–926, 2007.
- [KM81] Jean-Louis Krivine and Bernard Maurey. Espaces de Banach stables. *Israel J. Math.*, 39(4):273–295, 1981.
- [Köh95] Angela Köhler. Enveloping semigroups for flows. *Proceedings of the Royal Irish Academy. Section A*, 95A(2):179–191, 1995.
- [Kri72] J.-L. Krivine. Théorie des modèles et espaces L^p . *C. R. Acad. Sci. Paris Sér. A-B*, 275:A1207–A1210, 1972.
- [Kri74] J.-L. Krivine. Langages à valeurs réelles et applications. *Fund. Math.*, 81:213–253, 1974. Collection of articles dedicated to Andrzej Mostowski on the occasion of his sixtieth birthday, III.
- [Lot02] M. Lothaire. *Algebraic Combinatorics on Words*, volume 90 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 2002.
- [Lyu83] Mikhail Yu. Lyubich. Entropy properties of rational endomorphisms of the Riemann sphere. *Ergodic Theory and Dynamical Systems*, 3(3):351–385, 1983.
- [MH40] Marston Morse and Gustav A. Hedlund. Symbolic dynamics II. Sturmian trajectories. *American Journal of Mathematics*, 62:1–42, 1940.
- [Pap02] E. Pap, editor. *Handbook of measure theory. Vol. I, II*. North-Holland, Amsterdam, 2002.
- [Ros74] Haskell P. Rosenthal. A characterization of Banach spaces containing l^1 . *Proc. Nat. Acad. Sci. U.S.A.*, 71:2411–2413, 1974.
- [RPK19] M. Raissi, P. Perdikaris, and G. E. Karniadakis. Physics-informed neural networks: a deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. *J. Comput. Phys.*, 378:686–707, 2019.
- [Sau72] N. Sauer. On the density of families of sets. *J. Combinatorial Theory Ser. A*, 13:145–147, 1972.
- [She71] Saharon Shelah. Stability, the f.c.p., and superstability; model theoretic properties of formulas in first order theory. *Ann. Math. Logic*, 3(3):271–362, 1971.
- [She72] Saharon Shelah. A combinatorial problem; stability and order for models and theories in infinitary languages. *Pacific J. Math.*, 41:247–261, 1972.
- [She90] Saharon Shelah. *Classification theory and the number of nonisomorphic models*, volume 92 of *Studies in Logic and the Foundations of Mathematics*. North-Holland Publishing Co., Amsterdam, second edition, 1990.
- [Tal84] Michel Talagrand. *Pettis Integral and Measure Theory*, volume 51 of *Memoirs of the American Mathematical Society*. American Mathematical Society, Providence, RI, USA, 1984. Includes bibliography (pp. 220–224) and index.
- [Tka11] Vladimir V. Tkachuk. *A C_p -Theory Problem Book: Topological and Function Spaces*. Problem Books in Mathematics. Springer, 2011.

- [Tod97] Stevo Todorčević. *Topics in Topology*, volume 1652 of *Lecture Notes in Mathematics*. Springer Berlin, Heidelberg, 1997.
- [Tod99] Stevo Todorčević. Compact subsets of the first Baire class. *Journal of the American Mathematical Society*, 12(4):1179–1212, 1999.
- [Tod23] Stevo Todorčević. Dense metrizability. *Annals of Pure and Applied Logic*, 175:103327, 07 2023.
- [Val84] Leslie G. Valiant. A theory of the learnable. *Communications of the ACM*, 27(11):1134–1142, 1984.
- [VC71] Vladimir N Vapnik and Alexey Ya Chervonenkis. On the uniform convergence of relative frequencies of events to their probabilities. *Theory of Probability & Its Applications*, 16(2):264–280, 1971.
- [VC74] Vladimir N Vapnik and Alexey Ya Chervonenkis. *Theory of Pattern Recognition*. Nauka, Moscow, 1974. German Translation: Theorie der Zeichenerkennung, Akademie-Verlag, Berlin, 1979.