COMPLEXITY OF DEEP COMPUTATIONS VIA TOPOLOGY OF FUNCTION SPACES

EDUARDO DUEÑEZ[1] JOSÉ IOVINO[1] TONATIUH MATOS-WIEDERHOLD[2]
LUCIANO SALVETTI[2] FRANKLIN D. TALL[2]

[1]Department of Mathematics, University of Texas at San Antonio
[2]Department of Mathematics, University of Toronto

Abstract. We classify the limits of computations by Todorčević's trichotomy for Rosenthal compacta, we show that its alternatives measure how many observations are needed to tell these limits apart, and we realize every alternative by a classical algorithm. Limits of continuous computations along ultrafilters, the deep computations, need not be continuous. For computation structures with countably many features whose computations extend continuously to the type shards, the completions of prescribed invariant regions of bounded states, we prove that the absence of Shelah's independence property on these regions is equivalent to the requirement that every limit of the extended computations be of the first Baire class on each type shard, and that every such limit is then the limit of a sequence of computations. For a countable family, these limits form on each type shard a separable Rosenthal compactum, which, by Todorčević's trichotomy, is metrizable or contains the split Cantor set, the Alexandroff duplicate of the Cantor set, the extended duplicate of the split Cantor set, or the Alexandroff compactification of a discrete set of the cardinality of the continuum. An observation is a test, a pair of an input and a feature. The power method has a metrizable space of deep computations; Bresenham's line algorithm yields the split Cantor set in the plane and the extended duplicate of the split Cantor set in space; the group generated by the steps of the Euclidean algorithm yields the Alexandroff duplicate and, through exact equality tests, the Alexandroff compactification. A measure-theoretic counterpart, valid for arbitrary families of features, characterizes the absence of independence by the featurewise universal measurability of all limits of the extended computations on the type shards.

Introduction

In this paper we classify the limits of computations by Todorčević's trichotomy for Rosenthal compacta, we show that its alternatives measure how many observations are needed to tell these limits apart, and we realize every alternative by a classical algorithm. The computations act by continuous steps on states whose real-valued features are observed to finite precision, in the compositional computation

Date: November 14, 2026.

2020 *Mathematics Subject Classification.* Primary 68Q15, 03C45, 54H30; Secondary 03C98, 03D15, 05D10, 26A21, 37B05, 54C35, 68T07.

Key words and phrases. Deep computations, limit computations, complexity classes, deep learning, recurrent neural networks, deep equilibrium models, compositional computation structures, independence property, NIP, model theory, Rosenthal compacta, Baire class 1 functions, Todorčević's trichotomy, Argyros–Dodos–Kanellopoulos heptachotomy, Bourgain–Fremlin–Talagrand theorem, tame dynamical systems, infinite Ramsey theory.

This version: November 14, 2026. The most recent version of this paper is available at `https://utsa-quantum-math.github.io/resources/deep-computations-nip/`; it may supersede the version deposited on arXiv.

structures that Alva, Dueñez, Iovino and Walton introduced in the first paper of this series [ADIW26], which is to appear in *Mathematical Structures in Computer Science*. The *type* of a state records the values of all its features, and adjoining the limits of these records gives the *space of types*. The analysis takes place on *shards*, the sets of states on which each feature obeys a prescribed bound.[1] The closure of a shard in the space of types is its *type shard*, which includes its limit states. A *deep computation* is a pointwise limit of computations along an ultrafilter, and an *ultracomputation* is such a limit taken after the computations have been extended to the type shards. Limits of this kind are the outputs of iterative numerical methods, and they enter the recurrent, equilibrium and continuous-depth architectures of deep learning, from recurrent networks to the deep equilibrium models of Bai, Kolter and Koltun [BKK] and the neural ordinary differential equations of Chen, Rubanova, Bettencourt and Duvenaud [CRBD].

The first paper of this series treated the *stable* sets of computations, whose limits on the type shards remain continuous; by Grothendieck's double limit criterion [Gro52], every output feature of such a limit is then a uniform limit of polynomials in the input features. Familiar algorithms leave this regime at once. Newton's method for the square roots of a positive number, acting on the real projective line, has a limit that sends every positive number to the positive root and every negative number to the negative root, and is discontinuous at 0 and at ∞, although it is the limit of a sequence of iterates.

The regularity that survives is that of the first Baire class, whose compact sets of functions, the Rosenthal compacta [Ros77, God80], were classified by Todorčević in 1999 [Tod99]. Glasner and Megrelishvili brought his trichotomy into topological dynamics [GM22]: a countable system of continuous maps of a compact metrizable space is *tame* when every map in the pointwise closure of the system is of the first Baire class [Köh95, Gla06], and they distinguished tame systems by the metrizability, hereditary separability and first countability of their enveloping semigroups, with examples that separate these classes. Their setting is that of compact state spaces on which every type is realized, and the alternative of case (ii) of the trichotomy that they identify in the square of the double arrow space does not occur there (see the footnote to Theorem 4.10). In the present paper the trichotomy becomes a classification of limiting computations on type shards, each of its alternatives receives a meaning in terms of the observations needed to tell limiting computations apart, and each alternative, both alternatives of case (ii) included, is realized by a classical algorithm.

The gateway to the classification is a dividing line of Shelah's classification theory. A formula has the *independence property*, which Shelah isolated in 1971 [She71, Definition 4.1], when, for every natural number n, there are n parameters on which every pattern of truth and falsity of the formula is realized; the same line appeared, between 1968 and 1972, as the Sauer–Shelah dichotomy between polynomial and exponential growth of traces of families of sets [VC68, VC71, Sau72, She72]. For a sequence of real-valued functions, independence means that, for some lower and

[1] Real-world computations take place in shards because a machine represents every quantity with finitely many digits. The IEEE double-precision format, for instance, stores no number larger in magnitude than about $1.8 \cdot 10^{308}$, and a computation whose quantities exceed this bound overflows instead of continuing. A computation therefore commits in advance to a bound on the size of each quantity it handles, and the states it can reach lie in the set on which every feature obeys its bound (see Section 2.1.1).

upper thresholds and for any two disjoint finite sets of indices, some point sends the functions indexed by the first set below the lower threshold and those indexed by the second above the upper one. In 1978 Bourgain, Fremlin and Talagrand characterized the pointwise bounded families of continuous functions on a Polish space whose pointwise closure lies in the first Baire class [BFT78], and on a compact metrizable space their condition is the absence of independent sequences (see Theorem 1.6).

Theorem A (Theorem 3.1)**.** *Let $(L, \mathcal{P}, \Gamma)$ be a compositional computation structure with countably many features whose computations extend continuously to the type shards, and let $\Delta \subseteq \Gamma$ be a set of computations that preserves each shard of an exhaustive collection. Then every ultracomputation of Δ is of the first Baire class on the type shard of each shard of this collection if and only if, for each feature and each shard of the collection, the functions obtained by observing the feature at the outputs of the computations of Δ have no independent sequence. In that case, on each shard of the collection, every deep computation of Δ is the pointwise limit of a sequence of computations of Δ.*

The independence condition, the NIP, is finitary: it concerns finite patterns of high and low output values realized by computations on suitable inputs. For a countable set Δ with the NIP, the ultracomputations on the type shard of a shard of the collection form a separable Rosenthal compactum K, and Todorčević's trichotomy applies to K. We read its alternatives through *tests*: a test is a pair of an input and a feature, and applies the feature to the output on that input, the topology of K being the topology of pointwise convergence on tests. A member of K is *countably determined* when countably many tests separate it from every other member of K, and a subset of K is *finitely separated* when each member is separated from all the others, by a positive margin, on finitely many tests.

Theorem B (Theorem 3.3 and Proposition 3.4)**.** *Let Δ be a countable set of computations as in Theorem A with the NIP, and let K be the compactum of its ultracomputations on the type shard of a shard of the collection.*

(i) The compactum K is metrizable if and only if a single countable set of tests separates all its members.

(ii) The compactum K is hereditarily separable if and only if no finitely separated subset of K has the cardinality of the continuum; if K is hereditarily separable and not metrizable, then K contains a copy of the split Cantor set $S(2^{\mathbb{N}})$.

(iii) The compactum K is first countable if and only if every member of K is countably determined; if K is first countable and not hereditarily separable, then K contains a copy of the Alexandroff duplicate $D(2^{\mathbb{N}})$ of the Cantor set or of the extended duplicate $\hat{D}(S(2^{\mathbb{N}}))$ of the split Cantor set.

(iv) If K is not first countable, then K contains a copy of the Alexandroff compactification $A(2^{\mathbb{N}})$ of a discrete set of the cardinality of the continuum.

Accordingly, we call Δ $\mathrm{NIP}^{\mathrm{met}}$, $\mathrm{NIP}^{\mathrm{hs}}$ or $\mathrm{NIP}^{\mathrm{fc}}$ when its compacta of ultracomputations are metrizable, hereditarily separable or first countable, respectively, and obtain the hierarchy

$$\text{stable} \Rightarrow \mathrm{NIP}^{\mathrm{met}} \Rightarrow \mathrm{NIP}^{\mathrm{hs}} \Rightarrow \mathrm{NIP}^{\mathrm{fc}} \Rightarrow \mathrm{NIP}.$$

On compact state spaces this is the hierarchy of Glasner and Megrelishvili (see Proposition 4.1). The next theorem shows that every implication is strict and that

each canonical compactum of Theorem B arises from a classical algorithm. The algorithms are among the most widely used in computing. Newton's method and the power method, which underlies the ranking of web pages, are staples of scientific computation; Bresenham's algorithm, devised at IBM for a digital plotter, draws the lines of raster displays and printers; and the Euclidean algorithm, through continued fractions, yields the best rational approximations of real numbers.

Theorem C. *Every leaf of Figure 1 is realized by the algorithm placed at the leaf; more precisely:*

- *(i) Rotations of the circle by irrational angles are stable, and the power method, Newton's method for square roots on the real projective line and the logistic map $x \mapsto rx(1-x)$ for $1 < r < r_\infty$, below the Feigenbaum parameter r_∞, are NIP^{met} but not stable (Propositions 4.2, 4.3, 4.4 and 4.23).*
- *(ii) Bresenham's line algorithm in the plane is NIP^{hs} but not NIP^{met}: its deep computations other than the iterates form a double arrow circle, and the space of all its deep computations contains a copy of $S(2^{\mathbb{N}})$ whose isolated points are selected iterates (Theorem 4.5 and Proposition 4.6).*
- *(iii) Bresenham's line algorithm in space is NIP^{fc} but not NIP^{hs}, and its deep computations contain a copy of $\hat{D}(S(2^{\mathbb{N}}))$ and none of $D(2^{\mathbb{N}})$ (Theorem 4.10).*
- *(iv) A countable family $\Delta_{\mathfrak{C}}$ of elements of the group generated by the steps $x \mapsto x+1$ and $x \mapsto 1/x$ of the Euclidean algorithm is NIP^{fc} but not NIP^{hs}, and the closure of $\Delta_{\mathfrak{C}}$ is homeomorphic to the compactum $\hat{D}(2^{\mathbb{N}})$ that contains $D(2^{\mathbb{N}})$ (Proposition 4.15).*
- *(v) The group generated by the two steps of the Euclidean algorithm, acting on the real projective line, is NIP but not NIP^{fc}; its deep computations include exact equality tests, which, together with the constant map that they share as a baseline, form a copy of $A(2^{\mathbb{N}})$, and its space of deep computations contains a copy of $D(2^{\mathbb{N}})$ and none of $S(2^{\mathbb{N}})$ (Corollary 4.14 and Proposition 4.15).*

The examples show why the alternatives differ. In the power method a single test, one feature evaluated at one input, already separates all the deep computations. Bresenham's algorithm rounds at each step, and its limiting computations, which draw the line infinitely far away, come in pairs that agree everywhere except where the line passes exactly through a pixel corner, at which the two members of the pair apply opposite rounding conventions; any countable set of tests misses the corners of some pair, and the space is not metrizable, although each limiting computation is still determined by countably many tests of its own. In space, the limiting computations that break ties upward in both coordinates and whose two accumulated errors are opposite form a finitely separated set of the cardinality of the continuum. The Euclidean constructions turn on a baseline: an exact equality test sends one exceptional input to one value and every other input to a common baseline value. When the baseline is the same for uncountably many exceptional inputs, as in the full group, countably many tests cannot tell an equality test from the constant baseline map, and first countability fails; when the baseline moves with the exceptional input, as in $\Delta_{\mathfrak{C}}$, a few tests determine each limit, and the Alexandroff duplicate appears. The space of states belongs to the classification: Newton's method for the square roots of 1, which is NIP^{met} on the real projective line, has the independence property on the Riemann sphere.




FIGURE 1. Todorčević's trichotomy and its realizations.

Beyond the hierarchy lie the chaotic iterations: Newton's method on the Riemann sphere, the iteration of every rational map of degree at least two, the doubling map of the circle, the logistic family beyond the Feigenbaum parameter, and the shift of the Thue–Morse subshift all have the independence property (Propositions 4.20, 4.21, 4.23 and A.1). Single iterated maps realize the metrizable alternative, the split Cantor set and the extended duplicate of the split Cantor set; whether a single iterated map realizes the Alexandroff duplicate, or the failure of first countability, remains open (Questions 6.3 and 6.2), as does a finitary characterization of the three levels (Question 6.1).

Todorčević's trichotomy classifies the space of limits; the heptachotomy of Argyros, Dodos and Kanellopoulos [ADK08], which retains the indexing of a family by the binary tree, classifies the routes by which finite computations approach those limits, following a branch of the tree or departing from it below or above. The algorithms of Theorem C realize all seven canonical routes (Theorem 4.18). A convergent method erases the route, and a rotation recovers the point approached. Bresenham's algorithm in the plane records the side from which the point is approached, and the Euclidean algorithm decides an equality against a baseline that is fixed or, in the family $\Delta_{\mathfrak{C}}$, moves with the exceptional input. Bresenham's algorithm in space, finally, tells the two departures apart. The two alternatives of item (iii) of Theorem B are separated in exactly this way: departing from a branch on either side leads to one common limit for the family $\Delta_{\mathfrak{C}}$ and to two distinct limits for the selected iterates of Bresenham's algorithm in space (Section 4.6).

Finally, we pass from topology to measure, dispensing with the countability of the features. The measure-theoretic form of the theorem of Bourgain, Fremlin and Talagrand characterizes the NIP by the universal measurability of all ultracomputations on the type shards (Theorem 5.7). For every Radon probability measure on a type shard and every output feature, Luzin's theorem and the Stone–Weierstrass theorem then give uniform polynomial approximation in the input features outside a set of arbitrarily small measure; we call such deep computations *universally Monte Carlo computable*, and we show that, for countable sets of computations, universal Talagrand stability coincides with the NIP (Corollary 5.17).

The paper presupposes no acquaintance with advanced topology, model theory or the theory of computation; the necessary background is assembled in Section 1, beginning from undergraduate topology and measure theory.

A map of the classification. Figure 1, which accompanies Theorem C, draws Theorem B as three questions about the compactum of limits, with the algorithm that realizes each answer and, at each leaf, what tests can tell apart. Figure 2 draws the seven canonical families as patterns of the three routes, with the selected computations that realize them. The first figure sorts algorithms by the topology of the whole space of limits, and the second sorts selected families by the way in which they approach their limits.

Family	Level	Routes	Algorithm	What the limit retains
D_1	$\mathrm{NIP}^{\mathrm{met}}$		Newton's method for $\sqrt{a}$	nothing: one limit for all routes
D_2	$\mathrm{NIP}^{\mathrm{met}}$		rotation	the point approached
D_3	$\mathrm{NIP}^{\mathrm{hs}}$		Bresenham, plane	the side; the branch shares the lower one
D_4	$\mathrm{NIP}^{\mathrm{hs}}$		Bresenham, plane	the side; the branch shares the upper one
D_5	NIP		Euclid, fixed baseline	an equality; one baseline for all b
D_6	$\mathrm{NIP}^{\mathrm{fc}}$		Euclid, moving baseline	an equality; the baseline names b
D_7	$\mathrm{NIP}^{\mathrm{fc}}$		Bresenham, space	all three routes apart

FIGURE 2. The seven canonical families and their realizations. Arrows: departing below, along the branch, departing above; arrows ending at one dot share a limit, and an open dot is the same for every branch.

Organization of the paper. In Section 1 we present the topological and combinatorial preliminaries, including five canonical Rosenthal compacta (Section 1.4) and the role of NIP in model theory, combinatorics and learning (Section 1.5). In Section 2 we recall compositional computation structures and their stable sets of

computations; the section ends with two motivating examples, Newton's method for square roots and Newton's method on the Riemann sphere, whose different behaviors the classification is designed to separate (Section 2.5). Section 3 contains Theorems A and B, the hierarchy of levels and the seven canonical families of binary search. In Section 4 the alternatives of Todorčević's trichotomy are realized by classical algorithms, and chaotic iterations beyond the classification are described. In Section 5 we replace topology by measure and study deep computations that are universally measurable on each type shard. Section 6 contains the questions that remain open. In Appendix A we prove that the shift on the Thue–Morse subshift generates an action of $\mathbb{Z}$ that is not tame, a fact used in Section 4.7.

Contents

Introduction 1
A map of the classification 6
Organization of the paper 6
1. Topological preliminaries: From continuity to Baire class 1 8
1.1. From Rosenthal's dichotomy to the Bourgain–Fremlin–Talagrand dichotomy to Shelah's NIP 10
1.2. Rosenthal compacta 12
1.3. The special case $B_1(X, \mathbb{R}^{\mathcal{P}})$ with $\mathcal{P}$ countable 12
1.4. Five canonical Rosenthal compacta 14
1.5. NIP in model theory, combinatorics and learning: polynomial versus exponential growth 16
Historical remarks 18
2. Compositional computation structures: A structural approach to arbitrary-precision computations 18
2.1. Computation states structures 19
2.2. Compositional Computation Structures 20
2.3. Computability of deep computations and the Extendibility Axiom 22
2.4. Stability 23
2.5. Two motivating examples 24
3. Todorčević's trichotomy classifies deep computations 24
3.1. NIP, Rosenthal compacta, and deep computations 25
3.2. Todorčević's trichotomy and levels of NIP 26
3.3. Canonical families and binary search 28
4. Todorčević's trichotomy is realized by classical algorithms 31
4.1. Iterative algorithms and tame dynamics 31
4.2. Convergent methods and rotations have metrizable spaces of limits 33
4.3. Rounding produces the split Cantor set 36
4.4. Two rounding procedures produce the extended duplicate of the split Cantor set 39
4.5. Continued fractions produce exact equality tests and the Alexandroff duplicate 42
4.6. The same algorithms realize the seven canonical families 50
4.7. Chaotic iteration leads beyond the classification 53
4.8. Summary 60

5. Randomized versions of NIP and Monte Carlo computability of deep computations 60
5.1. NIP and Monte Carlo computability of deep computations 61
5.2. Talagrand stability and Monte Carlo computability of deep computations 64
6. Open questions 68
Appendix A. The Thue–Morse system is not tame 69
References 72

1. Topological preliminaries: From continuity to Baire class 1

In this section we assemble the topology of function spaces on which the rest of the paper rests. The paper pursues two themes: the approximation of a deep computation by a sequence of computations, and the complexity of the set of all deep computations. Deep computations are pointwise limits of continuous maps along ultrafilters, and both themes therefore belong to the theory of spaces of functions with the topology of pointwise convergence. We first recall Polish spaces, Baire category and the first Baire class. We then turn to Rosenthal's dichotomy and to the Bourgain–Fremlin–Talagrand dichotomy, from which the no-independence property (NIP) emerges. The Rosenthal compacta come next, and five of them serve as yardsticks for the classification. The section closes with the role of the NIP in model theory, in combinatorics and in statistical learning. All spaces are assumed to be Hausdorff unless explicitly stated otherwise.

We write $\mathbb{N}$ for the set $\{0, 1, 2, \dots\}$ of natural numbers, so that the semigroups of iterates considered below contain the identity. A countable union of closed subsets of a topological space is called an F_σ set, and a countable intersection of open subsets is called a G_δ set. In a metric space every open set is an F_σ set, and, equivalently, every closed set is a G_δ set.

A *Polish space* is a separable topological space whose topology is induced by a complete metric. Countable products of Polish spaces, with the product topology, are Polish. Besides the real line, the classical examples include the Cantor space $2^{\mathbb{N}}$ of all infinite binary sequences, the Baire space $\mathbb{N}^{\mathbb{N}}$ of all infinite sequences of natural numbers, and the space $\mathbb{R}^{\mathbb{N}}$ of all sequences of real numbers.

Fact 1.1. [Eng89, 4.3.24] *Let X be a Polish space. A subset of X is Polish in the subspace topology if and only if the subset is a G_δ set. In particular, every closed subset and every open subset of X is a Polish space.*

A subset A of a topological space X is *nowhere dense* if the closure of A has empty interior. The set A is of the *first category*, or *meager*, if A is contained in a countable union of closed nowhere dense sets, and of the *second category* otherwise. The complements of meager sets are called *comeager*. Since the closure of a nowhere dense set is nowhere dense, the meager subsets of X form the σ-ideal generated by the nowhere dense subsets of X. A topological space X is *Baire* if every comeager subset of X is dense, or, equivalently, if every countable intersection of dense open

sets is dense. The Baire category theorem states that compact Hausdorff spaces and completely metrizable spaces, Polish spaces in particular, are Baire.[2]

For topological spaces X and Y, we denote by $\mathrm{C_p}(X,Y)$ the set of all continuous functions $f\colon X \to Y$ with the topology of pointwise convergence, and we abbreviate $\mathrm{C_p}(X,\mathbb{R})$ to $\mathrm{C_p}(X)$. The topology of pointwise convergence is the one that $\mathrm{C_p}(X,Y)$ inherits as a subspace of the product Y^X, in which X serves merely as an index set.

A function $f\colon X \to Y$ between topological spaces is of *Baire class 1* if f is the pointwise limit of a sequence $(f_n)_{n\in\mathbb{N}}$ of continuous functions, that is, $f(x) = \lim_{n\to\infty} f_n(x)$ for all $x \in X$. When Y is metrizable, an elementary argument shows that a function of Baire class 1 is continuous at every point outside a set of the first category. We denote by $\mathrm{B_1}(X,Y)$ the set of functions $X \to Y$ of Baire class 1 with the topology of pointwise convergence, and we write $\mathrm{B_1}(X)$ for $\mathrm{B_1}(X,\mathbb{R})$; the inclusions $\mathrm{C_p}(X,Y) \subseteq \mathrm{B_1}(X,Y) \subseteq Y^X$ are topological. Relatively compact subsets of $\mathrm{C_p}(X)$ and of $\mathrm{B_1}(X)$, especially for Polish X, are central objects of study in functional analysis and in topological dynamics.

A topological space X is *perfectly normal* if X is normal and every closed subset of X is a G_δ set (equivalently, every open subset of X is an F_σ set). Every metrizable space (hence, every Polish space) is perfectly normal.

The following characterization of Baire class 1 functions is classical; for a proof, see [Tod97, Section 10].

Fact 1.2 (Lebesgue, Hausdorff, Baire[3])**.** *If X is perfectly normal, then the following conditions are equivalent for a function $f : X \to \mathbb{R}$:*

(1) *The function f is of Baire class 1.*
(2) *The preimage $f^{-1}[U]$ is an F_σ subset of X whenever $U \subseteq \mathbb{R}$ is open.*

If X is Polish, then (1) and (2) are equivalent to:

(3) *For every nonempty closed $F \subseteq X$, the restriction $f|_F$ has a point of continuity.*

Moreover, if X is Polish and $f \notin \mathrm{B_1}(X)$, then there exist nonempty countable $D_0, D_1 \subseteq X$ and reals $a < b$ such that

$$D_0 \subseteq f^{-1}(-\infty, a], \quad D_1 \subseteq f^{-1}[b,\infty), \quad \overline{D_0} = \overline{D_1}.$$

---

[2]More generally, any Čech-complete space is Baire [Eng89, Theorem 3.9.3].

[3]The implication (1)⇒(2) holds in every topological space. Indeed, if f is the pointwise limit of a sequence $(f_n)_{n\in\mathbb{N}}$ of continuous functions and a is a real number, then

$$f^{-1}(a,\infty) = \bigcup_{k\geq 1} \bigcup_{m\in\mathbb{N}} \bigcap_{n\geq m} f_n^{-1}[a + 1/k, \infty)$$

is a countable union of closed sets. The same is true of $f^{-1}(-\infty, a)$; since every open subset of $\mathbb{R}$ is a countable union of intervals $(a,b) = (a,\infty) \cap (-\infty, b)$, condition (2) follows. The reverse implication, (2)⇒(1), was established by Lebesgue for real-valued functions of finitely many real variables [Leb04] and extended by Hausdorff to real-valued functions on metric spaces [Kec95, Theorem 24.10]. The proof of (2)⇒(1) requires of X only that X be normal and that every open subset of X be an F_σ set, and these two requirements are precisely the defining properties of a perfectly normal space. The implication (1)⇒(3) holds whenever X is Polish. If f satisfies (1) and F is a nonempty closed subset of X, then the restriction $f|_F$ is of Baire class 1. As noted before Fact 1.2, the points of discontinuity of $f|_F$ therefore form a meager subset of F. Being closed in a Polish space, the set F is itself Polish, hence a Baire space, which the meager set of points of discontinuity cannot exhaust. Consequently, the restriction $f|_F$ has a point of continuity.

1.1. **From Rosenthal's dichotomy to the Bourgain–Fremlin–Talagrand dichotomy to Shelah's NIP.** The question of when a deep computation is the limit of a *sequence* of computations is answered by Rosenthal's dichotomy and by the Bourgain–Fremlin–Talagrand dichotomy, which we recall in this subsection; from the two dichotomies the NIP emerges in the theory of function spaces. In a metrizable space every point of the closure of a set is the limit of a sequence of points of the set. In spaces of functions with the topology of pointwise convergence, by contrast, the closure of a set need not admit this sequential description, and the two dichotomies decide when the description holds.

On the Cantor space $X = 2^{\mathbb{N}}$, the n-th coordinate map $\pi_n \colon X \to \{0,1\}$ is continuous for each $n \in \mathbb{N}$; however, the sequence $(\pi_n)_{n\in\mathbb{N}}$ has *no* convergent subsequence in $\mathbb{R}^X$ (see [Tod97, Chapter 1.1]). In a sense, this sequence exhibits the most extreme failure of sequential convergence. Indeed, the closure of $\{\pi_n : n \in \mathbb{N}\}$ in $\{0,1\}^X$, or in $\mathbb{R}^X$, is homeomorphic to the Stone–Čech compactification $\beta\mathbb{N}$ of the discrete space of natural numbers. The space $\beta\mathbb{N}$ has cardinality $2^{2^{\aleph_0}}$, the largest possible for a separable Hausdorff space, and the points of $\beta\mathbb{N}$ correspond to the ultrafilters on $\mathbb{N}$.[4]

In 1974 Haskell Rosenthal proved the following dichotomy for pointwise bounded sequences of continuous functions on a Polish space. His theorem shows that for these sequences a copy of $\beta\mathbb{N}$, such as the closure of the coordinate maps, is the only obstruction to the existence of a convergent subsequence.

Theorem 1.3 (Rosenthal's dichotomy [Ros74])**.** *If X is a Polish space and $(f_n)_{n\in\mathbb{N}}$ is a pointwise bounded sequence in $\mathrm{C_p}(X)$, then the sequence (f_n) has either a convergent subsequence or a subsequence whose closure in $\mathbb{R}^X$ is homeomorphic to $\beta\mathbb{N}$.*

Rosenthal's dichotomy grew out of *Rosenthal's ℓ_1-theorem*: every bounded sequence in a Banach space has a subsequence that is either weakly Cauchy or equivalent to the unit vector basis of ℓ_1, the space of absolutely summable sequences. In particular, a Banach space contains no isomorphic copy of ℓ_1 if and only if every bounded sequence in the space has a weakly Cauchy subsequence.

The Bourgain–Fremlin–Talagrand theorem, which concerns sets rather than sequences, characterizes the relative compactness in the larger space $\mathrm{B}_1(X)$ of a pointwise bounded set of continuous functions on a Polish space X. For such a set, relative compactness in $\mathrm{B}_1(X)$ means that every function in the pointwise closure of the set is of Baire class 1. The theorem is a counterpart of Rosenthal's dichotomy, although the usual statement is an equivalence rather than a dichotomy (see Theorem 1.4 below).

The reverse implication (3)⇒(1) is due to Baire, who proved the implication in his thesis for real-valued functions of real variables [Bai99]; for Polish spaces, see [Kec95, Theorem 24.15]. Finally, the last assertion of Fact 1.2 follows from the implication (3)⇒(1) together with the Baire category theorem.

[4]Addition on $\mathbb{N}$ extends to an associative operation on $\beta\mathbb{N}$, whose idempotent elements, the idempotent ultrafilters, exist by the Ellis–Numakura lemma. Alva, Dueñez, Iovino and Walton [ADIW26] use idempotent ultrafilters to prove the existence of *deep equilibria* in compositional computation structures. Deep equilibria are limits of the iterates of a single computation, taken along an ultrafilter on $\mathbb{N}$, that are idempotent under composition.

The statement of the theorem requires the following notation. For $f \in \mathbb{R}^X$ and a real number a, define

$$X^f_{\le a} \coloneqq f^{-1}(-\infty, a], \qquad X^f_{\ge a} \coloneqq f^{-1}[a, +\infty).$$

For a set $A \subseteq \mathbb{R}^X$ and a real number a, define, by a slight abuse of notation,

$$X^A_{\le a} \coloneqq \bigcap_{f\in A} X^f_{\le a} = \{x \in X : f(x) \le a \text{ for all } f \in A\},$$
$$X^A_{\ge a} \coloneqq \bigcap_{f\in A} X^f_{\ge a} = \{x \in X : f(x) \ge a \text{ for all } f \in A\};$$

in particular, we have $X^\emptyset_{\ge a} = X = X^\emptyset_{\le a}$. For a sequence $(f_n)_{n\in\mathbb{N}}$ in $\mathbb{R}^X$ and a set $I \subseteq \mathbb{N}$, define $I^\complement \coloneqq \mathbb{N} \setminus I$ and $f_I \coloneqq \{f_i : i \in I\}$.

Theorem 1.4 (The Bourgain–Fremlin–Talagrand dichotomy [BFT78])**.** *Let X be a Polish space, and let $A \subseteq \mathrm{C_p}(X)$ be pointwise bounded. The following are equivalent:*

(i) The set A is relatively compact in $\mathrm{B}_1(X)$.
(ii) For every sequence $(f_n)_{n\in\mathbb{N}}$ in A and all reals $a < b$ there is a set $I \subseteq \mathbb{N}$ such that

$$X^{f_I}_{\le a} \cap X^{f_{I^\complement}}_{\ge b} = \emptyset.$$

(In this form, the Bourgain–Fremlin–Talagrand dichotomy is a special case of the equivalence (ii)⇔(v) of Corollary 4G of Bourgain, Fremlin and Talagrand [BFT78].)

In condition (ii) of Theorem 1.4, the set $X^{f_I}_{\le a}$ is defined by the inequalities $f_i(x) \le a$ with $i \in I$, and the set $X^{f_{I^\complement}}_{\ge b}$ by the inequalities $f_j(x) \ge b$ with $j \in I^\complement$. Since I or $I^\complement$ is infinite, at least one of the two sets is defined by countably infinitely many inequalities. A computation, however, can check only finitely many inequalities at a time, and the following definitions accordingly impose only finitely many.

Definition 1.5. A set $A \subseteq \mathbb{R}^X$ of functions has the *finitary No-Independence Property (NIP)* if, for every sequence $(f_n)_{n\in\mathbb{N}}$ in A and all reals $a < b$, there exist finite disjoint sets $I, J \subseteq \mathbb{N}$ such that $X^{f_I}_{\le a} \cap X^{f_J}_{\ge b} = \emptyset$. We say that finite disjoint sets I, J with this property *witness finitary NIP for (f_n) and a, b.*

A set $A \subseteq \mathbb{R}^X$ has the *finitary Independence Property (IP)* if it does not have finitary NIP. Explicitly, finitary IP means that there exist a sequence $(f_n)_{n\in\mathbb{N}}$ in A and reals $a < b$ such that $X^{f_I}_{\le a} \cap X^{f_J}_{\ge b} \ne \emptyset$ for every pair of finite disjoint sets $I, J \subseteq \mathbb{N}$.

If the word "finite" is omitted in these definitions, we obtain *countable* NIP and *countable* IP. Countable NIP is weaker than finitary NIP, and countable IP is accordingly stronger than finitary IP. If, in addition, the witnesses are required to satisfy $J = I^\complement$, we obtain the properties that we call *BFT-NIP* and *BFT-IP*. Theorem 1.4 thus states that, on a Polish space X, a pointwise bounded set $A \subseteq \mathrm{C_p}(X)$ is relatively compact in $\mathrm{B}_1(X)$ if and only if it has BFT-NIP.

From now on, IP and NIP mean *finitary* IP and NIP unless otherwise specified.

Theorem 1.6. *Let X be a compact metrizable space, and let $A \subseteq \mathrm{C_p}(X)$ be pointwise bounded. The following are equivalent:*

(i) the set A is relatively compact in $\mathrm{B}_1(X)$;
(ii) the set A has BFT-NIP;

(iii) the set A has countable NIP;
(iv) the set A has finitary NIP.

(The equivalence *(ii)⇔(iv)* holds for every compact space X, even without the assumption that A is pointwise bounded, and the equivalence *(ii)⇔(iii)* holds for every set $A \subseteq \mathbb{R}^X$ on an arbitrary space X.)

Proof. Being compact and metrizable, the space X is Polish, and the equivalence (i)⇔(ii) is therefore Theorem 1.4. The implications (iv)⇒(iii) and (ii)⇒(iii) are trivial. The converse (iii)⇒(ii) holds because witnesses I and J of countable NIP satisfy $J \subseteq I^{\complement}$, whence I and $I^{\complement}$ witness BFT-NIP. To prove (ii)⇒(iv), suppose that (f_n) and $a < b$ witness finitary IP, and let $I \subseteq \mathbb{N}$ be given. The closed sets $X^{f_n}_{\leq a}$ $(n \in I)$ and $X^{f_m}_{\geq b}$ $(m \in I^{\complement})$ have the finite intersection property; by compactness, therefore, we obtain $X^{f_I}_{\leq a} \cap X^{f_{I^{\complement}}}_{\geq b} \neq \emptyset$. Since I is arbitrary, the set A does not have BFT-NIP, and (ii) fails. □

Theorem 1.6 can be cast as a dichotomy: on a compact metrizable space X, every pointwise bounded set $A \subseteq \mathrm{C_p}(X)$ either is relatively compact in $\mathrm{B}_1(X)$ or has IP in the finitary and, equivalently, in the countable sense.

The independence property was isolated by Saharon Shelah [She71, Definition 4.1] as one of the dividing lines of his classification of first-order theories [She90]. We return to Shelah's property in Section 1.5, where the property appears beside the corresponding notions of combinatorics and statistical learning.

1.2. **Rosenthal compacta.** The Bourgain–Fremlin–Talagrand dichotomy (Theorem 1.4) leads to the Rosenthal compacta, the compact spaces in which the deep computations of this paper are classified. By the dichotomy, if a pointwise bounded set of continuous functions on a Polish space has BFT-NIP, then the pointwise closure of the set is a compact set of functions of the first Baire class. Compact sets of this kind had appeared in Rosenthal's work on the ℓ_1-theorem, and in 1977 Rosenthal made them the subject of a study of their own [Ros77]. Godefroy [God80] named these compacta after Rosenthal in the following definition.

Definition 1.7. A *Rosenthal compactum* is a topological space homeomorphic to a compact subset of $\mathrm{B}_1(X)$, with the topology of pointwise convergence, for some Polish space X.

Rosenthal compacta have strong tameness properties; for instance, every point in the closure of a subset of a Rosenthal compactum is the limit of a sequence of points of the subset [BFT78, Theorem 3F]. In this paper Rosenthal compacta serve as spaces of deep computations. Theorem 3.1 below shows that for countably many features the deep computations of an NIP set of computations, extended to the limit states, form a Rosenthal compactum on each type shard. Here a type shard is a compact set of states and limit states on which every feature obeys a prescribed bound (see Section 2.1.1).

1.3. **The special case $\mathrm{B}_1(X, \mathbb{R}^{\mathcal{P}})$ with $\mathcal{P}$ countable.** This subsection is devoted to reducing the study of maps with values in a product of copies of the real line to that of real-valued functions. The reduction is needed because a state of a computation is described by the values of the features at the state, and a computation is therefore a map with values in the product of copies of the real line indexed by the

features. Fix a countable set $\mathcal{P}$ (in this paper, "countable" means finite or countably infinite), whose elements we call *predicate symbols* or *formal predicates*. Our goal is to characterize, for a Polish space X, the sets of continuous maps $X \to \mathbb{R}^{\mathcal{P}}$ whose restrictions to a compact subset of X form a relatively compact subset of the first Baire class.

Whenever $\mathcal{P}$ is regarded as a topological space, the set $\mathcal{P}$ carries the discrete topology, and the product $X \times \mathcal{P}$ is regarded either as a set or as a topological product, depending on the context. The spaces $\mathrm{C_p}(X, \mathbb{R}^{\mathcal{P}}) \subseteq \mathrm{B}_1(X, \mathbb{R}^{\mathcal{P}})$ are subspaces of the ambient space $(\mathbb{R}^{\mathcal{P}})^X$, which can be described in two other ways, through the natural homeomorphisms

$$(\mathbb{R}^{\mathcal{P}})^X \cong \mathbb{R}^{X \times \mathcal{P}} \cong (\mathbb{R}^X)^{\mathcal{P}},$$

given by

$$\mathbb{R}^{X \times \mathcal{P}} \to (\mathbb{R}^{\mathcal{P}})^X : \Phi \mapsto \hat{\Phi}$$
$$\mathbb{R}^{X \times \mathcal{P}} \to (\mathbb{R}^X)^{\mathcal{P}} : \Phi \mapsto \Phi\hat{},$$

where

$$\hat{\Phi}(x) \coloneqq \Phi(x, \cdot) \in \mathbb{R}^{\mathcal{P}}, \qquad \Phi\hat{}(P) \coloneqq \Phi(\cdot, P) \in \mathbb{R}^X.$$

These identifications treat X and $\mathcal{P}$ as mere sets; in particular, the topology of X plays no role in them.

The following proposition reduces the study of continuous functions, and of functions of Baire class 1, from X to $\mathbb{R}^{\mathcal{P}}$ to the case of real-valued functions.

Proposition 1.8. *The identification* $(\mathbb{R}^{\mathcal{P}})^X \cong \mathbb{R}^{X \times \mathcal{P}} \cong (\mathbb{R}^X)^{\mathcal{P}}$ *induces identifications*

$$\mathrm{C_p}(X, \mathbb{R}^{\mathcal{P}}) \cong \mathrm{C_p}(X \times \mathcal{P}) \cong \mathrm{C_p}(X)^{\mathcal{P}}, \quad \mathrm{B}_1(X, \mathbb{R}^{\mathcal{P}}) \cong \mathrm{B}_1(X \times \mathcal{P}) \cong \mathrm{B}_1(X)^{\mathcal{P}}.$$

(The cardinality of $\mathcal{P}$ plays no role here.)

Proof. The identification of the $\mathrm{C_p}$-spaces follows from the definition of the product topology and the discreteness of $\mathcal{P}$. By the former, a continuous map $X \to \mathbb{R}^{\mathcal{P}}$ is precisely a $\mathcal{P}$-indexed family of continuous functions $X \to \mathbb{R}$; by the latter, such families correspond to continuous functions $X \times \mathcal{P} \to \mathbb{R}$. For the spaces of functions of Baire class 1, recall that membership in the first Baire class is defined through pointwise limits of sequences of continuous functions. Since a sequence of $\mathcal{P}$-indexed families of continuous functions is the same as a $\mathcal{P}$-indexed family of sequences of continuous functions, the identification of these spaces follows as well. □

For $A \subseteq Y^X$ and $K \subseteq X$, we write $A|_K \coloneqq \{f|_K : f \in A\}$ for the set of restrictions to K of the functions in A. Similarly, for $A \subseteq (\mathbb{R}^{\mathcal{P}})^X$ and $P \in \mathcal{P}$, we write $\pi_P \circ A \coloneqq \{\pi_P \circ f : f \in A\}$, where $\pi_P \colon \mathbb{R}^{\mathcal{P}} \to \mathbb{R}$ is the projection onto the coordinate P. A set $A \subseteq \mathrm{C_p}(X, \mathbb{R}^{\mathcal{P}})$ is called *pointwise bounded* if the set $\pi_P \circ A \subseteq \mathrm{C_p}(X)$ is pointwise bounded for every $P \in \mathcal{P}$. The following theorem extends Theorem 1.6 to $\mathbb{R}^{\mathcal{P}}$-valued functions.

Theorem 1.9. *Assume that $\mathcal{P}$ is countable, that X is a Polish space, and that $A \subseteq \mathrm{C_p}(X, \mathbb{R}^{\mathcal{P}})$ is pointwise bounded. The following are equivalent for every compact $K \subseteq X$:*

(i) the set $A|_K$ is relatively compact in $\mathrm{B}_1(K, \mathbb{R}^{\mathcal{P}})$;
(ii) the set $\pi_P \circ A|_K$ has the NIP for every $P \in \mathcal{P}$.

Proof. By Proposition 1.8 and Tychonov's theorem, condition (i) holds if and only if $\pi_P \circ A|_K$ is relatively compact in $\mathrm{B}_1(K)$ for every $P \in \mathcal{P}$. Since K is compact metrizable, Theorem 1.6 shows that this is equivalent to (ii). □

In the applications, the NIP is verified on a set of states, whereas compactness is available only on the closure of the set. The following lemma shows that the NIP passes from a set to its closure.

Lemma 1.10. *Assume that X is Hausdorff and that $A \subseteq \mathrm{C_p}(X)$. The following are equivalent for every $L \subseteq X$:*

(i) the set $A|_L$ satisfies the NIP;
(ii) the set $A|_{\overline{L}}$ satisfies the NIP.

Proof. The implication (ii)⇒(i) is trivial. Conversely, suppose that a sequence (f_n) in A and reals $a < b$ witness the IP on $\overline{L}$, and choose a' and b' with $a < a' < b' < b$. For finite disjoint $I, J \subseteq \mathbb{N}$, the open set $\bigcap_{n\in I} f_n^{-1}(-\infty, a') \cap \bigcap_{n\in J} f_n^{-1}(b', \infty)$ contains a point of $\overline{L}$, namely a point at which $f_n \leq a$ for $n \in I$ and $f_n \geq b$ for $n \in J$; hence it meets L. Therefore the sequence (f_n) and the reals $a' < b'$ witness the IP on L. □

1.4. **Five canonical Rosenthal compacta.** Here we fix the notation for five Rosenthal compacta, all built on the Cantor set, that are the canonical members of the classes of Todorčević's trichotomy (see Theorem 3.3 below). By the trichotomy, the space of deep computations of each non-metrizable NIP example of Section 4 contains a copy of one of these compacta. The five compacta are distinguished from one another by three topological properties. Recall that a topological space is *separable* when the space has a countable dense subset, *hereditarily separable* when every subspace is separable, and *first countable* when every point has a countable base of neighborhoods.

Let $2^{\mathbb{N}}$ carry the product topology and the lexicographic order $<$, whose order topology is the product topology. The set of finite binary words is denoted $2^{<\mathbb{N}}$, and the length of a word $t \in 2^{<\mathbb{N}}$ is denoted $|t|$. For $a \in 2^{\mathbb{N}}$ and $n \in \mathbb{N}$, we write $a{\restriction}n$ for the word formed by the first n bits of a. Two words are *comparable* if one extends the other, and an *antichain* is a set of pairwise incomparable words. For a finite word $t \in 2^{<\mathbb{N}}$, let $t^\frown 0^\infty$ and $t^\frown 1^\infty$ be the sequences that begin with t and continue with all 0's and with all 1's, respectively. The set of sequences extending t is then the clopen interval $[t^\frown 0^\infty, t^\frown 1^\infty]$. The last of the five compacta below, like the families of Section 3.3, is built along a copy of the binary tree within itself. Such a copy is a *dyadic subtree* of $2^{<\mathbb{N}}$, that is, a subset of $2^{<\mathbb{N}}$ isomorphic to $2^{<\mathbb{N}}$ under the order of extension. We write a dyadic subtree as $\{w_t : t \in 2^{<\mathbb{N}}\}$, indexed through the canonical isomorphism, which preserves extension and the lexicographic order. A dyadic subtree is called *regular* if every level of the subtree is contained in a level of $2^{<\mathbb{N}}$. A sequence (u_k) of words *converges* to $a \in 2^{\mathbb{N}}$ if, for every $n \in \mathbb{N}$, all but finitely many of the words u_k extend $a{\restriction}n$.

Double brackets denote truth values: $[\![\ldots]\!]$ is 1 if the statement inside the brackets is true and 0 if the statement is false. We introduce the following functions $2^{\mathbb{N}} \to \{0, 1\}$:

- the *prefix test* $v_t := [\![\cdot{\restriction}|t| = t]\!]$, the characteristic function of the set of sequences extending a fixed word $t \in 2^{<\mathbb{N}}$;

- the *equality test* $\delta_a \coloneqq [\![\cdot = a]\!]$, the characteristic function of the singleton $\{a\}$ for fixed $a \in 2^{\mathbb{N}}$;
- the *threshold tests* $\chi_{[a} \coloneqq [\![a \leq \cdot]\!]$ and $\chi_{(a} \coloneqq [\![a < \cdot]\!]$, the characteristic functions of the final intervals $\{x : a \leq x\}$ (closed) and $\{x : a < x\}$ (open), for fixed $a \in 2^{\mathbb{N}}$; the special case $\mathbf{0} \coloneqq \chi_{(1^\infty}$ is the identically zero function.

Prefix tests are continuous, and so are the threshold tests $\chi_{[t^\frown 0^\infty}$ and $\chi_{(t^\frown 1^\infty}$, since the value of each of these functions at a sequence depends only on the first $|t|$ bits of the sequence. By contrast, equality tests and threshold tests at all other points are discontinuous functions of Baire class 1. All closures below are taken in the topology of pointwise convergence.

(1) The *Alexandroff compactification* $A(2^{\mathbb{N}}) = \{\delta_a : a \in 2^{\mathbb{N}}\} \cup \{\mathbf{0}\}$. Each δ_a is isolated, and every neighborhood of $\mathbf{0}$ contains all but finitely many of the functions δ_a. Thus the Alexandroff compactification is neither separable nor first countable at $\mathbf{0}$.

(2) The *extended Alexandroff compactification* $\hat{A}(2^{\mathbb{N}}) = \overline{\{v_t : t \in 2^{<\mathbb{N}}\}} = \{v_t : t \in 2^{<\mathbb{N}}\} \cup A(2^{\mathbb{N}})$. As $n \to \infty$, the prefix tests $v_{a\restriction n}$ converge to δ_a; along an antichain they converge to $\mathbf{0}$. Hence the extended Alexandroff compactification is separable but not first countable.

(3) The *split Cantor set* $S(2^{\mathbb{N}}) = \{\chi_{[a}, \chi_{(a} : a \in 2^{\mathbb{N}}\}$. The continuous thresholds $\chi_{[t^\frown 0^\infty}$, together with the isolated point $\mathbf{0} = \chi_{(1^\infty}$, form a countable dense subset: thresholds approaching a from below converge to $\chi_{[a}$, and from above to $\chi_{(a}$. It follows that the split Cantor set is separable, first countable and hereditarily separable, but not metrizable.

(4) The *Alexandroff duplicate* $D(2^{\mathbb{N}}) = \{(\mathbf{0}, a), (\delta_a, a) : a \in 2^{\mathbb{N}}\} \subseteq A(2^{\mathbb{N}}) \times 2^{\mathbb{N}}$. The points (δ_a, a) are isolated, whereas every neighborhood of $(\mathbf{0}, a)$ contains, for all b in some neighborhood of a, the point $(\mathbf{0}, b)$ and, with finitely many exceptions, the point (δ_b, b). Consequently, the Alexandroff duplicate is first countable but not separable; the duplicate is contained in the separable compactum $\hat{D}(2^{\mathbb{N}}) = \overline{\{(v_t, t^\frown 0^\infty) : t \in 2^{<\mathbb{N}}\}} = D(2^{\mathbb{N}}) \cup \{(v_t, t^\frown 0^\infty) : t \in 2^{<\mathbb{N}}\}$.

(5) The *extended duplicate of the split Cantor set*. Following Argyros, Dodos and Kanellopoulos [ADK08, 4.3.7], we define this compactum along a regular dyadic subtree $\{s_t : t \in 2^{<\mathbb{N}}\}$ of $2^{<\mathbb{N}}$, fixed once and for all. This subtree is required to satisfy $s_t^\frown 0^\infty \neq s_{t'}^\frown 0^\infty$ and $s_t^\frown 1^\infty \neq s_{t'}^\frown 1^\infty$ for all distinct $t, t' \in 2^{<\mathbb{N}}$, and the extended duplicate is then

$$\hat{D}(S(2^{\mathbb{N}})) = \overline{\{(v_{s_t}, \chi_{[s_t^\frown 0^\infty}) : t \in 2^{<\mathbb{N}}\}} \subseteq \hat{A}(2^{\mathbb{N}}) \times S(2^{\mathbb{N}}).$$

Call a sequence $a \in 2^{\mathbb{N}}$ a *branch* of the subtree if every initial segment $a\restriction n$ of a is extended by some word s_t; the branches form a perfect subset of $2^{\mathbb{N}}$. The compactum $\hat{D}(S(2^{\mathbb{N}}))$ consists of the isolated points $(v_{s_t}, \chi_{[s_t^\frown 0^\infty})$ and, for each branch a, of the point $(\delta_a, \chi_{[a})$ and of at most two points with first coordinate $\mathbf{0}$ [ADK08, 4.3.7]. The point $(\mathbf{0}, \chi_{[a})$ belongs to the compactum exactly when some sequence (u_k) of pairwise incomparable words of the subtree converges to a with $u_k^\frown 0^\infty$ increasing in k. Likewise, the point $(\mathbf{0}, \chi_{(a})$ belongs to the compactum exactly when some such sequence converges to a with $u_k^\frown 0^\infty$ decreasing in k. The points $(\delta_a, \chi_{[a})$ are reached along the branches. Indeed, if a is a branch and $u_1, u_2, \ldots$ are the words of the subtree that are initial segments of a, listed by increasing length,

then the sequences $u_k^\frown 0^\infty$ increase strictly to a. The points $(v_{u_k}, \chi_{[u_k^\frown 0^\infty)})$ then converge to $(\delta_a, \chi_{[a)})$ as $k \to \infty$. As the closure of a countable set, the extended duplicate of the split Cantor set is separable. This compactum is also first countable, since the compactum embeds into the Helly space of nondecreasing functions from $[0,1]$ into itself, which is first countable [ADK08, 4.3.7]. For each branch a, the point $(\delta_a, \chi_{[a)})$ is the only point $(\delta_b, \chi_{[b)})$, with b a branch, whose first coordinate takes the value 1 at a. The points $(\delta_a, \chi_{[a)})$, with a a branch, therefore form a discrete subspace, which has the cardinality of the continuum because the branches form a perfect set. Hence the extended duplicate of the split Cantor set is not hereditarily separable.

Figure 3 shows the five compacta as patterns of convergence of words of the binary tree $2^{<\mathbb{N}}$.




FIGURE 3. The five compacta as patterns of convergence in $2^{<\mathbb{N}}$.

The constructions of the Alexandroff compactification and of the Alexandroff duplicate make sense with $2^{\mathbb{N}}$ replaced by any compact metric space K; the resulting spaces are denoted $A(K)$ and $D(K)$, respectively. Since $A(K)$ is the one-point compactification of a discrete space of the cardinality of K, the space $A(K)$ depends, up to homeomorphism, only on the cardinality of K; for instance, $A([0,1])$ is homeomorphic to $A(2^{\mathbb{N}})$.

1.5. **NIP in model theory, combinatorics and learning: polynomial versus exponential growth.** We now trace the NIP of Definition 1.5 to the corresponding notions of model theory, combinatorics and statistical learning, and we make the relations among them precise. One of these relations is used in Section 4, where a bound on the Vapnik–Chervonenkis dimension of a family of arcs yields the NIP of Bresenham's algorithm. For families of $\{0,1\}$-valued functions (that is, of sets),

a combinatorial dividing line closely related to the NIP appeared independently in three fields between 1968 and 1972, some years before Theorem 1.4. In model theory the line arose in the work of Saharon Shelah [She71, She90], and in combinatorics in the work of Norbert Sauer [Sau72] and, again, of Shelah [She72, She90]. In statistical learning, finally, the same line appeared in the work of Vladimir Vapnik and Alexey Chervonenkis [VC68, VC71, VC74]. In each of the three fields, the line separates the "tame" families of sets from the "wild" ones. The number of traces of a tame family on an n-element set (the intersections of the members of the family with the set) is bounded by a polynomial in n. In a wild family, by contrast, for every $n \in \mathbb{N}$ some n-element set has all 2^n possible traces.

In model theory: Among the dividing lines of Shelah's classification theory [She90], the independence property is detected by counting types over finite sets of parameters. Let T be a complete first-order theory, and let $\varphi(x;y)$ be a formula in the language of T, in which x and y are finite tuples of variables of lengths $|x|$ and $|y|$. Finally, let B be a finite set of $|y|$-tuples of elements of a model M of T. A *φ-type* over B is a maximal set of formulas $\varphi(x;b)$ and $\neg\varphi(x;b)$, with $b \in B$, that is consistent with the theory of M with constants for the elements of M. Since B is finite, every φ-type over B is realized by an $|x|$-tuple a of elements of M, and the type is determined by the set $\{b \in B : M \models \varphi(a;b)\}$. Hence the set $S_\varphi(B)$ of φ-types over B is in bijection with the traces on B of the family of sets $\{b : M \models \varphi(a;b)\}$, with a ranging over $|x|$-tuples of elements of M. This bijection gives the bound $|S_\varphi(B)| \leq 2^{|B|}$. The formula φ has the *independence property* (IP) if, for every $n \in \mathbb{N}$, some set B of n tuples satisfies $|S_\varphi(B)| = 2^n$. Otherwise the formula is *NIP*, or *dependent*, and the theory T is NIP if every formula in the language of T is NIP. Since T is complete, these notions do not depend on the model M.

If φ is NIP, the Sauer–Shelah lemma below bounds $|S_\varphi(B)|$ by a polynomial in $|B|$. For a set A of parameters, we write $S_\varphi(A)$ for the set of φ-types over the set of $|y|$-tuples of elements of A. If φ has IP, then for every n some set A of at most $n|y|$ parameters satisfies $|S_\varphi(A)| \geq 2^n$ [Sim15, Section 2.3.4]. Consequently, φ is NIP if and only if $|S_\varphi(A)|$ is bounded by a polynomial in $|A|$ as A ranges over the finite sets of parameters. Over infinite sets of parameters, counting φ-types does not separate NIP from IP without further set-theoretic hypotheses. Indeed, every unstable formula, NIP or not, already has 2^κ φ-types over some set of κ parameters whenever κ is infinite and $2^\kappa = \kappa^+$ [Sim15, Section 2.3.4].

In combinatorics: Sauer [Sau72] and Shelah [She72] proved independently the following dichotomy, now known as the *Sauer–Shelah lemma*. Let $\mathscr{F} = \{C_i : i \in I\}$ be a family of subsets of a set Ω, indexed by a set I. Then either for every $n \in \mathbb{N}$ there is a set $B \subseteq \Omega$ with $|B| = n$ such that $|\{C_i \cap B : i \in I\}| = 2^n$, or there exists $N \in \mathbb{N}$ such that, for every $B \subseteq \Omega$ with $|B| \geq N$,

$$|\{C_i \cap B : i \in I\}| \leq \sum_{j=0}^{N-1} \binom{|B|}{j}.$$

In the first case $\mathscr{F}$ is said to have "exponential complexity", and in the second case "polynomial complexity"; every family of subsets of a *finite*

set Ω has polynomial complexity. The name "polynomial complexity" is justified by the following estimate, valid for fixed $N > 0$ and for $|B| = m > 0$, which bounds the number of traces by a polynomial in m:

$$\sum_{j=0}^{N-1} \binom{m}{j} \leq \sum_{j=0}^{N-1} \frac{m^j}{j!} \leq \left(\sum_{j=0}^{N-1} \frac{1}{j!} \right) \cdot m^{N-1} < e \cdot m^{N-1}.$$

In statistical learning: The Sauer–Shelah lemma is also the dichotomy that Vapnik and Chervonenkis had announced in 1968 [VC68], in their study of uniform convergence in statistics. Their proof of the dichotomy appeared in 1971 [VC71], a year before the papers of Sauer and Shelah (see also [VC74]). The largest size of a set $B \subseteq \Omega$ with $2^{|B|}$ traces $C_i \cap B$ is called the *Vapnik–Chervonenkis dimension*, or *VC dimension*, of $\mathscr{F}$. Equivalently, the VC dimension is the integer $N - 1$ for the least integer N as in the Sauer–Shelah lemma. If no such integer N exists, the VC dimension of $\mathscr{F}$ is infinite. Through the Vapnik–Chervonenkis theorem, the Sauer–Shelah lemma bounds the number of data points (the sample size) needed to "learn" a concept class $\mathscr{F}$ of finite VC dimension in a statistical sense. Here "learning" is understood up to a preset error and with a preset confidence. The Fundamental Theorem of Statistical Learning concerns PAC-learnability, where PAC stands for "Probably Approximately Correct" [Val84]. This theorem states that a class of $\{0, 1\}$-valued functions satisfying a mild measurability condition is PAC-learnable if and only if the class has finite VC dimension [BEHW89]; see also [SSBD14, Theorem 6.7].

For first-order formulas the three dichotomies coincide. To see this, fix a formula $\varphi(x; y)$ and consider the family of sets $\{b : M \models \varphi(a; b)\}$, with a ranging over the $|x|$-tuples of elements of a model M of T. By the description of φ-types given above, the formula φ has IP exactly when this family has infinite VC dimension. For an NIP formula, moreover, the Sauer–Shelah lemma gives the polynomial bound on the number of φ-types. For an arbitrary family of sets, finiteness of the VC dimension implies that the characteristic functions of its members have the NIP of Definition 1.5. The NIP does not, however, imply finite VC dimension: the finite subsets of $\mathbb{N}$ form a family of infinite VC dimension whose characteristic functions have the NIP.

Historical remarks. The Baire hierarchy of functions was introduced by René-Louis Baire in his 1899 doctoral thesis, *Sur les fonctions de variables réelles*. The theory of spaces of continuous functions with the topology of pointwise convergence, C_p-theory, was pioneered by A. V. Arhangel'skiĭ and his students in the 1970s and 1980s [Ark92]. This theory has since found numerous applications in model theory and in functional analysis; Tkachuk's problem book [Tka11] develops the subject systematically, through problems with complete solutions.

2. Compositional computation structures: A structural approach to arbitrary-precision computations

The theory of function spaces assembled in Section 1 is brought to bear on computation through the framework of compositional computation structures, to which the present section is devoted. A computation on real-valued data is observed through real-valued features of the states, each known to finite precision, and the

topology of this observation is the topology of pointwise convergence. Following the first paper of this series [ADIW26], we first describe states by types, the tuples of values of all the features. Computations act on states and are composed, and deep computations, the limits of computations along ultrafilters, become points of a space of maps into the space of types. Finally, we recall the stable sets of computations, studied in the first paper, and close the section with two examples that motivate the classification of Section 3.

2.1. **Computation states structures.** The first ingredient of the framework, which we introduce here, is the set of states on which computations act, together with the real-valued features through which the states are observed. A *computation states structure* is a pair $(L, \mathcal{P})$ consisting of a set L of *states* and a collection $\mathcal{P}$ of real-valued functions on L that we call *predicates* or *features*. Intuitively, the set L consists of the states of a computation, and the predicates are the primitives that are given and accepted as computable. For $P \in \mathcal{P}$ and $v \in L$, the number $P(v)$ is the value of the (observable) feature P at the state v. Maps $f\colon L \to L$ are called *transitions* of $(L, \mathcal{P})$; the computations of Definition 2.4 below are transitions.

The *type* of a state $v \in L$ is the indexed family[5]

$$\mathrm{tp}(v) = (P(v))_{P\in\mathcal{P}} \in \mathbb{R}^{\mathcal{P}}$$

of the values of the features at v.

We assume that each state $v \in L$ is determined by the type $\mathrm{tp}(v)$. Under this assumption we may identify L with the image $\mathrm{tp}[L] \subseteq \mathbb{R}^{\mathcal{P}}$ of L under the type map and endow L with the subspace topology. This topology is the topology of pointwise convergence induced by the product topology of $\mathbb{R}^{\mathcal{P}}$; in particular, for each $P \in \mathcal{P}$, the map $v \mapsto P(v)$ is continuous on L.

The principal examples are the state space $L = \mathbb{R}^{\mathbb{N}}$ with the predicates $P_i(v) = v_i$ for $i \in \mathbb{N}$ and, for a positive integer n, the state space $L = \mathbb{R}^n$ with the predicates $P_i(v) = v_i$ for $0 \le i < n$. Subspaces of these two spaces are the state spaces of primary interest in this paper. Indeed, once the predicates are enumerated, the identification of states with types makes every computation states structure with countably many predicates a subspace of $\mathbb{R}^{\mathbb{N}}$ or of some $\mathbb{R}^n$.

Definition 2.1. Let $(L, \mathcal{P})$ be a computation states structure. The type $\mathrm{tp}(v) \in \mathbb{R}^{\mathcal{P}}$ of a state $v \in L$ is called a *realized* (state) type. The closure of the set of realized types in $\mathbb{R}^{\mathcal{P}}$, with the topology of pointwise convergence, is called the *space of (state) types* of $(L, \mathcal{P})$ and is denoted $\mathcal{L}$. The elements $\mathbf{v}$ of $\mathcal{L}$ are called *state types*, and those in $\mathcal{L} \setminus L$ are called *unrealized* state types.

State types formalize the intuitive notion of a limit state, and the predicates extend continuously to state types. Indeed, for each $P \in \mathcal{P}$, the restriction to $\mathcal{L}$ of the P-th coordinate map $\mathbb{R}^{\mathcal{P}} \to \mathbb{R}$ given by $\mathbf{v} = (\mathbf{v}_Q)_{Q\in\mathcal{P}} \mapsto \mathbf{v}_P$ is a continuous extension of the predicate $P(\cdot)$ on L. Abusing notation, we still denote this extension by $P(\cdot)$, or just P.

[5]As in Proposition 1.8, where $\mathcal{P}$ is merely a set of symbols, we regard each predicate as named by a formal label P, a *predicate symbol*. The meaning of this symbol in the computation states structure is the real function $P(\cdot)\colon v \mapsto P(v)$ on L, the observable feature named by P. Whenever $\mathcal{P}$ is used as an index set, as in the product $\mathbb{R}^{\mathcal{P}}$ below, it is regarded as a set of symbols. The distinction between a symbol and its meaning is usually suppressed in mathematics, but the model-theoretic framework requires awareness of it. When necessary, we write $P(\cdot)$ for the function on L and P for the symbol; when no confusion can arise, we also write P for the function.

Remark 2.2. One may regard unrealized state types as "hyper-states", and those among them that are obtained as limits of sequences (or as ultralimits) of realized state types as "deep states". For readers versed in model theory we note that, at least for realizing state types with parameters in L, the type space $(\mathcal{L}, \mathcal{P})$ serves as a "poor man's" ultraproduct of $(L, \mathcal{P})$. Indeed, by the very definition of $\mathcal{L}$, every type $\mathbf{v}$ is realized in this type space.

2.1.1. *Bounds on precision and magnitude. Shards.* The analysis of this paper takes place on shards, the sets of states on which each feature obeys a prescribed bound; before defining them, we explain why actual computation imposes such bounds. Computation states structures are designed to model actual computation, in which operations on real-valued quantities can be carried out only approximately, to a preset degree of precision. A bound on the magnitude of the quantities involved is no less important in practice, for no machine can store, let alone compute with, arbitrarily large numbers. An actual computation therefore fixes at the outset bounds for its errors and for the absolute magnitudes of the quantities it handles. These bounds are fixed even when the collection $\mathcal{P}$ of predicates is infinite, possibly uncountable. Moreover, different parts of a computation may require different error bounds and may deal with quantities of very different sizes. Focusing on magnitudes, we consider sets of computations for which a bound for each predicate is fixed in advance. We are thereby led to the notion of *shard*, introduced in the first paper of this series [ADIW26].

Definition 2.3. Let $(L, \mathcal{P})$ be a computation states structure, with L identified with the set $\mathrm{tp}[L] \subseteq \mathbb{R}^{\mathcal{P}}$ of its realized types. A *sizer* is a family $r_\bullet = (r_P)_{P \in \mathcal{P}}$ of positive real numbers indexed by $\mathcal{P}$. For a sizer $r_\bullet$, let $\mathbb{R}[r_\bullet] \coloneqq \prod_{P \in \mathcal{P}} [-r_P, r_P]$, a compact space, and define the $r_\bullet$-*shard* of the state space L by

$$L[r_\bullet] = L \cap \mathbb{R}[r_\bullet].$$

The $r_\bullet$-*type shard* is the closure $\mathcal{L}[r_\bullet] = \overline{L[r_\bullet]}$, a closed and hence compact subset of $\mathbb{R}[r_\bullet]$.

The union of all type shards, as the sizer $r_\bullet$ varies, is denoted $\mathcal{L}_{\mathrm{sh}}$ and called the space of *shard-supported types*.

We have $\mathcal{L}_{\mathrm{sh}} \subseteq \mathcal{L}$, and the inclusion may be proper; equality holds in the important special case when $\mathcal{P}$ is countable (see [ADIW26]).

2.2. **Compositional Computation Structures.** In this subsection we recall the compositional computation structures of the first paper of this series [ADIW26], the framework in which deep computations are defined. Computations transform states, and two computations performed one after the other form a computation. The computations of a system therefore form a semigroup under composition, which we take to contain the identity, the computation that does nothing. Deep computations are the limits of computations along ultrafilters in the topology of pointwise convergence, taken in the space of types. Since an actual computation keeps the magnitudes that it handles within bounds, the shards of Section 2.1.1 enter through the notion of confinement. Throughout, the state space L of a computation states structure $(L, \mathcal{P})$ is identified with a subset of the space of types $\mathcal{L} \subseteq \mathbb{R}^{\mathcal{P}}$. Sets of maps $L \to L$ are topologized as subsets of the product L^L, in which, as usual, the exponent L serves merely as an index set, whereas the base $L \subseteq \mathbb{R}^{\mathcal{P}}$ carries the topology of pointwise convergence. The product L^L is thus a subspace

of $\mathcal{L}^L \subseteq (\mathbb{R}^{\mathcal{P}})^L$, and closures of sets of maps $L \to L$ are taken in $\mathcal{L}^L$. The following definition records the ingredients just described.

Definition 2.4. A *compositional computation structure* (CCS) is a triple $(L, \mathcal{P}, \Gamma)$, where

- the pair $(L, \mathcal{P})$ is a computation states structure, and
- the set $\Gamma \subseteq L^L$ is a monoid of transformations of L, that is, a semigroup under composition that contains the identity map id of L.

The elements of Γ are called the *computations* of $(L, \mathcal{P}, \Gamma)$. The elements ξ of the closure $\overline{\Gamma} \subseteq \mathcal{L}^L$ are called (real-valued) *deep computations*, and the elements of the closure $\overline{\Delta}$ of a set $\Delta \subseteq \Gamma$ are called the deep computations of Δ.[6]

A collection $\mathscr{R}$ of sizers is *exhaustive* if $L = \bigcup_{r_\bullet \in \mathscr{R}} L[r_\bullet]$. A computation $\gamma \in \Gamma$ is *$\mathscr{R}$-confined* if it maps $L[r_\bullet]$ into itself for every $r_\bullet \in \mathscr{R}$, and a subset $\Delta \subseteq \Gamma$ is *$\mathscr{R}$-confined* if each $\gamma \in \Delta$ is.

Unlike the subspace L^L, which is a semigroup under composition, the space $\mathcal{L}^L$ has no natural semigroup structure; deep computations are therefore not composable in general.

Proposition 2.5. *If $\Delta \subseteq \Gamma$ is confined by an exhaustive collection of sizers, then the closure $\overline{\Delta}$ is a compact subset of $\mathcal{L}^L_{\mathrm{sh}}$.*

Proof. Let $\mathscr{R}$ be an exhaustive collection of sizers that confines Δ, and choose, for each $v \in L$, a sizer $r_\bullet^{(v)} \in \mathscr{R}$ with $v \in L[r_\bullet^{(v)}]$. Since every $\gamma \in \Delta$ maps v into $L[r_\bullet^{(v)}]$, the set Δ lies in the compact space $\prod_{v \in L} \mathcal{L}[r_\bullet^{(v)}] \subseteq \mathcal{L}^L_{\mathrm{sh}}$, which therefore contains $\overline{\Delta}$. □

For a CCS $(L, \mathcal{P}, \Gamma)$ we regard the elements of Γ as "standard" finitary computations and the elements of $\overline{\Gamma}$, the deep computations, as possibly infinitary limits of standard computations.

Remarks 2.6. (For readers versed in model theory.)[7]

(1) Compositional computation structures, as defined above, are not literally real-valued structures, because nothing in the structure itself forces Γ to be realized as a subset of the product space L^L. However, as in their original definition [ADIW26], they may be regarded as real-valued structures $(L, \mathcal{P}, \Gamma, \circ, \mathrm{ev})$, where
   - the pair $(L, \mathcal{P})$ is a computation states structure,
   - the pair $(\Gamma, \circ)$ is a monoid,

[6] In the first paper of this series [ADIW26], the deep computations of Δ are the accumulation points of Δ that satisfy a confinement condition. Counting the members of Δ among them affects none of the properties studied in this paper. Indeed, under the Extendibility Axiom of Section 2.3, the members of Δ are continuous on the type shards, and those that are not accumulation points of Δ are isolated points of $\overline{\Delta}$.

[7] The real-valued structures used here are those of Keisler's model theory for real-valued structures [Kei23]; see also [DnI17, Section 6] for an introduction. In Keisler's structures every predicate comes with an *a priori bound*, a real number r such that the predicate takes values in $[-r, r]$. The features of a computation states structure need not be bounded on all of L, but on a shard $L[r_\bullet]$ each feature P takes values in $[-r_P, r_P]$. The computations that concern us, which are confined to shards, therefore fall within Keisler's framework.

- the map $\mathrm{ev} : \Gamma \times L \to L : (\gamma, v) \mapsto \mathrm{ev}(\gamma, v)$ is a faithful monoid action of Γ on L (i.e., $\mathrm{ev}(\gamma \circ \delta, v) = \mathrm{ev}(\gamma, \mathrm{ev}(\delta, v))$ for $\gamma, \delta \in \Gamma$ and $v \in L$, the identity element of Γ fixes every $v \in L$, and distinct members of Γ act differently).

The two definitions are reconciled by identifying each $\gamma \in \Gamma$ with the map $v \mapsto \mathrm{ev}(\gamma, v)$.

(2) Regarded as real-valued structures in this sense, the compositional computation structures with a fixed collection $\mathcal{P}$ of predicate symbols do not form an elementary class, because the requirement that each state be determined by its type is not preserved by ultrapowers. For instance, let $L = \{0,1\}^{\mathbb{N}}$ with the predicates $P_i(v) = v_i$, and let Γ consist of the permutations of coordinates with finite support. In an ultrapower along a nonprincipal ultrafilter on $\mathbb{N}$, the classes of the constant sequence with value 0 and of the sequence $(e_n)_{n\in\mathbb{N}}$ of unit vectors have the same type, yet the class of the sequence of transpositions of 0 and n sends them to states that differ at P_0. The model-theoretic remarks below concern the structures $(L, \mathcal{P}, \Gamma, \circ, \mathrm{ev})$ without that requirement. The notion of state type introduced above is essentially that of a quantifier-free type with parameters of sort L only. This notion may be enriched by further formulas, for instance with parameters in Γ or with quantifiers, and extended to types of sort Γ ("transform-types"). As usual, all types over a structure $\underline{C} = (L, \mathcal{P}, \Gamma, \circ, \mathrm{ev})$ are realized in some ultrapower $\underline{C}^{(\mathcal{U})}$ of $\underline{C}$, constructed as in the references above.[8] In Banach space theory, this approach to spaces of types goes back to Krivine's real-valued languages [Kri72, Kri74] and to the work of Krivine and Maurey on stable Banach spaces [KM81].

2.3. **Computability of deep computations and the Extendibility Axiom.** The rest of the paper depends on two notions, the computability of transforms and the Extendibility Axiom, both of which we recall in this subsection. The Extendibility Axiom requires every computation to have an extension that is continuous on each type shard. As for computability, a machine evaluates only sums and products of finitely many observed features, and only to within a preset error on a preset range of magnitudes. For this reason, computability is defined by uniform approximation on shards by polynomials in the features. Let $(L, \mathcal{P})$ be a computation states structure. A map $f\colon L \to \mathcal{L}$ from the state space to the space of types is called a *transform*. In the two definitions that follow, and in the characterization of computability after them, the collection $\mathcal{P}$ of predicates is assumed to be *countable*.

(1) A *computable predicate* is a function from L to $\mathbb{R}$ whose restriction to each shard is a uniform limit of polynomials, with real coefficients, in the features $P(\cdot)$.

(2) A transform $f\colon L \to \mathcal{L}$ is *computable* if, for each $Q \in \mathcal{P}$, the output feature $Q \circ f\colon L \to \mathbb{R}$ is a computable predicate.

Alva, Dueñez, Iovino and Walton [ADIW26] showed that a transform $f\colon L \to \mathcal{L}$ is computable if and only if it extends to a continuous map $\tilde{f}\colon \mathcal{L} \to \mathcal{L}$. This characterization motivates the following definition, in which continuity is required on each type shard.

[8] By contrast, no ultrapower is needed to realize state types in a computation states structure (see Remark 2.2).

A computation $\gamma \in \Gamma$ is *extendable* if γ has an extension $\tilde{\gamma}\colon \mathcal{L}_{\mathrm{sh}} \to \mathcal{L}_{\mathrm{sh}}$ such that, for every sizer $r_\bullet$, there is a sizer $s_\bullet$ for which the restriction $\tilde{\gamma}|_{\mathcal{L}[r_\bullet]}\colon \mathcal{L}[r_\bullet] \to \mathcal{L}[s_\bullet]$ is continuous. Such an extension is unique, since the extension is continuous on each type shard $\mathcal{L}[r_\bullet]$, in which $L[r_\bullet]$ is dense. For a set $\Delta \subseteq \Gamma$ of extendable computations, we write $\tilde{\Delta}$ for the set $\{\tilde{\gamma}\colon \gamma \in \Delta\}$.

Extendibility Axiom. *Every computation $\gamma \in \Gamma$ is extendable.*

The Extendibility Axiom is discussed at length in the first paper of this series [ADIW26]. *From now on, all compositional computation structures are assumed to satisfy the Extendibility Axiom.*

The following lemma extends deep computations to the type shards; it is used in Theorems 3.1 and 5.7.

Lemma 2.7. *Let $\mathscr{R}$ be an exhaustive collection of sizers and let $\Delta \subseteq \Gamma$ be $\mathscr{R}$-confined. Every deep computation $f \in \overline{\Delta}$ has an extension $\tilde{f} : \mathcal{L}_{\mathrm{sh}} \to \mathcal{L}_{\mathrm{sh}}$ that, for every $r_\bullet \in \mathscr{R}$, maps $\mathcal{L}[r_\bullet]$ into itself and satisfies $\tilde{f}|_{\mathcal{L}[r_\bullet]} \in \overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$.*

Proof. Write $f = \lim_{i,\mathcal{U}} \gamma_i$, where $(\gamma_i)_{i\in I}$ is a family in Δ and $\mathcal{U}$ is an ultrafilter on the index set I, and let $r_\bullet \in \mathscr{R}$ be given. Each $\tilde{\gamma}_i$ maps $L[r_\bullet]$ into itself and is continuous on $\mathcal{L}[r_\bullet]$; hence $\tilde{\gamma}_i$ maps the closure $\mathcal{L}[r_\bullet]$ of $L[r_\bullet]$ into itself. Since $\mathcal{L}[r_\bullet]$ is compact, the limit $\tilde{f}(v) \coloneqq \lim_{i,\mathcal{U}} \tilde{\gamma}_i(v)$ exists and lies in $\mathcal{L}[r_\bullet]$ for every $v \in \mathcal{L}[r_\bullet]$; in other words, the restriction $\tilde{f}|_{\mathcal{L}[r_\bullet]}$ equals $\lim_{i,\mathcal{U}} \tilde{\gamma}_i|_{\mathcal{L}[r_\bullet]}$. Since $\mathscr{R}$ is exhaustive, this defines $\tilde{f}$ on a set that contains L, and there $\tilde{f}$ agrees with f; at the remaining points of $\mathcal{L}_{\mathrm{sh}}$ we let $\tilde{f}$ be the identity. □

Lemma 2.7 connects deep computations with the second notion of limit introduced in the first paper of this series [ADIW26]. In that paper, this second notion is formulated in the space of maps from the space of types to itself, endowed with the topology of pointwise convergence. An *ultracomputation* of a set $\Delta \subseteq \Gamma$ of extendable computations is an accumulation point of $\tilde{\Delta}$ in this space of maps, subject to a confinement condition. Thus a deep computation sends states to types, whereas an ultracomputation sends types to types. Every deep computation is the restriction to L of an ultracomputation [ADIW26]. On the type shards $\mathcal{L}[r_\bullet]$ with $r_\bullet \in \mathscr{R}$, the ultralimit $\lim_{i,\mathcal{U}} \tilde{\gamma}_i$ in the proof of Lemma 2.7 provides such an extension of the deep computation $f = \lim_{i,\mathcal{U}} \gamma_i$ considered there. The classification of Section 3 is based on the sets $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$, whose members are the restrictions to $\mathcal{L}[r_\bullet]$ of such ultralimits of extended computations. Strictly speaking, the classification therefore concerns the restrictions of ultracomputations to type shards. For the dynamical compositional computation structures of Section 4.1, in which every type is realized and every computation is its own extension, deep computations and ultracomputations coincide.

2.4. **Stability.** The highest level of the hierarchy studied in this paper is occupied by the stable sets of computations, all of whose deep computations are computable when $\mathcal{P}$ is countable. We recall these sets here, before descending to the levels below in Section 3. By the theorem of Alva, Dueñez, Iovino and Walton recalled above, the computable transforms are exactly those that extend to continuous maps on the space of types. Accordingly, the stable sets are defined by the continuity of their deep computations on every type shard.

Definition 2.8. A set $\Delta \subseteq \Gamma$ of computations is *stable* if, for every sizer $r_\bullet$, every member of $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$ is continuous.

When $\mathcal{P}$ is countable, continuity on the type shards amounts to computability, by the theorem of Alva, Dueñez, Iovino and Walton recalled in Section 2.3. Stable sets of computations, the subject of the first paper of this series [ADIW26], are characterized there by Grothendieck's double limit criterion and identified with stability in the sense of model theory.

2.5. **Two motivating examples.** Before developing the classification, we describe two algorithms whose deep computations differ sharply in character, for the contrast between them indicates what the classification must distinguish. Both algorithms are treated in detail in Section 4.

Square roots. Newton's method for $p(x) = x^2 - a$, with $a > 0$, acts on the real projective line $\mathbb{RP}^1 = \mathbb{R} \cup \{\infty\}$ by the map $N_p(x) = (x^2 + a)/(2x)$, with $N_p(0) = N_p(\infty) = \infty$. The iterates of N_p are continuous and converge pointwise to a map $\mathfrak{f}$. This map sends every positive number to $\sqrt{a}$, every negative number to $-\sqrt{a}$, and both 0 and ∞ to ∞ (see Figure 4). The deep computations are the iterates together with $\mathfrak{f}$. Since the map $\mathfrak{f}$ is discontinuous, the map is not computable, and the set of iterates is not stable. Nevertheless, the map $\mathfrak{f}$ is the limit of a *sequence* of computations, and the space of deep computations is countable.

Newton's method on the Riemann sphere. For $q(z) = z^3 - 2z + 2$, Newton's method on the Riemann sphere $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ displays three distinct modes of behavior, each confined to a different part of the sphere (see Figure 9 in Section 4.7). The iterates converge on the basins of the three roots. On the basin of the cycle $0 \mapsto 1 \mapsto 0$, by contrast, the iterates do not converge, although the even iterates and the odd iterates converge there separately to different limits. Finally, on the common boundary of these basins, a fractal set, the iterates are chaotic. For the simpler polynomial $z^2 - 1$, chaos of this kind produces deep computations that are not even limits of sequences of computations (see Proposition 4.20 and Theorem 3.1).

The classification of Section 3 is designed to separate such behaviors. In Theorem 3.1 below we characterize, for countably many features, the sets of computations whose deep computations are of the first Baire class on every type shard. For these sets every deep computation is, on each shard, the limit of a sequence of computations, as in the case of square roots. Todorčević's trichotomy, which we recall in Section 3.2, then measures the complexity of the space of such deep computations.

3. Todorčević's trichotomy classifies deep computations

In this section we prove the first main theorem of the paper, which classifies sets of computations by the complexity of the deep computations they generate, and we derive from the theorem a hierarchy of complexity classes. The examples of Section 2.5 show why a classification is needed: some natural algorithms have discontinuous deep computations that are nevertheless limits of sequences of computations, whereas others have deep computations that are not even limits of sequences. The tools are the Bourgain–Fremlin–Talagrand dichotomy (Theorem 1.4) and Todorčević's trichotomy for Rosenthal compacta [Tod99], and our model is the work of Glasner and Megrelishvili on tame semigroups in topological dynamics [GM22].




FIGURE 4. Newton iterates for $x^2 - a$ on $\mathbb{RP}^1$, converging to $\mathfrak{f}$.

3.1. **NIP, Rosenthal compacta, and deep computations.** The classification rests on Theorem 3.1, which we prove in this subsection. This theorem characterizes, by a finitary condition and for countably many features, the sets of computations whose deep computations are of the first Baire class on every type shard. Analogously, the sets of computations whose deep computations are continuous were characterized in the first paper of this series [ADIW26] by Grothendieck's double limit criterion. Recall from Section 1 that $\pi_P \colon \mathbb{R}^{\mathcal{P}} \to \mathbb{R}$ denotes the projection onto the coordinate P. Condition (ii) of the theorem is finitary: unwound through Definition 1.5, the condition states that, for all $P \in \mathcal{P}$ and $r_\bullet \in \mathscr{R}$, all reals $a < b$ and every sequence $(\gamma_n)_{n\in\mathbb{N}}$ in Δ, there are finite disjoint sets $I, J \subseteq \mathbb{N}$ such that

$$L[r_\bullet] \cap \bigcap_{n\in I} (\pi_P \circ \gamma_n)^{-1}(-\infty, a] \cap \bigcap_{n\in J} (\pi_P \circ \gamma_n)^{-1}[b, \infty) = \emptyset.$$

Theorem 3.1. *Let $(L, \mathcal{P}, \Gamma)$ be a compositional computation structure satisfying the Extendibility Axiom, with $\mathcal{P}$ countable. Let $\mathscr{R}$ be an exhaustive collection of sizers, and suppose that $\Delta \subseteq \Gamma$ is $\mathscr{R}$-confined. Then the following conditions are equivalent:*

(i) the inclusion $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}} \subseteq \mathrm{B}_1(\mathcal{L}[r_\bullet], \mathcal{L}[r_\bullet])$ holds for all $r_\bullet \in \mathscr{R}$;
(ii) for all $P \in \mathcal{P}$ and $r_\bullet \in \mathscr{R}$, the set $\pi_P \circ \Delta|_{L[r_\bullet]}$ satisfies the NIP.

Furthermore, if these conditions hold, then every deep computation $f \in \overline{\Delta}$ has an extension $\tilde{f} \colon \mathcal{L}_{sh} \to \mathcal{L}_{sh}$ such that $\tilde{f}|_{\mathcal{L}[r_\bullet]} \in \mathrm{B}_1(\mathcal{L}[r_\bullet], \mathcal{L}[r_\bullet])$ for all $r_\bullet \in \mathscr{R}$. In particular, for every $r_\bullet \in \mathscr{R}$ and every deep computation $f \in \overline{\Delta}$, there is a sequence $(\gamma_n)_{n\in\mathbb{N}}$ in Δ such that $f(v) = \lim_{n\to\infty} \gamma_n(v)$ for every $v \in L[r_\bullet]$.

Proof. Since $\mathcal{P}$ is countable, each type shard $\mathcal{L}[r_\bullet]$ is a compact metrizable, hence Polish, subspace of $\mathbb{R}^{\mathcal{P}}$. Let $r_\bullet \in \mathscr{R}$ be given; by the Extendibility Axiom and confinement, every $\tilde{\gamma}$ with $\gamma \in \Delta$ maps $\mathcal{L}[r_\bullet]$ continuously into itself (see the proof of Lemma 2.7). Hence, for every $P \in \mathcal{P}$, the set $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$ consists of continuous functions and is pointwise bounded. By Theorem 1.9 and Lemma 1.10, condition (ii) holds at $r_\bullet$ if and only if $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}} \subseteq \mathrm{B}_1(\mathcal{L}[r_\bullet], \mathbb{R}^{\mathcal{P}})$, which (i) implies. Conversely, suppose that this inclusion holds. For a Polish space X, every compact subset of

$\mathrm{B}_1(X)$ is Fréchet–Urysohn (see [BFT78, Theorem 3F] or [Deb13, Theorem 4.1]). In a Fréchet–Urysohn space, every point in the closure of a subset is the limit of a sequence of points of the subset. By Proposition 1.8, the closure $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$ may be regarded as a compact subset of $\mathrm{B}_1(\mathcal{L}[r_\bullet] \times \mathcal{P})$, where the space $\mathcal{L}[r_\bullet] \times \mathcal{P}$ is Polish because $\mathcal{P}$ is countable and discrete. Every member of this closure is therefore the pointwise limit of a sequence of maps $\tilde{\gamma}_n|_{\mathcal{L}[r_\bullet]}$ with $\gamma_n \in \Delta$, which are continuous maps from $\mathcal{L}[r_\bullet]$ to itself; hence (i) holds at $r_\bullet$. Finally, if (i) holds, then the extension $\tilde{f}$ of Lemma 2.7 satisfies $\tilde{f}|_{\mathcal{L}[r_\bullet]} \in \overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}} \subseteq \mathrm{B}_1(\mathcal{L}[r_\bullet], \mathcal{L}[r_\bullet])$ for all $r_\bullet \in \mathscr{R}$. By the argument just given, the restriction $\tilde{f}|_{\mathcal{L}[r_\bullet]}$ is the pointwise limit of a sequence of maps $\tilde{\gamma}_n|_{\mathcal{L}[r_\bullet]}$ with $\gamma_n \in \Delta$, and restriction to $L[r_\bullet]$ gives the last assertion. □

3.2. **Todorčević's trichotomy and levels of NIP.** Todorčević's trichotomy now enables us to stratify the NIP sets of computations into three levels. The trichotomy concerns separable Rosenthal compacta, and, under the hypotheses of Theorem 3.1, the deep computations of a countable NIP set of computations form such a compactum on each type shard. We measure the complexity of such a compactum by three topological properties, namely metrizability, hereditary separability and first countability. The three properties form a chain: every compact metrizable space is hereditarily separable, and R. Pol proved that every hereditarily separable Rosenthal compactum is first countable (see [Deb13, Section 10]). These properties define three levels of NIP.

Definition 3.2. Let $(L, \mathcal{P}, \Gamma)$ be a CCS satisfying the Extendibility Axiom, with $\mathcal{P}$ countable, and let $\mathscr{R}$ be an exhaustive collection of sizers. Let $\Delta \subseteq \Gamma$ be an $\mathscr{R}$-confined countable set of computations that satisfies condition (ii) of Theorem 3.1. We say that:

(i) the set Δ is $\mathrm{NIP}^{\mathrm{fc}}$ if $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$ is first countable for every $r_\bullet \in \mathscr{R}$;
(ii) the set Δ is $\mathrm{NIP}^{\mathrm{hs}}$ if $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$ is hereditarily separable for every $r_\bullet \in \mathscr{R}$;
(iii) the set Δ is $\mathrm{NIP}^{\mathrm{met}}$ if $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$ is metrizable for every $r_\bullet \in \mathscr{R}$.

The superscripts abbreviate first countable, hereditarily separable and metrizable. We avoid numerical subscripts because in model theory NIP_k denotes Shelah's k-dependence, a condition that weakens as k grows.


By the chain of properties above, the levels satisfy $\mathrm{NIP}^{\mathrm{met}} \Rightarrow \mathrm{NIP}^{\mathrm{hs}} \Rightarrow \mathrm{NIP}^{\mathrm{fc}} \Rightarrow$ NIP. Moreover, a countable stable set of computations (see Definition 2.8) that is $\mathscr{R}$-confined, which has the NIP by Theorem 3.1, is $\mathrm{NIP}^{\mathrm{met}}$. Indeed, on each type shard $\mathcal{L}[r_\bullet]$ with $r_\bullet \in \mathscr{R}$, confinement makes the deep computations of such a set a compact set and stability makes them continuous, so that they form a separable compact subset of a countable product of spaces $\mathrm{C}_{\mathrm{p}}(\mathcal{L}[r_\bullet])$, and hence a separable Eberlein compactum. Separable Eberlein compacta, in turn, are metrizable [Ark92]. The five compacta of Section 1.4 show that none of the implications between the three topological properties can be reversed for separable Rosenthal compacta. Specifically, the extended Alexandroff compactification $\hat{A}(2^{\mathbb{N}})$ is separable but not first countable. The extended duplicate $\hat{D}(S(2^{\mathbb{N}}))$ of the split Cantor set and the compactum $\hat{D}(2^{\mathbb{N}})$ that contains the Alexandroff duplicate are separable and first countable but not hereditarily separable. Finally, the split Cantor set

$S(2^{\mathbb{N}})$ is hereditarily separable but not metrizable. Todorčević proved that these examples are canonical [Tod99], in the following sense.

Theorem 3.3 (Todorčević's trichotomy [Tod99]; see also [ADK08, Theorem 3])**.** *Let K be a separable Rosenthal compactum.*

(i) If K is hereditarily separable but non-metrizable, then the split Cantor set $S(2^{\mathbb{N}})$ embeds into K.
(ii) If K is first countable but not hereditarily separable, then either the Alexandroff duplicate $D(2^{\mathbb{N}})$ or the extended duplicate $\hat{D}(S(2^{\mathbb{N}}))$ of the split Cantor set embeds into K.
(iii) If K is not first countable, then the Alexandroff compactification $A(2^{\mathbb{N}})$ embeds into K.

Together with the chain of properties above, the trichotomy yields the following classification of separable Rosenthal compacta:



The three levels admit a second description, in terms of the number of tests required to distinguish deep computations from one another. Let Δ and $\mathscr{R}$ be as in Definition 3.2, fix $r_\bullet \in \mathscr{R}$, and write $K_{r_\bullet} = \overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$. We identify each $f \in K_{r_\bullet}$ with the real-valued function $(x, P) \mapsto P(f(x))$ on $\mathcal{L}[r_\bullet] \times \mathcal{P}$, whose points we call *tests*: the test (x, P) applies the feature P to the output on the input x. The topology of pointwise convergence on $K_{r_\bullet}$ is then the topology of pointwise convergence on tests. A set $\mathcal{T}$ of tests *separates* a set of deep computations if any two distinct members of the set differ on some test in $\mathcal{T}$. A member f of $K_{r_\bullet}$ is *countably determined* if some countable set $\mathcal{T}_f$ of tests separates f from every other member of $K_{r_\bullet}$. A set $E \subseteq K_{r_\bullet}$ is *finitely separated* if each $f \in E$ admits finitely many tests and some $\varepsilon_f > 0$ such that every $g \in E \setminus \{f\}$ differs from f by at least ε_f on one of these tests. Since the topology of $K_{r_\bullet}$ is the topology of pointwise convergence on tests, a subset of $K_{r_\bullet}$ is finitely separated exactly when the subset is a discrete subspace of $K_{r_\bullet}$.

Proposition 3.4. *Let Δ and $\mathscr{R}$ be as in Definition 3.2, let $r_\bullet \in \mathscr{R}$ be given, and let $K_{r_\bullet} = \overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$.*

(i) The compactum $K_{r_\bullet}$ is metrizable if and only if one countable set of tests separates $K_{r_\bullet}$.
(ii) The compactum $K_{r_\bullet}$ is first countable if and only if every member of $K_{r_\bullet}$ is countably determined.

(iii) The compactum $K_{r_\bullet}$ is hereditarily separable if and only if no finitely separated subset of $K_{r_\bullet}$ has cardinality $2^{\aleph_0}$.

Proof. (i) If a countable set $\mathcal{T}$ of tests separates $K_{r_\bullet}$, then restriction to $\mathcal{T}$ is a continuous injection of the compact space $K_{r_\bullet}$ into the metrizable space $\mathbb{R}^{\mathcal{T}}$. This injection is therefore a homeomorphism onto the image. Conversely, if $K_{r_\bullet}$ is metrizable, then the square $K_{r_\bullet} \times K_{r_\bullet}$ is second countable, and hence its open subset $\{(f,g) : f \neq g\}$ is Lindelöf. This open set is covered by the open sets $\{(f,g) : f(t) \neq g(t)\}$ for tests t, and a countable subcover gives the required set $\mathcal{T}$. (ii) If f is countably determined, the singleton $\{f\}$ is a countable intersection of open sets, and in a compact Hausdorff space such a point has a countable base of neighborhoods. Conversely, let $(U_k)_{k\in\mathbb{N}}$ be a countable base of neighborhoods of f, and choose in each U_k a basic neighborhood of f defined by a finite set F_k of tests. The union $\mathcal{T}_f = \bigcup_k F_k$ then separates f from every other member of $K_{r_\bullet}$. (iii) As noted before the proposition, the finitely separated subsets of $K_{r_\bullet}$ are the discrete subspaces; by Theorem 3.1, moreover, the compactum $K_{r_\bullet}$ is a separable Rosenthal compactum. A hereditarily separable space has no uncountable discrete subspace, since a separable discrete space is countable. Conversely, suppose that $K_{r_\bullet}$ is not hereditarily separable; then Theorem 3.3 provides a copy of $D(2^{\mathbb{N}})$, of $\hat{D}(S(2^{\mathbb{N}}))$ or of $A(2^{\mathbb{N}})$ inside $K_{r_\bullet}$. Each of these three compacta contains a discrete subspace of cardinality $2^{\aleph_0}$ (see Section 1.4). The existence of such a subspace in every separable Rosenthal compactum that is not hereditarily separable is Pol's theorem (see [ADK08] and [GM22, Theorem 0.4]). □

Thus the set Δ is $\mathrm{NIP}^{\mathrm{met}}$ when, on each type shard, a single countable set of tests separates all the deep computations of Δ. It is $\mathrm{NIP}^{\mathrm{hs}}$ when no type shard carries a finitely separated set of deep computations of the cardinality of the continuum. Finally, the set Δ is $\mathrm{NIP}^{\mathrm{fc}}$ when, on each type shard, each deep computation is separated from all the others by countably many tests of its own.

In Section 4 we exhibit computations at each level of the hierarchy and show that every implication between the classes stable, $\mathrm{NIP}^{\mathrm{met}}$, $\mathrm{NIP}^{\mathrm{hs}}$, $\mathrm{NIP}^{\mathrm{fc}}$ and NIP is strict. The definitions of $\mathrm{NIP}^{\mathrm{fc}}$, $\mathrm{NIP}^{\mathrm{hs}}$ and $\mathrm{NIP}^{\mathrm{met}}$ are topological, and the descriptions of Proposition 3.4 still involve infinitely many tests. Whether the three levels nevertheless admit a finitary, combinatorial characterization, like that of NIP in Theorem 3.1, is Question 6.1.

3.3. **Canonical families and binary search.** Todorčević's trichotomy organizes the hierarchy of the preceding subsection by the topology of the whole space of deep computations. The heptachotomy of Argyros, Dodos and Kanellopoulos [ADK08], which we now recall, works at a finer grain, on families of computations indexed by the binary tree. Recall from Section 1.4 that a dyadic subtree of the binary tree $2^{<\mathbb{N}}$ is written $\{w_t : t \in 2^{<\mathbb{N}}\}$, indexed through the canonical isomorphism preserving extension and the lexicographic order.

Definition 3.5. Let X and Y be Polish spaces.

(1) If I is countable and $\{f_i : i \in I\} \subseteq \mathbb{R}^X$, $\{g_i : i \in I\} \subseteq \mathbb{R}^Y$ are two families indexed by I, we say that they are *equivalent* if the map $f_i \mapsto g_i$ extends to a homeomorphism from $\overline{\{f_i : i \in I\}}$ to $\overline{\{g_i : i \in I\}}$.

(2) If $\{f_t : t \in 2^{<\mathbb{N}}\} \subseteq \mathbb{R}^X$ is a pointwise bounded family, we say that it is *minimal* if, for every dyadic subtree $\{w_t : t \in 2^{<\mathbb{N}}\}$ of $2^{<\mathbb{N}}$, the family $\{f_{w_t} : t \in 2^{<\mathbb{N}}\}$ is equivalent to $\{f_t : t \in 2^{<\mathbb{N}}\}$.

The seven canonical minimal families in the classification of Argyros, Dodos and Kanellopoulos [ADK08] are described below. We retain their notation and the tree indexing, which distinguishes families even when their closures are homeomorphic. The description is carried out along the regular dyadic subtree $\{s_t : t \in 2^{<\mathbb{N}}\}$ fixed in Section 1.4 for the definition of the extended duplicate of the split Cantor set. This subtree satisfies $s_t^\frown 0^\infty \neq s_{t'}^\frown 0^\infty$ and $s_t^\frown 1^\infty \neq s_{t'}^\frown 1^\infty$ whenever $t \neq t'$. The families are built from the prefix tests v_t and the threshold tests $\chi_{[a}$, $\chi_{(a}$ of Section 1.4. Two further conventions are taken from the paper of Argyros, Dodos and Kanellopoulos [ADK08]. First, a point $\sigma \in 2^{\mathbb{N}}$, identified with a point of the Cantor set in $[0,1]$, is regarded as the constant function on $2^{\mathbb{N}}$ with value σ. Second, a pair (f,g) of functions on $2^{\mathbb{N}}$ is regarded as a single function on the disjoint union $2^{\mathbb{N}} \sqcup 2^{\mathbb{N}}$.

(1) The family $D_1 = \{\frac{1}{|t|+1} v_t : t \in 2^{<\mathbb{N}}\}$ has closure $D_1 \cup \{\mathbf{0}\}$, the one-point compactification of a countable discrete set.
(2) The family $D_2 = \{s_t^\frown 0^\infty : t \in 2^{<\mathbb{N}}\}$ is a discrete subset of its closure $\overline{D_2}$, which Argyros, Dodos and Kanellopoulos denote by $2^{\leq \mathbb{N}}$. This closure is compact and metrizable, and its accumulation points form a copy of $2^{\mathbb{N}}$.
(3) The family $D_3 = \{\chi_{[s_t^\frown 0^\infty} : t \in 2^{<\mathbb{N}}\}$ is a discrete subset of its closure $\overline{D_3}$, a closed subspace of $S(2^{\mathbb{N}})$.
(4) The family $D_4 = \{\chi_{(s_t^\frown 1^\infty} : t \in 2^{<\mathbb{N}}\}$ is a discrete subset of its closure $\overline{D_4}$, a closed subspace of $S(2^{\mathbb{N}})$.
(5) The family $D_5 = \{v_t : t \in 2^{<\mathbb{N}}\}$ is a discrete subset of its closure $\overline{D_5} = \hat{A}(2^{\mathbb{N}})$.
(6) The family $D_6 = \{(v_t, t^\frown 0^\infty) : t \in 2^{<\mathbb{N}}\}$ is a discrete subset of its closure $\overline{D_6} = \hat{D}(2^{\mathbb{N}})$.
(7) The family $D_7 = \{(v_{s_t}, \chi_{[s_t^\frown 0^\infty}) : t \in 2^{<\mathbb{N}}\}$ is a discrete subset of its closure $\overline{D_7} = \hat{D}(S(2^{\mathbb{N}}))$.

Proposition 3.6 (Canonical minimal families [ADK08, Section 4.3])**.** *The seven families $D_1, \ldots, D_7$ defined above are relatively compact in the first Baire class on their respective Polish domains. The seven families are minimal and pairwise nonequivalent under the tree-indexed equivalence just defined.*

Theorem 3.7 (The heptachotomy of Argyros, Dodos and Kanellopoulos [ADK08, Theorem 2])**.** *Let X be a Polish space, and let $\{f_t : t \in 2^{<\mathbb{N}}\}$ be a family of pairwise distinct real-valued functions on X, relatively compact in $\mathrm{B}_1(X)$. Then there is a regular dyadic subtree $\{w_t : t \in 2^{<\mathbb{N}}\}$ of $2^{<\mathbb{N}}$ such that the family $\{f_{w_t} : t \in 2^{<\mathbb{N}}\}$ is equivalent to exactly one of the families $D_1, \ldots, D_7$.*

The heptachotomy and the trichotomy answer different questions. The trichotomy concerns the closure of a family as a topological space, and the indexing of the family plays no role in the trichotomy. The heptachotomy keeps the indexing by the binary tree and classifies, after passage to a subtree, the routes by which the family approaches its limits: along a branch, or departing from the branch below or above. Through their closures, the seven families fall into four levels of the trichotomy [ADK08, Section 4.3]. The closures of D_1 and D_2 are metrizable, and

those of D_3 and D_4 are hereditarily separable and not metrizable. The closures of D_6 and D_7 are first countable and not hereditarily separable, and the closure of D_5 is not first countable. Families at the same level may therefore follow different routes, and Section 4.6 shows that the difference between the routes reflects a concrete feature of an algorithm.

The building blocks of the families $D_1, \ldots, D_7$ admit a computational interpretation as primitives of binary search on $2^{\mathbb{N}}$, the space of binary expansions. A node $t \in 2^{<\mathbb{N}}$ determines the dyadic interval $[t^\frown 0^\infty, t^\frown 1^\infty]$, the set of inputs that round to t at precision $|t|$. The prefix test v_t asks whether the input rounds to t, and the point $t^\frown 0^\infty$ is the rounded value itself. The comparison $\chi_{[t^\frown 0^\infty}$ asks whether the input lies at or to the right of the left endpoint of the interval. Likewise, the comparison $\chi_{(t^\frown 1^\infty}$ asks whether the input lies strictly to the right of the right endpoint. The resulting reading of the seven families is recorded in Table 1.

Family	Function at node t	Limits
D_1	prefix test weighted by $1/(\|t\|+1)$	the test that never succeeds: confidence vanishes with depth
D_2	return the rounded value $t^\frown 0^\infty$	the exact input: successive approximation
D_3	compare with the left endpoint	comparison with a limit of left endpoints, with ties broken upward and, at limits approached from above, also downward
D_4	compare with the right endpoint	comparison with a limit of right endpoints, with ties broken downward and, at limits approached from below, also upward
D_5	prefix test	exact equality, or no match
D_6	prefix test and rounded value	a dictionary lookup that returns its key
D_7	prefix test and comparison	a three-way comparison: below, inside or above

TABLE 1. The minimal families of Argyros, Dodos and Kanellopoulos as primitives of binary search.

The translation to a concrete computation is made after fixing the magnitude bounds specified by the parameters of the computation. Let $r_\bullet$ be the resulting sizer, and let $\{\gamma_t : t \in 2^{<\mathbb{N}}\}$ be a family of computations in a CCS satisfying the Extendibility Axiom and having countably many features. Suppose that the family preserves the shard $L[r_\bullet]$ and has the NIP there for every feature. For a chosen feature P, the argument of Theorem 3.1 supplies a relatively compact family of scalar functions $\{P \circ \tilde{\gamma}_t|_{\mathcal{L}[r_\bullet]} : t \in 2^{<\mathbb{N}}\}$ in the first Baire class on the type shard $\mathcal{L}[r_\bullet]$. Restriction to a shard or observation through a single feature can identify distinct computations; a constant feature, for example, gives a constant family. Accordingly, the seven families are used here as canonical patterns of observable limit behavior, whose occurrence in a particular computation requires a separate identification.

4. Todorčević's trichotomy is realized by classical algorithms

In this section we realize every alternative of Todorčević's trichotomy (Theorem 3.3) by a classical algorithm, and with these realizations we show that every implication of the hierarchy of Section 3 is strict. By selecting computations along the binary tree, we also realize in the same algorithms all seven canonical families of the heptachotomy (Theorem 3.7 and Section 4.6). The section follows the alternatives of the trichotomy in the order of Theorem B of the Introduction. Convergent methods and rotations have metrizable spaces of deep computations; Bresenham's algorithm in the plane yields the split Cantor set; Bresenham's algorithm in space yields the extended duplicate of the split Cantor set; and the steps of the Euclidean algorithm yield the Alexandroff duplicate and, through exact equality tests, the Alexandroff compactification. For each algorithm we describe the computations and their features, the deep computations, the canonical compactum that they contain, the invariant that decides the level, and the tests that do or do not tell the deep computations apart (see Proposition 3.4). At all levels but one, the computations are the powers of a single step map; the Euclidean constructions use the group generated by two steps, and a countable family of its elements. The section closes with chaotic iterations, Newton's method on the Riemann sphere among them, which lie beyond both classifications. For the algorithms of this section the hierarchy coincides with the Glasner–Megrelishvili hierarchy of tame dynamical systems [GM22], and we begin with this coincidence (see Proposition 4.1); Table 3 at the end of the section summarizes the examples.

4.1. **Iterative algorithms and tame dynamics.** The algorithms of this section act by continuous steps on a compact space of states. Within this setting, which we now isolate, our hierarchy coincides with the Glasner–Megrelishvili hierarchy of tame dynamical systems. The coincidence allows the results of Glasner and Megrelishvili to be brought to bear on the examples that follow. A *dynamical CCS* is a compositional computation structure $(L, \mathcal{P}, \Gamma)$ in which L is a compact metrizable space and the predicates in $\mathcal{P}$ are continuous and separate the points of L. Moreover, the family $\mathcal{P}$ and the semigroup Γ are countable, and the members of Γ are continuous maps $L \to L$, among them the identity. A dynamical CCS is a *cascade CCS* if $\Gamma = \{\gamma^n : n \in \mathbb{N}\}$ is the semigroup of iterates of a single continuous map $\gamma\colon L \to L$. Since $\mathcal{P}$ separates the points of the compact space L, the map $v \mapsto \mathrm{tp}(v)$ is a homeomorphism of L onto the space of types. Consequently a single shard covers L, every type is realized, and the Extendibility Axiom holds with $\tilde{\gamma} = \gamma$ for every $\gamma \in \Gamma$. The deep computations of a dynamical CCS form the pointwise closure

$$E(L, \Gamma) = \overline{\Gamma} \subseteq L^L,$$

which is the *enveloping semigroup*, introduced by Ellis, of the dynamical system (L, Γ); for a cascade CCS we write $E(L, \gamma)$. The system (L, Γ) is called *tame* if, for every continuous function $f\colon L \to \mathbb{R}$, the family $\{f \circ \gamma : \gamma \in \Gamma\}$ has the NIP of Definition 1.5 [Köh95, Gla06]. We say that a dynamical CCS is *stable* when the set Γ of computations has the property of Definition 2.8. Similarly, the CCS is called NIP when Γ satisfies condition (ii) of Theorem 3.1 with respect to a single sizer $r_\bullet$ with $L[r_\bullet] = L$. With respect to the same sizer, it is called $\mathrm{NIP}^{\mathrm{fc}}$, $\mathrm{NIP}^{\mathrm{hs}}$ or $\mathrm{NIP}^{\mathrm{met}}$ when Γ has the corresponding property of Definition 3.2.

A map $\xi\colon L \to L$ is *of Baire class* 1 if $f \circ \xi$ is of Baire class 1 for every continuous $f\colon L \to \mathbb{R}$. It suffices that $P \circ \xi$ be of Baire class 1 for every $P \in \mathcal{P}$. Indeed, since the type map embeds L into $\mathbb{R}^{\mathcal{P}}$ with $\mathcal{P}$ countable, every open subset of L is a countable union of finite intersections of sets $P^{-1}(U)$ with $P \in \mathcal{P}$ and $U \subseteq \mathbb{R}$ open. By Fact 1.2, the preimage of every open set under ξ is then an F_σ set, and so is the preimage of every open set under $f \circ \xi$. A second appeal to Fact 1.2 shows that the function $f \circ \xi$ is of Baire class 1. In general, the definition of Baire class 1 given above for maps from L to L is less restrictive than the one adopted in Section 1. The latter requires a map from L to L to be the pointwise limit of a sequence of continuous maps from L to L. For example, if $L = [0,1] \cup \{2\} \subseteq \mathbb{R}$, then the map that sends $[0, 1/2]$ to 0 and the rest of L to 2 is of Baire class 1 in the sense defined above. This map, however, is not the pointwise limit of a sequence of continuous maps from L to L, because each such map sends the connected set $[0,1]$ into $[0,1]$ or to 2. For the deep computations of a dynamical CCS, by contrast, the two definitions agree. If every deep computation is of Baire class 1 in the sense defined above, then the CCS is NIP, as we show in Proposition 4.1 below. In that case the last assertion of Theorem 3.1, applied to the single sizer $r_\bullet$ with $L = L[r_\bullet]$, shows that every deep computation is the pointwise limit of a sequence of computations. Since computations are continuous maps from L to L, every deep computation is then of Baire class 1 in the sense of Section 1 as well.

Proposition 4.1. *A dynamical CCS is NIP if and only if the dynamical system (L, Γ) is tame, and if and only if every deep computation is of Baire class* 1*. In that case the CCS is NIP^{fc}, NIP^{hs} or NIP^{met} if and only if $E(L, \Gamma)$ is first countable, hereditarily separable or metrizable, respectively.*


Proof. The single sizer $r_\bullet$ with $L = L[r_\bullet]$ forms an exhaustive collection that confines Γ. If (L, Γ) is tame, the CCS is NIP: take $f = P$ for $P \in \mathcal{P}$. If the CCS is NIP, then Theorem 3.1 and Proposition 1.8 show that the function $P \circ \xi$ is of Baire class 1 for every $\xi \in E(L, \Gamma)$ and $P \in \mathcal{P}$. Every deep computation is therefore of Baire class 1. Finally, suppose that every deep computation is of Baire class 1, and let $f\colon L \to \mathbb{R}$ be continuous. The map $\xi \mapsto f \circ \xi$ is continuous on L^L, so the closure of $\{f \circ \gamma : \gamma \in \Gamma\}$ is the compact set $\{f \circ \xi : \xi \in E(L, \Gamma)\}$, which lies in $\mathrm{B}_1(L)$. By Theorem 1.6, the family $\{f \circ \gamma : \gamma \in \Gamma\}$ has the NIP, so (L, Γ) is tame. The last statement is Definition 3.2, since $\overline{\widetilde{\Gamma}|_L} = E(L, \Gamma)$. □

For metrizable systems, Glasner and Megrelishvili call a tame system *tame*$_1$ when its enveloping semigroup is first countable, *tame*$_2$ when it is hereditarily separable, and *hereditarily non-sensitive* when it is metrizable. They prove the inclusions between these classes [GM22, Proposition 0.6] and show by examples that the inclusions are proper. It follows from Proposition 4.1 that a dynamical CCS is NIP^{fc}, NIP^{hs} or NIP^{met} exactly when the dynamical system (L, Γ) is tame$_1$, tame$_2$ or hereditarily non-sensitive, respectively.

The simplest iterative methods are those whose iterates converge pointwise. In Proposition 4.2 we show that all of them are NIP^{met}, and that the continuity of the limit decides whether they are stable.

Proposition 4.2. *Let $(L, \mathcal{P}, \Gamma)$ be a cascade CCS, with $\Gamma = \{\gamma^n : n \in \mathbb{N}\}$ for a continuous map $\gamma\colon L \to L$, and suppose that the iterates γ^n converge pointwise to a map $\gamma_\infty\colon L \to L$. Then $E(L, \gamma) = \{\gamma^n : n \in \mathbb{N}\} \cup \{\gamma_\infty\}$, and the CCS is NIP^{met}. The CCS is stable if and only if γ_∞ is continuous.*

Proof. The set of iterates together with their pointwise limit γ_∞ is a compact subset of the Hausdorff space L^L, hence closed, and the enveloping semigroup $E(L,\gamma)$ therefore consists of the iterates and γ_∞. For each $P \in \mathcal{P}$, the map $P \circ \gamma_\infty$ is the pointwise limit of the continuous functions $P \circ \gamma^n$ and is therefore of Baire class 1; by Proposition 4.1, the CCS is NIP. The space $E(L,\gamma)$ is a countable compact Hausdorff space, hence metrizable, so the CCS is $\mathrm{NIP}^{\mathrm{met}}$. The members of $E(L,\gamma)$ other than γ_∞ are continuous, which gives the statement about stability. □



4.2. **Convergent methods and rotations have metrizable spaces of limits.** Our descent through the hierarchy starts at the top two levels, stable and $\mathrm{NIP}^{\mathrm{met}}$, at which the deep computations form a metrizable space. A convergent method forgets how its limit is approached, and a rotation recovers the point approached but not the side; selected iterates of the two realize the canonical families D_1 and D_2 (Section 4.6). At these two levels we place a rotation of the circle, Newton's method for square roots, the power method and the logistic map for $0 < r \leq 3$. By Proposition 4.2, every convergent method lies at one of these two levels. Convergence, however, is not necessary for stability, as the first example shows.

A stable example: rotation. Let the space of states be the circle $L = \mathbb{T} = \mathbb{R}/\mathbb{Z}$, with the predicates $\cos 2\pi x$ and $\sin 2\pi x$, and let the step be the rotation $\gamma(x) = x+\alpha$ by an irrational number α. Since a pointwise limit of rotations is a rotation and the points $n\alpha$ with $n \in \mathbb{N}$ are dense in $\mathbb{T}$, the deep computations are all the rotations $x \mapsto x + c$ with $c \in \mathbb{T}$. These rotations are continuous, so the rotation CCS is stable, and the space of deep computations of the rotation CCS is the circle group, a metrizable compactum.

Square roots: $\mathrm{NIP}^{\mathrm{met}}$ but not stable. We return to the first motivating example of Section 2.5, Newton's method on the real projective line for the polynomial $p(x) = x^2 - a$ with real $a > 0$, whose iterates approximate the square roots $\pm\sqrt{a}$. As for every rational function with real coefficients, the Newton map N_p maps the real projective line $\mathbb{RP}^1 = \mathbb{R} \cup \{\infty\}$ into itself. Newton's method for p, the iteration of its Newton map $N_p(x) = x - p(x)/p'(x)$, may therefore be regarded as a CCS with state space $\mathbb{RP}^1$.

Let $\mathbb{R} \to S^1$, $x \mapsto (P_1(x), P_3(x))$, be the inverse stereographic projection of the real line onto $S^1 \subseteq \mathbb{R}^2$, given by

$$P_1(x) = \frac{2x}{x^2+1}, \qquad P_3(x) = \frac{x^2-1}{x^2+1},$$

and extended to a homeomorphism of the real projective line $\mathbb{RP}^1 = \mathbb{R} \cup \{\infty\}$ onto S^1 by $\infty \mapsto (P_1(\infty), P_3(\infty)) = (0,1)$. These coordinates are the restrictions to $\mathbb{RP}^1$ of two of the coordinates of the Riemann sphere used in Section 4.7, whose names they keep. As predicates, the two coordinates make $(\mathbb{RP}^1, (P_1, P_3))$ a computation states structure with a compact state space. The Newton map N_p is

$$x \mapsto x - \frac{x^2-a}{2x} = \begin{cases} \frac{x^2+a}{2x}, & x \neq 0, \infty; \\ \infty, & x = 0, \infty. \end{cases}$$

From every initial point $x_0 \neq 0, \infty$, the Newton steps converge to one of the square roots $\pm\sqrt{a}$. More precisely, the iterates of N_p converge pointwise to the map

$\mathfrak{f}\colon \mathbb{RP}^1 \to \mathbb{RP}^1$ given by

$$\mathfrak{f}(x) = \lim_{n\to\infty} N_p^n(x) = \begin{cases} \sqrt{a}, & x \in (0,\infty); \\ -\sqrt{a}, & x \in (-\infty, 0); \\ \infty, & x = 0, \infty. \end{cases}$$

The map $\mathfrak{f}$ is therefore discontinuous at 0 and at ∞ (see Figure 4). Proposition 4.2 therefore shows that the square-root CCS is NIP^{met} but not stable.

The power method: NIP^{met} but not stable. The power method computes a dominant eigenvector of a matrix by repeated multiplication and normalization. Let $d \geq 2$ be an integer, and let $A = \operatorname{diag}(\lambda_1, \ldots, \lambda_d)$ be a diagonal matrix with $\lambda_1 > \lambda_2 > \cdots > \lambda_d > 0$. The space of states is the real projective space $L = \mathbb{RP}^{d-1}$ of lines through the origin in $\mathbb{R}^d$, in which $[v]$ denotes the line spanned by a nonzero vector v. The family $\mathcal{P}$ consists of the predicates $P_{ij}([v]) = v_i v_j / \|v\|^2$ with $1 \leq i, j \leq d$, which are the entries of the orthogonal projection onto the line $[v]$. The step of the algorithm is $\gamma([v]) = [Av]$. For $[v] \in \mathbb{RP}^{d-1}$, let $i(v)$ be the least index i with $v_i \neq 0$, and define $\ell([v]) = [e_{i(v)}]$, where $e_1, \ldots, e_d$ is the standard basis of $\mathbb{R}^d$.

Proposition 4.3. *The deep computations of the power method are $E(L,\gamma) = \{\gamma^n : n \in \mathbb{N}\} \cup \{\ell\}$. Hence the power method is NIP^{met}. The deep computation ℓ, the only one that is not an iterate, is of Baire class* 1 *and discontinuous, so the power method is not stable.*

Proof. For $i = i(v)$, we have $\lambda_i^{-n} A^n v = v_i e_i + \sum_{j>i} (\lambda_j/\lambda_i)^n v_j e_j \xrightarrow{n\to\infty} v_i e_i$, so the iterates γ^n converge pointwise to ℓ, and Proposition 4.2 applies. The map ℓ is discontinuous at every $[v]$ with $i(v) > 1$, since $[v + te_1] \xrightarrow{t\to 0} [v]$ while $\ell([v + te_1]) = [e_1] \neq \ell([v])$. □

The deep computation ℓ sends each input to the eigenline of the largest eigenvalue whose eigenvector has a nonzero component in the input. Being discontinuous on the space of types L, the map ℓ is not computable in the sense of Section 2. The map ℓ is nevertheless the limit of a sequence of computations, and the space of deep computations of the power method is countable and metrizable. A single test, one feature evaluated at one input, separates all these deep computations (see Proposition 3.4(i)). Take the input $[e_1 + e_2]$ and the feature P_{22}. At the iterate γ^n the test returns $P_{22}([\lambda_1^n e_1 + \lambda_2^n e_2]) = \lambda_2^{2n}/(\lambda_1^{2n} + \lambda_2^{2n})$, which decreases strictly with n, and at the limit ℓ it returns 0, since $\ell([e_1 + e_2]) = [e_1]$. Distinct members of $E(L,\gamma)$ therefore return distinct values at this one test.

Both levels: the logistic map. The logistic map $f_r(x) = rx(1-x)$, with $0 < r \leq 4$, maps $[0,1]$ into itself. Its CCS has states $L = [0,1]$, the single predicate $x \mapsto x$, and the semigroup of iterates of f_r (see Figure 5). For $0 < r \leq 1$, let ℓ_r be the constant map 0; for $1 < r \leq 3$, let ℓ_r be the map with $\ell_r(x) = 1 - 1/r$ for $0 < x < 1$ and $\ell_r(0) = \ell_r(1) = 0$.

Proposition 4.4. *For $0 < r \leq 3$, the iterates of f_r converge pointwise to ℓ_r. Consequently the logistic CCS is stable for $0 < r \leq 1$, and NIP^{met} but not stable for $1 < r \leq 3$.*

Proof. Write $f = f_r$ and $x^* = 1 - 1/r$, and note that $f(x)/x = r(1-x)$ for $x > 0$. If $r \leq 1$, then $f(x) < x$ for $0 < x \leq 1$, so every orbit decreases to a fixed point of f, and 0 is the only fixed point in $[0,1]$.

Suppose that $1 < r \le 2$, in which case $0 < x^* \le \frac{1}{2}$. The inequality $f(x) > x$ holds for $0 < x < x^*$, and the reverse inequality for $x^* < x \le 1$. For $0 < x < 1$ we have $f(x) \in (0, \frac{r}{4}] \subseteq (0, \frac{1}{2}]$, and the function f, which fixes x^*, is increasing on $[0, \frac{1}{2}]$. Hence the orbit of $f(x)$ is increasing and bounded by x^* if $f(x) < x^*$, and decreasing and bounded below by x^* if $f(x) > x^*$. In both cases, as well as trivially when $f(x) = x^*$, the orbit of $f(x)$ converges to a fixed point of f in $(0, \frac{1}{2}]$, namely to x^*.

Suppose that $2 < r \le 3$, and let $b = f(c)$, where $c = \frac{r}{4}$ is the maximum of f. We have $x^* < c$, since $c - x^* = (r-2)^2/(4r)$, and therefore $b < f(x^*) = x^*$, since f is strictly decreasing on $[\frac{1}{2}, 1]$. We also have $b \ge 1/r$, because $b - 1/r = (r^3(4-r) - 16)/(16r)$ and the polynomial $r^3(4-r)$ increases from 16 on $[2, 3]$. The concave function f takes the value x^* at $1/r$ and at x^*, so $f \ge x^*$ on $[1/r, x^*]$. Hence the function f maps $J_- = [b, x^*]$ into $J_+ = [x^*, c]$, and, being decreasing on J_+, it maps J_+ onto J_-. Fix x with $0 < x < 1$; the orbit of $f(x)$ then lies in $(0, c]$, and it meets $J_- \cup J_+ = [b, c]$. Indeed, if the orbit did not meet $[b, c]$, all its points would lie in $(0, b)$, where $f(y) > y$ because $b < x^*$. The orbit would then increase to a fixed point of f in $(0, b]$. No such fixed point exists, since the only positive fixed point of f is x^*, which exceeds b. Since f maps $J_- \cup J_+$ into itself, it remains to be shown that the orbit of every $y \in J_- \cup J_+$ converges to x^*. The points of period 2 of f that are not fixed are the roots of $r^2y^2 - r(r+1)y + (r+1)$, whose discriminant $r^2(r+1)(r-3)$ is negative for $r < 3$; for $r = 3$ the only root is x^*. Hence the function $f(f(y)) - y$ vanishes on $[0, 1]$ only at 0 and x^*, and it is negative on $(x^*, 1]$, since it is negative at 1. For $y \in J_+$ with $y > x^*$, the point $f(f(y))$ lies in J_+, so $x^* \le f(f(y)) < y$. The sequence $(f^{2n}(y))$ therefore decreases to a fixed point of $f \circ f$ in $[x^*, y]$, that is, to x^*, whence the odd iterates $f^{2n+1}(y)$ converge to $f(x^*) = x^*$. Every $y \in J_-$ satisfies $f(y) \in J_+$, so its orbit converges to x^* as well.

Finally, the equalities $f(0) = f(1) = 0$ account for the endpoints, and in all cases the iterates therefore converge pointwise to ℓ_r. The map ℓ_r is continuous for $r \le 1$, and discontinuous at 0 for $r > 1$, so Proposition 4.2 gives the remaining statements. □





FIGURE 5. The logistic family: long-run values (left) and two orbits for $r = 2.8$ (right).

We return to the logistic family for $3 < r \leq 4$ at the end of the section, where we complete its classification (see Proposition 4.23 and Remark 4.24).



4.3. **Rounding produces the split Cantor set.** Bresenham's line algorithm in the plane, which we analyze in this subsection, lies at the level $\mathrm{NIP}^{\mathrm{hs}}$ but not $\mathrm{NIP}^{\mathrm{met}}$, immediately below the metrizable level. Below the metrizable level, no countable set of tests separates the deep computations (see Proposition 3.4), a test being a pair of an input and a feature (see Section 3.2). Rounding gives rise to spaces of deep computations of this kind, since an algorithm that rounds at each step takes a new discrete decision at each step. The decision bit records the side from which a limit is approached, and the same bit lets selected iterates realize the canonical families D_3 and D_4 (Remark 4.7).

Drawing a line. Fix an irrational number α with $0 < \alpha < 1$. To draw the line $y = \alpha x + \rho$ on a screen, Bresenham's algorithm [Bre65] moves one pixel to the right at each step and decides whether to move one pixel up. The n-th decision is $b_n(\rho) = \lfloor (n+1)\alpha + \rho \rfloor - \lfloor n\alpha + \rho \rfloor = [\![\{n\alpha + \rho\} \in [1-\alpha, 1)]\!]$, where $\{\cdot\}$ denotes the fractional part, and the state of the algorithm is the error term ρ modulo 1. The step $\rho \mapsto \rho + \alpha$ is a rotation of the circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$, but the decision bit is not a continuous function of ρ. The standard remedy [MH40, Lot02] is to take the decision sequences themselves as states.

Let $\mathcal{O} = \{-k\alpha : k \in \mathbb{Z}\} \subseteq \mathbb{T}$, a countable dense set containing both endpoints 0 and $1-\alpha$ of the decision interval. The *split circle* $\mathbb{T}_{\mathcal{O}}$ is obtained from $\mathbb{T}$ by replacing each $t \in \mathcal{O}$ by two points $t^- < t^+$, with the circular order topology; the split circle is a compact metrizable space, homeomorphic to the Cantor set. The rotation by α preserves $\mathcal{O}$ and lifts to a homeomorphism R of $\mathbb{T}_{\mathcal{O}}$ that preserves the labels $\pm$. The decision interval becomes clopen in $\mathbb{T}_{\mathcal{O}}$, and the coding map $\mathbb{T}_{\mathcal{O}} \to 2^{\mathbb{Z}}$, $y \mapsto ([\![R^n y \in [1-\alpha, 1)]\!])_{n \in \mathbb{Z}}$, is a homeomorphism onto the *Sturmian shift* X_α that conjugates R to the shift σ. The *Bresenham CCS* has states $L = X_\alpha$, the coordinate predicates $P_j(y) = y_j$ for $j \in \mathbb{Z}$, and the semigroup $\Gamma = \{\sigma^n : n \in \mathbb{N}\}$.

A point of $\mathbb{T}_{\mathcal{O}}$ lies *over* $t \in \mathbb{T}$ if the natural projection $\mathbb{T}_{\mathcal{O}} \to \mathbb{T}$ maps the point to t. For $c \in \mathbb{T}$, let R_c^+ (respectively R_c^-) be the map of $\mathbb{T}_{\mathcal{O}}$ that sends a point over t to $(t+c)^+$ (respectively $(t+c)^-$) when $t + c \in \mathcal{O}$, and to $t + c$ otherwise. Through the coding map, we regard R_c^+ and R_c^- as maps of X_α. The *double arrow circle* $\mathbb{D}$ is $\mathbb{T} \times \{-, +\}$ with the lexicographic circular order and the topology of this order, and we also write c^- and c^+ for the points $(c, -)$ and $(c, +)$ of $\mathbb{D}$. The double arrow circle is compact, first countable, hereditarily separable and not metrizable [Eng89, 3.10.C].

Theorem 4.5. *The Bresenham CCS is NIP, and the space of deep computations of the CCS is*

$$E(X_\alpha, \sigma) = \{\sigma^n : n \in \mathbb{N}\} \cup \{R_c^+, R_c^- : c \in \mathbb{T}\},$$

in which each σ^n is isolated. The map $c^\pm \mapsto R_c^\pm$ is a homeomorphism of $\mathbb{D}$ onto $\{R_c^+, R_c^- : c \in \mathbb{T}\}$. In particular, the Bresenham CCS is $\mathrm{NIP}^{\mathrm{hs}}$ but not $\mathrm{NIP}^{\mathrm{met}}$.

Proof. To prove that the CCS is NIP, note that the predicates $P_j \circ \sigma^n$ are $\{0,1\}$-valued. Moreover, the j-th coordinate of $\sigma^n y$ is 1 exactly when $R^{n+j} y$ lies in the clopen arc $[1-\alpha, 1)$ of $\mathbb{T}_{\mathcal{O}}$, that is, when y lies in an arc of $\mathbb{T}_{\mathcal{O}}$. A family of arcs of a circularly ordered set has VC dimension at most 3. By the remark at the end of Section 1.5, the characteristic functions of the members of such a family therefore have the NIP.

For the deep computations, let γ be the limit of σ^{n_i} along an ultrafilter $\mathcal{U}$, and let $c \in \mathbb{T}$ be the limit of $n_i\alpha$ along $\mathcal{U}$. If n_i is constant on a member of $\mathcal{U}$, then γ is a power of σ. Otherwise, since the points $n\alpha$ are distinct, the ultrafilter decides whether $n_i\alpha$ approaches c from the right or from the left. In the first case, for every point y over t, the points $R^{n_i}y$ over $t + n_i\alpha$ approach $t + c$ strictly from the right. These points converge in $\mathbb{T}_{\mathcal{O}}$ to $(t + c)^+$ if $t + c \in \mathcal{O}$ and to $t + c$ otherwise, whence $\gamma = R_c^+$; the second case gives $\gamma = R_c^-$. Conversely, for every N, every c is approached from both sides by the points $n\alpha$ with $n \geq N$; hence every $R_c^\pm$ is a deep computation. The maps R_c^+ and R_c^- are distinct, since the two maps differ at the points over $d - c$ with $d \in \mathcal{O}$. The same argument shows that, whenever a net $(c_i^{\varepsilon_i})$ converges to c^+ in $\mathbb{D}$, the maps $R_{c_i}^{\varepsilon_i}$ converge pointwise to R_c^+, and symmetrically for c^-. The map $c^\pm \mapsto R_c^\pm$ is therefore a continuous bijection from the compact space $\mathbb{D}$ onto a Hausdorff space, hence a homeomorphism. Finally, fix $d \in \mathcal{O}$ and consider the two points $(d - n\alpha)^\pm$ of $\mathbb{T}_{\mathcal{O}}$, which σ^n sends to d^- and d^+. Every map $R_c^\pm$ sends the two points to the same point, and so does every pointwise limit of maps σ^{m_i} with m_i tending to infinity, by the argument above. Hence no net of such maps, and no net of maps σ^m with $m \neq n$ and m bounded, converges to σ^n, so σ^n is isolated. Thus $E(X_\alpha, \sigma)$ is the union of a countable set and a copy of $\mathbb{D}$, and the space $E(X_\alpha, \sigma)$ is therefore hereditarily separable and not metrizable. □

The two deep computations R_c^+ and R_c^- are the two ways of drawing the line infinitely far away, where the accumulated error has reached c. The two maps agree except at the pixels where the line passes exactly through a corner, and at those pixels they apply the two opposite rounding conventions. Figure 6 shows the two conventions.

The topology of Theorem 4.5 has a direct reading in terms of tests (see Proposition 3.4). The maps R_c^+ and R_c^- differ only at the points of X_α that lie over $\mathcal{O} - c$, the inputs at which the line passes exactly through a pixel corner. A countable set of tests involves the inputs over countably many points $t_1, t_2, \ldots$ of $\mathbb{T}$, and every c outside the countable set $\bigcup_i (\mathcal{O} - t_i)$ gives two distinct deep computations R_c^+ and R_c^- that agree on all these tests. No countable set of tests therefore separates the deep computations, and this is the nonmetrizability of $E(X_\alpha, \sigma)$. Each deep computation is nevertheless determined by countably many tests of its own. The coordinates of the output at one fixed input y_0 determine the point $R_c^\pm(y_0)$, which lies over $\pi(y_0) + c$, where $\pi\colon \mathbb{T}_{\mathcal{O}} \to \mathbb{T}$ is the natural projection, and hence determine c. The coordinate P_0 at the inputs over $-c$, of which there are two when $-c \in \mathcal{O}$, then tells R_c^+, which sends these inputs to 0^+, from R_c^-, which sends them to 0^-, and both from an iterate, which keeps the two inputs apart. The iterates themselves are isolated, and finitely many tests determine each of them. Determination by countably many tests does not mean isolation; by Proposition 3.4(ii), it means that each deep computation has a countable base of neighborhoods.

By Theorem 3.1, the space $E(X_\alpha, \sigma)$, the closure of a countable set of computations of an NIP dynamical CCS, is a separable Rosenthal compactum. Being hereditarily separable and not metrizable, the compactum contains, by Todorčević's trichotomy (see Theorem 3.3), a copy of the split Cantor set $S(2^{\mathbb{N}})$. Such a copy can be exhibited explicitly, with iterates as the dense set of isolated points, by placing the error terms of the selected iterates in the gaps of a Cantor set.

Let C be the affine copy $\frac{1}{4} + \frac{1}{4}C_0$ of the middle-thirds Cantor set $C_0 \subseteq [0, 1]$, constructed from the interval $I_\varnothing = [\frac{1}{4}, \frac{1}{2}]$. For $t \in 2^{<\mathbb{N}}$, let I_t be the interval




FIGURE 6. Bresenham's algorithm (left) and the two tie-breaking conventions $R_c^{\pm}$ (right).

of length $\ell_t = \frac{1}{4}\,3^{-|t|}$ indexed by t in the construction of C, whose left and right thirds are $I_{t^\frown 0}$ and $I_{t^\frown 1}$, and let G_t be the open middle third of I_t, the gap removed from I_t. We write z_t and x_t for the left and right endpoints of I_t. For $b \in 2^{\mathbb{N}}$, let x_b be the point of C that lies in every interval $I_{b\restriction n}$. The map $b \mapsto x_b$ is an order-preserving homeomorphism of $2^{\mathbb{N}}$, with the lexicographic order, onto C, and the gap G_u separates the points $x_{u^\frown 0^\frown 1^\infty}$ and $x_{u^\frown 1^\frown 0^\infty}$, which code adjacent sequences. In $S(2^{\mathbb{N}})$, adjacent sequences give one function, $\chi_{(u^\frown 0^\frown 1^\infty} = \chi_{[u^\frown 1^\frown 0^\infty}$, and these functions, together with $\mathbf{1} = \chi_{[0^\infty}$ and $\mathbf{0} = \chi_{(1^\infty}$, are the isolated points of $S(2^{\mathbb{N}})$.

Proposition 4.6. *Choose pairwise distinct natural numbers n_u, for $u \in 2^{<\mathbb{N}}$, and m^-, m^+ such that, modulo 1, the point $n_u\alpha$ lies in the gap G_u, the point $m^-\alpha$ in $(0, \frac{1}{4})$ and the point $m^+\alpha$ in $(\frac{1}{2}, 1)$. Then the map that sends $\chi_{[u^\frown 1^\frown 0^\infty}$ to σ^{n_u}, the function $\mathbf{1}$ to σ^{m^-} and the function $\mathbf{0}$ to σ^{m^+} extends to a homeomorphism of $S(2^{\mathbb{N}})$ onto the closure of these iterates in $E(X_\alpha, \sigma)$. The homeomorphism sends $\chi_{[a}$ to $R^-_{x_a}$ whenever a has infinitely many digits 1, and $\chi_{(a}$ to $R^+_{x_a}$ whenever a has infinitely many digits 0.*

Proof. The functions $\chi_{[u^\frown 1^\frown 0^\infty}$, $\mathbf{1}$ and $\mathbf{0}$ are isolated in $S(2^{\mathbb{N}})$ and form a dense subset, and the iterates are isolated in $E(X_\alpha, \sigma)$ by Theorem 4.5. By a lemma of Argyros, Dodos and Kanellopoulos [ADK08, Lemma 19], the map extends to a homeomorphism between the two closures as soon as, for every sequence (u_k) of pairwise distinct words, the functions $\chi_{[u_k^\frown 1^\frown 0^\infty}$ converge pointwise exactly when the iterates $\sigma^{n_{u_k}}$ do; the finitely many remaining indices play no role.

Write $a_u = u^\frown 1^\frown 0^\infty$, and let $p_u \in G_u$ be the point $n_u\alpha$ modulo 1. For every $a \in 2^{\mathbb{N}}$ distinct from a_u, the inequality $a_u < a$ holds exactly when $p_u < x_a$. Indeed, if $a_u < a$, then $p_u < x_{a_u} < x_a$; if $a < a_u$, then $a \le u^\frown 0^\frown 1^\infty$, because no sequence lies strictly between the two adjacent sequences, and therefore $x_a \le x_{u^\frown 0^\frown 1^\infty} < p_u$. Moreover, the distance from p_u to x_{a_u} is less than the length $\ell_u/3$ of G_u, which tends to 0 as $|u| \to \infty$.


As recalled in Section 1.4, the thresholds $\chi_{[a_{u_k}}$ converge pointwise exactly when the sequences a_{u_k} converge to some $a \in 2^{\mathbb{N}}$ and eventually lie on one side of a, the limit being $\chi_{[a}$ from below and $\chi_{(a}$ from above. By the proof of Theorem 4.5, and because the maps $R_c^{\pm}$ are pairwise distinct, the iterates $\sigma^{n_{u_k}}$ converge exactly when the points p_{u_k} converge to some $c \in \mathbb{T}$ and eventually lie on one side of c, the limit being R_c^- from the left and R_c^+ from the right. Since $|u_k| \to \infty$, the points

p_{u_k} converge to c exactly when c lies in C and the sequences a_{u_k} converge to the code of c; the sides agree by the equivalence above. The two kinds of convergence therefore coincide, and the limit $\chi_{[a}$ corresponds to $R^-_{x_a}$, the limit $\chi_{(a}$ to $R^+_{x_a}$. □

The copy of $S(2^{\mathbb{N}})$ in Proposition 4.6 thus consists of the selected iterates, of the maps R^-_c for the points $c \in C$ that are limits of points of C from the left, and of the maps R^+_c for the points $c \in C$ that are limits of points of C from the right. Each gap of C contributes one isolated iterate, just as each pair of adjacent sequences contributes one isolated threshold to the split Cantor set.

Remark 4.7. The same Cantor set realizes the canonical families D_3 and D_4 of Section 3.3, which differ in the rounding convention that the limit along a branch shares with the departures from the branch. By Kronecker's theorem, there are pairwise distinct natural numbers n^+_t, for $t \in 2^{<\mathbb{N}}$, such that the point $n^+_t\alpha$ lies in $(x_t, x_t + \ell_t/3)$ modulo 1, and pairwise distinct natural numbers n^-_t such that the point $n^-_t\alpha$ lies in $(z_t - \ell_t/3, z_t)$. Both intervals lie outside C, immediately to the right and to the left of I_t. Fix $b \in 2^{\mathbb{N}}$, and call a word t that is not an initial segment of b a *left departure* from b if $t^\frown 0^\infty < b$, and a *right departure* if $t^\frown 0^\infty > b$. If u is the longest common initial segment of t and b, the points $n^+_t\alpha$ and $n^-_t\alpha$ lie within distance $\frac{4}{3}\ell_u$ of x_b. Both points lie to the left of x_b when t is a left departure and to the right when t is a right departure, whereas along the initial segments of b the first point lies to the right of x_b and the second to the left. Consequently, as t converges to b, the iterates $\sigma^{n^+_t}$ converge to $R^+_{x_b}$ along initial segments and right departures, and to $R^-_{x_b}$ along left departures. Symmetrically, the iterates $\sigma^{n^-_t}$ converge to $R^-_{x_b}$ along initial segments and left departures, and to $R^+_{x_b}$ along right departures. In either family, a sequence of pairwise distinct iterates converges exactly when the indexing words converge to some b and the corresponding points of $\mathbb{T}$ eventually lie on one side of x_b; in particular, initial segments may be interleaved with the departures that share their limit. The patterns are those of D_4 and of D_3, and since the iterates are isolated in $E(X_\alpha, \sigma)$, the lemma of Argyros, Dodos and Kanellopoulos [ADK08, Lemma 19] shows that the family $\{\sigma^{n^+_t} : t \in 2^{<\mathbb{N}}\}$ is equivalent to D_4 and the family $\{\sigma^{n^-_t} : t \in 2^{<\mathbb{N}}\}$ to D_3. In terms of drawing, iterates whose accumulated error approaches the Cantor set from the right realize D_4, and iterates whose accumulated error approaches the Cantor set from the left realize D_3: along a branch, the line is drawn in the limit with the tie-breaking convention of the departures on the same side.

For the action of $\mathbb{Z}$ generated by the shift, Glasner and Megrelishvili proved that Sturmian systems are tame_2 but not hereditarily non-sensitive [GM22, Corollary 8.3].



4.4. **Two rounding procedures produce the extended duplicate of the split Cantor set.** When two rounding algorithms run in parallel, the computation descends a further level. We establish this descent here in the case of Bresenham's algorithm for lines in three-dimensional space, which lies at the level NIP^{fc} but not NIP^{hs}. Of the two alternatives that the trichotomy offers at this level, the spatial algorithm realizes the extended duplicate of the split Cantor set: its two coordinates tell apart the departures from a branch on the two sides, as in the canonical family D_7. At this level each deep computation is still separated from all the others by countably many

tests chosen for the deep computation in question (see Proposition 3.4). The same proposition shows, however, that some finitely separated set of deep computations has the cardinality of the continuum.

Drawing a line in space. Let α and β be irrational numbers in $(0,1)$ such that $1, \alpha, \beta$ are linearly independent over $\mathbb{Q}$. To draw a line in $\mathbb{Z}^3$ with direction $(1, \alpha, \beta)$, the three-dimensional Bresenham algorithm runs the planar algorithm in two coordinates at once. The CCS of the algorithm has states $L = X_\alpha \times X_\beta$, the coordinate predicates of both factors, and the semigroup generated by $\sigma \times \sigma$.

The analysis of this algorithm uses a property of the double arrow circle $\mathbb{D}$ of Section 4.3 that keeps every compact metrizable subset of $\mathbb{D} \times \mathbb{D}$ countable. Recall that a *network* of a space is a family of subsets such that every neighborhood of a point contains a member of the family that contains the point. In particular, every base is a network.

Lemma 4.8. *Every subspace of the double arrow circle $\mathbb{D}$ that has a countable network, in particular every metrizable subspace, is countable.*

Proof. A metrizable subspace of $\mathbb{D}$ is separable, since $\mathbb{D}$ is hereditarily separable, and the subspace therefore has a countable base, which is a countable network. Let $M \subseteq \mathbb{D}$ have a countable network $\{N_k : k \in \mathbb{N}\}$. For each $c \in \mathbb{T}$ with $c^+ \in M$, the open arc $[c^+, (c + \frac{1}{2})^-)$ is a neighborhood of c^+ in $\mathbb{D}$, and we may choose $k(c) \in \mathbb{N}$ with $c^+ \in N_{k(c)} \subseteq [c^+, (c + \frac{1}{2})^-)$. If $k(c) = k(c')$, then the point c'^+ lies in $[c^+, (c + \frac{1}{2})^-)$ and the point c^+ in $[c'^+, (c' + \frac{1}{2})^-)$, whence both $c' - c$ and $c - c'$ lie in $[0, \frac{1}{2})$ modulo 1, which forces $c = c'$. Thus the map $c \mapsto k(c)$ is injective, and the set M contains only countably many points c^+. The same argument, applied to the arcs $((c - \frac{1}{2})^+, c^-]$, shows that M contains only countably many points c^-. □

Corollary 4.9. *Every compact metrizable subspace of $\mathbb{D} \times \mathbb{D}$ is countable.*

Proof. Let $C \subseteq \mathbb{D} \times \mathbb{D}$ be compact and metrizable, and let $\mathcal{N}$ be a countable base of C. Since the two projections are continuous, the images of the members of $\mathcal{N}$ under each projection form a countable network of the image of C. By Lemma 4.8, both images are countable, and so is C, which is contained in their product. □

For $c' \in \mathbb{T}$, we also write $R^+_{c'}$ and $R^-_{c'}$ for the maps of X_β defined as in Section 4.3, with β in place of α.

Theorem 4.10. *The three-dimensional Bresenham CCS is NIP^{fc} but not NIP^{hs}. The space $E = E(X_\alpha \times X_\beta, \sigma \times \sigma)$ of deep computations of the CCS is the union of the set of iterates $(\sigma \times \sigma)^n$, $n \in \mathbb{N}$, and of the closed subspace*

$$\{(R^\varepsilon_c, R^{\varepsilon'}_{c'}) : c, c' \in \mathbb{T},\ \varepsilon, \varepsilon' \in \{+, -\}\},$$

which is homeomorphic to $\mathbb{D} \times \mathbb{D}$. Each of these iterates is isolated in E. The space E contains a copy of $\hat{D}(S(2^{\mathbb{N}}))$, and E contains no copy of $D(2^{\mathbb{N}})$, of $\hat{D}(2^{\mathbb{N}})$ or of $A(2^{\mathbb{N}})$.

Proof. Each predicate is a coordinate of one factor, so the CCS is NIP by Theorem 4.5. Every deep computation is the limit ξ of the iterates $(\sigma \times \sigma)^n$ along an ultrafilter $\mathcal{U}$ on $\mathbb{N}$. The coordinates of ξ are then the limits along $\mathcal{U}$ of the iterates σ^n on X_α and on X_β. If $\mathcal{U}$ is principal, the deep computation ξ is an iterate; otherwise the proof of Theorem 4.5 shows that ξ is a pair $(R^\varepsilon_c, R^{\varepsilon'}_{c'})$. By Kronecker's theorem, the points $(n\alpha, n\beta)$ with $n \geq N$ are dense in $\mathbb{T}^2$ for every N. Every point (c, c') is


therefore approached by these points from inside each of the four open quadrants at (c, c'). As in the proof of Theorem 4.5, it follows that every pair $(R_c^{\varepsilon}, R_{c'}^{\varepsilon'})$ with $\varepsilon, \varepsilon' \in \{+, -\}$ is a deep computation. By Theorem 4.5, applied to each factor, the set of these pairs is homeomorphic to $\mathbb{D} \times \mathbb{D}$, and it is therefore compact and closed in E. Each iterate $(\sigma \times \sigma)^n$ is isolated in E, since σ^n is isolated in $E(X_\alpha, \sigma)$ and is the first coordinate of no other deep computation. The space E is a closed subspace of $E(X_\alpha, \sigma) \times E(X_\beta, \sigma)$, which is first countable as a product of two first countable spaces. Hence the CCS is $\mathrm{NIP}^{\mathrm{fc}}$, and no copy of the space $A(2^{\mathbb{N}})$, which is not first countable, lies in E. Consider the pairs (R_c^+, R_{-c}^+), for $c \in \mathbb{T}$. For $0 < \delta < \frac{1}{2}$, the set $[c^+, (c+\delta)^-) \times [(-c)^+, (-c+\delta)^-)$ is a neighborhood of $(c^+, (-c)^+)$ in $\mathbb{D} \times \mathbb{D}$ that contains $(s^+, (-s)^+)$ only if $c \le s < c + \delta$ and $-c \le -s < -c + \delta$, that is, only if $s = c$. These pairs therefore form an uncountable discrete subspace, and the space of deep computations is not hereditarily separable; thus the CCS is not $\mathrm{NIP}^{\mathrm{hs}}$. A compact metrizable subspace of E meets the copy of $\mathbb{D} \times \mathbb{D}$ in a compact metrizable set, which is countable by Corollary 4.9, and the subspace itself is therefore countable. Since the non-isolated points of $D(2^{\mathbb{N}})$ form a copy of the Cantor set, neither $D(2^{\mathbb{N}})$ nor the compactum $\hat{D}(2^{\mathbb{N}})$ that contains it embeds into E. Finally, by Theorem 3.1 the space E, being the closure of a countable set, is a separable Rosenthal compactum that is first countable but not hereditarily separable. By Theorem 3.3, this compactum contains a copy of $D(2^{\mathbb{N}})$ or of $\hat{D}(S(2^{\mathbb{N}}))$, necessarily of the latter. □

The uncountable discrete set consists of the deep computations that break ties upward in both coordinates at points where the two accumulated errors are opposite. Each of the two planar algorithms is $\mathrm{NIP}^{\mathrm{hs}}$, and drawing in two directions at once leaves the class. In just the same way, the Sorgenfrey plane shows that a product of two hereditarily separable spaces need not be hereditarily separable. Glasner and Megrelishvili show similarly that a product of two tame_2 systems need not be tame_2 [GM22, Proposition 4.1].[9] A second example at this level arises from the Euclidean algorithm (see Proposition 4.15). This example is a countable family of elements of the group generated by the steps of the Euclidean algorithm, and the deep computations of the family include a copy of $D(2^{\mathbb{N}})$. Whether a single iterated map can realize the level with such a copy is Question 6.3.

Remark 4.11. A copy of $\hat{D}(S(2^{\mathbb{N}}))$ in E whose dense set of isolated points consists of iterates can be exhibited explicitly with the Cantor set C of Proposition 4.6. With the lengths ℓ_t and the right endpoints x_t of the intervals I_t fixed there, let $y_t = \frac{3}{2}\ell_t - x_t$ for $t \in 2^{<\mathbb{N}}$. By Kronecker's theorem, there are pairwise distinct $n_t \in \mathbb{N}$ such that, modulo 1, the point $n_t\alpha$ lies in $(x_t, x_t + \ell_t/3)$ and the point $n_t\beta$ in $(y_t, y_t + \ell_t/3)$. As t runs through the initial segments of b, the iterates $(\sigma \times \sigma)^{n_t}$

[9] The double arrow space is the set of the points $p^- < p^+$, for $p \in [0, 1]$, with its order topology. Glasner and Megrelishvili [GM22, Remark 4.2] assert that the square of this space contains a copy of $D(2^{\mathbb{N}})$, namely the set of the points $(p^+, (1-p)^+)$ and $(p^-, (1-p)^+)$ with p in the Cantor set. The set of these points is not closed, and hence not compact. For instance, if p is the limit of a decreasing sequence $(x_k)_{k \in \mathbb{N}}$ of points of the Cantor set, then the points $(x_k^-, (1-x_k)^+)$ of this set converge to $(p^+, (1-p)^-)$, a point outside it. Moreover, the square of the double arrow space contains no copy of $D(2^{\mathbb{N}})$ at all. Indeed, with intervals in place of arcs, the arguments of Lemma 4.8 and Corollary 4.9 apply to the double arrow space, and the non-isolated points of $D(2^{\mathbb{N}})$ form an uncountable compact metrizable set. Proposition 4.1 of Glasner and Megrelishvili does not depend on their Remark 4.2.

converge to $(R^+_{x_b}, R^+_{-x_b})$. Along words t that converge to b, are not initial segments of b and satisfy $t^\frown 0^\infty < b$, the iterates converge to $(R^-_{x_b}, R^+_{-x_b})$. Finally, along words that converge to b and satisfy $t^\frown 0^\infty > b$, the iterates converge to $(R^+_{x_b}, R^-_{-x_b})$. The three convergences just described are the convergence patterns of the family D_7 of Section 3.3. Since the iterates are isolated in E, a lemma of Argyros, Dodos and Kanellopoulos [ADK08, Lemma 19] shows that the family $\{(\sigma \times \sigma)^{n_t} : t \in 2^{<\mathbb{N}}\}$ is equivalent to D_7 and therefore has a closure homeomorphic to $\hat{D}(S(2^{\mathbb{N}}))$. These three patterns are drawn in the left panel of Figure 8; the right panel shows the copy of $\hat{D}(2^{\mathbb{N}})$ that maps built from the Euclidean algorithm produce in Section 4.5.



4.5. **Continued fractions produce exact equality tests and the Alexandroff duplicate.** The last tame level, NIP but not NIP^{fc}, is reached by the Euclidean algorithm, acting on the real projective line through the group generated by the steps of the algorithm. Exact equality tests that share one baseline witness the failure of first countability; when the baseline moves with the exceptional input, the same group realizes the Alexandroff duplicate of the preceding level, and the two choices give the canonical families D_5 and D_6 (Section 4.6). At this level some deep computation is separated from the others by no countable set of tests, that is, of pairs of an input and a feature (see Proposition 3.4). A constant deep computation has this property whenever uncountably many exact equality tests, maps each of which agrees with the constant map at every input but one, are also deep computations. Indeed, countably many tests involve only countably many inputs and therefore cannot distinguish the constant map from an exact equality test whose exceptional input is not among those inputs. The Euclidean algorithm gives rise to equality tests of precisely this kind among the deep computations of the Euclidean CCS defined below.

Let $L = \mathbb{RP}^1 = \mathbb{R} \cup \{\infty\}$, with the predicates P_1, P_3 of Section 4.2, and let $\Gamma = \mathrm{PGL}(2, \mathbb{Z})$ be the group of Möbius maps $x \mapsto (ax + b)/(cx + d)$ with integer coefficients and $ad - bc = \pm 1$. The group Γ is generated by the steps $x \mapsto x + 1$ and $x \mapsto 1/x$ of the Euclidean algorithm, and the maps that the continued fraction algorithm composes are elements of Γ. We call $(\mathbb{RP}^1, (P_1, P_3), \Gamma)$ the *Euclidean CCS*. The Euclidean CCS is a dynamical CCS whose semigroup is a group, and the deep computations of the CCS form $E(\mathbb{RP}^1, \Gamma)$.

For an irrational number $\eta = [a_0; a_1, a_2, \dots]$, let p_k/q_k be the convergents of η, with the convention $p_{-1} = 1$ and $q_{-1} = 0$, and let $\eta_{k+1} = [a_{k+1}; a_{k+2}, \dots]$ be its complete quotients. Let M_k be the Möbius map $t \mapsto (p_k t + p_{k-1})/(q_k t + q_{k-1})$, which belongs to Γ because $p_k q_{k-1} - p_{k-1} q_k = (-1)^{k+1}$. Writing $A_j(t) = a_j + 1/t$, we have $M_k = A_0 \circ A_1 \circ \cdots \circ A_k$, and hence the inverse map is $M_k^{-1} = A_k^{-1} \circ \cdots \circ A_0^{-1}$, with $A_j^{-1}(t) = 1/(t - a_j)$. Thus M_k^{-1} is the map that the first $k + 1$ steps of the continued fraction algorithm for η apply. In these terms, the classical identity $\eta = M_k(\eta_{k+1})$ states that these steps send η to the complete quotient η_{k+1}.

Lemma 4.12. *Let η be an irrational number, and let $x \in \mathbb{RP}^1$ be distinct from η. Then $M_k^{-1}(x) + q_{k-1}/q_k \xrightarrow{k\to\infty} 0$, while $M_k^{-1}(\eta) = \eta_{k+1} > 1$ for every k.*

Proof. Solving $M_k(t) = x$ gives $M_k^{-1}(\infty) = -q_{k-1}/q_k$ and, for real x,

$$M_k^{-1}(x) = \frac{q_{k-1}x - p_{k-1}}{p_k - q_k x} = -\frac{q_{k-1}}{q_k} \cdot \frac{x - p_{k-1}/q_{k-1}}{x - p_k/q_k}$$

for $k \geq 1$ (for such k we have $q_{k-1} \geq 1$), where the value is ∞ if $x = p_k/q_k$. The convergents of an irrational number are pairwise distinct, so x equals p_k/q_k for at most one k; for all larger k the formula gives a real number. Both convergents tend to $\eta \neq x$, so the last fraction tends to 1, and $0 < q_{k-1}/q_k \leq 1$ for $k \geq 1$; hence $M_k^{-1}(x) + q_{k-1}/q_k \xrightarrow{k\to\infty} 0$. The identity $M_k^{-1}(\eta) = \eta_{k+1}$ is the classical one, and $\eta_{k+1} > a_{k+1} \geq 1$ because η_{k+1} is irrational. □

Let $\varphi = (1 + \sqrt{5})/2$ be the golden ratio. Let G be the set of irrational numbers $\eta = [0; a_1, a_2, \dots]$ whose sequence of partial quotients is a concatenation $1^1 d_1 1^2 d_2 1^3 d_3 \cdots$, where 1^n denotes n consecutive ones and each $d_n \in \{1, 2\}$. The block 1^n occupies the indices $s_n, \dots, s_n + n - 1$ with $s_n = n(n+1)/2$. Distinct choices of (d_n) give distinct numbers, by the uniqueness of continued fraction expansions, so G has the cardinality of the continuum. For $\eta \in G$, let δ_η be the map that sends η to φ and every other point of $\mathbb{RP}^1$ to $-1/\varphi$, and let c be the constant map with value $-1/\varphi$.

Proposition 4.13. *(i) For every $\eta \in G$, the map δ_η is a deep computation of the Euclidean CCS.*

(ii) The map c is a deep computation, and c has no countable base of neighborhoods in $E(\mathbb{RP}^1, \Gamma)$.

Consequently $E(\mathbb{RP}^1, \Gamma)$ is not first countable.

Proof. (i) Let $\eta \in G$ be given; for each n, let $k_n = s_n + \lfloor n/2 \rfloor$ be the middle index of the block 1^n, and write $m_n = \lceil n/2 \rceil - 1$. Then $a_{k_n} = a_{k_n - 1} = \cdots = a_{k_n - m_n} = 1$ and $a_{k_n+1} = \cdots = a_{k_n+m_n} = 1$, while $m_n \to \infty$. The two quantities that occur in Lemma 4.12 are continued fractions: $q_{k-1}/q_k = [0; a_k, a_{k-1}, \dots, a_1]$ and $\eta_{k+1} = [a_{k+1}; a_{k+2}, \dots]$. Suppose that two continued fractions, finite or infinite, $[c_0; c_1, \dots]$ and $[c_0'; c_1', \dots]$ satisfy $c_j = c_j'$ for $j < m$, with $c_1 = \cdots = c_{m-1} = 1$. Then the two continued fractions share the convergents of index at most $m - 1$, and both lie between the convergents of index $m - 2$ and $m - 1$ (or equal one of these two convergents). It follows that the two continued fractions differ by at most $1/(q_{m-2}q_{m-1}) \leq 1/(F_{m-1}F_m)$, whatever their remaining partial quotients are. Here F_j is the jth Fibonacci number, with $F_1 = F_2 = 1$. The hypothesis holds with $m = m_n$ for the pair $[0; a_{k_n}, \dots, a_1]$, $[0; 1, 1, \dots]$ and for the pair $[a_{k_n+1}; a_{k_n+2}, \dots]$, $[1; 1, 1, \dots]$. Hence $q_{k_n-1}/q_{k_n} \xrightarrow{n\to\infty} [0; 1, 1, \dots] = 1/\varphi$ and $\eta_{k_n+1} \xrightarrow{n\to\infty} [1; 1, 1, \dots] = \varphi$, and Lemma 4.12 shows that the maps $M_{k_n}^{-1} \in \Gamma$ converge pointwise to δ_η, at $x = \infty$ as well.

(ii) Let (η_j) be a sequence of distinct elements of G. For each $x \in \mathbb{RP}^1$, at most one j has $\eta_j = x$, so $\delta_{\eta_j}(x) = -1/\varphi$ for all but at most one j. Hence the maps δ_{η_j} converge pointwise to c, and $c \in E(\mathbb{RP}^1, \Gamma)$ because this set is closed. Now let $(U_i)_{i\in\mathbb{N}}$ be open neighborhoods of c in $E(\mathbb{RP}^1, \Gamma)$. Each U_i contains a basic neighborhood $\{f : d(f(x), c(x)) < \varepsilon_i \text{ for } x \in F_i\}$ with $F_i \subseteq \mathbb{RP}^1$ finite, $\varepsilon_i > 0$ and d a metric on $\mathbb{RP}^1$. Since the set $F = \bigcup_i F_i$ is countable, some $\eta \in G$ lies outside F; the map δ_η then agrees with c on F and hence lies in $\bigcap_i U_i$, although $\delta_\eta \neq c$. Thus no countable family of neighborhoods of c has intersection $\{c\}$. In a Hausdorff space, the intersection of a base of neighborhoods of c is $\{c\}$, so c has no countable base of neighborhoods. □




FIGURE 7. The maps M_k^{-1} (left) and their limits, exact equality tests δ_η (right).

Corollary 4.14. *The Euclidean CCS is NIP but not NIP*^{fc}*. The set* $\{\delta_\eta : \eta \in G\} \cup \{c\}$ *of deep computations of the CCS is homeomorphic to the Alexandroff compactification* $A(2^{\mathbb{N}})$*.*

Proof. Ellis proved that the action of $\mathrm{GL}(2,\mathbb{R})$ on the projective line is tame [Ell93] (see [GM22, Example 5.7]). For every continuous $f\colon \mathbb{RP}^1 \to \mathbb{R}$, the family $\{f \circ \gamma : \gamma \in \Gamma\}$ is contained in the corresponding family for $\mathrm{GL}(2,\mathbb{R})$, so it has the NIP. Hence the action of Γ is tame, and, by Proposition 4.1, the Euclidean CCS is NIP. By Proposition 4.13, it is not NIP^{fc}. Each δ_η is isolated in $\{\delta_\eta : \eta \in G\} \cup \{c\}$, since δ_η is the only member of this set that sends η near φ. Moreover, as in the proof of Proposition 4.13, every neighborhood of c contains all but finitely many of the maps δ_η. The set $\{\delta_\eta : \eta \in G\} \cup \{c\}$ is therefore the one-point compactification of a discrete set of cardinality $2^{\aleph_0}$, and hence homeomorphic to $A(2^{\mathbb{N}})$ (see Section 1.4). □

The deep computation δ_η is an exact equality test (see Figure 7), which takes one value if the input is exactly η and another value otherwise. The map δ_η is the pointwise limit of maps that the continued fraction algorithm applies to η, when these maps are applied to all inputs. The mechanism is that of convergence groups. These are groups of homeomorphisms of a compact space in which every sequence of distinct elements has a subsequence converging to a constant map uniformly on the compact subsets of the complement of some point. By Lemma 4.12, the maps M_k^{-1} draw the images of all points other than η together, near $-q_{k-1}/q_k \in [-1,0)$, while sending η itself into $(1,\infty)$. In the language of convergence groups, every irrational number η is thus a conical limit point of Γ: some sequence in Γ sends η to one limit and every other point to another. The choice of G makes the two limits the same for uncountably many η. Glasner and Megrelishvili use the same device to prove that an infinite countable uniform convergence group acting on a compact metric space with infinite limit set is tame but not tame$_1$ [GM22, Proposition 6.1 and Theorem 6.5]. Their theorem does not cover Γ, which contains parabolic elements and is not a uniform convergence group on $\mathbb{RP}^1$. Ellis's example, the projective action of the whole group $\mathrm{GL}(2,\mathbb{R})$, is also tame but not tame$_1$ [GM22, Example 5.7]. The Euclidean CCS shows that the countable group generated by the two steps of the algorithm already reaches this level.

Iterating a single element of Γ cannot produce δ_η. For $g \in \Gamma$, the closure of $\{g^n : n \in \mathbb{Z}\}$ in $(\mathbb{RP}^1)^{\mathbb{RP}^1}$ is countable, hence metrizable, and the closure does not contain δ_η. Indeed, if g has finite order, the set of its powers is finite. If g is parabolic, with fixed point x_0, then g^n converges pointwise to the constant map x_0 as $n \to \pm\infty$. If g is hyperbolic, with attracting and repelling fixed points x_a and x_r, then g^n converges as $n \to +\infty$ to the map that fixes x_r and sends every other point to x_a. As $n \to -\infty$, the two fixed points exchange their roles, and g^n converges to the map that fixes x_a and sends every other point to x_r. If g reverses orientation and has infinite order, then the trace t of g is nonzero, whence the square g^2 has trace $t^2 + 2 > 2$ and is a hyperbolic map with the same fixed points as g. The odd powers $g^{2m+1} = g \circ g^{2m}$ have the same two limits as the even ones. In every case the closure consists of the powers of g, which are homeomorphisms, together with at most two limit maps. Each of these limit maps is constant or fixes the unique point at which the map differs from a constant. The map δ_η is neither a power of g nor such a limit map, since δ_η is not injective and $\delta_\eta(\eta) = \varphi \neq \eta$.

A copy of the Alexandroff duplicate. Among the deep computations of the Euclidean CCS, the device of Proposition 4.13 also produces a copy of the Alexandroff duplicate $D(2^{\mathbb{N}})$. With this copy comes a countable set of computations at the level $\mathrm{NIP}^{\mathrm{fc}}$ but not $\mathrm{NIP}^{\mathrm{hs}}$. Instead of sending uncountably many irrational numbers to the same pair of values, we now let the value that a deep computation takes away from the exceptional point vary continuously with the exceptional point. Let $\mathcal{C}$ be the set of irrational numbers $[0; \omega_1, \omega_2, \dots]$ with $\omega_n \in \{1, 2\}$ for every $n \geq 1$, a Cantor set homeomorphic to $\{1, 2\}^{\mathbb{N}}$ through continued fraction expansions. For $\omega = (\omega_1, \omega_2, \dots) \in \{1, 2\}^{\mathbb{N}}$ we write $[0; \omega]$ for the point $[0; \omega_1, \omega_2, \dots]$ of $\mathcal{C}$. For a finite or infinite sequence ω over the alphabet $\{1, 2\}$ and for each positive integer n not exceeding its length, let $\Lambda_n(\omega)$ be the block $(\omega_n, \omega_{n-1}, \dots, \omega_1, 1^n)$. This block consists of the first n terms of ω in reverse order, followed by n ones. For $\omega \in \{1, 2\}^{\mathbb{N}}$, let $\eta(\omega)$ be the irrational number in $(0, 1)$ whose partial quotients after the initial 0 form the concatenation $\Lambda_1(\omega)\Lambda_2(\omega)\Lambda_3(\omega)\cdots$. Since the terms $\omega_1, \dots, \omega_n$ can be recovered from $\Lambda_n(\omega)$, the map $\omega \mapsto \eta(\omega)$ is injective. For a finite word $\mathsf{w} = (\mathsf{w}_1, \dots, \mathsf{w}_k)$ of positive integers, let M_{w} be the Möbius map $x \mapsto [0; \mathsf{w}_1, \dots, \mathsf{w}_{k-1}, \mathsf{w}_k + 1/x]$, and let $M_\emptyset(x) = 1/x$ for the empty word. For an irrational number $[0; a_1, a_2, \dots]$ in $(0, 1)$, the map M_k introduced before Lemma 4.12 is then $M_{(a_1, \dots, a_k)}$. Write $\{1, 2\}^{<\mathbb{N}}$ for the set of finite words over $\{1, 2\}$. For a word $t = (t_1, \dots, t_m)$ in this set, let $t^\sharp$ be the word $\Lambda_1(t) \cdots \Lambda_{m-1}(t)\,(t_m, t_{m-1}, \dots, t_1)$ of length m^2, let the empty word be its own image, and write $\gamma_t = M_{t^\sharp}^{-1} \in \Gamma$. Finally, for $\omega \in \{1, 2\}^{\mathbb{N}}$, let τ_ω be the map that sends $\eta(\omega)$ to φ and every other point of $\mathbb{RP}^1$ to $-[0; \omega]$, and let c_ω be the constant map with value $-[0; \omega]$. For the constant sequence $\omega = (1, 1, 1, \dots)$, the map c_ω is the map c of Proposition 4.13.

Proposition 4.15. *(i) The closure of the countable set $\Delta_{\mathcal{C}} = \{\gamma_t : t \in \{1, 2\}^{<\mathbb{N}}\}$ is $\Delta_{\mathcal{C}} \cup \{\tau_\omega : \omega \in \{1, 2\}^{\mathbb{N}}\} \cup \{c_\omega : \omega \in \{1, 2\}^{\mathbb{N}}\}$, and this closure is homeomorphic to $\hat{D}(2^{\mathbb{N}})$ through a homeomorphism that maps $\{\tau_\omega : \omega \in \{1, 2\}^{\mathbb{N}}\} \cup \{c_\omega : \omega \in \{1, 2\}^{\mathbb{N}}\}$ onto $D(2^{\mathbb{N}})$. In particular, $E(\mathbb{RP}^1, \Gamma)$ contains a copy of $D(2^{\mathbb{N}})$, and the set $\Delta_{\mathcal{C}}$ of computations of the Euclidean CCS is $\mathrm{NIP}^{\mathrm{fc}}$ but not $\mathrm{NIP}^{\mathrm{hs}}$.*

(ii) Every hereditarily separable compact subset of $E(\mathbb{RP}^1, \Gamma)$ is metrizable. In particular, $E(\mathbb{RP}^1, \Gamma)$ contains no copy of $S(2^{\mathbb{N}})$ and none of $\hat{D}(S(2^{\mathbb{N}}))$.

Proof. (i) We use two facts from the proofs of Lemma 4.12 and Proposition 4.13. First, consider two continued fractions, finite or infinite, with positive integer partial quotients that agree in their terms of index less than m. Both lie between two consecutive convergents of index $m-2$ and $m-1$, and hence differ by at most $1/(F_{m-1}F_m)$. Second, the formula for M_k^{-1} in the proof of Lemma 4.12 uses only the terms $a_0,\dots,a_k$: if $k\geq 1$ and p_i/q_i are the convergents of $[a_0;a_1,\dots,a_k]$, the map $M(x)=[a_0;a_1,\dots,a_{k-1},a_k+1/x]$ satisfies

$$M^{-1}(x) = -\frac{q_{k-1}}{q_k}\cdot\frac{x-p_{k-1}/q_{k-1}}{x-p_k/q_k} \quad (x\in\mathbb{R}), \qquad M^{-1}(\infty) = -\frac{q_{k-1}}{q_k}, \tag{4.16}$$

where $q_{k-1}/q_k = [0;a_k,a_{k-1},\dots,a_1]$.

Limits along initial segments. Let $\omega\in\{1,2\}^{\mathbb{N}}$ be given, and write $\eta=\eta(\omega)=[0;a_1,a_2,\dots]$. For $m\geq 1$, the word $(\omega{\restriction}m)^\sharp$ consists of the first m^2 partial quotients of η, whence $\gamma_{\omega\restriction m}=M_{m^2}^{-1}$. The terms $a_{m^2},a_{m^2-1},\dots,a_{m^2-m+1}$ are $\omega_1,\omega_2,\dots,\omega_m$, and $a_{m^2+1}=\cdots=a_{m^2+m}=1$. By the first fact, as $m\to\infty$, the quotient $q_{m^2-1}/q_{m^2}=[0;a_{m^2},\dots,a_1]$ tends to $[0;\omega]$ and the complete quotient η_{m^2+1} tends to $[1;1,1,\dots]=\varphi$. Lemma 4.12 therefore shows that the maps $\gamma_{\omega\restriction m}$ converge pointwise to τ_ω.

Limits along other words. Let $(t_j)_{j\in\mathbb{N}}$ be a sequence of words that converges to ω, none of which is an initial segment of ω. For each j, let n_j be the length of the longest common initial segment of t_j and ω, and observe that n_j tends to infinity because the words t_j converge to ω. To show that the maps γ_{t_j} converge pointwise to c_ω, we write $t_j^\sharp=(\mathsf{w}_1,\dots,\mathsf{w}_k)$, where $k=k_j=|t_j|^2$, and we let $\ell=\ell_j=n_j(n_j+1)+1$. The blocks $\Lambda_1(t_j),\dots,\Lambda_{n_j}(t_j)$ coincide with $\Lambda_1(\omega),\dots,\Lambda_{n_j}(\omega)$, and the next term of $t_j^\sharp$ is the term of index n_j+1 of t_j. Hence $\mathsf{w}_i=a_i$ for $i<\ell$, while w_ℓ and a_ℓ are distinct elements of $\{1,2\}$. Moreover, since $t_j^\sharp$ ends with the terms of t_j in reverse order, the terms $\mathsf{w}_k,\mathsf{w}_{k-1},\dots$ begin with $\omega_1,\dots,\omega_{n_j}$, and $k-\ell\geq(n_j+1)^2-\ell=n_j$. Let $x\in\mathbb{RP}^1$ be distinct from η. The convergents of $[0;\mathsf{w}_1,\dots,\mathsf{w}_k]$ of index at least $\ell-1$ agree with η in their terms of index less than ℓ. By the first fact, the convergents of index $k-1$ and k therefore tend to η as $j\to\infty$, while $q_{k-1}/q_k=[0;\mathsf{w}_k,\dots,\mathsf{w}_1]$ tends to $[0;\omega]$. Formula (4.16) then shows that $\gamma_{t_j}(x)$ converges to $-[0;\omega]$ for every such x. At the point η we write γ_{t_j} as the composite of the map $x\mapsto 1/x$ followed by the maps $x\mapsto 1/(x-\mathsf{w}_i)$, for $i=1,\dots,k$ in this order. We then apply these $k+1$ factors to η one at a time. The first ℓ factors send η to its complete quotient $\eta_\ell=[a_\ell;a_{\ell+1},\dots]$, which lies in $(a_\ell,a_\ell+1)$. The next factor sends η_ℓ to $z_j=1/(\eta_\ell-\mathsf{w}_\ell)$, which lies in $(\frac12,1)$ if $(a_\ell,\mathsf{w}_\ell)=(2,1)$ and in $(-\infty,-1)$ if $(a_\ell,\mathsf{w}_\ell)=(1,2)$; in either case $z_j<1$. The remaining factors form the inverse of the map $M'(x)=[\mathsf{w}_{\ell+1};\mathsf{w}_{\ell+2},\dots,\mathsf{w}_{k-1},\mathsf{w}_k+1/x]$, and therefore $\gamma_{t_j}(\eta)=M'^{-1}(z_j)$. Let p_i'/q_i' be the convergents of $[\mathsf{w}_{\ell+1};\mathsf{w}_{\ell+2},\dots,\mathsf{w}_k]$, the last of which has index $i=k-\ell-1\geq n_j-1$. Every convergent of index at least 1 has the form $\mathsf{w}_{\ell+1}+1/y$ with $1\leq y\leq 3$, since all its partial quotients lie in $\{1,2\}$. Each of these convergents is therefore at least $\frac43$, and its distance to z_j exceeds $\frac13$. The convergents of index $i-1$ and i differ by $1/(q_{i-1}'q_i')\leq 1/(F_iF_{i+1})$, and $q_{i-1}'/q_i'=[0;\mathsf{w}_k,\dots,\mathsf{w}_{\ell+2}]$ tends to $[0;\omega]$, since the first n_j-1 partial quotients of this fraction are $\omega_1,\dots,\omega_{n_j-1}$. By (4.16), applied to M', we conclude that $\gamma_{t_j}(\eta)$ converges to $-[0;\omega]$, and hence that the maps γ_{t_j} converge pointwise to c_ω.

The closure. By Corollary 4.14 and Theorem 3.1, the space $E(\mathbb{RP}^1,\Gamma)$ is a Rosenthal compactum. Consequently, every point of the closure of $\Delta_{\mathfrak{C}}$ that does not

belong to $\Delta_{\mathcal{C}}$ is the limit of a sequence (γ_{t_j}) with pairwise distinct words t_j [BFT78, Theorem 3F]. A compactness argument yields a subsequence of (t_j) that converges to some $\omega \in \{1,2\}^{\mathbb{N}}$. A further subsequence consists either of initial segments of ω or of words that are not initial segments of ω. By the two preceding paragraphs, the corresponding maps converge to τ_ω in the first case and to c_ω in the second. Both kinds of limit occur for every ω, and the closure of $\Delta_{\mathcal{C}}$ is therefore the set described in (i).

The homeomorphism. Identify $\{1,2\}^{\mathbb{N}}$ with $2^{\mathbb{N}}$ and $\{1,2\}^{<\mathbb{N}}$ with $2^{<\mathbb{N}}$ by replacing the letters 1 and 2 with 0 and 1. Under this identification, define a map Θ on $\hat{D}(2^{\mathbb{N}})$ by $\Theta(v_t, t^\frown 0^\infty) = \gamma_t$, $\Theta(\delta_\omega, \omega) = \tau_\omega$ and $\Theta(\mathbf{0}, \omega) = c_\omega$. The map Θ is surjective by the preceding paragraph. To see that Θ is injective, note first that the constant maps c_ω are pairwise distinct, because distinct sequences ω give distinct numbers $[0;\omega]$. For the same reason, the maps τ_ω are pairwise distinct, and none of them is constant. Neither kind of map is injective, whereas the maps γ_t are homeomorphisms. Finally, the maps γ_t are pairwise distinct, because $t^\sharp$ determines t and the Möbius map M_{w} determines the word w. Indeed, a rational number in $[0,1]$ has at most two expansions $[0;\mathsf{w}_1,\dots,\mathsf{w}_k]$, whose lengths differ by one. The number $M_{\mathsf{w}}(\infty) = [0;\mathsf{w}_1,\dots,\mathsf{w}_k]$ and the determinant $(-1)^{k+1}$ of a matrix that represents M_{w} therefore determine w. Since $\hat{D}(2^{\mathbb{N}})$ is first countable, the continuity of Θ may be tested on sequences. A sequence that converges to a point (δ_ω, ω) consists, apart from finitely many terms and from terms equal to that point, of points $(v_t, t^\frown 0^\infty)$ with t an initial segment of ω, whose lengths tend to infinity. By the paragraph on limits along initial segments, the image of this sequence under Θ converges to τ_ω. A sequence that converges to a point $(\mathbf{0}, \omega)$ consists, apart from finitely many terms and from terms equal to that point, of points of at most three kinds. Points of the first kind have the form $(\mathbf{0}, \omega')$, and points of the second kind the form $(\delta_{\omega'}, \omega')$, with $\omega' \to \omega$ and $\omega' \neq \omega$. Points of the third kind have the form $(v_t, t^\frown 0^\infty)$, with t converging to ω and eventually not an initial segment of ω. The images of the points of the first kind converge to c_ω because $[0;\omega']$ depends continuously on ω'. The same holds for the images of the points of the third kind, by the paragraph on limits along other words. For the second kind, observe that the equality $\tau_{\omega'}(x) = -[0;\omega']$ fails at a given point $x \in \mathbb{RP}^1$ only if $\eta(\omega') = x$, which happens for at most one ω'. The images of the points of the second kind therefore converge to c_ω as well. Hence Θ is a continuous bijection of a compact space onto a Hausdorff space, and therefore a homeomorphism; the map Θ sends $D(2^{\mathbb{N}})$ onto $\{\tau_\omega : \omega \in \{1,2\}^{\mathbb{N}}\} \cup \{c_\omega : \omega \in \{1,2\}^{\mathbb{N}}\}$. The set $\Delta_{\mathcal{C}}$ is countable, and $\Delta_{\mathcal{C}}$ has the NIP because Γ does (see Corollary 4.14). The closure of $\Delta_{\mathcal{C}}$, a copy of $\hat{D}(2^{\mathbb{N}})$, is first countable but not hereditarily separable (see Section 3.2), and $\Delta_{\mathcal{C}}$ is therefore $\mathrm{NIP}^{\mathrm{fc}}$ but not $\mathrm{NIP}^{\mathrm{hs}}$.

(ii) We first describe the deep computations that are not computations. For $y, z, w \in \mathbb{RP}^1$, let $\zeta_{y,z,w}$ be the map that sends z to w and every other point to y. Let $\xi \in E(\mathbb{RP}^1, \Gamma) \setminus \Gamma$ be given; since $E(\mathbb{RP}^1, \Gamma)$ is a Rosenthal compactum, the map ξ is the limit of a sequence $(g_n)_{n\in\mathbb{N}}$ of pairwise distinct elements of Γ. Represent each g_n by an integer matrix $\mathbf{G}_n$ with determinant ± 1. Since only finitely many integer matrices have norm below a given bound, the norms $\|\mathbf{G}_n\|$ tend to infinity. Passing to a subsequence, we may assume that the normalized matrices $\mathbf{G}_n/\|\mathbf{G}_n\|$ converge to a matrix $\mathbf{G}$ of norm 1. Since $\det(\mathbf{G}_n/\|\mathbf{G}_n\|) = \pm\|\mathbf{G}_n\|^{-2} \to 0$, the matrix $\mathbf{G}$ has rank one, and $\mathbf{G}\mathbf{e} = \langle \mathbf{e}, \mathbf{e}' \rangle \, \mathbf{e}''$ for all $\mathbf{e} \in \mathbb{R}^2$, for some nonzero vectors

$\mathbf{e}', \mathbf{e}'' \in \mathbb{R}^2$. If $x \in \mathbb{RP}^1$ is the line spanned by a vector $\mathbf{e}$ with $\langle \mathbf{e}, \mathbf{e}' \rangle \neq 0$, then $\mathbf{G}_n\mathbf{e}/\|\mathbf{G}_n\| \xrightarrow{n\to\infty} \langle \mathbf{e}, \mathbf{e}' \rangle \, \mathbf{e}'' \neq 0$, and $g_n(x)$ converges to the line y spanned by $\mathbf{e}''$. Thus, if z is the line orthogonal to $\mathbf{e}'$ and $w = \xi(z)$, then $\xi = \zeta_{y,z,w}$.

Now let $K \subseteq E(\mathbb{RP}^1, \Gamma)$ be compact and hereditarily separable, and let Z_K be the set of points $z \in \mathbb{RP}^1$ for which some map $\zeta_{y,z,w}$ with $y \neq w$ belongs to K; such a map determines z. Choose, for each $z \in Z_K$, one such map $\zeta_z = \zeta_{y_z,z,w_z} \in K$. We claim that the set $\{\zeta_z : z \in Z_K\}$ is discrete. Indeed, fix a metric d on $\mathbb{RP}^1$ and a point $z \in Z_K$, let $\varepsilon = d(y_z, w_z)/2$, and choose distinct points $x, x' \in \mathbb{RP}^1 \setminus \{z\}$. Consider the neighborhood of ζ_z formed by the maps g with $d(g(z), w_z) < \varepsilon$, $d(g(x), y_z) < \varepsilon$ and $d(g(x'), y_z) < \varepsilon$. Given $z' \in Z_K \setminus \{z\}$, one of the points x and x' differs from z', and we may suppose that x does. Then $\zeta_{z'}(z) = \zeta_{z'}(x) = y_{z'}$, since neither z nor x equals z'. Were $\zeta_{z'}$ in the neighborhood of ζ_z just described, we would have $d(y_z, w_z) \leq d(y_z, y_{z'}) + d(y_{z'}, w_z) < 2\varepsilon$, which is impossible. Since K is hereditarily separable, the discrete set $\{\zeta_z : z \in Z_K\}$ is countable, and so is Z_K. Evaluation at the points of the countable set $Z_K \cup \mathbb{Q}$ separates the members of K. To see this, note first that two members of Γ that agree on the dense set $\mathbb{Q}$ agree everywhere. A member of Γ, moreover, is injective, whereas a map $\zeta_{y,z,w}$ takes the value y at all points of $\mathbb{Q}$ but at most one. Two maps $\zeta_{y,z,w}$ and $\zeta_{y',z',w'}$ with $y \neq y'$ differ at a point of $\mathbb{Q} \setminus \{z, z'\}$. Lastly, let $\zeta_{y,z,w}$ and $\zeta_{y,z',w'}$ be two distinct maps, labelled so that the first satisfies $w \neq y$. These two maps differ at the point $z \in Z_K$, at which the second takes the value y if $z' \neq z$ and the value $w' \neq w$ if $z' = z$. Hence restriction to $Z_K \cup \mathbb{Q}$ is a continuous injection of the compact space K into the metrizable space $(\mathbb{RP}^1)^{Z_K \cup \mathbb{Q}}$, and K is metrizable.

Finally, the split Cantor set $S(2^{\mathbb{N}})$ is hereditarily separable and not metrizable, and, as we now show, so is the closed subspace of $\hat{D}(S(2^{\mathbb{N}}))$ formed by the points of $\hat{D}(S(2^{\mathbb{N}}))$ with first coordinate $\mathbf{0}$. The subspace just described is hereditarily separable, since the projection onto the second coordinate maps this subspace homeomorphically onto a subspace of $S(2^{\mathbb{N}})$. To see that this subspace is not metrizable, let $\varpi \in 2^{\mathbb{N}}$ have infinitely many zeros, and let a be the branch of the subtree $\{s_t : t \in 2^{<\mathbb{N}}\}$ of Section 1.4 that extends the words $s_{\varpi\restriction n}$. Let $u_1, u_2, \ldots$ be the words $s_{(\varpi\restriction n)^\frown 1}$ with $\varpi(n) = 0$, listed by increasing n. These words are pairwise incomparable and converge to a, and the sequences $u_k^\frown 0^\infty$ decrease, since the canonical isomorphism preserves the lexicographic order. By the description of $\hat{D}(S(2^{\mathbb{N}}))$ in Section 1.4, the point $(\mathbf{0}, \chi_{(a})$ belongs to the subspace just described, and distinct sequences ϖ give distinct branches. In $S(2^{\mathbb{N}})$, each point $\chi_{(a}$ has the neighborhoods $\{\chi_{(a}\} \cup \{\chi_{[x}, \chi_{(x} : a < x < b\}$, for $b > a$, and the argument of Lemma 4.8 shows that a metrizable subspace of $S(2^{\mathbb{N}})$ contains only countably many of the points $\chi_{(a}$. By the preceding paragraph, therefore, neither $S(2^{\mathbb{N}})$ nor $\hat{D}(S(2^{\mathbb{N}}))$ embeds into $E(\mathbb{RP}^1, \Gamma)$. □

Together with Corollary 4.14, Proposition 4.15 shows that the space of deep computations of the Euclidean CCS contains copies of $A(2^{\mathbb{N}})$ and of $D(2^{\mathbb{N}})$, but no copy of $S(2^{\mathbb{N}})$ or of $\hat{D}(S(2^{\mathbb{N}}))$. At the level $\mathrm{NIP}^{\mathrm{fc}}$ but not $\mathrm{NIP}^{\mathrm{hs}}$, the set $\Delta_{\mathcal{C}}$ thus realizes the alternative of Todorčević's trichotomy opposite to the one realized by the three-dimensional Bresenham algorithm (see Theorem 4.10). The two alternatives are drawn side by side in Figure 8.

The Euclidean constructions differ in a single feature, the baseline. Every map δ_η of Proposition 4.13 takes the value φ at its exceptional input η and the common baseline value $-1/\varphi$ at every other input. A countable set of tests involves countably many inputs, and an exceptional input $\eta \in G$ outside them gives a map δ_η that agrees with the constant map c on all these tests; no countable set of tests determines c, and first countability fails. In the family $\Delta_{\mathcal{C}}$ the baseline $-[0;\omega]$ of τ_ω and of c_ω moves with ω, and distinct sequences ω give distinct baselines. Three fixed distinct inputs x_1, x_2, x_3 therefore determine ω for every member of the closure of $\Delta_{\mathcal{C}}$ that is not a computation: each such member differs from a constant map at one input at most, so at least one of the three inputs is ordinary for any two of them, and there their baselines must agree. One further input, $\eta(\omega)$, tells τ_ω from c_ω, and no computation γ_t, which is injective, agrees with such a member at all three inputs. With the two predicates P_1 and P_3 at each of these four inputs, eight tests separate each member of the closure of $\Delta_{\mathcal{C}}$ that is not a computation from all the others, and the closure is first countable. Exact determination by finitely many tests is not isolation: the rational tolerances on these tests give countably many neighborhoods whose intersection is the chosen member, and compactness turns them into a countable base of neighborhoods.

Both constructions at the level NIP^{fc} but not NIP^{hs} follow the binary tree, and the difference between them appears when the words depart from a branch. Along the initial segments of a branch, the selected iterates of the spatial Bresenham algorithm converge to a map $(R^+_{x_b}, R^+_{-x_b})$ (see Remark 4.11), and the maps γ_t to a map τ_ω. Departing from the branch below and above, the Bresenham iterates converge to the two distinct maps $(R^-_{x_b}, R^+_{-x_b})$ and $(R^+_{x_b}, R^-_{-x_b})$, whereas the maps γ_t converge from either side to the single constant map c_ω. Two companions for each limit along a branch give the extended duplicate of the split Cantor set, and one companion gives the Alexandroff duplicate; these are the patterns of the families D_7 and D_6 of Section 3.3.




FIGURE 8. The two alternatives of case (ii) of the trichotomy: spatial Bresenham (left) and the maps γ_t (right).

Free reduction. A second example at the level NIP but not NIP^{fc} is the free group F_2 on two generators a, b, acting on its boundary ∂F_2. This boundary, a Cantor set, is the space of infinite reduced words in a, a^{-1}, b, b^{-1}. A generator acts

on an infinite word by prepending a letter and cancelling the letter against the first letter of the word if the two letters are inverse, as a stack does. With the predicates $w \mapsto [\![w_n = x]\!]$, for $n \in \mathbb{N}$ and each letter x, the boundary ∂F_2 and the action of F_2 form a dynamical CCS. Glasner and Megrelishvili proved that this system is tame but not tame$_1$ [GM22, Corollary 6.6 and Example 6.7]. By Proposition 4.1, this CCS is therefore NIP but not NIP^{fc}.

An open question. Both examples at this level are actions of groups with two generators. Whether a single map can reach this level, that is, whether some cascade CCS is NIP but not NIP^{fc}, is Question 6.2. For actions of $\mathbb{Z}$, and more generally of abelian groups, the existence of a tame metrizable system whose enveloping semigroup is not first countable is Question 6.8 of Glasner and Megrelishvili [GM22]. To our knowledge, this question remains open.

4.6. **The same algorithms realize the seven canonical families.** Todorčević's trichotomy, realized in the preceding subsections, describes the space of deep computations as a whole: the trichotomy decides whether the space is metrizable and, if not, which canonical compactum the space contains. The heptachotomy of Argyros, Dodos and Kanellopoulos (Theorem 3.7) describes instead the routes by which a family of computations indexed by the binary tree approaches its deep computations: along a branch, or departing from the branch below or above. In this subsection we show that the algorithms of this section realize all seven canonical families, and we compare the realizations in pairs. Each comparison isolates one feature of an algorithm that decides the route, and several of these features are invisible to the trichotomy, which places the two members of a pair at the same level.

The only family not yet realized is D_5, whose closure is the extended Alexandroff compactification $\hat{A}(2^{\mathbb{N}})$. The Euclidean algorithm realizes this family with the device of Proposition 4.13, in which all exact equality tests share one baseline, arranged along the binary tree as in Proposition 4.15. For $\omega \in \{1,2\}^{\mathbb{N}}$, let $\theta(\omega)$ be the irrational number in $(0,1)$ whose partial quotients after the initial 0 form the concatenation of the blocks $(\omega_n, 1^{2n})$, for $n = 1, 2, \dots$, and let ε_ω be the map that sends $\theta(\omega)$ to φ and every other point of $\mathbb{RP}^1$ to $-1/\varphi$. For a word $t = (t_1, \dots, t_m)$ in $\{1,2\}^{<\mathbb{N}}$, let $t^\natural$ be the word $(t_1, 1^2, t_2, 1^4, \dots, t_{m-1}, 1^{2m-2}, t_m, 1^m)$, which ends with half of the block that follows t_m in the expansion of $\theta(\omega)$ for every ω extending t; the empty word is again its own image, and we write $\beta_t = M_{t^\natural}^{-1} \in \Gamma$. As before, c denotes the constant map with value $-1/\varphi$.

Proposition 4.17. *The closure of* $\{\beta_t : t \in \{1,2\}^{<\mathbb{N}}\}$ *in* $E(\mathbb{RP}^1, \Gamma)$ *is* $\{\beta_t : t \in \{1,2\}^{<\mathbb{N}}\} \cup \{\varepsilon_\omega : \omega \in \{1,2\}^{\mathbb{N}}\} \cup \{c\}$. *With the letters* 1 *and* 2 *replaced by* 0 *and* 1, *the family* $\{\beta_t : t \in \{1,2\}^{<\mathbb{N}}\}$ *is equivalent to* D_5, *and the closure of the family is homeomorphic to* $\hat{A}(2^{\mathbb{N}})$.

Proof. The proof follows the proof of Proposition 4.15(i), and we use the two facts recalled at its beginning. Fix $\omega \in \{1,2\}^{\mathbb{N}}$, and write $\theta(\omega) = [0; a_1, a_2, \dots]$.

Limits along initial segments. For $m \geq 1$, the word $(\omega{\restriction}m)^\natural$ consists of the first $k = m^2 + m$ partial quotients of $\theta(\omega)$, whence $\beta_{\omega\restriction m} = M_k^{-1}$. The fraction $q_{k-1}/q_k = [0; a_k, \dots, a_1]$ begins with m partial quotients equal to 1, and the complete quotient $\theta(\omega)_{k+1}$ begins with the remaining m ones of the block 1^{2m}. By the first fact, q_{k-1}/q_k tends to $1/\varphi$ and $\theta(\omega)_{k+1}$ to φ as $m \to \infty$, and Lemma 4.12 shows that the maps $\beta_{\omega\restriction m}$ converge pointwise to ε_ω.

Limits along other words. Let (t_j) be a sequence of words that converges to ω, none of which is an initial segment of ω, and let n_j be the length of the longest common initial segment of t_j and ω, which tends to infinity. Write $t_j^\natural = (\mathsf{w}_1, \dots, \mathsf{w}_k)$ and $\ell = (n_j+1)^2$. Then $\mathsf{w}_i = a_i$ for $i < \ell$, whereas w_ℓ and a_ℓ are the distinct terms of index n_j+1 of t_j and of ω. The word $t_j^\natural$ ends with at least n_j+1 terms equal to 1, and these terms follow the index ℓ. For $x \in \mathbb{RP}^1$ distinct from $\theta(\omega)$, the first fact shows that the convergents of index $k-1$ and k of $[0; \mathsf{w}_1, \dots, \mathsf{w}_k]$ tend to $\theta(\omega)$ and that $q_{k-1}/q_k = [0; \mathsf{w}_k, \dots, \mathsf{w}_1]$ tends to $1/\varphi$; by formula (4.16), $\beta_{t_j}(x)$ tends to $-1/\varphi$. At the point $\theta(\omega)$, the argument of Proposition 4.15 applies word for word: the first $\ell+1$ factors of β_{t_j} send $\theta(\omega)$ to a point $z_j < 1$, and the remaining factors form the inverse of a Möbius map M' whose partial quotients lie in $\{1, 2\}$, whose convergents of positive index exceed $z_j + \frac{1}{3}$, and whose reversed fraction q'_{i-1}/q'_i tends to $1/\varphi$. Hence $\beta_{t_j}(\theta(\omega))$ also tends to $-1/\varphi$, and the maps β_{t_j} converge pointwise to c.

The closure and the equivalence. The maps β_t are pairwise distinct homeomorphisms, since $t^\natural$ determines t and M_{w} determines w, whereas the maps ε_ω and c are not injective; the maps ε_ω are pairwise distinct because θ is injective. As in the proof of Proposition 4.15, every point of the closure outside the family is the limit of a sequence (β_{t_j}) with pairwise distinct words, a subsequence of the words converges to some ω, and a further subsequence consists of initial segments of ω or of words that are not; the two preceding paragraphs give the limit ε_ω or c. Hence the closure is the set described, and each map β_t is isolated in the closure. The same argument shows that, for a sequence (t_j) of pairwise distinct words, the maps β_{t_j} converge exactly when either all but finitely many of the words t_j are initial segments of one sequence ω, with limit ε_ω, or every ω has only finitely many of the words t_j as initial segments, with limit c. The prefix tests v_{t_j}, which are isolated in $\hat{A}(2^{\mathbb{N}})$, converge under exactly the same condition, with limits δ_ω and $\mathbf{0}$, and the lemma of Argyros, Dodos and Kanellopoulos [ADK08, Lemma 19] shows that the two families are equivalent. □

With Proposition 4.17, every canonical family is realized. A canonical family belongs to a selected family of computations indexed by the binary tree, and not to an algorithm. Bresenham's algorithm in the plane realizes both D_3 and D_4, and the Euclidean group realizes both D_5 and D_6; the space of all deep computations of the group is not first countable, whereas the closure of the selected family $\Delta_{\mathfrak{C}}$ is. Changing the selected computations, or the information retained in the output, can therefore change the canonical family, and the routes of Table 2 are the convergence patterns of the selected families.

Theorem 4.18. *Each of the seven canonical families $D_1, \dots, D_7$ of Argyros, Dodos and Kanellopoulos is equivalent to a family of computations of one of the algorithms of this section, as follows.*

- *(i) The family D_1: iterates of Newton's method for the square roots of a positive number, on the real projective line.*
- *(ii) The family D_2: iterates of a rotation of the circle by an irrational angle.*
- *(iii) The families D_3 and D_4: iterates of Bresenham's algorithm in the plane.*
- *(iv) The family D_5: maps of the Euclidean CCS with a fixed baseline, the family $\{\beta_t\}$ of Proposition 4.17.*

(v) The family D_6: maps of the Euclidean CCS with a moving baseline, the family $\Delta_{\mathfrak{C}}$ of Proposition 4.15.
(vi) The family D_7: iterates of Bresenham's algorithm in space.

Proof. (i) Let N_p be the Newton map of the square-root CCS, and let $\mathfrak{f}$ be the pointwise limit of its iterates. The iterates N_p^n are pairwise distinct rational maps, of degree 2^n, and $\mathfrak{f}$ is discontinuous, so the closure of every infinite set of iterates consists of these iterates, each isolated, together with $\mathfrak{f}$. For any injective assignment $t \mapsto n_t$, every sequence of pairwise distinct maps $N_p^{n_t}$ converges to $\mathfrak{f}$, just as every sequence of pairwise distinct members of D_1 converges to $\mathbf{0}$, and the lemma of Argyros, Dodos and Kanellopoulos [ADK08, Lemma 19] shows that the families $\{N_p^{n_t}\}$ and D_1 are equivalent.

(ii) Let rot_c denote the rotation $x \mapsto x+c$ of $\mathbb{T}$, and let n_t^+ be the natural numbers of Remark 4.7, with $n_t^+\alpha$ in $(x_t, x_t+\ell_t/3)$ modulo 1. The rotations $\mathrm{rot}_{n_t^+\alpha}$ converge pointwise exactly when the points $n_t^+\alpha$ converge in $\mathbb{T}$. A sequence of these points, indexed by pairwise distinct words, converges exactly when the words converge to some $b \in 2^{\mathbb{N}}$, the limit being x_b: the point $n_t^+\alpha$ lies within $\ell_t/3$ of the point x_t of C, and the points of C in the interval I_t, of length ℓ_t, are the points coded by the extensions of t. Since the points $n_t^+\alpha$ lie outside the closed set C, which contains all their accumulation points, each rotation $\mathrm{rot}_{n_t^+\alpha}$ is isolated in the closure of the family. The members of D_2 converge exactly when the words t converge, and the lemma of Argyros, Dodos and Kanellopoulos gives the equivalence.

(iii) This is Remark 4.7. (iv) This is Proposition 4.17. (v) The homeomorphism Θ of Proposition 4.15 sends $(v_t, t^\frown 0^\infty)$ to γ_t, and is therefore an equivalence of D_6 with $\Delta_{\mathfrak{C}}$. (vi) This is Remark 4.11. □

Table 2 records the limits of the seven realizations along the three routes. For a sequence b, the departures below and above are the words t, converging to b and not initial segments of b, with $t^\frown 0^\infty < b$ and $t^\frown 0^\infty > b$, as in Remark 4.7; for the Euclidean families, b is a sequence over $\{1, 2\}$, ordered after the letters 1 and 2 are replaced by 0 and 1.

As recalled after Theorem 3.7, the trichotomy places the closures of the seven families at four levels. Four comparisons show what the finer classification adds.

Forgetting the route and recovering the point: D_1 and D_2. Both families lie at the metrizable level. A convergent method forgets everything about the route: every sequence of distinct iterates of Newton's method has the same limit, whatever words index the sequence. A rotation recovers the point approached: the limit rot_{x_b} determines b, although no route leaves a trace in the limit beyond b.

Rounding records the side: D_2, D_3 and D_4. The rotation of (ii) and the family $\{\sigma^{n_t^+}\}$ of Remark 4.7 use the same exponents n_t^+, and the natural projection of the Sturmian shift onto the circle carries the shift to the rotation by α and both maps $R_c^{\pm}$ to rot_c. The decision bits of Bresenham's algorithm, which the projection discards, are what records the side from which the accumulated error approaches the limit, and they alone turn D_2 into D_4. Whether the limit along the branch shares the convention of the departures below or above depends on the side from which the selected error terms approach the Cantor set, and distinguishes D_3 from D_4.

A fixed and a moving baseline: D_5 and D_6. Both families consist of maps generated by the steps of the Euclidean algorithm, and in both the limit along a branch is an exact equality test, while the departures on either side lead to a

Family	Algorithm	Along the branch	Departing below	Departing above
D_1	Newton, square roots	∮	∮	∮
D_2	rotation	rot_{x_b}	rot_{x_b}	rot_{x_b}
D_3	Bresenham, plane	$R^-_{x_b}$	$R^-_{x_b}$	$R^+_{x_b}$
D_4	Bresenham, plane	$R^+_{x_b}$	$R^-_{x_b}$	$R^+_{x_b}$
D_5	Euclid, fixed baseline	ε_b	c	c
D_6	Euclid, moving baseline	τ_b	c_b	c_b
D_7	Bresenham, space	$(R^+_{x_b}, R^+_{-x_b})$	$(R^-_{x_b}, R^+_{-x_b})$	$(R^+_{x_b}, R^-_{-x_b})$

TABLE 2. Limits of the realizations along the three routes.

constant map. In D_5 the constant is the same for every branch, and no countable set of tests determines it: first countability fails. In D_6 the constant c_b, the baseline of the equality test τ_b, moves with the branch, and eight tests determine every limit. The trichotomy places the two families at different levels, although the algorithm, the group and the routes coincide; only the baseline differs.

One or two companions: D_6 and D_7. Both families have closures that are first countable and not hereditarily separable, and the trichotomy distinguishes them only by the canonical compactum that each closure contains. The routes make the difference visible: departures from a branch on the two sides lead to one common limit for the Euclidean maps and to two distinct limits for the selected iterates of Bresenham's algorithm in space, whose two accumulated errors tend to opposite limits (see Figure 8).

4.7. **Chaotic iteration leads beyond the classification.** Chaotic iteration leads beyond the hierarchy, to sets of computations with the independence property. By Theorem 3.1, some deep computation of such a set then fails to be of the first Baire class on some type shard, and neither classification applies: Todorčević's trichotomy is a theorem about Rosenthal compacta, and the heptachotomy a theorem about families relatively compact in the first Baire class. In a set of computations with the independence property, the computations realize, for some feature and on suitable inputs, every finite pattern of high and low values. We close the section with examples of chaotic iteration, among them Newton's method on the Riemann sphere, the doubling map of the circle and the logistic map beyond the Feigenbaum parameter.

We begin by formalizing the Newton–Raphson root-finding method as a CCS [Gal00], [BFB16, Section 2.3]. Let $\hat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ be the extended complex plane, that is, the Alexandroff (one-point) compactification of $\mathbb{C}$. We identify $\hat{\mathbb{C}}$ with the unit sphere $S^2 \subseteq \mathbb{R}^3$, the Riemann sphere, through the inverse stereographic projection. This projection is the homeomorphism $\hat{\mathbb{C}} \to S^2$ given by $z \mapsto (P_1(z), P_2(z), P_3(z))$, where $(P_1(\infty), P_2(\infty), P_3(\infty)) = (0, 0, 1)$ and, for

$z \neq \infty$,

$$P_1(z) = \frac{2\,\Re z}{|z|^2+1}, \qquad P_2(z) = \frac{2\,\Im z}{|z|^2+1}, \qquad P_3(z) = \frac{|z|^2-1}{|z|^2+1}.$$

The extended complex plane $\hat{\mathbb{C}}$ thus carries the real predicates P_1, P_2, P_3 with values in $[-1, 1]$. Through the same identification, the topology of $\hat{\mathbb{C}}$ is metrized by the pullback of the Euclidean metric of S^2.

Newton CCSs: steps and fractals. Here we describe the compositional computation structure generated by the Newton map, the single step that Newton's method iterates to approximate a root of a function. The structure so obtained is the common setting of the examples that follow. We then recall the Fatou and Julia sets, on which the iterates of the Newton map behave regularly and chaotically, respectively.

Let $p = p(z)$ be a nonconstant rational function with complex coefficients, that is, a nonconstant meromorphic map $\hat{\mathbb{C}} \to \hat{\mathbb{C}}$. The *Newton step*, or *Newton map*, of p is the rational map

$$N_p : z \mapsto z - \frac{p(z)}{p'(z)}. \tag{4.19}$$

The iterates of the Newton map, that is, the compositional powers N_p^n of N_p, may be regarded as the computations of a CCS with state space $L = \hat{\mathbb{C}}$. The natural predicates are the coordinates P_1, P_2, P_3 of the inverse stereographic projection, since the real and imaginary parts $z \mapsto \Re z$ and $z \mapsto \Im z$ are not defined at $z = \infty$.

Let $\Gamma_p \coloneqq \{N_p^n : n \in \mathbb{N}\}$ be the semigroup of iterates of the Newton map (4.19). The CCS $(\hat{\mathbb{C}}, (P_1, P_2, P_3), \Gamma_p)$ is called *Newton's method for $p(z)$*, or the *Newton CCS for $p(z)$*. Since the state space $\hat{\mathbb{C}}$ is compact and the Newton map N_p is continuous on $\hat{\mathbb{C}}$, every Newton CCS satisfies the Extendibility Axiom.

Newton's method succeeds at an initial estimate $z_0 \in \hat{\mathbb{C}}$ when the sequence $(z_n)_{n\in\mathbb{N}}$ of Newton steps, where $z_n \coloneqq N_p^n(z_0)$, converges to a root of p.

Fatou and Julia sets. Recall that the *Fatou set* $\mathcal{F}_g$ of a rational map $g\colon \hat{\mathbb{C}} \to \hat{\mathbb{C}}$ is the set of points near which the iterates of g behave regularly. More precisely, a point belongs to $\mathcal{F}_g$ if the point has a neighborhood on which every sequence of iterates of g has a subsequence that converges uniformly with respect to the spherical metric. The complement $\mathcal{J}_g \coloneqq \hat{\mathbb{C}} \setminus \mathcal{F}_g$ of the Fatou set is the *Julia set* of g, a closed set that is typically a fractal. The Julia set of the Newton map N_p is called the *Newton fractal* of p.

The Newton CCS for the cubic $q(z) = z^3 - 2z + 2$. We begin with the second motivating example of Section 2.5, a cubic polynomial whose Newton map has, besides the superattracting fixed points at the three roots, an attracting cycle of period two. Newton's method therefore fails on an open set of initial estimates, on which the iterates of the Newton map do not converge. For the same reason, the deep computations of the method include two distinct limits of the iterates: one along an ultrafilter that contains the even numbers, the other along an ultrafilter that contains the odd numbers. The polynomial $q(z) = z^3 - 2z + 2$ has Newton map

$$N_q : z \mapsto z - \frac{z^3 - 2z + 2}{3z^2 - 2} = \frac{2z^3 - 2}{3z^2 - 2}.$$

Denote the real root of q by r_0, and the roots in the upper and the lower half plane by r_+ and r_-. Starting from $z_0 = 0$, the Newton steps alternate $0, 1, 0, 1, \ldots$, and neither 0 nor 1 is a root of q.

Let B_0, B_+, B_- be the basins of the roots r_0, r_+, r_-, and let B_{01} be the basin of the superattracting cycle $\{0, 1\}$. The Fatou set of N_q is $B_{01} \cup B_0 \cup B_+ \cup B_-$, and each of these four sets is nonempty and open. Since B_{01} is nonempty, the iterates N_q^n do not converge pointwise. Fix nonprincipal ultrafilters $\mathcal{U}_0$ and $\mathcal{U}_1$ on $\mathbb{N}$ that contain the even and the odd numbers, respectively. The limits of the iterates N_q^n along $\mathcal{U}_0$ and along $\mathcal{U}_1$ are deep computations, which we denote by f_0 and f_1, respectively. On the Fatou set the even and the odd iterates converge pointwise, whence the restrictions of the two limits to the Fatou set do not depend on the choice of the ultrafilters. Both limits send each basin of a root to that root; on B_{01}, by contrast, they take only the values 0 and 1 and exchange them. The exchange of the two values follows from the identity $f_1 = N_q \circ f_0$, which holds on the Fatou set by the continuity of N_q. In particular, the two deep computations are distinct.[10] Figure 9 depicts f_0 and f_1, together with the Newton fractal of q, on which the iterates behave chaotically.




FIGURE 9. The deep computations f_0 and f_1 of Newton's method for $z^3 - 2z + 2$, colored by basin.

The Newton CCS for the cubic $u(z) = z^3 - 1$. For contrast, we turn to $u(z) = z^3 - 1$, whose roots are the cube roots of unity $r_0 = 1$ and $r_\pm = \exp(\pm 2\pi i/3)$. The Newton map of u is

$$N_u : z \mapsto z - \frac{z^3 - 1}{3z^2} = \frac{2z^3 + 1}{3z^2}.$$

The critical points of this rational map of degree three are the three roots of u, which are superattracting fixed points, and the point 0. The map N_u sends the point 0 to ∞, a repelling fixed point of multiplier $3/2$. The immediate basin of every attracting or parabolic cycle contains a critical point whose orbit converges to the cycle [Mil06, Sections 8 and 10]. Unlike N_q, the map N_u therefore has no attracting cycle other than its three attracting fixed points, the roots of u, and it has no parabolic cycle. The boundary of a Siegel disk or of a Herman ring, moreover, is infinite and lies in the closure of the critical orbits [Mil06, Sections 11 and 15]. Since every critical orbit of N_u is finite, the map N_u has no such domain.

[10]The point ∞ is a repelling fixed point of N_q: if $|z|$ is large, then $N_q(z) = \frac{2}{3}z + O(1)$. Hence the point ∞ lies on the Newton fractal of q, and the two deep computations satisfy $f_0(\infty) = \infty = f_1(\infty)$.

Sullivan's theorem excludes wandering domains, and the classification of periodic Fatou components then shows that the Fatou set of N_u is the union of the basins of the three roots [Mil06, Section 16 and Appendix F]. On each of these basins, the iterates of N_u converge to the corresponding root. Figure 10 depicts Newton's method for u. To render the iterates themselves, and not merely their limits, we color each point z_0 by blending the colors of the three roots with relative weights $|z_n - r|^{-3}$ for $r \in \{r_0, r_+, r_-\}$, where $z_n := N_u^n(z_0)$. This coloring is a continuous function of z_0, blurred for small n. As n increases, the coloring sharpens into large regions of nearly constant color, separated by narrow regions in which the color varies so rapidly that the coloring is nearly discontinuous.




FIGURE 10. Newton's method for $z^3 - 1$: one and three iterations, and the limit on the Fatou set.

The chaos of Newton's method on the Julia set of the Newton map is visible to the predicates P_1, P_2, P_3. In Propositions 4.20 and 4.21 below we show that, for the polynomials $z^2 - 1$, q and u, this chaos prevents the Newton CCS on the Riemann sphere from being NIP.

Newton's method on the Riemann sphere. On the Julia set of the Newton map the iteration is chaotic, and the chaos is visible to the predicates of the CCS. We exhibit the independence in the simplest case, the square roots of 1. Let $p(z) = z^2 - 1$, whose Newton map on $\hat{\mathbb{C}}$ is $N(z) = \frac{1}{2}\big(z + \frac{1}{z}\big)$, and recall the second coordinate P_2 of the inverse stereographic projection $\hat{\mathbb{C}} \to S^2$ defined above.

Proposition 4.20. *The functions $P_2 \circ N^{3k}$, $k \in \mathbb{N}$, form an independent sequence: for all finite disjoint sets $I, J \subseteq \mathbb{N}$ there is $z \in \hat{\mathbb{C}}$ with $P_2(N^{3k}z) \geq \frac{1}{2}$ for $k \in I$ and $P_2(N^{3k}z) \leq -\frac{1}{2}$ for $k \in J$. In particular, the Newton CCS of p is not NIP.*

Proof. The Möbius transformation $\psi(z) = (z-1)/(z+1)$ conjugates N to the squaring map: $\psi(N(z)) = \psi(z)^2$. The map ψ sends the imaginary axis, together with ∞, onto the unit circle, with $\psi^{-1}(e^{i\theta}) = i\cot(\theta/2)$. Hence, for $z = i\cot(\theta/2)$, we have $N^n(z) = i\cot(2^n\theta/2)$ and

$$P_2\big(N^n(z)\big) = \frac{2\cot(2^{n-1}\theta)}{\cot^2(2^{n-1}\theta) + 1} = \sin(2^n\theta).$$

Writing $\theta = 2\pi\sum_{j\geq 1} x_j 2^{-j}$ with bits x_j, we find that the fractional part of $2^{3k}\theta/2\pi$ has binary expansion $0.x_{3k+1}x_{3k+2}x_{3k+3}\cdots$. If $(x_{3k+1}, x_{3k+2}, x_{3k+3}) = (0,1,0)$,

this fractional part lies in $[\frac{1}{4}, \frac{3}{8}]$, and hence $\sin(2^{3k}\theta) \geq \sin\frac{3\pi}{4} > \frac{1}{2}$. If the block is $(1,1,0)$, the fractional part lies in $[\frac{3}{4}, \frac{7}{8}]$, whence $\sin(2^{3k}\theta) < -\frac{1}{2}$. Choosing the block $(0,1,0)$ for $k \in I$ and $(1,1,0)$ for $k \in J$ gives the required z. □

The proof also shows that the doubling map $x \mapsto 2x$ of the circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$, with the predicates $\cos 2\pi x$ and $\sin 2\pi x$, is not NIP: with $x = \theta/2\pi$, the functions $\sin(2\pi\, 2^{3k}x)$ form an independent sequence. The doubling map shifts the binary digits of x, and the independence comes from the freedom to choose those digits.

On the real projective line, by contrast, the same algorithm is tame. The Julia set of N, the imaginary axis together with ∞, meets $\mathbb{RP}^1$ only at 0 and ∞, and on $\mathbb{RP}^1$ the iterates converge to the deep computation of Section 4.2 for $a = 1$. By Proposition 4.2, the corresponding CCS is therefore $\mathrm{NIP}^{\mathrm{met}}$ (see Section 4.2). Whether Newton's method is tame is thus decided by the space of states, and not by the formula that defines the Newton step.

Rational maps of the sphere. The chaos exhibited in Proposition 4.20 is not peculiar to the square roots of 1. Chaos of this kind, and with it the independence property, is present on the Julia set of every rational map of degree at least 2. The proof of Proposition 4.21 below rests on two classical facts [Mil06, Sections 4 and 14] about the Julia set $\mathcal{J}_g$ of such a map g (see Section 4.7). The first is that $\mathcal{J}_g$ is a nonempty compact set without isolated points that satisfies $g^{-1}(\mathcal{J}_g) = \mathcal{J}_g$. The second is that, for every neighborhood U of a point of $\mathcal{J}_g$, the image $g^m(U)$ contains $\mathcal{J}_g$ for all sufficiently large m. This second fact is a consequence of Montel's theorem and of the density of repelling cycles in $\mathcal{J}_g$.

Proposition 4.21. *Let $g\colon \hat{\mathbb{C}} \to \hat{\mathbb{C}}$ be a rational map of degree at least 2, and consider the CCS with states $\hat{\mathbb{C}}$, the predicates P_1, P_2, P_3 of Section 4, and the semigroup of iterates of g. There are $j \in \{1,2,3\}$, a positive integer n and reals $a < b$ such that the functions $P_j \circ g^{nk}$, $k \in \mathbb{N}$, form an independent sequence: for all finite disjoint sets $I, J \subseteq \mathbb{N}$ there is $z \in \hat{\mathbb{C}}$ with $P_j(g^{nk}(z)) \leq a$ for $k \in I$ and $P_j(g^{nk}(z)) \geq b$ for $k \in J$. In particular, the CCS is not NIP, and neither are the Newton CCSs for $q(z) = z^3 - 2z + 2$ and for $u(z) = z^3 - 1$.*

Proof. Choose distinct points $z_0, z_1 \in \mathcal{J}_g$ and an index j with $P_j(z_0) \neq P_j(z_1)$, say $P_j(z_0) < P_j(z_1)$, and reals $a < b$ strictly between these two values. By continuity, the points z_0 and z_1 have closed neighborhoods U_0 and U_1 such that $P_j \leq a$ on U_0 and $P_j \geq b$ on U_1, and we may choose n with $g^n(U_0) \supseteq \mathcal{J}_g$ and $g^n(U_1) \supseteq \mathcal{J}_g$. Let $I, J \subseteq \mathbb{N}$ be finite and disjoint, and let m exceed all their elements; let $U^{(k)}$ denote U_0 for $k \in I$ and U_1 for $k \notin I$. Let $C_m = U^{(m)} \cap \mathcal{J}_g$ and, for $k = m-1, \ldots, 0$, let $C_k = U^{(k)} \cap \mathcal{J}_g \cap g^{-n}(C_{k+1})$. Each C_k is nonempty: since $g^n(U^{(k)}) \supseteq \mathcal{J}_g \supseteq C_{k+1}$, some point of $U^{(k)}$ is mapped by g^n into C_{k+1}, and that point lies in $\mathcal{J}_g$ because $g^{-1}(\mathcal{J}_g) = \mathcal{J}_g$. Every $z \in C_0$ satisfies $g^{nk}(z) \in U^{(k)}$ for $k \leq m$, and hence $P_j(g^{nk}(z)) \leq a$ for $k \in I$ and $P_j(g^{nk}(z)) \geq b$ for $k \in J$. For the last assertion, note that the Newton maps $N_q(z) = (2z^3 - 2)/(3z^2 - 2)$ and $N_u(z) = (2z^3 + 1)/(3z^2)$ of Section 4 have degree 3, since in each quotient the numerator and the denominator have no common zero. □

The logistic map at $r = 4$. The doubling map also governs the logistic map $f_4(x) = 4x(1-x)$ of Section 4.2, as Figure 11 illustrates.

Proposition 4.22. *The functions f_4^{3k}, $k \in \mathbb{N}$, form an independent sequence on $[0,1]$: for all finite disjoint sets $I, J \subseteq \mathbb{N}$ there is $x \in [0,1]$ with $f_4^{3k}(x) \geq \frac{3}{4}$ for*

$k \in I$ and $f_4^{3k}(x) \leq \frac{1}{4}$ for $k \in J$. In particular, the logistic CCS for $r = 4$ is not NIP.

Proof. Let $h(\theta) = \sin^2(\theta/2)$ for $\theta \in \mathbb{R}$, and note the identity $f_4(h(\theta)) = 4\sin^2(\theta/2)\cos^2(\theta/2) = \sin^2\theta = h(2\theta)$. Iterated, this identity yields $f_4^n(h(\theta)) = h(2^n\theta) = \sin^2(2^{n-1}\theta)$. Writing $\theta = 2\pi\sum_{j\geq 1} x_j 2^{-j}$ with bits x_j, we find that the fractional part of $2^{3k}\theta/2\pi$ has binary expansion $0.x_{3k+1}x_{3k+2}x_{3k+3}\cdots$. Moreover, the angle $2^{3k-1}\theta$ differs from π times this fractional part by an integer multiple of π. If $(x_{3k+1}, x_{3k+2}, x_{3k+3}) = (0,1,1)$, this fractional part lies in $[\frac{3}{8}, \frac{1}{2}]$, and hence $f_4^{3k}(h(\theta)) = \sin^2(2^{3k-1}\theta) \geq \sin^2\frac{3\pi}{8} > \frac{3}{4}$. If the block is $(0,0,0)$, the fractional part lies in $[0, \frac{1}{8}]$, whence $f_4^{3k}(h(\theta)) \leq \sin^2\frac{\pi}{8} < \frac{1}{4}$. Choosing the block $(0,1,1)$ for $k \in I$ and $(0,0,0)$ for $k \in J$, and taking $x = h(\theta)$, gives the required point. □




FIGURE 11. Independence for the doubling map and for f_4.

Thus the logistic family, which is stable for $0 < r \leq 1$ and NIP^{met} but not stable for $1 < r \leq 3$ (see Proposition 4.4), leaves the hierarchy at $r = 4$. Let r_∞ be the supremum of the parameters $r \in (0,4]$ for which every periodic point of f_r has a power of 2 as period; this number is the Feigenbaum parameter $r_\infty \approx 3.5699$, the limit of the cascade of period doublings. The passage out of the hierarchy takes place at r_∞, and the family skips the intermediate levels. Indeed, in Proposition 4.23 below we show that the family remains NIP^{met} below r_∞ and has the independence property beyond r_∞, periodic windows included. Moreover, in Remark 4.24 we sketch the argument that the family is still NIP^{met} at r_∞. The arguments rest on the kneading theory of Milnor and Thurston [MT88] and on the theory of interval maps all of whose periodic points have periods that are powers of 2 [BC92, dMvS93].

Proposition 4.23. *(i) For $3 < r < r_\infty$, the logistic CCS is NIP^{met} but not stable.*

(ii) For $r_\infty < r \leq 4$, there are a positive integer n and reals $a < b$ such that the functions f_r^{nk}, $k \in \mathbb{N}$, form an independent sequence: for all finite disjoint sets $I, J \subseteq \mathbb{N}$ there is $x \in [0,1]$ with $f_r^{nk}(x) \leq a$ for $k \in I$ and $f_r^{nk}(x) \geq b$ for $k \in J$. In particular, the logistic CCS is not NIP.

Proof. Writing f for f_r, we use three classical facts [MT88, BC92, dMvS93]. First, by the monotonicity of the kneading invariant in the logistic family, every periodic point of f has a power of 2 as period if and only if $r \leq r_\infty$. Moreover, for $r < r_\infty$ only finitely many periods occur, whence f has only finitely many periodic points. Second, for $2 < r < 4$, every limit point of an orbit that starts in $(0,1)$ lies in

the interval $[f^2(\frac{1}{2}), f(\frac{1}{2})]$, which does not contain 0. Third, a continuous map of a compact interval with only finitely many periodic points has every orbit asymptotic to a periodic orbit [BC92].

(i) Suppose that $3 < r < r_\infty$, and let N be a common multiple of the periods of f, which are powers of 2. If the orbit of x is asymptotic to the periodic orbit $\mathcal{Q}$, then $f^{Nk}(x)$ converges to a point of $\mathcal{Q}$. Indeed, the map f^N fixes $\mathcal{Q}$ pointwise, and by uniform continuity the point of $\mathcal{Q}$ nearest to $f^i(x)$ is eventually carried by f to the point of $\mathcal{Q}$ nearest to $f^{i+1}(x)$. Let $\ell_r(x)$ be the limit of the sequence $(f^{Nk}(x))_{k\in\mathbb{N}}$. Then, for $0 \le i < N$, the maps f^{Nk+i} converge pointwise to $f^i \circ \ell_r$ as k tends to infinity, and every limit of the iterates along a nonprincipal ultrafilter on $\mathbb{N}$ is one of these N maps. Hence $E([0,1], f)$ is countable, and therefore metrizable. Each of the maps $f^i \circ \ell_r$ is of Baire class 1, being the pointwise limit of a sequence of iterates, and Proposition 4.1 therefore shows that the CCS is $\mathrm{NIP}^{\mathrm{met}}$. The map ℓ_r, which fixes 0 and maps $(0,1)$ into the interval $[f^2(\frac{1}{2}), f(\frac{1}{2})]$, is thus discontinuous at 0, and the CCS is not stable.

(ii) Suppose that $r_\infty < r \le 4$; by the first fact, the map f then has a periodic point of period not a power of 2. Hence, by a theorem of Block and Coppel [BC92, Chapter II], some iterate f^n is strictly turbulent. Strict turbulence of f^n means that there are disjoint closed intervals H_0 and H_1, with H_0 to the left of H_1, such that $f^n(H_0)$ and $f^n(H_1)$ both contain $H_0 \cup H_1$. Let $a = \max H_0$ and $b = \min H_1$, let $I, J \subseteq \mathbb{N}$ be finite and disjoint, and let m exceed all their elements; let $H^{(k)}$ denote H_0 for $k \in I$ and H_1 for $k \notin I$. Let $C_m = H^{(m)}$ and, for $k = m-1, \dots, 0$, let $C_k = H^{(k)} \cap f^{-n}(C_{k+1})$. Each set C_k is nonempty, since $f^n(H^{(k)}) \supseteq C_{k+1}$. Every $x \in C_0$ satisfies $f^{nk}(x) \in H^{(k)}$ for $k \le m$, and hence $f^{nk}(x) \le a$ for $k \in I$ and $f^{nk}(x) \ge b$ for $k \in J$. □

Remark 4.24. At the Feigenbaum parameter itself the logistic CCS is again $\mathrm{NIP}^{\mathrm{met}}$ but not stable; we sketch the argument, which rests on the structure of infinitely renormalizable maps [BC92, dMvS93]. For $r = r_\infty$ and every $m \in \mathbb{N}$, the map $f = f_r$ has a cycle of 2^m pairwise disjoint closed intervals, permuted cyclically by f and contained in the intervals of the cycle of level $m-1$. Moreover, the lengths of the intervals of level m tend to 0 as $m \to \infty$. Finally, every orbit that starts in $(0,1)$ either lands on a periodic orbit or enters the cycle of every level. Label the intervals of the cycle of level m by the residues modulo 2^m, in such a way that f adds 1 to the label. Let $\mathcal{U}$ be a nonprincipal ultrafilter on $\mathbb{N}$, and let ς be the limit of n along $\mathcal{U}$ in the ring $\mathbb{Z}_2$ of 2-adic integers. If the orbit of x lands on a periodic orbit, of period 2^m, then $f^n(x)$ is eventually determined by the residue of n modulo 2^m. Otherwise, for each m, the point $f^n(x)$ eventually lies in the interval of level m whose label differs from n by a certain residue modulo 2^m. This residue depends on x and m but not on n, and the intervals obtained for successive m shrink to a point. In either case, the points $f^n(x)$ converge along $\mathcal{U}$ to a limit $\ell^\varsigma(x)$ that depends only on ς. The map ℓ^ς is the pointwise limit of the iterates f^{n_j} along any sequence of natural numbers n_j that tends to infinity and converges to ς in $\mathbb{Z}_2$. Each map ℓ^ς is therefore of Baire class 1, and $E([0,1], f)$ consists of the iterates and the maps ℓ^ς, $\varsigma \in \mathbb{Z}_2$. The rational points of $[0,1]$, among them the point $\frac{1}{2}$, separate the members of $E([0,1], f)$: distinct iterates are continuous and differ at a rational point. Moreover, an iterate f^n and a map ℓ^ς differ at every rational point $x > 0$ so small that $f^n(x) < f^2(\frac{1}{2})$, since ℓ^ς maps $(0,1)$ into $[f^2(\frac{1}{2}), f(\frac{1}{2})]$. Finally, two maps ℓ^ς and $\ell^{\varsigma'}$ with $\varsigma \ne \varsigma'$ differ at $\frac{1}{2}$, whose orbit lands on no periodic orbit.

Indeed, whenever $\varsigma \not\equiv \varsigma'$ modulo 2^m, the points $\ell^\varsigma(\frac{1}{2})$ and $\ell^{\varsigma'}(\frac{1}{2})$ lie in distinct intervals of level m. As in the proof of Proposition 3.4(i), the space $E([0,1], f)$ is therefore metrizable, and by Proposition 4.1 the CCS is $\mathrm{NIP}^{\mathrm{met}}$. The CCS is not stable, since each map ℓ^ς fixes 0 and is discontinuous there.

The Thue–Morse sequence. Even a sequence produced by a simple substitution rule can be wild. The Thue–Morse sequence t_n is the parity of the number of ones in the binary expansion of n. The same sequence is the limit of the words obtained from 0 by the substitution $0 \mapsto 01$, $1 \mapsto 10$. The two-sided sequences all of whose finite blocks occur in (t_n) form the Thue–Morse subshift $X_{TM} \subseteq 2^{\mathbb{Z}}$, on which the shift σ is a homeomorphism. Glasner and Megrelishvili observed that the Morse minimal system, which consists of the subshift X_{TM} and the action of $\mathbb{Z}$ generated by σ, is not tame [GM18]. In Appendix A we give a self-contained proof, which relies only on the structure theorem for tame minimal systems. By Proposition 4.1, the CCS with states X_{TM}, the coordinate predicates and the group $\{\sigma^n : n \in \mathbb{Z}\}$ is therefore not NIP. The cascade CCS of σ, whose computations are only the iterates σ^n with $n \in \mathbb{N}$, is not NIP either. Indeed, the reversal of two-sided sequences maps X_{TM} onto itself and conjugates σ to its inverse. The reversal therefore turns an independent sequence of functions $f \circ \sigma^{n_k}$ with negative exponents n_k into one with positive exponents. Both facts are stated, with complete proofs, in Proposition A.1.

4.8. **Summary.** Table 3 collects the examples of this section by the levels of the hierarchy, that is, by the alternatives of Todorčević's trichotomy; Table 2 collects them by the routes of the heptachotomy. The Euclidean algorithm and free reduction are actions of groups with two generators, and the family $\Delta_{\mathcal{C}}$ consists of elements of the first of these groups. Each of the remaining entries of the table is a single map iterated.

5. Randomized versions of NIP and Monte Carlo computability of deep computations

Replacing topology by measure, we dispense in this section with the countability of the set $\mathcal{P}$ of predicates, on which the theory of Sections 3–4 depends. In those sections we studied deep computations as *topological* limits of computations, and we showed that, for countable $\mathcal{P}$ and under the NIP, each deep computation is of the first Baire class on the type shards and, on each shard, is the limit of a sequence of computations. The theory of those sections rests on the type shards being Polish, and hence on the countability of $\mathcal{P}$, which is used in the proof of Theorem 3.1. In place of membership in the first Baire class, we now ask that a deep computation extend to the type shards as a *universally measurable* map, that is, as a map measurable with respect to every Radon probability measure on the type shards. By Luzin's theorem and the Stone–Weierstrass theorem, each output feature of such an extension is then, for every such measure, a uniform limit of polynomials in the features outside a set of arbitrarily small measure. We call a deep computation with an extension of this kind *universally Monte Carlo computable* (see Definition 5.6). The role of the Bourgain–Fremlin–Talagrand dichotomy is then taken by the measure-theoretic form of the dichotomy (see Theorem 5.5), and the role of the NIP is shared with *Talagrand stability*, a measure-theoretic condition introduced by Talagrand in his study of Pettis integration (see Definition 5.9).

Level	Computation	Deep computations; source
stable	rotation of the circle	the circle group
	logistic map, $0 < r \leq 1$	iterates and one continuous limit; Prop. 4.4
$\mathrm{NIP}^{\mathrm{met}}$, not stable	power method	countable; Prop. 4.3
	Newton's method for $x^2 - a$ on $\mathbb{RP}^1$	countable; Prop. 4.2
	logistic map, $1 < r \leq r_\infty$	countable for $r < r_\infty$, metrizable for $r = r_\infty$; Props. 4.4, 4.23, Rem. 4.24
$\mathrm{NIP}^{\mathrm{hs}}$, not $\mathrm{NIP}^{\mathrm{met}}$	Bresenham, plane (Sturmian shift)	isolated iterates and a double arrow circle; Thm. 4.5
$\mathrm{NIP}^{\mathrm{fc}}$, not $\mathrm{NIP}^{\mathrm{hs}}$	Bresenham, space (two Sturmian shifts)	isolated iterates and a copy of $\mathbb{D} \times \mathbb{D}$; contains $\hat{D}(S(2^{\mathbb{N}}))$, not $D(2^{\mathbb{N}})$; Thm. 4.10
	the family $\Delta_{\mathfrak{C}}$ in $\mathrm{PGL}(2, \mathbb{Z})$	a copy of $\hat{D}(2^{\mathbb{N}})$; Prop. 4.15
	a single map, with $D(2^{\mathbb{N}})$	open; Question 6.3
NIP, not $\mathrm{NIP}^{\mathrm{fc}}$	Euclidean algorithm: $\mathrm{PGL}(2, \mathbb{Z})$ on $\mathbb{RP}^1$ (group)	contains $A(2^{\mathbb{N}})$, of exact equality tests, and $D(2^{\mathbb{N}})$; Cor. 4.14, Prop. 4.15
	free reduction: F_2 on ∂F_2 (group)	not first countable; [GM22]
	a single map	open; Question 6.2
not NIP	doubling map of the circle	Prop. 4.20
	Newton's method for $z^2 - 1$ on $\hat{\mathbb{C}}$	Prop. 4.20
	Newton's method for q and u on $\hat{\mathbb{C}}$	Prop. 4.21
	logistic map, $r_\infty < r \leq 4$	Props. 4.22, 4.23
	Thue–Morse shift	Prop. A.1

TABLE 3. Computations at each level of the hierarchy and beyond it, with the cases left open.

The first main result of the section, Theorem 5.7, characterizes the NIP by the universal measurability of the restrictions of ultracomputations to the type shards, and derives Monte Carlo computability from the NIP thus characterized. The second main result, Corollary 5.15, derives the same conclusion from universal Talagrand stability. By Theorem 5.16, these two hypotheses, the NIP and universal Talagrand stability, coincide for countable sets of computations (see Corollary 5.17).

5.1. **NIP and Monte Carlo computability of deep computations.** Of the two routes to universal Monte Carlo computability followed in this section, the first starts from the NIP and passes through universal measurability, which we introduce here. Universal measurability is the measure-theoretic counterpart of membership

in the first Baire class. This first route leads to Theorem 5.7, by which the deep computations of a confined set of computations with the NIP on shards and features are universally Monte Carlo computable. By Fact 1.2, a function $f\colon X \to \mathbb{R}$ on a perfectly normal space X is of Baire class 1 if and only if $f^{-1}[U]$ is an F_σ subset of X for every open $U \subseteq \mathbb{R}$. The following definition replaces the F_σ condition by measurability with respect to every Radon measure.

Definition 5.1. Let X be a Hausdorff space, and let (Y, Σ) be a measurable space. A map $f : X \to Y$ is *universally measurable* (with respect to Σ) if $f^{-1}(E)$ is measurable with respect to the completion of μ for every Radon measure μ on X and every $E \in \Sigma$.

When $Y = \mathbb{R}$, we always take Σ to be the Borel σ-algebra $\mathcal{B}(\mathbb{R})$ of $\mathbb{R}$.

If X is a compact Hausdorff space, then Radon measures on X are finite, and every nonzero Radon measure on X becomes a probability measure after normalization. Moreover, the sets $E \subseteq \mathbb{R}$ whose preimages under a given function are μ-measurable form a σ-algebra, which contains the Borel sets as soon as it contains the open sets. Together, these two observations yield the following simplification of Definition 5.1 for compact spaces.

Remark 5.2. If X is compact, then a function $f : X \to \mathbb{R}$ is universally measurable if and only if $f^{-1}(U)$ is μ-measurable for every Radon probability measure μ on X and every open set $U \subseteq \mathbb{R}$.

An account of universally measurable sets and functions may be found in the *Handbook of measure theory* [Pap02, Chapters 1 and 2].

Notation 5.3. Following Bourgain, Fremlin and Talagrand [BFT78], we denote by $M_r(X)$ the collection of all universally measurable real-valued functions on X. For a Radon measure μ on X, we further denote by $\mathscr{M}^0(X, \mu)$ the collection of all real-valued functions on X measurable with respect to the completion of μ. A set $A \subseteq \mathbb{R}^X$ is *relatively countably compact* in $\mathscr{M}^0(X, \mu)$ if every countably infinite subset of A has an accumulation point in $\mathscr{M}^0(X, \mu)$.

The maps that arise in the study of deep computations, transitions and deep computations themselves among them, send a state space $L \subseteq \mathbb{R}^{\mathcal{P}}$ into $\mathbb{R}^{\mathcal{P}}$. The product space $\mathbb{R}^{\mathcal{P}}$ carries two natural σ-algebras: the Borel σ-algebra, generated by the open subsets of $\mathbb{R}^{\mathcal{P}}$, and the cylinder σ-algebra. The latter, generated by the subbasic open sets $\pi_P^{-1}(U)$ with $P \in \mathcal{P}$ and $U \subseteq \mathbb{R}$ open, is in general strictly smaller than the Borel σ-algebra, but the two σ-algebras coincide when $\mathcal{P}$ is countable. We use the cylinder σ-algebra to define universally measurable maps into $\mathbb{R}^{\mathcal{P}}$, because of the following characterization.

Proposition 5.4. *Let X be a Hausdorff space and $Y = \prod_{i \in I} Y_i$ be any product of measurable spaces (Y_i, Σ_i) for $i \in I$. Let Σ_Y be the cylinder σ-algebra generated by the measurable spaces (Y_i, Σ_i), and let $\pi_i\colon Y \to Y_i$ be the projections. The following are equivalent for $f : X \to Y$:*

(i) The map $f : X \to Y$ is universally measurable (with respect to Σ_Y).
(ii) The map $\pi_i \circ f : X \to Y_i$ is universally measurable (with respect to Σ_i) for all $i \in I$.

Proof. The implication (i)⇒(ii) holds because each projection π_i is measurable and $(\pi_i \circ f)^{-1}(E) = f^{-1}(\pi_i^{-1}(E))$ for every $E \in \Sigma_i$. To prove (ii)⇒(i), fix a Radon

measure μ on X, and note that the sets $C \subseteq Y$ for which $f^{-1}(C)$ is μ-measurable form a σ-algebra. This σ-algebra contains every measurable cylinder $C = \prod_{i \in I} C_i$: if J is the finite set of indices $i \in I$ with $C_i \neq Y_i$, then $f^{-1}(C) = \bigcap_{i \in J} (\pi_i \circ f)^{-1}(C_i)$ is μ-measurable by (ii). Since the measurable cylinders generate Σ_Y, the set $f^{-1}(C)$ is μ-measurable for every $C \in \Sigma_Y$. □

By Proposition 5.4, the universal measurability of a map into $\mathbb{R}^{\mathcal{P}}$ can be checked feature by feature, just as membership in the first Baire class can (see Proposition 1.8).

We saw in Section 1 that the NIP corresponds to relative compactness in the space of functions of Baire class 1 (see Theorems 1.4, 1.6 and 1.9). In Section 3 we saw, moreover, that this correspondence extends to deep computations when the set $\mathcal{P}$ of predicates is countable (see Theorem 3.1). The following theorem, which holds on every Hausdorff space, characterizes the NIP in terms of universal measurability. Applied to the type shards, the theorem therefore requires no countability of $\mathcal{P}$.

Theorem 5.5 (Bourgain–Fremlin–Talagrand [BFT78, Theorem 2F])**.** *Let X be a Hausdorff space, and let $A \subseteq \mathrm{C_p}(X)$ be pointwise bounded. The following are equivalent:*

(i) The closure $\overline{A}$ is contained in $M_r(X)$.
(ii) For every compact $K \subseteq X$, the restriction $A|_K$ satisfies the NIP.
(iii) For every Radon measure μ on X, the set A is relatively countably compact in $\mathscr{M}^0(X, \mu)$.

Theorem 5.5 leads to a measure-theoretic counterpart of computability, the universal Monte Carlo computability of Definition 5.6 below, whose motivation lies in two classical facts.

(1) Littlewood's second principle, that every Lebesgue measurable function is "nearly continuous", is made precise by Luzin's theorem. Let (X, Σ, μ) be a complete finite Radon measure space, and let Y be a second-countable topological space (e.g., $Y = \mathbb{R}^{\mathcal{P}}$ with $\mathcal{P}$ countable), equipped with its Borel σ-algebra. Then a function $f\colon X \to Y$ is measurable if and only if, for every $E \in \Sigma$ and every $\varepsilon > 0$, there is a closed set $F \subseteq E$ such that the restriction $f|_F$ is continuous and $\mu(E \setminus F) < \varepsilon$.
(2) Computability of a transform is characterized by the continuous extendibility of the transform to the space of types (see Section 2.3); this characterization is at the core of the first paper of this series [ADIW26].

Taken together, the two facts justify Definition 5.6 below. Indeed, suppose that an extension of a transform is universally measurable on a type shard, and fix a Radon probability measure on the shard. By Luzin's theorem, each output feature of the extension is then continuous on a closed subset of the shard whose measure is as close to 1 as desired. On this subset, by the Stone–Weierstrass theorem, the output feature in question is a uniform limit of polynomials in the features.

Definition 5.6. Let $(L, \mathcal{P}, \Gamma)$ be a CCS, and let $\mathscr{R}$ be a collection of sizers. A transform $f\colon L \to \mathcal{L}_{\mathrm{sh}}$ is *universally Monte Carlo computable on $\mathscr{R}$* if f has an extension $\tilde{f}\colon \mathcal{L}_{\mathrm{sh}} \to \mathcal{L}_{\mathrm{sh}}$ such that, for every $r_\bullet \in \mathscr{R}$, there is a sizer $s_\bullet$ for which the restriction $\tilde{f}|_{\mathcal{L}[r_\bullet]}\colon \mathcal{L}[r_\bullet] \to \mathcal{L}[s_\bullet]$ is universally measurable.

The deep computations of a set of computations confined by an exhaustive collection of sizers are transforms $L \to \mathcal{L}_{\mathrm{sh}}$ (see Proposition 2.5). Since each type shard

$\mathcal{L}[r_\bullet]$ is compact, Proposition 5.4 and Remark 5.2 show that the universal measurability required in Definition 5.6 means the following: for every Radon probability measure μ on $\mathcal{L}[r_\bullet]$ and every $P \in \mathcal{P}$, the map $\pi_P \circ \tilde{f}|_{\mathcal{L}[r_\bullet]} \colon \mathcal{L}[r_\bullet] \to [-s_P, s_P]$ is μ-measurable.

Theorem 5.7. *Let $(L, \mathcal{P}, \Gamma)$ be a CCS satisfying the Extendibility Axiom, let $\mathscr{R}$ be an exhaustive collection of sizers, and let $\Delta \subseteq \Gamma$ be $\mathscr{R}$-confined. The following are equivalent:*

(i) for all $P \in \mathcal{P}$ and all $r_\bullet \in \mathscr{R}$, the set $\pi_P \circ \Delta|_{L[r_\bullet]}$ satisfies the NIP;

(ii) for every $r_\bullet \in \mathscr{R}$, every member of $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$ is universally measurable, as a map into $\mathbb{R}^{\mathcal{P}}$ with the cylinder σ-algebra.

If these conditions hold, then every deep computation in $\overline{\Delta}$ is universally Monte Carlo computable on $\mathscr{R}$.

Proof. Fix $r_\bullet \in \mathscr{R}$ and $P \in \mathcal{P}$, and write $\mathcal{K} = \overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$, a compact subset of $\mathcal{L}[r_\bullet]^{\mathcal{L}[r_\bullet]}$, and $A_P = \pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$. By the Extendibility Axiom and confinement, the set A_P consists of continuous functions on the compact space $\mathcal{L}[r_\bullet]$ and is pointwise bounded. Moreover, since the map $g \mapsto \pi_P \circ g$ is continuous, the closure of A_P is the compact set $\{\pi_P \circ g : g \in \mathcal{K}\}$. Assume first that (i) holds; then the restriction of A_P to $L[r_\bullet]$, namely $\pi_P \circ \Delta|_{L[r_\bullet]}$, has the NIP. Since $L[r_\bullet]$ is dense in $\mathcal{L}[r_\bullet]$, Lemma 1.10 shows that the set A_P itself has the NIP, and so does the restriction of A_P to every compact subset of $\mathcal{L}[r_\bullet]$. Theorem 5.5 then gives the inclusion $\overline{A_P} \subseteq M_r(\mathcal{L}[r_\bullet])$, that is, every member of $\overline{A_P}$ is universally measurable. Since P is arbitrary, Proposition 5.4 shows that every member of $\mathcal{K}$ is universally measurable, and (ii) follows. Conversely, if (ii) holds, then every member of $\overline{A_P}$ is universally measurable by Proposition 5.4. By the implication (i)⇒(ii) of Theorem 5.5, applied to the compact set $\mathcal{L}[r_\bullet]$ itself, the set A_P has the NIP. Consequently, so does its restriction $\pi_P \circ \Delta|_{L[r_\bullet]}$, and (i) holds. Finally, let $f \in \overline{\Delta}$, and let $\tilde{f}$ be the extension of Lemma 2.7. For every $r_\bullet \in \mathscr{R}$, the restriction $\tilde{f}|_{\mathcal{L}[r_\bullet]}$ belongs to $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$, and it is therefore universally measurable when (ii) holds. Since $\tilde{f}$ maps $\mathcal{L}[r_\bullet]$ into itself, the deep computation f satisfies Definition 5.6 with $s_\bullet = r_\bullet$ for every $r_\bullet \in \mathscr{R}$. □

Remark 5.8. Condition (ii) of Theorem 5.7 concerns all the members of the sets $\overline{\tilde{\Delta}|_{\mathcal{L}[r_\bullet]}}$, that is, the restrictions of ultracomputations to the type shards. By contrast, the equivalence of the theorem fails if (ii) is weakened to the universal Monte Carlo computability of every deep computation, which requires only that some extension be universally measurable. To see this, let $L = \mathbb{Q}/\mathbb{Z} \subseteq \mathbb{T}$ carry the predicates $\cos 2\pi x$ and $\sin 2\pi x$, and let Γ consist of the iterates of the doubling map $x \mapsto 2x$. The space of types is the circle $\mathbb{T}$, and each computation extends to the continuous map $x \mapsto 2^n x$ of $\mathbb{T}$. Thus the Extendibility Axiom holds, and the single sizer with both bounds equal to 1 forms an exhaustive collection that confines Γ. By the paragraph that follows Proposition 4.20, the iterates of the doubling map of $\mathbb{T}$ do not have the NIP; by Lemma 1.10, neither do their restrictions to the dense set L. Every deep computation $f \colon L \to \mathbb{T}$, however, extends to the map that agrees with f on L and takes the value 0 on $\mathbb{T} \setminus L$. Since L is countable, this extension is Borel, whence every deep computation is universally Monte Carlo computable.

5.2. **Talagrand stability and Monte Carlo computability of deep computations.** The second route to universal Monte Carlo computability starts from

Talagrand stability, a notion that Talagrand introduced in his study of Pettis integration [Tal84] and that is closely related to the NIP. We follow this route here and compare the two routes at the end of the subsection. Let X be a compact Hausdorff space, let $A \subseteq \mathbb{R}^X$ be a set of functions, and let μ be a Radon probability measure on X. For a μ-measurable set $E \subseteq X$, a positive integer k and real numbers $a < b$, we write the points of E^{2k} as $x = (x_0, \dots, x_{2k-1})$ and define

$$D_k(A, E, a, b) = \bigcup_{f \in A} \{x \in E^{2k} : f(x_{2i}) \leq a,\ f(x_{2i+1}) \geq b \ \text{ for all } i < k\}.$$

Definition 5.9. The set A is *Talagrand μ-stable* if, for every μ-measurable set $E \subseteq X$ of positive measure and all reals $a < b$, there is an integer $k \geq 1$ such that

$$(\mu^{2k})^*(D_k(A, E, a, b)) < (\mu(E))^{2k}.$$

Here $(\mu^{2k})^*$ denotes the outer measure associated with the product measure μ^{2k} on X^{2k}. The outer measure is needed because $D_k(A, E, a, b)$ need not be measurable. The set $D_k(A, E, a, b)$ is measurable when A is a countable set of μ-measurable functions, since $D_k(A, E, a, b)$ is then a countable union of measurable sets.

Remarks 5.10. Let X, μ, $A \subseteq \mathbb{R}^X$ and $E \subseteq X$ be as above, and let $k \geq 1$ be an integer and $a < b$ be reals. The following hold.

(1) If $A \subseteq B \subseteq \mathbb{R}^X$, then

$$D_k(A, E, a, b) \subseteq D_k(B, E, a, b).$$

(2) If $a < a' < b' < b$, then

$$D_k(\overline{A}, E, a, b) \subseteq D_k(A, E, a', b').$$

[*Proof*: If $x \in D_k(\overline{A}, E, a, b)$, then there is $f \in \overline{A}$ such that $f(x_{2i}) \leq a < a'$ and $f(x_{2i+1}) \geq b > b'$ for all $i < k$. By the definition of the topology of pointwise convergence, there is $g \in A$ such that $g(x_{2i}) < a'$ and $g(x_{2i+1}) > b'$ for all $i < k$; hence $x \in D_k(A, E, a', b')$.]

(3) Every subset of a Talagrand μ-stable set is Talagrand μ-stable, by (1).

(4) If A is Talagrand μ-stable, then so is $\overline{A}$, by (2).

In Corollary 5.15 below we consider a confined set Δ of computations such that, for every feature P and every type shard $\mathcal{L}[r_\bullet]$, the set $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$ is universally Talagrand stable, that is, Talagrand μ-stable for every Radon probability measure μ on the shard (see Definition 5.13). We show there that the deep computations of every such set are universally Monte Carlo computable. The proof of the corollary rests on Lemma 5.12, by which the pointwise closure of a Talagrand μ-stable set consists of μ-measurable functions. The lemma, in turn, rests on a characterization of measurable functions (compare Fact 1.2), for which we recall four properties of a measure space (X, Σ, μ). The space is *complete* if every subset of a set of measure zero belongs to Σ, and *semi-finite* if every $E \in \Sigma$ with $\mu(E) = \infty$ contains some $F \in \Sigma$ with $0 < \mu(F) < \infty$. The space is *saturated* if a set $E \subseteq X$ belongs to Σ whenever $E \cap F \in \Sigma$ for all $F \in \Sigma$ with $\mu(F) < \infty$. Finally, the space is *inner regular* with respect to a collection $\mathcal{K} \subseteq \Sigma$ if $\mu(E) = \sup\{\mu(K) : K \in \mathcal{K} \text{ and } K \subseteq E\}$ for every $E \in \Sigma$. For a function $f \colon X \to \mathbb{R}$, a set $K \subseteq X$ and reals $a < b$, we write $K_{\leq a} = \{x \in K : f(x) \leq a\}$ and $K_{\geq b} = \{x \in K : f(x) \geq b\}$.

Fact 5.11 ([Fre03, Lemma 413G])**.** *Let (X, Σ, μ) be a complete, semi-finite and saturated measure space that is inner regular with respect to a collection $\mathcal{K} \subseteq \Sigma$. A function $f\colon X \to \mathbb{R}$ is measurable if and only if, for every $K \in \mathcal{K}$ with $0 < \mu(K) < \infty$ and all reals $a < b$, either $\mu^*(K_{\leq a}) < \mu(K)$ or $\mu^*(K_{\geq b}) < \mu(K)$.*

For instance, let X be a compact Hausdorff space, and let μ be a Radon probability measure on X. If Σ is the completion by μ of the Borel σ-algebra of X and $\mathcal{K}$ is the collection of compact subsets of X, then all four conditions of Fact 5.11 hold.

We first show that Talagrand stability passes to the pointwise closure and forces measurability.

Lemma 5.12. *Let X be a compact Hausdorff space, let μ be a Radon probability measure on X, and let $A \subseteq \mathbb{R}^X$ be a set of functions. If A is Talagrand μ-stable, then $\overline{A}$ is also Talagrand μ-stable and $\overline{A} \subseteq \mathscr{M}^0(X, \mu)$.*

Proof. By Remarks 5.10, the closure $\overline{A}$ is Talagrand μ-stable, and hence so is $\{f\}$ for every $f \in \overline{A}$. It therefore suffices to show that if $\{f\}$ is Talagrand μ-stable, then f is μ-measurable. Suppose that f is not μ-measurable. Then Fact 5.11 gives a compact set E of positive measure and reals $a < b$ with $\mu^*(E_{\leq a}) = \mu^*(E_{\geq b}) = \mu(E)$, where $E_{\leq a} = \{x \in E : f(x) \leq a\}$ and $E_{\geq b} = \{x \in E : f(x) \geq b\}$. For every $k \geq 1$ we have $(E_{\leq a} \times E_{\geq b})^k \subseteq D_k(\{f\}, E, a, b)$, and this product has outer measure $(\mu(E))^{2k}$ (see [Tal84]); hence $\{f\}$ is not Talagrand μ-stable. □

Definition 5.13. Let X be a compact Hausdorff space. A set $A \subseteq \mathbb{R}^X$ is *universally Talagrand stable* if it is Talagrand μ-stable for every Radon probability measure μ on X.

For pointwise bounded sets of continuous functions on a compact space, universal Talagrand stability implies the NIP.

Proposition 5.14. *Let X be a compact Hausdorff space, and let $A \subseteq \mathrm{C_p}(X)$ be pointwise bounded. If A is universally Talagrand stable, then A satisfies the NIP.*

Proof. By Lemma 5.12, the closure $\overline{A}$ is contained in $\mathscr{M}^0(X, \mu)$ for every Radon probability measure μ on X, that is, $\overline{A} \subseteq M_r(X)$. By the implication (i)⇒(ii) of Theorem 5.5, applied with $K = X$, the set A satisfies the NIP. □

Corollary 5.15. *Let $(L, \mathcal{P}, \Gamma)$ be a CCS satisfying the Extendibility Axiom, let $\mathscr{R}$ be an exhaustive collection of sizers, and let $\Delta \subseteq \Gamma$ be $\mathscr{R}$-confined. If $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$ is universally Talagrand stable for all $P \in \mathcal{P}$ and all $r_\bullet \in \mathscr{R}$, then every deep computation in $\overline{\Delta}$ is universally Monte Carlo computable on $\mathscr{R}$.*

Proof. Let $P \in \mathcal{P}$ and $r_\bullet \in \mathscr{R}$ be given. By the Extendibility Axiom and confinement, the set $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$ is a pointwise bounded family of continuous functions on the compact space $\mathcal{L}[r_\bullet]$. Proposition 5.14 therefore shows that this set has the NIP, and hence so does its restriction $\pi_P \circ \Delta|_{L[r_\bullet]}$. Theorem 5.7 now completes the proof. □

In the context of deep computations we have thus found two routes to universal Monte Carlo computability, the NIP and universal Talagrand stability, and the second implies the first (see Proposition 5.14). For countable sets of *continuous* functions on a compact space, the converse also holds: the NIP implies universal

Talagrand stability (see Theorem 5.16 below). Consequently, since the Extendibility Axiom makes computations continuous on the type shards, the two routes to universal Monte Carlo computability coincide for countable sets of computations. After a reduction to a compact metrizable space, and after composition with an increasing homeomorphism of $\mathbb{R}$ onto a bounded interval, which makes the family uniformly bounded and preserves both the NIP and Talagrand stability, this implication for countable sets of continuous functions can be derived from two results on the Glivenko–Cantelli property of a family of functions. The Glivenko–Cantelli property is the almost sure convergence of empirical means to expectations, uniformly over the family. The first result is a theorem of van Handel [vH13, Theorem 1.3], which concerns separable, uniformly bounded families of measurable functions on a standard Borel space. According to this theorem, such a family has the Glivenko–Cantelli property for every probability measure if and only if the family contains no Boolean σ-independent sequence. The second result is a theorem of Talagrand [Tal87], which identifies, among uniformly bounded families, the Glivenko–Cantelli families for a measure μ with the Talagrand μ-stable ones (see also [Fre03, §465]). We give instead a direct proof, which reduces the question to a compact metrizable space and then applies Fact 1.2 to a cluster point of the reduced family.

Theorem 5.16. *Let X be a compact Hausdorff space, and let $A \subseteq \mathrm{C_p}(X)$ be countable and pointwise bounded. If A satisfies the NIP, then A is universally Talagrand stable.*

Proof. Suppose that A satisfies the NIP, and let μ be a Radon probability measure on X. Towards a contradiction, suppose further that A is not Talagrand μ-stable. Then there are a μ-measurable set $E \subseteq X$ of positive measure and reals $a < b$ such that $\mu^{2k}(D_k(A, E, a, b)) = (\mu(E))^{2k}$ for every $k \geq 1$; the sets $D_k(A, E, a, b)$ are measurable, since A is a countable set of continuous functions. We first transfer the situation to a compact metrizable space. The map $\Psi\colon X \to \mathbb{R}^A$ given by $\Psi(x) = (f(x))_{f\in A}$ is continuous, and the image $Y = \Psi(X)$ is compact and metrizable because A is countable. Moreover, each $f \in A$ factors as $f = \hat{f}\circ\Psi$, where $\hat{f}\colon Y \to \mathbb{R}$ is the restriction of the coordinate projection. The set $\hat{A} = \{\hat{f} : f \in A\} \subseteq \mathrm{C_p}(Y)$ is pointwise bounded, and $\hat{A}$ satisfies the NIP, because $Y^{\hat{f}_I}_{\leq a'} \cap Y^{\hat{f}_J}_{\geq b'} = \Psi(X^{f_I}_{\leq a'} \cap X^{f_J}_{\geq b'})$ for every sequence $(f_n)_{n\in\mathbb{N}}$ in A, all finite sets $I, J \subseteq \mathbb{N}$ and all reals $a' < b'$. Let ν be the Borel probability measure on Y given by $\nu(B) = \mu(E \cap \Psi^{-1}(B))/\mu(E)$. Since $D_k(A, E, a, b) = E^{2k} \cap (\Psi^{2k})^{-1}(D_k(\hat{A}, Y, a, b))$, where Ψ^{2k} applies Ψ to each coordinate, we obtain $\nu^{2k}(D_k(\hat{A}, Y, a, b)) = 1$ for every $k \geq 1$.

Now consider the product measure $\nu^{\mathbb{N}}$ on $Y^{\mathbb{N}}$, under which the coordinates y_n of a sequence $(y_n)_{n\in\mathbb{N}}$ are independent, each with distribution ν. Let Z denote the support of ν, which is a nonempty closed, and hence Polish, subspace of Y. For each $k \geq 1$, almost every sequence admits some $\hat{f} \in \hat{A}$ with $\hat{f}(y_{2i}) \leq a$ and $\hat{f}(y_{2i+1}) \geq b$ for all $i < k$, since this event has probability $\nu^{2k}(D_k(\hat{A}, Y, a, b)) = 1$. Moreover, since Y has a countable base and every basic open set that meets Z has positive measure, both $\{y_{2i} : i \in \mathbb{N}\}$ and $\{y_{2i+1} : i \in \mathbb{N}\}$ are dense subsets of Z for almost every sequence. Fix a sequence (y_n) with all these properties, and, for each $k \geq 1$, choose $\hat{f}_k \in \hat{A}$ with $\hat{f}_k(y_{2i}) \leq a$ and $\hat{f}_k(y_{2i+1}) \geq b$ for all $i < k$. Since $\hat{A}$ is pointwise bounded, the sequence $(\hat{f}_k)$ has a cluster point g in $\mathbb{R}^Y$, and $g \in \mathrm{B}_1(Y)$ by Theorem 1.6. For every i, the inequalities $\hat{f}_k(y_{2i}) \leq a$ and $\hat{f}_k(y_{2i+1}) \geq b$ hold for

all $k > i$, and therefore $g(y_{2i}) \leq a$ and $g(y_{2i+1}) \geq b$. By condition (3) of Fact 1.2, applied to the Polish space Y and its closed subset Z, the restriction $g|_Z$ has a point of continuity z. Every neighborhood of z in Z, however, contains points y_{2i} with $g(y_{2i}) \leq a$ and points y_{2i+1} with $g(y_{2i+1}) \geq b$; this contradicts the continuity of $g|_Z$ at z. □

Corollary 5.17. *Let $(L, \mathcal{P}, \Gamma)$ be a CCS satisfying the Extendibility Axiom, let $\mathscr{R}$ be an exhaustive collection of sizers, and let $\Delta \subseteq \Gamma$ be countable and $\mathscr{R}$-confined. For all $P \in \mathcal{P}$ and $r_\bullet \in \mathscr{R}$, the set $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$ is universally Talagrand stable if and only if $\pi_P \circ \Delta|_{L[r_\bullet]}$ satisfies the NIP.*

Proof. By the Extendibility Axiom and confinement, the set $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$ is a countable, pointwise bounded family of continuous functions on the compact space $\mathcal{L}[r_\bullet]$, in which $L[r_\bullet]$ is dense. If this set is universally Talagrand stable, it has the NIP by Proposition 5.14, and so does its restriction to $L[r_\bullet]$. Conversely, suppose that $\pi_P \circ \Delta|_{L[r_\bullet]}$ has the NIP. By Lemma 1.10, so does the set $\pi_P \circ \tilde{\Delta}|_{\mathcal{L}[r_\bullet]}$, to which Theorem 5.16 therefore applies. □

6. Open questions

We close with the questions that remain open, in the order in which they arise from the sections of the paper. The first concerns the hierarchy of Section 3.2, whose levels were defined through the topology of the space of deep computations. Proposition 3.4 describes these levels by the number of tests needed to tell deep computations apart, but the descriptions so obtained still involve infinitely many tests.

Question 6.1. Is there a finitary, combinatorial characterization of $\mathrm{NIP}^{\mathrm{fc}}$, $\mathrm{NIP}^{\mathrm{hs}}$ and $\mathrm{NIP}^{\mathrm{met}}$, in the spirit of the characterization of NIP by finite patterns in Theorem 3.1?

In Section 4 we place a single iterated map at every level of the hierarchy except one, and at three of the five leaves of Figure 1; Figure 12 marks the two leaves that remain open.

The exception is the level NIP but not $\mathrm{NIP}^{\mathrm{fc}}$, at which both of our examples are actions of groups with two generators (see Section 4.5).

Question 6.2. Is there a cascade CCS that is NIP but not $\mathrm{NIP}^{\mathrm{fc}}$? Equivalently, is there a continuous map γ of a compact metrizable space L such that (L, γ) is tame and $E(L, \gamma)$ is not first countable?

For actions of $\mathbb{Z}$, and more generally of abelian groups, the analogous question is Question 6.8 of Glasner and Megrelishvili [GM22].

At the level $\mathrm{NIP}^{\mathrm{fc}}$ but not $\mathrm{NIP}^{\mathrm{hs}}$, the deep computations of the three-dimensional Bresenham algorithm contain a copy of $\hat{D}(S(2^{\mathbb{N}}))$ and none of $D(2^{\mathbb{N}})$ (see Theorem 4.10). By contrast, the countable family $\Delta_{\mathcal{C}}$ of elements of the group generated by the steps of the Euclidean algorithm realizes the other alternative of Todorčević's trichotomy (see Proposition 4.15).

Question 6.3. Is there a cascade CCS that is $\mathrm{NIP}^{\mathrm{fc}}$ but not $\mathrm{NIP}^{\mathrm{hs}}$ and whose space of deep computations contains a copy of $D(2^{\mathbb{N}})$?




FIGURE 12. Leaves realized by a single iterated map; dashed leaves are open.

APPENDIX A. THE THUE–MORSE SYSTEM IS NOT TAME

In this appendix we prove that the shift on the Thue–Morse subshift generates an action of $\mathbb{Z}$ that is not tame, a fact observed by Glasner and Megrelishvili [GM18]. From this fact we derive that the two dynamical CCSs defined by this shift, which we use in Section 4.7 as examples beyond the hierarchy, are not NIP. Our proof is self-contained except for the structure theorem for tame minimal systems under actions of abelian groups, due to Huang, to Kerr and Li, and to Glasner. We recall the theorem below and combine the theorem with an elementary computation of the correlations of the Thue–Morse sequence.

We first recall the objects involved. The Thue–Morse sequence $(t_n)_{n\in\mathbb{N}}$ is defined by letting t_n be the parity of the number of ones in the binary expansion of n, whence $t_{2k} = t_k$ and $t_{2k+1} = 1 - t_k$ for every $k \in \mathbb{N}$. The same sequence is the limit of the words W_m, for $m \in \mathbb{N}$, obtained from 0 by m applications of the substitution $0 \mapsto 01$, $1 \mapsto 10$. The two-sided sequences all of whose finite blocks occur in (t_n) form the Thue–Morse subshift $X_{TM} \subseteq 2^{\mathbb{Z}}$, a closed subset of the compact metrizable space $2^{\mathbb{Z}}$. On this subshift the shift σ, defined by $\sigma(x)_n = x_{n+1}$, is a homeomorphism. As in Section 4.1, a system (L, Γ) consists of a compact metrizable space L and a countable semigroup Γ of continuous maps $L \to L$. Such a system is *tame*, in the sense of Section 4.1, if for every continuous function $f\colon L \to \mathbb{R}$ the family $\{f \circ \gamma : \gamma \in \Gamma\}$ has the NIP of Definition 1.5. We call a sequence $(f \circ \gamma_k)_k$ of members of the family *independent* if, for some reals $a < b$, the sequence witnesses the finitary independence property of Definition 1.5. Hence the system (L, Γ) is

tame exactly when no independent sequence exists. The CCS with states X_{TM}, the coordinate predicates $x \mapsto x_n$ for $n \in \mathbb{Z}$, and the group $\{\sigma^n : n \in \mathbb{Z}\}$ is a dynamical CCS in the sense of Section 4.1. The cascade CCS of σ, in turn, is the dynamical CCS with the same states and predicates but with the semigroup $\{\sigma^n : n \in \mathbb{N}\}$. By Proposition 4.1, each of these two CCSs is NIP if and only if its dynamical system is tame.

That the dynamics of σ on X_{TM} is wild follows from the structure theory of tame minimal systems under actions of abelian groups. Huang [Hua06] and, independently, Kerr and Li [KL07] proved that every metrizable minimal tame system under an action of an abelian group is uniquely ergodic. Unique ergodicity means that the system carries exactly one invariant Borel probability measure. The same authors proved, in addition, that such a system is an almost one-to-one extension of the maximal equicontinuous factor of the system. Glasner [Gla07] showed, moreover, that the factor map onto this factor is an isomorphism of measure-preserving systems. For a minimal homeomorphism, the maximal equicontinuous factor is a rotation $y \mapsto y + \alpha$ of a compact abelian group Y in which the multiples of α are dense, equipped with the Haar measure of Y. The characters of Y are eigenfunctions of this rotation and span a dense subspace of the space of square-integrable functions on Y. Consider now a minimal system given by a homeomorphism σ that generates a tame action. The Koopman operator of this system sends each function that is square-integrable with respect to the invariant measure to the composite of the function with σ. By the results just recalled, every such function is a limit in mean square of linear combinations of eigenfunctions of the Koopman operator. The Thue–Morse subshift fails this test, because the correlations of the Thue–Morse sequence are, on average, too small.

Proposition A.1. *Let $X_{TM} \subseteq 2^{\mathbb{Z}}$ be the Thue–Morse subshift, and let σ be the shift on X_{TM}. The action of $\mathbb{Z}$ on X_{TM} generated by σ is not tame. Consequently, neither the CCS with states X_{TM}, the coordinate predicates and the group $\{\sigma^n : n \in \mathbb{Z}\}$ nor the cascade CCS of σ is NIP.*

Proof. We first show that X_{TM} is minimal. Every finite block w of (t_n) occurs in some word W_m. If $\overline{W}_m$ denotes the word obtained from 1 by m applications of the substitution, then the words $W_{m+1} = W_m\overline{W}_m$ and $\overline{W}_{m+1} = \overline{W}_m W_m$ both contain W_m. Moreover, since $t_{2k} = t_k$ and $t_{2k+1} = 1 - t_k$, the block of (t_n) of length 2^{m+1} that begins at the index $2^{m+1}j$ is W_{m+1} or $\overline{W}_{m+1}$ according as t_j is 0 or 1. Hence the block w occurs in every block of length 2^{m+2} of (t_n), and therefore in every point of X_{TM}; every orbit of σ is thus dense in X_{TM}.

Suppose now that the action of $\mathbb{Z}$ generated by σ is tame. By the structure theorem recalled above, the subshift X_{TM} carries a unique σ-invariant Borel probability measure μ. The eigenfunctions of the unitary operator V on $L^2(\mu)$ that sends each function to the composite of the function with σ span, by the same theorem, a dense subspace of $L^2(\mu)$. Let $\vartheta(x) = (-1)^{x_0}$ for $x \in X_{TM}$, and let $c(n) = \int \vartheta \cdot (\vartheta \circ \sigma^n)\, d\mu$ for $n \in \mathbb{N}$. By minimality and compactness, the subshift X_{TM} contains a point x with $x_j = t_j$ for every $j \geq 0$. Every weak* limit of the measures $\frac{1}{K}\sum_{k<K} \delta_{\sigma^k x}$ is σ-invariant and hence equal to μ, whence

$$c(n) = \lim_{K\to\infty} \frac{1}{K} \sum_{k<K} (-1)^{t_k + t_{k+n}}.$$

Splitting the sum according to the parity of k and using $t_{2k} = t_k$ and $t_{2k+1} = 1 - t_k$, we obtain $c(2n) = c(n)$ and $c(2n+1) = -\frac{1}{2}\big(c(n) + c(n+1)\big)$ for every $n \in \mathbb{N}$, and in particular $c(1) = -\frac{1}{3}$. For $K \geq 1$, let $A(K) = \sum_{n<K} c(n)^2$ and $B(K) = \sum_{n<K} c(n)c(n+1)$. For every $K \geq 1$, the recursions yield the identities and inequalities

$$B(2K) = -\tfrac{1}{2} \sum_{n<K} \big(c(n) + c(n+1)\big)^2 \leq 0,$$

$$A(2K) = A(K) + \tfrac{1}{4} \sum_{n<K} \big(c(n) + c(n+1)\big)^2 \leq \tfrac{3}{2}A(K) + \tfrac{1}{2}B(K),$$

where the last inequality follows by expanding the squares, because $c(0) = 1 \geq c(K)^2$. Hence we have $A(4K) \leq \frac{3}{2}A(2K)$ for every $K \geq 1$, and, since $A(2) = 1 + c(1)^2 \leq 2$, it follows that $A(2^k) \leq 2\big(\frac{3}{2}\big)^{k-1}$ for every $k \geq 1$. Since the function A is nondecreasing, we conclude that $A(K)/K \xrightarrow{K\to\infty} 0$ (see Figure 13). Now let λ be a complex number of modulus 1, and let Π_λ be the orthogonal projection of $L^2(\mu)$ onto the kernel of $V - \lambda I$. This kernel consists of the eigenfunctions of V with eigenvalue λ and the function 0. By von Neumann's mean ergodic theorem, applied to the unitary operator $\bar{\lambda}V$, the averages $\frac{1}{K}\sum_{n<K} \bar{\lambda}^n\, \vartheta \circ \sigma^n$ converge to $\Pi_\lambda \vartheta$ in $L^2(\mu)$. Consequently, since $c(n)$ is the inner product of $\vartheta \circ \sigma^n$ and ϑ, the Cauchy–Schwarz inequality yields

$$\|\Pi_\lambda \vartheta\|^2 = \lim_{K\to\infty} \Big|\frac{1}{K} \sum_{n<K} \bar{\lambda}^n c(n)\Big| \leq \lim_{K\to\infty} \Big(\frac{A(K)}{K}\Big)^{1/2} = 0.$$

Thus the function ϑ is orthogonal to every eigenfunction of V, and therefore to every function in $L^2(\mu)$, since these eigenfunctions span a dense subspace of $L^2(\mu)$. This conclusion is absurd, since $|\vartheta| = 1$. The action is therefore not tame, and, by Proposition 4.1, the CCS with the group $\{\sigma^n : n \in \mathbb{Z}\}$ is not NIP.




FIGURE 13. Thue–Morse correlations $c(n)$ (left) and the averages $A(K)/K$ (right).

For the cascade CCS of σ, we use a symmetry of the Thue–Morse sequence. The words W_{2m} are palindromes, and every finite block of (t_n) occurs in one of them. Consequently, the reversal ι of two-sided sequences, defined by $\iota(x)_n = x_{-n}$, maps X_{TM} onto itself and satisfies $\iota \circ \sigma = \sigma^{-1} \circ \iota$. Since the action of $\mathbb{Z}$ is not tame, there are a continuous function $f\colon X_{TM} \to \mathbb{R}$ and pairwise distinct integers n_k such that the sequence $(f \circ \sigma^{n_k})_k$ is independent. Moreover, infinitely many of the indices n_k

have the same sign. If these indices are positive, the corresponding terms already form an independent sequence in the cascade. If these indices are negative, the functions $f \circ \sigma^{n_k} \circ \iota = (f \circ \iota) \circ \sigma^{-n_k}$ form an independent sequence with positive exponents, because the reversal ι is a bijection of X_{TM}. In either case the cascade of σ is not tame, and, by Proposition 4.1, its CCS is not NIP. □

References

[ADIW26] Samson Alva, Eduardo Dueñez, José Iovino, and Claire Walton, *Approximability of deep computations*, Mathematical Structures in Computer Science (2026), To appear; arXiv:2409.06064.

[ADK08] Spiros A. Argyros, Pandelis Dodos, and Vassilis Kanellopoulos, *A classification of separable Rosenthal compacta and its applications*, Dissertationes Mathematicae **449** (2008), 1–55.

[Ark92] A. V. Arkhangel'skii, *Topological function spaces*, Mathematics and its Applications (Soviet Series), vol. 78, Kluwer Academic Publishers Group, Dordrecht, 1992, Translated from the Russian by R. A. M. Hoksbergen. MR 1144519

[Bai99] René Baire, *Sur les fonctions de variables réelles*, Annali di Matematica Pura ed Applicata (3) **3** (1899), 1–123.

[BC92] Louis S. Block and William A. Coppel, *Dynamics in one dimension*, Lecture Notes in Mathematics, vol. 1513, Springer, Berlin, 1992.

[BEHW89] Anselm Blumer, Andrzej Ehrenfeucht, David Haussler, and Manfred K. Warmuth, *Learnability and the Vapnik–Chervonenkis dimension*, J. Assoc. Comput. Mach. **36** (1989), no. 4, 929–965.

[BFB16] Richard L. Burden, J. Douglas Faires, and Annette M. Burden, *Numerical analysis*, 10th ed., Cengage Learning, Boston, MA, 2016.

[BFT78] J. Bourgain, D. H. Fremlin, and M. Talagrand, *Pointwise compact sets of Baire-measurable functions*, American Journal of Mathematics **100** (1978), no. 4, 845–886.

[BKK] Shaojie Bai, J. Zico Kolter, and Vladlen Koltun, *Deep equilibrium models*, Preprint. https://arxiv.org/abs/1909.01377, https://implicit-layers-tutorial.org/.

[Bre65] Jack E. Bresenham, *Algorithm for computer control of a digital plotter*, IBM Systems Journal **4** (1965), no. 1, 25–30.

[CRBD] Ricky T. Q. Chen, Yulia Rubanova, Jesse Bettencourt, and David Duvenaud, *Neural ordinary differential equations*, Preprint. https://arxiv.org/abs/1806.07366.

[Deb13] Gabriel Debs, *Descriptive aspects of Rosenthal compacta*, Recent Progress in General Topology III, Springer, 2013, pp. 205–227.

[dMvS93] Welington de Melo and Sebastian van Strien, *One-dimensional dynamics*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), vol. 25, Springer, Berlin, 1993.

[DnI17] Eduardo Dueñez and José N. Iovino, *Model theory and metric convergence I: Metastability and dominated convergence*, Beyond First Order Model Theory, Volume I, CRC Press, Boca Raton, FL, 2017, pp. 131–187. MR 3729326

[Ell93] Robert Ellis, *The enveloping semigroup of projective flows*, Ergodic Theory and Dynamical Systems **13** (1993), no. 4, 635–660.

[Eng89] Ryszard Engelking, *General topology*, second ed., Sigma Series in Pure Mathematics, vol. 6, Heldermann Verlag, Berlin, 1989, Translated from the Polish by the author. MR MR1039321 (91c:54001)

[Fre03] David H. Fremlin, *Measure theory, volume 4: Topological measure spaces*, Torres Fremlin, Colchester, UK, 2003.

[Gal00] A. Galántai, *The theory of Newton's method*, Journal of Computational and Applied Mathematics **124** (2000), no. 1, 25–44, Numerical Analysis 2000. Vol. IV: Optimization and Nonlinear Equations.

[Gla06] Eli Glasner, *On tame dynamical systems*, Colloquium Mathematicum **105** (2006), no. 2, 283–295.

[Gla07] ———, *The structure of tame minimal dynamical systems*, Ergodic Theory and Dynamical Systems **27** (2007), no. 6, 1819–1837.

[GM18] Eli Glasner and Michael Megrelishvili, *More on tame dynamical systems*, Ergodic theory and dynamical systems in their interactions with arithmetics and combinatorics, Lecture Notes in Mathematics, vol. 2213, Springer, Cham, 2018, pp. 351–392. MR 3821724

[GM22] ———, *Todorčević' trichotomy and a hierarchy in the class of tame dynamical systems*, Transactions of the American Mathematical Society **375** (2022), no. 7, 4513–4548.

[God80] Gilles Godefroy, *Compacts de Rosenthal*, Pacific J. Math. **91** (1980), no. 2, 293–306. MR 615679

[Gro52] A. Grothendieck, *Critères de compacité dans les espaces fonctionnels généraux*, Amer. J. Math. **74** (1952), 168–186. MR 47313

[Hua06] Wen Huang, *Tame systems and scrambled pairs under an abelian group action*, Ergodic Theory and Dynamical Systems **26** (2006), no. 5, 1549–1567.

[Kec95] Alexander S. Kechris, *Classical descriptive set theory*, Graduate Texts in Mathematics, vol. 156, Springer-Verlag, 1995.

[Kei23] H. Jerome Keisler, *Model theory for real-valued structures*, Beyond First Order Model Theory, Volume II (José Iovino, ed.), CRC Press, Boca Raton, FL, 2023.

[KL07] David Kerr and Hanfeng Li, *Independence in topological and C^*-dynamics*, Mathematische Annalen **338** (2007), no. 4, 869–926.

[KM81] Jean-Louis Krivine and Bernard Maurey, *Espaces de Banach stables*, Israel J. Math. **39** (1981), no. 4, 273–295. MR 83a:46030

[Köh95] Angela Köhler, *Enveloping semigroups for flows*, Proceedings of the Royal Irish Academy. Section A **95A** (1995), no. 2, 179–191.

[Kri72] J.-L. Krivine, *Théorie des modèles et espaces L^p*, C. R. Acad. Sci. Paris Sér. A-B **275** (1972), A1207–A1210. MR 46 #7879

[Kri74] ———, *Langages à valeurs réelles et applications*, Fund. Math. **81** (1974), 213–253, Collection of articles dedicated to Andrzej Mostowski on the occasion of his sixtieth birthday, III. MR 50 #1873

[Leb04] Henri Lebesgue, *Une propriété caractéristique des fonctions de classe 1*, Bulletin de la Société Mathématique de France **32** (1904), 229–242.

[Lot02] M. Lothaire, *Algebraic combinatorics on words*, Encyclopedia of Mathematics and its Applications, vol. 90, Cambridge University Press, Cambridge, 2002.

[MH40] Marston Morse and Gustav A. Hedlund, *Symbolic dynamics II. Sturmian trajectories*, American Journal of Mathematics **62** (1940), 1–42.

[Mil06] John Milnor, *Dynamics in one complex variable*, third ed., Annals of Mathematics Studies, vol. 160, Princeton University Press, Princeton, NJ, 2006.

[MT88] John Milnor and William Thurston, *On iterated maps of the interval*, Dynamical Systems (College Park, MD, 1986–87), Lecture Notes in Mathematics, vol. 1342, Springer, Berlin, 1988, pp. 465–563.

[Pap02] E. Pap (ed.), *Handbook of measure theory. Vol. I, II*, North-Holland, Amsterdam, 2002. MR 1953489

[Ros74] Haskell P. Rosenthal, *A characterization of Banach spaces containing l^1*, Proc. Nat. Acad. Sci. U.S.A. **71** (1974), 2411–2413. MR 358307

[Ros77] ———, *Point-wise compact subsets of the first Baire class*, Amer. J. Math. **99** (1977), no. 2, 362–378.

[Sau72] N. Sauer, *On the density of families of sets*, J. Combinatorial Theory Ser. A **13** (1972), 145–147. MR 307902

[She71] Saharon Shelah, *Stability, the f.c.p., and superstability; model theoretic properties of formulas in first order theory*, Ann. Math. Logic **3** (1971), no. 3, 271–362. MR 317926

[She72] ———, *A combinatorial problem; stability and order for models and theories in infinitary languages*, Pacific J. Math. **41** (1972), 247–261. MR 307903

[She90] ———, *Classification theory and the number of nonisomorphic models*, second ed., Studies in Logic and the Foundations of Mathematics, vol. 92, North-Holland Publishing Co., Amsterdam, 1990.

[Sim15] Pierre Simon, *A guide to nip theories*, Lecture Notes in Logic, vol. 44, Cambridge University Press, 2015.

[SSBD14] Shai Shalev-Shwartz and Shai Ben-David, *Understanding machine learning: From theory to algorithms*, Cambridge University Press, Cambridge, 2014.

[Tal84] Michel Talagrand, *Pettis integral and measure theory*, Memoirs of the American Mathematical Society, vol. 51, American Mathematical Society, Providence, RI, USA, 1984, Includes bibliography (pp. 220–224) and index.

[Tal87] ———, *The Glivenko-Cantelli problem*, The Annals of Probability **15** (1987), no. 3, 837–870.

[Tka11] Vladimir V. Tkachuk, *A C_p-theory problem book: Topological and function spaces*, Problem Books in Mathematics, Springer, 2011.

[Tod97] Stevo Todorčević, *Topics in topology*, Lecture Notes in Mathematics, vol. 1652, Springer Berlin, Heidelberg, 1997.

[Tod99] Stevo Todorčević, *Compact subsets of the first Baire class*, Journal of the American Mathematical Society **12** (1999), no. 4, 1179–1212.

[Val84] Leslie G. Valiant, *A theory of the learnable*, Communications of the ACM **27** (1984), no. 11, 1134–1142.

[VC68] Vladimir N. Vapnik and Alexey Ya. Chervonenkis, *The uniform convergence of frequencies of the appearance of events to their probabilities*, Dokl. Akad. Nauk SSSR **181** (1968), no. 4, 781–783, In Russian; English translation in Soviet Math. Dokl. **9** (1968), 915–918.

[VC71] Vladimir N Vapnik and Alexey Ya Chervonenkis, *On the uniform convergence of relative frequencies of events to their probabilities*, Theory of Probability & Its Applications **16** (1971), no. 2, 264–280.

[VC74] ———, *Theory of pattern recognition*, Nauka, Moscow, 1974, German Translation: Theorie der Zeichenerkennung, Akademie-Verlag, Berlin, 1979.

[vH13] Ramon van Handel, *The universal Glivenko–Cantelli property*, Probability Theory and Related Fields **155** (2013), no. 3–4, 911–934, arXiv:1009.4434.