


# FACT OF THE DAY



Today, October 6, Richard Dedekind  
would have turned 191 years old

Slides available at:  
[tonellicueto.xyz/pdf/8IMM\\_slides.pdf](http://tonellicueto.xyz/pdf/8IMM_slides.pdf)

Why

does the DESCARTES Solver work?

Alperen A.  
ERGÜR



The University of Texas at San Antonio™

Josué  
TONELLI-CUETO

Elias  
TSIGARIDAS  
*Inria*





Photo while working on this project

# Real Root Isolation I: The Problem

# Real Root Isolation I: The Problem

INPUT:

# Real Root Isolation I: The Problem

INPUT:

$$g \in \mathbb{Z}[x]$$

# Real Root Isolation I: The Problem

INPUT:

$$g \in \mathbb{Z}[x]$$

OUTPUT:

# Real Root Isolation I: The Problem

INPUT:

$$g \in \mathbb{Z}[x]$$

OUTPUT:

Intervals  $J_1, \dots, J_k$  s.t.

# Real Root Isolation I: The Problem

INPUT:

$$g \in \mathbb{Z}[x]$$

OUTPUT:

Intervals  $J_1, \dots, J_k$  s.t.

0)  $J_i = (a_i, b_i)$  with  $a_i, b_i \in \mathbb{Q}$

# Real Root Isolation I: The Problem

INPUT:

$$g \in \mathbb{Z}[x]$$

OUTPUT:

Intervals  $J_1, \dots, J_k$  s.t.

0)  $J_i = (a_i, b_i)$  with  $a_i, b_i \in \mathbb{Q}$

1)  $Z(g) \cap \mathbb{R} \subseteq \bigcup_{i=1}^k J_i$

# Real Root Isolation I: The Problem

INPUT:

$$g \in \mathbb{Z}[x]$$

OUTPUT:

Intervals  $J_1, \dots, J_k$  s.t.

0)  $J_i = (a_i, b_i)$  with  $a_i, b_i \in \mathbb{Q}$

1)  $Z(g) \cap \mathbb{R} \subseteq \bigcup_{i=1}^k J_i$

2)  $\forall i, \# Z(g) \cap J_i = 1$

# Real Root Isolation I: The Problem

INPUT:

$$g \in \mathbb{Z}[x]$$

OUTPUT:

Intervals  $J_1, \dots, J_k$  s.t.

0)  $J_i = (a_i, b_i)$  with  $a_i, b_i \in \mathbb{Q}$

1)  $Z(g) \cap \mathbb{R} \subseteq \bigcup_{i=1}^k J_i$

2)  $\forall i, \# Z(g) \cap J_i = 1$

INPUT SIZE PARAMETERS:

# Real Root Isolation I: The Problem

INPUT:

$$g \in \mathbb{Z}[x]$$

OUTPUT:

Intervals  $J_1, \dots, J_k$  s.t.

0)  $J_i = (a_i, b_i)$  with  $a_i, b_i \in \mathbb{Q}$

1)  $Z(g) \cap \mathbb{R} \subseteq \bigcup_{i=1}^k J_i$

2)  $\forall i, \# Z(g) \cap J_i = 1$

INPUT SIZE PARAMETERS:

$d$ : degree of  $g$

# Real Root Isolation I: The Problem

INPUT:

$$g \in \mathbb{Z}[x]$$

OUTPUT:

Intervals  $J_1, \dots, J_k$  s.t.

0)  $J_i = (a_i, b_i)$  with  $a_i, b_i \in \mathbb{Q}$

1)  $Z(g) \cap \mathbb{R} \subseteq \bigcup_{i=1}^k J_i$

2)  $\forall i, \# Z(g) \cap J_i = 1$

INPUT SIZE PARAMETERS:

$d$ : degree of  $g$

$\gamma$ : bit-size of coefficients of  $g$

# Real Root Isolation I: The Problem

INPUT:

$$g \in \mathbb{Z}[x]$$

OUTPUT:

Intervals  $J_1, \dots, J_k$  s.t.

0)  $J_i = (a_i, b_i)$  with  $a_i, b_i \in \mathbb{Q}$

1)  $Z(g) \cap \mathbb{R} \subseteq \bigcup_{i=1}^k J_i$

2)  $\forall i, \# Z(g) \cap J_i = 1$

INPUT SIZE PARAMETERS:

$d$ : degree of  $g$

$\gamma$ : bit-size of coefficients of  $g$

MEASURE OF RUN-TIME

# Real Root Isolation I: The Problem

INPUT:

$$g \in \mathbb{Z}[x]$$

OUTPUT:

Intervals  $J_1, \dots, J_k$  s.t.

0)  $J_i = (a_i, b_i)$  with  $a_i, b_i \in \mathbb{Q}$

1)  $Z(g) \cap \mathbb{R} \subseteq \bigcup_{i=1}^k J_i$

2)  $\forall i, \# Z(g) \cap J_i = 1$

INPUT SIZE PARAMETERS:

$d$ : degree of  $g$

$\gamma$ : bit-size of coefficients of  $g$

MEASURE OF RUN-TIME

Bit complexity

# Real Root Isolation II: The State of the Art

# Real Root Isolation II: The State of the Art

STURM SOLVER

$$\tilde{\mathcal{O}}_B(d^4 \gamma^2)$$

# Real Root Isolation II: The State of the Art

STURM SOLVER

$$\tilde{\mathcal{O}}_B(d^4 \gamma^2)$$

DESCARTES SOLVER

$$\tilde{\mathcal{O}}_B(d^4 \gamma^2)$$

# Real Root Isolation II: The State of the Art

STURM SOLVER

$$\tilde{\mathcal{O}}_B(d^4\gamma^2)$$

DESCARTES SOLVER

$$\tilde{\mathcal{O}}_B(d^4\gamma^2)$$

ANewDsc

$$\tilde{\mathcal{O}}_B(d^3 + d^2\gamma)$$

(Sagraloff & Mehlhorn; 2016)

# Real Root Isolation II: The State of the Art

STURM SOLVER

$$\tilde{O}_B(d^4 \gamma^2)$$

DESCARTES SOLVER

$$\tilde{O}_B(d^4 \gamma^2)$$

ANewDsc

$$\tilde{O}_B(d^3 + d^2 \gamma)$$

(Sagraloff & Mehlhorn; 2016)

PAN'S ALGORITHM

$$\tilde{O}_B(d^2 \gamma)$$

(Pan; 2002)

# Real Root Isolation II: The State of the Art

STURM SOLVER

$$\tilde{O}_B(d^4 \gamma^2)$$

DESCARTES SOLVER

$$\tilde{O}_B(d^4 \gamma^2)$$

ANewDsc

$$\tilde{O}_B(d^3 + d^2 \gamma)$$

(Sagraloff & Mehlhorn; 2016)

PAN'S ALGORITHM

$$\tilde{O}_B(d^2 \gamma)$$

(Pan; 2002)

Q: Can we beat the champion?

# Real Root Isolation III:

What do we wish?

# Real Root Isolation III:

What do we wish?

$$\tilde{G}_B(d\gamma)$$

# Real Root Isolation III:

What do we wish?

$$\tilde{O}_B(d\gamma)$$

We wish to find real roots  
almost as fast as we read the polynomial!

# DESCARTES SOLVER I:

## Rule of Signs

# DESCARTES SOLVER I:

## Rule of Signs

$V(g) := \#$  sign variations of  $g_0, g_1, \dots$

# DESCARTES SOLVER I:

## Rule of Signs

$V(g) := \# \text{ sign variations of } g_0, g_1, \dots$

THM (Descartes' rule of signs)

$$\#\mathcal{Z}(g, R) \leq V(g)$$



Portrait by Frans Hals  
Source: Wikimedia Commons

# DESCARTES SOLVER I:

## Rule of Signs

$V(g) := \# \text{ sign variations of } g_0, g_1, \dots$

THM (Descartes' rule of signs)

$$\#\mathcal{Z}(g, \mathbb{R}_+) \leq V(g)$$

Moreover,

$$V(g) \leq 1 \Rightarrow \text{Equality}$$



Portrait by Frans Hals  
Source: Wikimedia Commons

# DESCARTES SOLVER I:

## Rule of Signs

$V(g) := \# \text{ sign variations of } g_0, g_1, \dots$

THM (Descartes' rule of signs)

$$\#\mathcal{Z}(g, \mathbb{R}_+) \leq V(g)$$

Moreover,

$$V(g) \leq 1 \Rightarrow \text{Equality}$$

COR

$$\#\mathcal{Z}(g, (a, b)) \leq V(g, (a, b)) := V\left((x+1)^d g\left(\frac{bx+a}{x+1}\right)\right)$$



Portrait by Frans Hals  
Source: Wikimedia Commons

# DESCARTES SOLVER I:

## Rule of Signs

$V(g) := \# \text{sign variations of } g_0, g_1, \dots$

THM (Descartes' rule of signs)

$$\#\mathcal{Z}(g, \mathbb{R}_+) \leq V(g)$$

Moreover,

$$V(g) \leq 1 \Rightarrow \text{Equality}$$

COR

$$\#\mathcal{Z}(g, (a, b)) \leq V(g, (a, b)) := V\left((x+1)^d g\left(\frac{bx+a}{x+1}\right)\right)$$

$$(0, \infty) \xrightarrow{\text{bijection}} (a, b)$$



Portrait by Frans Hals  
Source: Wikimedia Commons

DESCARTES SOLVER II:

The Descartes' Oracle

# DESCARTES SOLVER II:

The Descartes' Oracle

1) Overcounting:

# DESCARTES SOLVER II:

The Descartes' Oracle

1) Overcounting:  $\#Z(g, J) \leq V(g, J)$

# DESCARTES SOLVER II:

The Descartes' Oracle

- 1) Overcounting:  $\#Z(g, J) \leq V(g, J)$
- 2) Exactness I:

# DESCARTES SOLVER II:

The Descartes' Oracle

- 1) Overcounting:  $\#Z(g, J) \leq V(g, J)$
- 2) Exactness I:  $V(g, J) \leq 1 \Rightarrow$  Equality

# DESCARTES SOLVER II:

The Descartes' Oracle

- 1) Overcounting:  $\#Z(g, J) \leq V(g, J)$
- 2) Exactness I:  $V(g, J) \leq 1 \Rightarrow$  Equality
- 3) Exactness II:

# DESCARTES SOLVER II:

The Descartes' Oracle

- 1) Overcounting:  $\#Z(g, J) \leq V(g, J)$
- 2) Exactness I:  $V(g, J) \leq 1 \Rightarrow$  Equality
- 3) Exactness II:

$$\#Z(g, D(m(J), c_w(J))) \leq K \Rightarrow V(g, J) \leq K$$

# DESCARTES SOLVER II:

The Descartes' Oracle

- 1) Overcounting:  $\#Z(g, J) \leq V(g, J)$
- 2) Exactness I:  $V(g, J) \leq 1 \Rightarrow$  Equality
- 3) Exactness II:

$$\#Z(g, D(m(J), c_w(J))) \leq K \Rightarrow V(g, J) \leq K$$

Obreshkoff's Thm: DESCARTES sees the complex roots around!

# DESCARTES SOLVER II:

The Descartes' Oracle

- 1) Overcounting:  $\#Z(g, J) \leq V(g, J)$
- 2) Exactness I:  $V(g, J) \leq 1 \Rightarrow$  Equality
- 3) Exactness II:

$$\#Z(g, D(m(J), c_w(J))) \leq K \Rightarrow V(g, J) \leq K$$

Obreshkoff's Thm: DESCARTES sees the complex roots around!

- 4) Subadditivity:

# DESCARTES SOLVER II:

The Descartes' Oracle

- 1) Overcounting:  $\#Z(g, J) \leq V(g, J)$
- 2) Exactness I:  $V(g, J) \leq 1 \Rightarrow$  Equality
- 3) Exactness II:

$$\#Z(g, D(m(J), c_w(J))) \leq K \Rightarrow V(g, J) \leq K$$

Obreshkoff's Thm: DESCARTES sees the complex roots around!

- 4) Subadditivity:

$$\bigcup_{i=1}^n J_i \subseteq J$$

# DESCARTES SOLVER II:

The Descartes' Oracle

- 1) Overcounting:  $\#Z(g, J) \leq V(g, J)$
- 2) Exactness I:  $V(g, J) \leq 1 \Rightarrow$  Equality

3) Exactness II:

$$\#Z(g, D(m(J), c_w(J))) \leq K \Rightarrow V(g, J) \leq K$$

Obreshkoff's Thm: DESCARTES sees the complex roots around!

4) Subadditivity:


$$\bigcup_{i=1}^n J_i \subseteq J \Rightarrow \sum V(g, J_i) \leq V(g, J)$$

DESCARTES SOLVER III:

The Algorithm


# DESCARTES SOLVER III:

The Algorithm




# DESCARTES SOLVER III:

## The Algorithm




# DESCARTES SOLVER III:

## The Algorithm




# DESCARTES SOLVER III:

The Algorithm




# DESCARTES SOLVER III:

## The Algorithm




# DESCARTES SOLVER III:

## The Algorithm




# DESCARTES SOLVER III:

## The Algorithm




# DESCARTES SOLVER III:

## The Algorithm




# DESCARTES SOLVER III:

## The Algorithm




# DESCARTES SOLVER III:

## The Algorithm




# DESCARTES SOLVER III:

## The Algorithm



# DESCARTES SOLVER III:


## The Algorithm



Subdivide until for all  $J$ ,  
 $V(g, J) \leq 1$ !

# DESCARTES SOLVER III:

## The Algorithm



Subdivide until for all  $J$ ,  
 $V(g, J) \leq 1!$

# Real Root Isolation IV:

Are we being pessimistic?

# Real Root Isolation IV:

Are we being pessimistic?

Worst-case complexity:

# Real Root Isolation IV:

Are we being pessimistic?

Worst-case complexity:

$$\max \{ \text{cost}(\text{SOLVER}, g) \mid \text{bit-size}(g) \leq \tau, \deg(g) \leq d \}$$

# Real Root Isolation IV:

Are we being pessimistic?

Worst-case complexity:

$$\max \{ \text{cost}(\text{SOLVER}, g) \mid \text{bit-size}(g) \leq \tau, \deg(g) \leq d \}$$



Pessimistic in practice

# Real Root Isolation IV:

Are we being pessimistic?

Worst-case complexity:

$$\max \{ \text{cost}(\text{SOLVER}, g) \mid \text{bit-size}(g) \leq \tau, \deg(g) \leq d \}$$



Pessimistic in practice

DESCARTES SOLVER

seems to behave faster in practice!

# Real Root Isolation IV:

Are we being pessimistic?

Worst-case complexity:

$$\max \{ \text{cost}(\text{SOLVER}, g) \mid \text{bit-size}(g) \leq \tau, \deg(g) \leq d \}$$



Pessimistic in practice

DESCARTES SOLVER

seems to behave faster in practice!

Can we explain this?

Real Root Isolation V:

Beyond pessimism

Real Root Isolation V:

Beyond pessimism

Worst-case complexity:

# Real Root Isolation V:

Beyond pessimism

Worst-case complexity:

$$\max \{ \text{cost}(\text{SOLVER}, g) \mid \text{bit-size}(g) \leq \tau, \deg(g) \leq d \}$$

# Real Root Isolation V:

Beyond pessimism

Worst-case complexity:

$$\max \{ \text{cost}(\text{SOLVER}, g) \mid \text{bit-size}(g) \leq \tau, \deg(g) \leq d \}$$

(Goldstine & von Neumann, 1951)  
(Demmel, 1988) (Smale; 1985, 1997)

↓  
(Roughgarden, 2021)

# Real Root Isolation V:

Beyond pessimism

Worst-case complexity:

$$\max \{ \text{cost}(\text{SOLVER}, g) \mid \text{bit-size}(g) \leq \tau, \deg(g) \leq d \}$$

(Goldstine & von Neumann, 1951)  
(Demmel, 1988) (Smale; 1985, 1997)

(Roughgarden, 2021)

Probabilistic complexity

# Real Root Isolation V:

Beyond pessimism

Worst-case complexity:

$$\max \{ \text{cost}(\text{SOLVER}, g) \mid \text{bit-size}(g) \leq \tau, \deg(g) \leq d \}$$

(Goldstine & von Neumann, 1951)  
(Demmel, 1988) (Smale; 1985, 1997)

(Roughgarden, 2021)

Probabilistic complexity

$$\mathbb{E} \{ \text{cost}(\text{SOLVER}, g)^l \mid \text{bit-size}(g) \leq \tau, \deg(g) \leq d \}$$

# Real Root Isolation V:

Beyond pessimism

Worst-case complexity:

$$\max \{ \text{cost}(\text{SOLVER}, g) \mid \text{bit-size}(g) \leq \tau, \deg(g) \leq d \}$$

(Goldstine & von Neumann, 1951)  
(Demmel, 1988) (Smale; 1985, 1997)

(Roughgarden, 2021)

Probabilistic complexity

$$\mathbb{E} \{ \text{cost}(\text{SOLVER}, g)^l \mid \text{bit-size}(g) \leq \tau, \deg(g) \leq d \}$$

What's a 'good' random model for  $g$ ?

# Real Root Isolation V:

Beyond pessimism

Worst-case complexity:

$$\max \{ \text{cost}(\text{SOLVER}, g) \mid \text{bit-size}(g) \leq \tau, \deg(g) \leq d \}$$

(Goldstine & von Neumann, 1951)  
(Demmel, 1988) (Smale; 1985, 1997)

(Roughgarden, 2021)

Probabilistic complexity

$$\mathbb{E} \{ \text{cost}(\text{SOLVER}, g)^l \mid \text{bit-size}(g) \leq \tau, \deg(g) \leq d \}$$

What's a 'good' random model for  $g$ ?

↑  
Many choices of randomness 😱

Beyond pessimism I:  
Uniform Random Bit Polynomials  
& A SIMPLE MAIN THEOREM

Beyond pessimism I:

Uniform Random Bit Polynomials

& A SIMPLE MAIN THEOREM

$$F = \sum_{k=0}^d F_k X^k$$

# Beyond pessimism I: Uniform Random Bit Polynomials & A SIMPLE MAIN THEOREM

$$F = \sum_{k=0}^d F_k X^k$$

s.t.  $F_k \sim \mathcal{U}([-2^\gamma, 2^\gamma] \cap \mathbb{Z})$  independent

# Beyond pessimism I: Uniform Random Bit Polynomials & A SIMPLE MAIN THEOREM

$$F = \sum_{k=0}^d F_k X^k$$

s.t.  $F_k \sim \mathcal{U}([-2^\gamma, 2^\gamma] \cap \mathbb{Z})$  independent

SIMPLE MAIN THM (Ergür, T-C, Tsigaridas)

$$\mathbb{E} \text{cost}(\text{DESCARTES}, F) = \tilde{\mathcal{O}}_B(d^2 + d\gamma)$$

# Beyond pessimism I: Uniform Random Bit Polynomials & A SIMPLE MAIN THEOREM

$$F = \sum_{k=0}^d F_k X^k$$

s.t.  $F_k \sim \mathcal{U}([-2^\gamma, 2^\gamma] \cap \mathbb{Z})$  independent

SIMPLE MAIN THM (Ergür, T-C, Tsigaridas)

$$\mathbb{E} \text{cost}(\text{DESCARTES}, F) = \tilde{\mathcal{O}}_B(d^2 + d\gamma)$$

On average, DESCARTES is almost optimal!

# Beyond pessimism II: Random Bit Polynomials

# Beyond pessimism II: Random Bit Polynomials

$$f = \sum_{k=0}^d f_k x^k \in \mathbb{Z}[x]$$

# Beyond pessimism II:

## Random Bit Polynomials

$$F = \sum_{k=0}^d f_k X^k \in \mathbb{Z}[X]$$

s.t.  $f_k$  independent

# Beyond pessimism II:

## Random Bit Polynomials

$$F = \sum_{k=0}^d f_k X^k \in \mathbb{Z}[X]$$

bit-size of  $F$ :

s.t.  $f_k$  independent

# Beyond pessimism II:

## Random Bit Polynomials

$$F = \sum_{k=0}^d f_k X^k \in \mathbb{Z}[X]$$

bit-size of  $F$ : s.t.  $f_k$  independent

$$\gamma(F) := \min\{\gamma \mid \forall K, P(|f_k| \leq 2^\gamma) = 1\}$$

# Beyond pessimism II:

## Random Bit Polynomials

$$F = \sum_{k=0}^d f_k X^k \in \mathbb{Z}[X]$$

bit-size of  $F$ : s.t.  $f_k$  independent

$$\gamma(F) := \min\{\gamma \mid \forall K, P(|f_k| \leq 2^\gamma) = 1\}$$

weight of  $F$ :

# Beyond pessimism II:

## Random Bit Polynomials

$$F = \sum_{k=0}^d f_k X^k \in \mathbb{Z}[X]$$

bit-size of  $F$ : s.t.  $f_k$  independent

$$\gamma(F) := \min\{\gamma \mid \forall K, P(|f_k| \leq 2^\gamma) = 1\}$$

weight of  $F$ :

$$w(F) := \max \{P(f_k = c) \mid c \in \mathbb{R}, k \in \{0, 1, d-1, d\}\}$$

# Beyond pessimism II:

## Random Bit Polynomials

$$F = \sum_{k=0}^d f_k X^k \in \mathbb{Z}[X]$$

bit-size of  $F$ : s.t.  $f_k$  independent

$$\gamma(F) := \min\{\gamma \mid \forall K, P(|f_k| \leq 2^\gamma) = 1\}$$

weight of  $F$ :

$$w(F) := \max \left\{ P(f_k = c) \mid c \in \mathbb{R}, k \in \{0, 1, \downarrow d-1, d\} \right\}$$

No middle indexes!

# Beyond pessimism II:

## Random Bit Polynomials

$$F = \sum_{k=0}^d f_k X^k \in \mathbb{Z}[X]$$

bit-size of  $F$ : s.t.  $f_k$  independent

$$\gamma(F) := \min\{\gamma \mid \forall K, P(|f_k| \leq 2^\gamma) = 1\}$$

weight of  $F$ : No middle indexes!

$$w(F) := \max \left\{ P(f_k = c) \mid c \in \mathbb{R}, k \in \{0, 1, \downarrow d-1, d\} \right\}$$

uniformity of  $F$ :

# Beyond pessimism II:

## Random Bit Polynomials

$$F = \sum_{k=0}^d F_k X^k \in \mathbb{Z}[X]$$

bit-size of  $F$ : s.t.  $F_k$  independent

$$\gamma(F) := \min\{\gamma \mid \forall K, P(|F_k| \leq 2^\gamma) = 1\}$$

weight of  $F$ : No middle indexes!

$$w(F) := \max \left\{ P(F_k = c) \mid c \in \mathbb{R}, k \in \{0, 1, \downarrow d-1, d\} \right\}$$

uniformity of  $F$ :  $u(F) := \ln(w(F)(1 + 2^{\gamma(F)+1}))$

# Beyond pessimism III: MAIN THEOREM

# Beyond pessimism III:

MAIN THEOREM

MAIN THM (Ergür, T-C, Tsigaridas)

$$\mathbb{E} \text{cost}(\text{DESCARTES}, F) = \tilde{\mathcal{O}}_B(d^2 + d\gamma)(1+u(F))^4$$

# Beyond pessimism III:

## MAIN THEOREM

MAIN THM (Ergür, T-C, Tsigaridas)

$$\mathbb{E} \text{cost}(\text{DESCARTES}, F) = \tilde{\mathcal{O}}_B(d^2 + d\gamma)(1 + u(F))^4$$

Note:  $F$  uniform  $\Rightarrow u(F) = 0$

# Beyond pessimism III:

## MAIN THEOREM

MAIN THM (Ergür, T-C, Tsigaridas)

$$\mathbb{E} \text{cost}(\text{DESCARTES}, F) = \tilde{\mathcal{O}}_B(d^2 + d\gamma)(1 + u(F))^4$$

Note:  $F$  uniform  $\Rightarrow u(F) = 0$

Claim: For many cases,  $u(F) = O(1)$

# Beyond pessimism III:

## MAIN THEOREM

MAIN THM (Ergür, T-C, Tsigaridas)

$$\mathbb{E} \text{cost}(\text{DESCARTES}, F) = \tilde{\mathcal{O}}_B(d^2 + d\gamma)(1 + u(F))^4$$

Note:  $F$  uniform  $\Rightarrow u(F) = 0$

Claim: For many cases,  $u(F) = O(1)$

IF  $\gamma = \Omega(d)$ , almost like reading!

# Beyond pessimism III:

## MAIN THEOREM

MAIN THM (Ergür, T-C, Tsigaridas)

$$\mathbb{E} \text{cost}(\text{DESCARTES}, F) = \tilde{O}_B(d^2 + d\gamma)(1 + u(F))^4$$

Note:  $F$  uniform  $\Rightarrow u(F) = 0$

Claim: For many cases,  $u(F) = O(1)$

IF  $\gamma = \Omega(d)$ , almost like reading!

On average, DESCARTES is almost optimal!

Beyond pessimism IV:

Examples of Random Bit Polynomials I

## Beyond pessimism IV:

### Examples of Random Bit Polynomials I

- Support control  $\{0, 1, d-1, d\} \subseteq A$

## Beyond pessimism IV:

### Examples of Random Bit Polynomials I

- Support control  $\{0, 1, d-1, d\} \subseteq A$

$$f = \sum_{k \in A} f_k X^k \quad \text{with } f_k \sim \mathcal{U}([-2^\gamma, 2^\gamma] \cap \mathbb{Z})$$

## Beyond pessimism IV:

### Examples of Random Bit Polynomials I

- Support control  $\{0, 1, d-1, d\} \subseteq A$

$$f = \sum_{k \in A} f_k X^k \quad \text{with } f_k \sim \mathcal{U}([-2^\gamma, 2^\gamma] \cap \mathbb{Z})$$

... then  $u(f) = 0$

## Beyond pessimism IV:

### Examples of Random Bit Polynomials I

- Support control  $\{0, 1, d-1, d\} \subseteq A$

$$f = \sum_{k \in A} f_k X^k \quad \text{with } f_k \sim \mathcal{U}([-2^\gamma, 2^\gamma] \cap \mathbb{Z})$$

... then  $u(f) = 0$

- Sign control  $\sigma \in \{-1, +1\}^{\{0, \dots, d\}}$

## Beyond pessimism IV:

### Examples of Random Bit Polynomials I

- Support control  $\{0, 1, d-1, d\} \subseteq A$

$$f = \sum_{k \in A} f_k X^k \quad \text{with } f_k \sim \mathcal{U}([-2^\gamma, 2^\gamma] \cap \mathbb{Z})$$

... then  $u(f) = 0$

- Sign control  $\sigma \in \{-1, +1\}^{\{0, \dots, d\}}$

$$F = \sum_{k=1}^d f_k X^k \quad \text{with } f_k \sim \mathcal{U}(\sigma_k ([1, 2^\gamma] \cap \mathbb{N}))$$

## Beyond pessimism IV:

### Examples of Random Bit Polynomials I

- Support control  $\{0, 1, d-1, d\} \subseteq A$

$$f = \sum_{k \in A} f_k X^k \quad \text{with } f_k \sim \mathcal{U}([-2^\gamma, 2^\gamma] \cap \mathbb{Z})$$

... then  $u(f) = 0$

- Sign control  $\sigma \in \{-1, +1\}^{\{0, \dots, d\}}$

$$F = \sum_{k=1}^d f_k X^k \quad \text{with } f_k \sim \mathcal{U}(\sigma_k ([1, 2^\gamma] \cap \mathbb{N}))$$

... then  $u(F) \leq \ln 3$

Beyond pessimism V:

Examples of Random Bit Polynomials II

## Beyond pessimism V:

### Examples of Random Bit Polynomials II

- Exact bitsize

## Beyond pessimism V:

### Examples of Random Bit Polynomials II

- Exact bitsize

$$F = \sum_{k=1}^d f_k X^k \quad \text{with} \quad f_k \sim \mathcal{U}\left(\{n \in \mathbb{Z} \mid \lfloor \log n \rfloor = r\}\right)$$

## Beyond pessimism V:

### Examples of Random Bit Polynomials II

- Exact bitsize

$$F = \sum_{k=1}^d f_k X^k \quad \text{with } f_k \sim \mathcal{U}\left(\{n \in \mathbb{Z} \mid \lfloor \log n \rfloor = r\}\right)$$

... then  $u(F) \leq \ln 3$

# Beyond pessimism V:

## Examples of Random Bit Polynomials II

- Exact bitsize

$$F = \sum_{k=1}^d f_k X^k \quad \text{with } f_k \sim \mathcal{U}\left(\{n \in \mathbb{Z} \mid \lfloor \log n \rfloor = r\}\right)$$

... then  $u(F) \leq \ln 3$

+ their combinations

## Beyond pessimism V:

### Examples of Random Bit Polynomials II


- Exact bitsize

$$F = \sum_{k=1}^d f_k X^k \quad \text{with } f_k \sim \mathcal{U}\left(\{n \in \mathbb{Z} \mid \lfloor \log n \rfloor = r\}\right)$$

... then  $u(F) \leq \ln 3$

+ their combinations

Our random model is flexible!



LOTS OF DETAILS  
WILL BE OMITTED

How to bound  
the run-time of DESCARTES?

How to bound  
the run-time of DESCARTES?

$\gamma(g) \leftarrow$  DESCARTES' Computation Tree


How to bound  
the run-time of DESCARTES?

$\gamma(g) \leftarrow$  DESCARTES' Computation Tree

$\langle -1, 1 \rangle$


How to bound  
the run-time of DESCARTES?

$\gamma(g) \leftarrow$  DESCARTES' Computation Tree




How to bound  
the run-time of DESCARTES?

$\gamma(g) \leftarrow$  DESCARTES' Computation Tree




How to bound  
the run-time of DESCARTES?

$\gamma(g) \leftarrow$  DESCARTES' Computation Tree




# How to bound the run-time of DESCARTES?

$\gamma(g) \leftarrow$  DESCARTES' Computation Tree




# How to bound the run-time of DESCARTES?

$\gamma(g) \leftarrow$  DESCARTES' Computation Tree



# How to bound the run-time of DESCARTES?

$\gamma(g) \leftarrow$  DESCARTES' Computation Tree




# How to bound the run-time of DESCARTES?

$\gamma(g) \leftarrow$  DESCARTES' Computation Tree

PROP.

$\text{cost}(\text{DESCARTES}, g)$



# How to bound the run-time of DESCARTES?


$\gamma(g) \leftarrow$  DESCARTES' Computation Tree

PROP.

$\text{cost}(\text{DESCARTES}, g)$


$\wedge$

$$\mathcal{O}(\text{d} \cdot \text{width} \gamma(g) \cdot \text{depth} \gamma(g) + d^2 \cdot \text{width} \gamma(g) \cdot \text{depth}^2 \gamma(g))$$



# How to bound the run-time of DESCARTES?

$\gamma(g) \leftarrow$  DESCARTES' Computation Tree



PROP.  
 $\text{cost}(\text{DESCARTES}, g)$   
 $\propto$   
 $O(d \gamma \text{width} \gamma(g) \text{depth} \gamma(g))$   
 $+ d^2 \text{width} \gamma(g) \text{depth}^2 \gamma(g))$

I.e. size of  $\gamma(g)$  bounds  
run-time of DESCARTES!

# The Ingredients of the Analysis I: Condition Number

# The Ingredients of the Analysis I: Condition Number

$$C(g) := \max_{x \in [-1, 1]} \frac{\sum_{k=0}^d |g_k|}{\max \{ |g(x)|, |g'(x)|/d \}}$$

# The Ingredients of the Analysis I: Condition Number

$$C(g) := \frac{\sum_{k=0}^d |g_k|}{\max \{ |g(x)|, |g'(x)|/d \}}$$

$x \in [-1, 1]$

$$C(g) = \infty$$

# The Ingredients of the Analysis I: Condition Number

$$C(g) := \frac{\sum_{k=0}^d |g_k|}{\max_{x \in [-1,1]} \{ |g(x)|, |g'(x)|/d \}}$$

$C(g) = \infty \Leftrightarrow g$  has a singular root in  $[-1,1]$

# The Ingredients of the Analysis I: Condition Number

$$C(g) := \frac{\sum_{k=0}^d |g_k|}{\max_{x \in [-1,1]} \{ |g(x)|, |g'(x)|/d \}}$$

$C(g) = \infty \Leftrightarrow g$  has a singular root in  $[-1,1]$

$\frac{1}{C(g)} \sim \left\{ \begin{array}{l} \text{How much I have to perturb} \\ \text{the coefficients of } g \text{ so that} \\ \tilde{g} \text{ has a singular root in } [-1,1] \end{array} \right.$

Bounding depth  $\gamma(g)$

Bounding depth  $\gamma(g)$

PROP.

$$\text{depth } \gamma(g) \leq 5 + \log d + \log C(g)$$

# Bounding depth $\gamma(g)$

PROP.

$$\text{depth } \gamma(g) \leq 5 + \log d + \log C(g)$$



sep. between complex roots of  $g$   
nearby  $[-1, 1]$

# Bounding depth $\gamma(g)$

PROP.

$$\text{depth } \gamma(g) \leq 5 + \log d + \log C(g)$$



$$\text{sep. between complex roots of } g \geq \frac{1}{12 d C(g)}$$

nearby  $[-1, 1]$

# Bounding depth $\gamma(g)$

PROP.

$$\text{depth } \gamma(g) \leq 5 + \log d + \log C(g)$$



$$\text{sep. between complex roots of } g \geq \frac{1}{12dc(g)}$$

nearby  $[-1, 1]$

11

1/2  $\text{depth } \gamma(g)$

# Bounding depth $\gamma(g)$

PROP.

$$\text{depth } \gamma(g) \leq 5 + \log d + \log C(g)$$



$$\text{sep. between complex roots of } g \geq \frac{1}{12dC(g)}$$

$\wedge$

$$\frac{1}{2} \text{depth } \gamma(g)$$

Here it is  
essential for  
roots being near  $[-1, 1]$ !

# Bounding depth $\gamma(g)$

PROP.

$$\text{depth } \gamma(g) \leq 5 + \log d + \log C(g)$$



$$\text{sep. between complex roots of } g \geq \frac{1}{12dC(g)}$$

↑

nearby  $[-1, 1]$

Here  $\rightarrow 1/\lambda$

no

$$\frac{1}{2} \text{ depth } \gamma(g)$$

Here it is  
essential for  
roots being near  $[-1, 1]$ !

Bounding width  $\gamma(8)$

Bounding width  $\gamma(8)$

PROP

width  $\gamma(8)$

Bounding width  $\gamma(g)$

PROP

width  $\gamma(g)$

$\leq \#$  complex roots of  $g$  nearby  $[-1, 1]$

Bounding width  $\gamma(8)$

PROP

width  $\gamma(8)$

$\leq \#$  complex roots of  $f$  nearby  $[-1, 1]$



Important  
For good bound  
of the RHS!

# Bounding width $\gamma(8)$

PROP

width  $\gamma(8)$

$\leq \#$  complex roots of  $f$  nearby  $[-1, 1]$

↑  
Important  
For good bound  
of the RHS!

Secret tool to bound RHS:

Bounding width  $\gamma(8)$

PROP

width  $\gamma(8)$

$\leq \#$  complex roots of  $f$  nearby  $[-1, 1]$



Important  
For good bound  
of the RHS!

Secret tool to bound RHS:

Titchmarsh's thm.

# The Ingredients of the Analysis II: Cont. Probabilistic Toolbox

# The Ingredients of the Analysis II: Cont. Probabilistic Toolbox

We can handle

$$C(F)$$

for a wide class of random  $F$

# The Ingredients of the Analysis II: Cont. Probabilistic Toolbox

We can handle

$$C(F)$$

for a wide class of random  $F$   
as long as continuous distribution

# The Ingredients of the Analysis II: Cont. Probabilistic Toolbox

We can handle

$$C(F) \left( \text{ & # complex roots of } F \text{ nearby } [-1, 1] \right)$$

for a wide class of random  $F$   
as long as continuous distribution

# The Ingredients of the Analysis II: Cont. Probabilistic Toolbox

We can handle

$$C(F) \left( \text{ & # complex roots of } F \text{ nearby } [-1, 1] \right)$$

for a wide class of random  $F$   
as long as continuous distribution  
— using geometric Functional analysis

# The Ingredients of the Analysis II: Cont. Probabilistic Toolbox

We can handle

$$C(F) \left( \text{ & # complex roots of } F \text{ nearby } [-1, 1] \right)$$

for a wide class of random  $F$   
as long as continuous distribution  
— using geometric functional analysis

... but we don't have a continuous dist.

# The Ingredients of the Analysis III: Ball's Smoothing Trick

# The Ingredients of the Analysis III: Ball's Smoothing Trick

$x \in \mathbb{Z}^N$  discrete random variable

# The Ingredients of the Analysis III: Ball's Smoothing Trick

$x \in \mathbb{Z}^N$  discrete random variable

$y \in \mathbb{R}^N$  s.t.  $y_i \sim \mathcal{U}((-1/2, 1/2))$  i.i.d.

# The Ingredients of the Analysis III: Ball's Smoothing Trick

$x \in \mathbb{Z}^N$  discrete random variable  
 $y \in \mathbb{R}^N$  s.t.  $y_i \sim \mathcal{U}(-\frac{1}{2}, \frac{1}{2})$  i.i.d.

$\rightarrow x + y$  continuous random var.

# The Ingredients of the Analysis III: Ball's Smoothing Trick

$x \in \mathbb{Z}^N$  discrete random variable  
 $y \in \mathbb{R}^N$  s.t.  $y_i \sim \mathcal{U}(-\frac{1}{2}, \frac{1}{2})$  i.i.d.

→  $x + y$  continuous random var.

We can use our old cont. toolbox!

Summing up:

SUMMING UP:

DESCARTES' SOLVER

IS ALMOST OPTIMAL ON AVERAGE!

SUMMING UP:

DESCARTES' SOLVER

IS ALMOST OPTIMAL ON AVERAGE!

... AND THAT'S WHY DESCARTES WORKS SO WELL

Muito Obrigado

pela Atenção!

Muchas Gracias

por su Atención !