

How many Real Zeros  
does  
a random sparse polynomial system  
have?

Algebra Seminar  
UPV/EHU

Josué TONELLI-CUETO  
(Johns Hopkins University)

# Joint work with...



Alperen A. ERGÜR



Mate L. TELEK



Elias TSIGARIDAS

$$\left\{ \begin{array}{l} a_1 + b_1 X + c_1 Y + d_1 X Y Z^d = 0 \\ a_2 + b_2 X + c_2 Y + d_2 X Y Z^d = 0 \\ a_3 + b_3 X + c_3 Y + d_3 X Y Z^d = 0 \end{array} \right.$$

How many...

Complex Zeros?

$$\left\{ \begin{array}{l} a_1 + b_1 X + c_1 Y + d_1 XYZ^d = 0 \\ a_2 + b_2 X + c_2 Y + d_2 XYZ^d = 0 \\ a_3 + b_3 X + c_3 Y + d_3 XYZ^d = 0 \end{array} \right.$$

How many...

Complex Zeros?

$$\left\{ \begin{array}{l} a_1 + b_1 X + c_1 Y + d_1 XYZ^d = 0 \\ a_2 + b_2 X + c_2 Y + d_2 XYZ^d = 0 \\ a_3 + b_3 X + c_3 Y + d_3 XYZ^d = 0 \end{array} \right.$$

*d complex  
zeros  
(generically)*

How many...

Real Zeros?

$$\left\{ \begin{array}{l} a_1 + b_1 X + c_1 Y + d_1 XYZ^d = 0 \\ a_2 + b_2 X + c_2 Y + d_2 XYZ^d = 0 \\ a_3 + b_3 X + c_3 Y + d_3 XYZ^d = 0 \end{array} \right.$$

How many...

Real Zeros?

$$\left\{ \begin{array}{l} a_1 + b_1 X + c_1 Y + d_1 XYZ^d = 0 \\ a_2 + b_2 X + c_2 Y + d_2 XYZ^d = 0 \\ a_3 + b_3 X + c_3 Y + d_3 XYZ^d = 0 \end{array} \right.$$

$\leq 2$   
real zeros  
(generically)

How many...

Positive Zeros?

$$\left\{ \begin{array}{l} a_1 + b_1 X + c_1 Y + d_1 XYZ^d = 0 \\ a_2 + b_2 X + c_2 Y + d_2 XYZ^d = 0 \\ a_3 + b_3 X + c_3 Y + d_3 XYZ^d = 0 \end{array} \right.$$

How many...

Positive Zeros?

$$\left\{ \begin{array}{l} a_1 + b_1 X + c_1 Y + d_1 XYZ^d = 0 \\ a_2 + b_2 X + c_2 Y + d_2 XYZ^d = 0 \\ a_3 + b_3 X + c_3 Y + d_3 XYZ^d = 0 \end{array} \right.$$

0 or 1  
positive zeros  
(generically)

General Phenomenon...

Many Complex Zeros,

but a lot fewer Real Zeros

Why do we care about Real Zeros?

[Applied answer]

Why do we care about Real Zeros?

- Chemical Reaction Networks

[Applied answer]

Why do we care about Real Zeros?

- Chemical Reaction Networks
- Computer Vision

[Applied answer]

Why do we care about Real Zeros?

- Chemical Reaction Networks
- Computer Vision
- Phylogenetic Trees

[Applied answer]

Why do we care about Real Zeros?

In applications,

the Real & Positive Zeros

are the ones that matter

[Applied Answer]

Why do we care about Real Zeros?

[Pure answer]

Why do we care about Real Zeros?

The number of Complex zeros  
is a geometric quantity:

[Pure answer]

Why do we care about Real Zeros?

The number of Complex zeros  
is a geometric quantity:  
degree, (mixed) volume of Newton polytopes

[Pure answer]

Why do we care about Real Zeros?

The number of Complex zeros  
is a geometric quantity:  
degree, (mixed) volume of Newton polytopes

The number of Real zeros  
is a more combinatorial quantity:

[Pure answer]

Why do we care about Real Zeros?

The number of Complex zeros  
is a geometric quantity:  
degree, (mixed) volume of Newton polytopes

The number of Real zeros  
is a more combinatorial quantity:  
size of the support

[Pure answer]

# Khovanskii's Theorem on Fewnomial Systems



Kiil's Theorem  
Fewnomial Systems

i?



Kiil's Theorem  
Fewnomial Systems


Polynomial



# Kiil's Theorem

## Fewnomial Systems

i?



Polynomial  $\rightarrow$  poly-nomial



Polynomial → poly-nomial → МНОГО-ЧЛЕН



# Kiil's Theorem

## Fewnomial Systems

i?

Polynomial  $\rightarrow$  poly-nomial  $\rightarrow$  МНОГО-ЧЛЕН  
 $\rightarrow$  МНОГОЧЛЕН  
"many monomial"



# Kiil's Theorem

## Fewnomial Systems

i?

Polynomial → poly-nomial → МНОГО-ЧЛЕН  
→ МНОГОЧЛЕН → МАЛОЧЛЕН  
"many monomial"      "few monomials"



# Kiil's Theorem

## Fewnomial Systems

i?

Polynomial → poly-nomial → МНОГО-ЧЛЕН  
→ МНОГОЧЛЕН → МАЛОЧЛЕН  
"many monomial"      "few monomials"  
→ МАЛО-ЧЛЕН



# Kiil's Theorem

## Fewnomial Systems

Polynomial → poly-nomial → МНОГО-ЧЛЕН  
→ МНОГОЧЛЕН → МАЛОЧЛЕН  
"many monomial"      "few monomials"  
→ МАЛО-ЧЛЕН → Few-nomial



# Kiil's Theorem

## Fewnomial Systems

i?

Polynomial  $\rightarrow$  poly-nomial  $\rightarrow$  МНОГО-ЧЛЕН  
 $\rightarrow$  МНОГОЧЛЕН  $\rightarrow$  МАЛОЧЛЕН  
"many monomial"      "few monomials"  
 $\rightarrow$  МАЛО-ЧЛЕН  $\rightarrow$  Few-nomial  $\rightarrow$  Fewnomial



# Kiil's Theorem

## Fewnomial Systems

i?

Polynomial  $\rightarrow$  poly-nomial  $\rightarrow$  МНОГО-ЧЛЕН  
 $\rightarrow$  МНОГОЧЛЕН  $\rightarrow$  МАЛОЧЛЕН  
"many monomial"      "few monomials"  
 $\rightarrow$  МАЛО-ЧЛЕН  $\rightarrow$  Few-nomial  $\rightarrow$  Fewnomial  
 $\rightsquigarrow$  Gutxinomia



# Kiil's Theorem

## Fewnomial Systems

i?

Polynomial → poly-nomial → МНОГО-ЧЛЕН  
→ МНОГОЧЛЕН → МАЛОЧЛЕН  
"many monomial"      "few monomials"  
→ МАЛО-ЧЛЕН → Few-nomial → Fewnomial  
↔ Gutxinomia      ↔ Poconomio

# Khovanskii's Theorem on Fewnomial Systems

$g \in \mathbb{R}[x_1, \dots, x_n]^n$  fewnomial system

# Khovanskii's Theorem on Fewnomial Systems

$g \in \mathbb{R}[X_1, \dots, X_n]^n$  fewnomial system  
with at most  $t$  exponents

# Khovanskii's Theorem on Fewnomial Systems

$g \in \mathbb{R}[x_1, \dots, x_n]^n$  fewnomial system  
with at most  $t$  exponents

where  $\mathcal{X}_v(g, \mathbb{R}_+^n) := \{x \in \mathbb{R}_+^n \mid g(x) = 0, D_x g \text{ non-sing.}\}$

# Khovanskii's Theorem on Fewnomial Systems

$g \in \mathbb{R}[x_1, \dots, x_n]^n$  fewnomial system  
with at most  $t$  exponents

$$\# \mathcal{Z}_r(g, \mathbb{R}^n_+) \leq 2^{\binom{t-1}{2}} (n+1)^{t-1}$$

where  $\mathcal{Z}_r(g, \mathbb{R}^n_+) := \{x \in \mathbb{R}^n_+ \mid g(x) = 0, D_x g \text{ non-sing.}\}$

# Khovanskii's Theorem on Fewnomial Systems

$g \in \mathbb{R}[x_1, \dots, x_n]^n$  fewnomial system  
with at most  $t$  exponents

No degree!

$$\# \mathcal{Z}_r(g, \mathbb{R}_+^n) \leq 2^{\binom{t-1}{2}} \binom{t-1}{n+1}$$

where  $\mathcal{Z}_r(g, \mathbb{R}_+^n) := \{x \in \mathbb{R}_+^n \mid g(x) = 0, D_x g \text{ non-sing.}\}$

# Kushnirenko's Hypothesis

# Kushnirenko's Hypothesis

$g \in \mathbb{R}[x_1, \dots, x_n]^n$  Fewnomial system

where  $g_i$  has  $t_i$  terms

# Kushnirenko's Hypothesis

$g \in \mathbb{R}[x_1, \dots, x_n]^n$  Fewnomial system

where  $g_i$  has  $t_i$  terms

Is there universal  $c > 0$  s.t.

$$\#\mathcal{Z}_r(g, \mathbb{R}_+^n) \leq (t_1 \cdots t_n)^c ?$$

State of the Art (Zero dimensional case)

State of the Art (Zero dimensional case)

(Descartes, XVII) if  $n=1$ ,  $\#\mathcal{Z}(g, \mathbb{R}_+) \leq t-1$

State of the Art (Zero dimensional case)

(Descartes, XVII) if  $n=1$ ,  $\#\mathcal{Z}(g, \mathbb{R}_+) \leq t-1$

(Khovanskii, 1991)

$$\#\mathcal{Z}(g, \mathbb{R}_+^n) \leq 2^{\binom{t-1}{2}} (n+1)^{t-1}$$

# State of the Art (Zero dimensional case)

(Descartes, XVII) if  $n=1$ ,  $\#\mathcal{Z}(g, \mathbb{R}_+) \leq t-1$

(Khovanskii, 1991)

$$\#\mathcal{Z}_r(g, \mathbb{R}_+^n) \leq 2^{\binom{t-1}{2}} (n+1)^{t-1}$$

(Bihan, Ottile; 2007)

$$\#\mathcal{Z}_r(g, \mathbb{R}_+^n) \leq 3 \cdot 2^{\binom{t-n-1}{2}} n^{t-n-1}$$

# State of the Art (Zero dimensional case)

(Descartes, XVII) if  $n=1$ ,  $\#\mathcal{Z}(g, \mathbb{R}_+) \leq t-1$

(Khovanskii, 1991)

$$\#\mathcal{Z}_r(g, \mathbb{R}_+^n) \leq 2^{\binom{t-1}{2}} (n+1)^{t-1}$$

(Bihan, Ottile; 2007)

$$\#\mathcal{Z}_r(g, \mathbb{R}_+^n) \leq 3 \cdot 2^{\binom{t-n-1}{2}} \cdot n^{t-n-1}$$

(Sevastyanov, 1978; Koiran, Portier, Tavenas; 2015)

$$\text{if } n=2, \quad \#\mathcal{Z}_r(g, \mathbb{R}_+^2) \leq O(d_2^3 t_1 + d_2^2 t_1^3)$$

State of the Art (Zero dimensional case)

Even the bivariate case  
 $(n=2)$

is widely open !

A Probabilistic Approach

# A Probabilistic Approach

Randomize  $F_j$

# A Probabilistic Approach

Randomize  $F_i$ ,

what can we say about

$$\mathbb{E} \# Z_r(F, \mathbb{R}_+^n) ?$$

# Probabilistic State of the Art I

(Bügisser, Ergür, Tonelli-Cueto; 2018)

# Probabilistic State of the Art I

(Bügisser, Ergür, Tonelli-Cueto; 2018)

$$F_i = \sum_{\alpha \in A} F_{i,\alpha} X^\alpha$$

$$t := \#A$$

# Probabilistic State of the Art I

(Bügisser, Ergür, Tonelli-Cueto; 2018)

$$f_i = \sum_{\alpha \in A} f_{i,\alpha} X^\alpha \quad \text{with } f_{i,\alpha} \text{ ind. cent. Gaussian}$$

$$t := \#A$$

# Probabilistic State of the Art I

(Bügisser, Ergür, Tonelli-Cueto; 2018)

$$F_i = \sum_{\alpha \in A} F_{i,\alpha} X^\alpha \quad \text{with } F_{i,\alpha} \text{ ind. cent. Gaussian}$$

s.t.  $\text{Var}(F_{i,\alpha})$  only depends on  $\alpha$

$$t := \#A$$

# Probabilistic State of the Art I

(Bügisser, Ergür, Tonelli-Cueto; 2018)

$$F_i = \sum_{\alpha \in A} F_{i,\alpha} X^\alpha \quad \text{with } F_{i,\alpha} \text{ ind. cent. Gaussian}$$

s.t.  $\text{Var}(F_{i,\alpha})$  only depends on  $\alpha$

$$t := \#A$$

$$\mathbb{E} \# Z_r(f, \mathbb{R}_+^n) \leq \frac{1}{2^{n-1}} \binom{t}{n}$$

# Probabilistic State of the Art II

(Bürgisser, ISSAC'2023)

$$F_i = \sum_{\alpha \in A_i} F_{i,\alpha} X^\alpha \quad \text{with } F_{i,\alpha} \text{ i.i.d. standard Gaussian}$$

$$t_i := \#A_i$$

$$\mathbb{E} \# Z_r(f, \mathbb{R}_+^n) \leq \frac{1}{(2\pi)^{\frac{n}{2}}} V\left(\sum_{i=1}^n P_i\right) t_1 \cdots t_n$$

where  $P_i := \text{conv}(A_i)$  &  $V(P) := \# \text{vertices of } P$

# Probabilistic State of the Art II

(Bürgisser, ISSAC'2023)

$$F_i = \sum_{\alpha \in A_i} F_{i,\alpha} X^\alpha \quad \text{with } F_{i,\alpha} \text{ i.i.d. standard Gaussian}$$

$$t_i := \#A_i$$

$$\mathbb{E} \# Z_r(f, \mathbb{R}_+^n) \leq \frac{1}{(2\pi)^{\frac{n}{2}}} V\left(\sum_{i=1}^n P_i\right) t_1 \cdots t_n$$

where  $P_i := \text{conv}(A_i)$  &  $V(P) := \# \text{vertices of } P$

# Probabilistic State of the Art II

(Bürgisser, ISSAC'2023)

$$F_i = \sum_{\alpha \in A_i} F_{i,\alpha} X^\alpha \quad \text{with } F_{i,\alpha} \text{ i.i.d. standard Gaussian}$$

$t_i := \#A_i$  Call the same variance

$$\mathbb{E} \# Z_r(f, \mathbb{R}_+^n) \leq \frac{1}{(2\pi)^{\frac{n}{2}}} V\left(\sum_{i=1}^n P_i\right) t_1 \cdots t_n$$

where  $P_i := \text{conv}(A_i)$  &  $V(P) := \# \text{vertices of } P$

# Main Result I (Ergür, Telek, Tonelli-Cueto; 2023+)

# Main Result I (Ergür, Telek, Tonelli-Cueto; 2023+)

$$f \in \mathbb{R}[x_1, \dots, x_n]^n$$

random Fewnomial system

# Main Result I (Ergür, Telek, Tonelli-Cueto; 2023+)

$f \in \mathbb{R}[x_1, \dots, x_n]^n$  random Fewnomial system  
such that the coefficients are ind. cent. Gaussians

# Main Result I (Ergür, Telek, Tonelli-Cueto; 2023+)

$f \in \mathbb{R}[x_1, \dots, x_n]^n$  random Fewnomial system  
such that the coefficients are ind. cent. Gaussians

$$\mathbb{E} \# Z_r(f, \mathbb{R}_+^n) \leq \frac{1}{4^n} t_1(t_1-1) \cdots t_n(t_n-1)$$

where  $t_i$  is # monomials of  $i$ ;

# Main Result II (Ergür, Telek, Tonelli-Cueto; 2023+)

# Main Result II (Ergür, Telek, Tonelli-Cueto; 2023+)

$$f_i = \sum_{\alpha \in A_i} f_{i,\alpha} X^\alpha \quad \text{with } f_{i,\alpha} \text{ ind. cent. Gaussians}$$

$$t_i := \#A_i$$

# Main Result II (Ergür, Telek, Tonelli-Cueto; 2023+)

$$F_i = \sum_{\alpha \in A_i} F_{i,\alpha} X^\alpha \quad \text{with } F_{i,\alpha} \text{ ind. cent. Gaussians}$$

$$t_i := \#A_i \quad \& \quad P_i := \text{conv}(A_i)$$

# Main Result II (Ergür, Telek, Tonelli-Cueto; 2023+)

$$F_i = \sum_{\alpha \in A_i} F_{i,\alpha} X^\alpha \quad \text{with } F_{i,\alpha} \text{ ind. cent. Gaussians}$$

$$t_i := \#A_i \quad \& \quad P_i := \text{conv}(A_i)$$

$$\text{IF } \text{Var}(F_{i,\alpha}) \leq 1,$$

# Main Result II (Ergür, Telek, Tonelli-Cueto; 2023+)

$$F_i = \sum_{\alpha \in A_i} F_{i,\alpha} X^\alpha \quad \text{with } F_{i,\alpha} \text{ ind. cent. Gaussians}$$

$$t_i := \#A_i \quad \& \quad P_i := \text{conv}(A_i)$$

IF  $\text{Var}(F_{i,\alpha}) \leq 1$ ,

with equality whenever  $\alpha$  is a vertex  $P_i$

# Main Result II (Ergür, Telek, Tonelli-Cueto; 2023+)

$$f_i = \sum_{\alpha \in A_i} f_{i,\alpha} X^\alpha \quad \text{with } f_{i,\alpha} \text{ ind. cent. Gaussians}$$

$$t_i := \#A_i \quad \& \quad P_i := \text{conv}(A_i)$$

IF  $\text{Var}(f_{i,\alpha}) \leq 1$ ,  
with equality whenever  $\alpha$  is a vertex  $P_i$

$$\mathbb{E} \# Z_r(f, \mathbb{R}_+^n) \leq \frac{1}{4^n} V\left(\sum_{i=1}^n P_i\right) t_1 \cdots t_n$$

where  $V(P) := \# \text{vertices of } P$

# Main Result III (Ergür, Telek, Tonelli-Cueto; 2023+)

# Main Result III (Ergür, Telek, Tonelli-Cueto; 2023+)

$$f_i = \sum_{\alpha \in A_i} f_{i,\alpha} X^\alpha \quad \text{with } f_{i,\alpha} \text{ ind. cent. Gaussians}$$

# Main Result III (Ergür, Telek, Tonelli-Cueto; 2023+)

$$f_i = \sum_{\alpha \in A_i} f_{i,\alpha} X^\alpha \quad \text{with } f_{i,\alpha} \text{ ind. cent. Gaussians}$$

$$t_i := \#A_i$$

# Main Result III (Ergür, Telek, Tonelli-Cueto; 2023+)

$$f_i = \sum_{\alpha \in A_i} f_{i,\alpha} X^\alpha \quad \text{with } f_{i,\alpha} \text{ ind. cent. Gaussians}$$

$$t_i := \#A_i \quad \& \quad \pi_i : A_i \rightarrow \mathbb{R} \quad \text{s.t. } \pi_i(\alpha) := \log \text{Var}(f_{i,\alpha})$$

# Main Result III (Ergür, Telek, Tonelli-Cueto; 2023+)

$$F_i = \sum_{\alpha \in A_i} F_{i,\alpha} X^\alpha \quad \text{with } F_{i,\alpha} \text{ ind. cent. Gaussians}$$

$$t_i := \#A_i \quad \& \quad \pi_i : A_i \rightarrow \mathbb{R} \quad \text{s.t.} \quad \pi_i(\alpha) := \log \text{Var}(F_{i,\alpha})$$

$$\mathcal{L}(A, \pi) := \text{conv} \left\{ (\pi_\alpha^{(\alpha)-s}) \mid \alpha \in A, s \geq 0 \right\}$$

# Main Result III (Ergür, Telek, Tonelli-Cueto; 2023+)

$F_i = \sum_{\alpha \in A_i} F_{i,\alpha} X^\alpha$  with  $F_{i,\alpha}$  ind. cent. Gaussians

$t_i := \#A_i$  &  $\pi_i : A_i \rightarrow \mathbb{R}$  s.t.  $\pi_i(\alpha) := \log \text{Var}(F_{i,\alpha})$

$\mathcal{L}(A, \pi) := \text{conv} \left\{ (\pi_\alpha^{(\alpha)-s}) \mid \alpha \in A, s \geq 0 \right\}$

$$\mathbb{E} \# Z_r(F, \mathbb{R}_+^n) \leq \frac{1}{4^n} V \left( \sum_{i=1}^n \mathcal{L}(A_i, \pi_i) \right) t_1 \cdots t_n$$

where  $V(P) := \# \text{vertices of } P$

# Two Tools I: Rice's Formula

# Two Tools I: Rice's Formula

$$\mathbb{E} \# \mathcal{Z}(F, \mathbb{R}_+^n) = \int_{\mathbb{R}_+^n} \mathbb{E}(|\det D_x F| \mid F(x) = 0) p_{F(x)}(0) dx$$

# Two Tools II: Cauchy-Binet Formula

## Two Tools II: Cauchy-Binet Formula

$A, B$   $m \times n$  matrices

$$\det A B^T$$

## Two Tools II: Cauchy-Binet Formula

$A, B$   $m \times n$  matrices

$$\det AB^T = \sum_{\substack{I \subseteq \{1, \dots, n\} \\ |I|=m}} \det A_I \det B_I$$

where  $X_I$  is the submatrix of  $X$   
• whose columns are indexed by  $I$

What about the variance?

What about the variance?

THM (Tsigaridas, Tonelli-Cueto; 2024+)

What about the variance?

THM (Tsigeridas, Tonelli-Cueto; 2024+)

$f \in \mathbb{R}[X_1, \dots, X_n]^n$  random Fewnomial system

What about the variance?

THM (Tsigaridas, Tonelli-Cueto; 2024+)

$f \in \mathbb{R}[X_1, \dots, X_n]^n$  random Fewnomial system  
with  $\sum P_i$  simple, where  $P_i = \text{conv}(A_i)$   
 $\& A_i$  support of  $A_i$

What about the variance?

THM (Tsigaridas, Tonelli-Cueto; 2024+)

$f \in \mathbb{R}[X_1, \dots, X_n]^n$  random Fewnomial system

with  $\sum P_i$  simple, where  $P_i = \text{conv}(A_i)$

&  $A_i$  support of  $A_i$

Under robust prob. assump.

$$\left( \mathbb{E} \# \mathcal{Z}_r(f, \mathbb{R}_+^n) \right)^{1/e} \leq V\left(\sum_{i=1}^n P_i\right) O(\log^2 D + n \log D)^n e^n$$

where  $V(P) := \# \text{vertices of } P$  &  $D = \max \text{ degree}$

## Future Work:

Can we answer deterministically

Kushnirenko's question

probabilistically?

Eskerrik  
asko!