

TU DRESDEN

ADVANCED PRACTICAL COURSE

LAB REPORT

Nuclear Magnetic Resonance

Authors:

Toni EHMCKE
Christian SIEGEL

Supervisor:

Samata CHAUDHUR

Dresden, November 29, 2015

Date of experimental procedure: November 20, 2015

Contents

1	Introduction	2
1.1	Motivation	2
2	Experimental procedure	2
3	Data Analysis	2
4	Discussion and conclusions	2

1 Introduction

1.1 Motivation

Nuclear Magnetic Resonance is a physical phenomenon that can be observed while placing an ensemble of nuclei into a static magnetic field and stimulate it with a high-frequent alternating field. A necessary condition for this effect is that the atoms of the sample have a *nuclear spin* different from zero. It is the central concept that is used for *NMR-Spectroscopy*, a standard methodology for the investigation of the structure and interaction of complex molecules and solid state bodies by measuring local magnetic fields, and the *magnetic resonance tomography* which is an imaging technique used in clinical diagnostics for describing the morphologic and physiologic build-up of tissues and organs. For all of those applications we need to find out some central parameters of particular physical compensation-processes, the so called *relaxation times* T_1 and T_2 . In the following experiment exactly those material-characteristic obserables are determined for an ensemble of ^{57}Fe -nuclei. But at first some basic knowledge.

1.2 Nuclear Zeeman-Effect

2 Experimental procedure

3 Data Analysis

4 Discussion and conclusions

References

- [01] S. Diez. *Advanced practical course - Experiment MMC*, Dresden, 2015
- [02] <https://en.wikipedia.org/wiki/Histogram> [November 20, 2015]
- [03] <https://en.wikipedia.org/wiki/Kinesin> [November 22, 2015]
- [04] S. Ferbrugge et al. *Novel ways to determine kinesin-1's run length and randomness using fluorescence microscopy*. Amsterdam. October 2009
- [05] S. Klumpp, R. Lipowsky. *Cooperative cargo transport by several molecular motors*. Potsdam. November 2005