

Dynamics Cheat Sheet

Linear Momentum L

Definition: total force acting on a particle is equal to the time rate of change of its linear momentum.

Equation: $\mathbf{L} = m\mathbf{v}$

- for a system of particles (G : center of mass):

$$\mathbf{L} = \sum_i m_i \mathbf{v}_i = m \mathbf{v}_G$$

Time rate of change: $\dot{\mathbf{L}} = \mathbf{F}$

- for a system of particles (G : center of mass):

$$\dot{\mathbf{L}} = \sum_i m_i \mathbf{a}_i = \sum_i \mathbf{F}_i = \underbrace{\mathbf{F}}_{\text{external forces}} = m \mathbf{a}_G$$

Angular Momentum H

Definition: "moment" of the particle's linear momentum \mathbf{L} about a point O .

Equation: $\mathbf{H}_O = \mathbf{r} \times \mathbf{L} = \mathbf{r} \times m\mathbf{v}$

Equation: $\mathbf{H}_O = \mathbf{r}_G \times m\mathbf{v}_G + \mathbf{H}_G$

- for a system of particles (about fixed point O):

$$\mathbf{H}_O = \sum_i (\mathbf{r}_i \times m_i \mathbf{v}_i)$$

Equation: taken about the center of mass G :

$$\mathbf{H}_G = \sum_i (\mathbf{r}'_i \times m\mathbf{v}_i) \quad | \quad \mathbf{v}_i = \dot{\mathbf{r}}_i = \dot{\mathbf{r}}_G + \dot{\mathbf{r}}'_i = \mathbf{v}_G + \mathbf{v}'_i \quad (1)$$

$$= \sum_i (\mathbf{r}'_i \times m(\mathbf{v}_G + \mathbf{v}'_i))$$

$$= \sum_i (\mathbf{r}'_i \times m\mathbf{v}_G) + \sum_i (\mathbf{r}'_i \times m\mathbf{v}'_i) \quad | \quad -\mathbf{v}_G \times \sum_i m_i \mathbf{r}'_i = 0$$

$$\mathbf{H}_G = \sum_i (\mathbf{r}'_i \times m\mathbf{v}'_i) \quad (2)$$

Equation (1) is the **absolute angular momentum**, whereas equation (2) is the **relative angular momentum**; when taken about G , these quantities are identical.

Time rate of change: wrt fixed point O

$$\begin{aligned} \frac{d\mathbf{H}_O}{dt} &= \cancel{\dot{\mathbf{r}} \times m\dot{\mathbf{v}}}^0 + \mathbf{r} \times m\dot{\mathbf{v}} \quad | \quad \dot{\mathbf{r}} = \mathbf{v} \\ &= \mathbf{r} \times m\mathbf{a} \quad | \quad \mathbf{F} = m\mathbf{a} \\ &= \mathbf{r} \times \mathbf{F} \\ &= \mathbf{M}_O \end{aligned}$$

- for a system of particles (about fixed point O):

$$\begin{aligned} \dot{\mathbf{H}}_O &= \sum_i \left(\cancel{\dot{\mathbf{r}}_i \times m_i \dot{\mathbf{v}}_i}^0 \right) + \sum_i (\mathbf{r}_i \times m_i \mathbf{a}_i) \\ &= \sum_i (\mathbf{r}_i \times \mathbf{F}_i) + \sum_i M_i \quad | \quad M_i = \text{external moments} \\ &= \mathbf{M}_O \end{aligned}$$

Time rate of change: about CoM G :

$$\dot{\mathbf{H}}_G = \sum_i (\mathbf{r}'_i \times m\dot{\mathbf{v}}'_i) = \sum_i (\mathbf{r}'_i \times \mathbf{F}_i) + \sum_i M_i = \mathbf{M}_G,$$

where \mathbf{M}_G is the total moment about G of the applied external forces plus any external moments. Expression valid for any movement of G !

Angular Impulse

$$\int_{t_1}^{t_2} \mathbf{M}_O dt = \int_{t_1}^{t_2} \dot{\mathbf{H}}_O dt = \mathbf{H}_O(t_2) - \mathbf{H}_O(t_1) = \Delta \mathbf{H}_O$$

"Useful when dealing with impulsive forces. Possible to calculate the integrated effect of a force on a particle without knowing in detail the actual value of the force as a function of time."

Kinetic Energy T

Equation: for a system of particles:

$$\begin{aligned} T &= \sum_i T_i = \sum_i \left(\frac{1}{2} m_i \mathbf{v}_i \cdot \mathbf{v}_i \right) \quad | \quad \text{decompose into relative and } G \\ &= \frac{1}{2} m v_G^2 + \sum_i \frac{1}{2} m_i v_i'^2 \end{aligned}$$

Center of Mass

$$\mathbf{r}_G = \frac{1}{m} \left(\sum_i m_i \mathbf{r}_i \right); m = \sum_i m_i$$

Tangential Velocity

$$\mathbf{v} = \boldsymbol{\omega} \times \mathbf{r}$$

Coreolis Theorem

Space and Body Cones