Perception: Camera Modeling

Kostas Daniilidis

Can our aerial robots control their flight like gannets?



How can gannets' eyes and brain estimate distance to water!

Nature Vol. 293 24 September 1981

- Spiess, F. N. et al. Science 207, 1421-1433 (1980).
- Corliss, J. B. Science 203, 1073-1083 (1979).
- 7. Edmond, J. M. et al. Earth planet. Sci. Lett. 46, 19-30 (1979).
- 8. Ruby, E. G., Wirsen, C. O. & Jannasch, H. W. Appl. envir. Microbiol. (in the press).
- 9. Cavanaugh, C. M., Gardiner, S. L., Jones, M. L., Jannasch, H. W. & Waterbury, J. B. Science 213, 340-342 (1981).
- 10. Felbeck, H. Science 213, 336-338 (1981).
- 11. Jones, M. L. Science 213, 333-336 (1981).
- 12. Peck, H. D. Jr Enzymes 10, 651-669 (1974).
- 13. Latzko, E. & Gibbs, M. Pl. Physiol. 44, 295-300 (1969).
- 14. Reid, R. G. B. Can. J. Zool. 58, 386-393 (1980).
- 15. Los Angeles County (California) Sanitation District files.
- 16. Rau, G. H. Science 213, 338-340 (1981). 17. Rau, G. H. Nature 289, 484-485 (1981).
- 18. Rau, G. H. & Hedges, J. I. Science 203, 648-649 (1979).
- 19. Emery, K. O. & Hulsemann, J. Deep-Sea Res. 8, 165-180 (1962).
- 20. Hartman, O. & Barnard, J. L. Allan Hancock Pacific Exped. 22, (1958).
- 21. Nicholas, D. J. D., Ferrante, J. V. & Clarke, G. R. Analyt. Biochem. 95, 24-31 (1979).
- Lonsdale, P. Nature 281, 531-534 (1979).


Plummeting gannets: a paradigm of ecological optics

David N. Lee & Paul E. Reddish

Department of Psychology, University of Edinburgh, Edinburgh EH8 9JZ, UK



Quadrotor "sees" with a camera



In robotics we use all kinds of cameras!





In robotics we use all kinds of cameras!






In robotics we use all kinds of cameras!






A camera is an imaging chip and a lens




A camera is an imaging chip and a lens




Magnifying glass






Rays from on object point P converge on a point p on the image plane




Rays from a point P in the scene converge into a point in the image plane



Rays parallel to the optical axis meet the focus after leaving the lens. Rays through center of the lens do not change direction.


$$\frac{1}{f} = \frac{1}{a} + \frac{1}{b}$$



These rays meet at one point if

$$\frac{1}{f} = \frac{1}{a} + \frac{1}{b}$$


What happens when we move b, the image plane



Moving the image plane is what we call (de-) focusing! Image starts blurring!

$$\frac{1}{f} \neq \frac{1}{a} + \frac{1}{b}$$


Perspective projection: size of object image



If you look only at the ray going through the center of the lens

$$\frac{Y}{a} = \frac{y}{b}$$


Perspective projection: size of object image



If you look only at the ray going through the center of the lens

$$\frac{Y}{a} = \frac{y}{b}$$


Perspective projection




- An object of the same size coming closer results on a larger image
- A point moving on the same ray does not change its image

$$\frac{Y}{a} = \frac{y}{b}$$

Perspective projection = Pinhole Model



Pinhole model




 $rac{Y}{Z} = rac{y}{b}$ Z = a

If we replace b with f and include a minus because object image is upside down....

 $y = -f\frac{1}{2}$

Pinhole model used in computer vision and robotics



... amd assume that image plane is in front of the lens

$$y = f \frac{Y}{Z}$$

What is the effect of f=b?

- Theoretically, we expect an offset in the x and y coordinates caused by the error (f-b).
- If the object is on focus:

$$\frac{b-f}{f} = \frac{b}{Z}$$

Relative error depends on the ratio of focal length to depth!

This would matter if we would actually use the f from specs of the came

In practice we use a process called calibration, yielding the f that best satisfies

$$y = f \frac{Y}{Z}$$