

Introduction to Statistical Method

Estimation

- An **estimator** for a population parameter θ is a statistic and denoted by $\hat{\theta}$.
- Any given value of $\hat{\theta}$ is an **estimate**.

Desirable Properties of a Point Estimator

- The expected value of $\hat{\theta}$ should be θ .
- $\hat{\theta}$ should have **small variance** for **large sample size**.

Bias

- The difference $\theta - E[\hat{\theta}]$ is the bias of an estimator $\hat{\theta}$ for a population parameter θ .
- $E[\hat{\theta}] = \theta$ means $\hat{\theta}$ is unbiased.

Mean Square Error of Estimator

- The **mean square error** of $\hat{\theta}$ is defined as $MSE(\hat{\theta}) := E[(\hat{\theta} - \theta)^2]$.
- The mean square error measures the **overall quality of an estimator**.

$$\begin{aligned} MSE(\hat{\theta}) &= E[(\hat{\theta} - E[\hat{\theta}] + E[\hat{\theta}] - \theta)^2] = E[(\hat{\theta} - E[\hat{\theta}])^2 + (E[\hat{\theta}] - \theta)^2 + 2(\hat{\theta} - E[\hat{\theta}])(E[\hat{\theta}] - \theta)] \\ &= Var(\hat{\theta}) + (bias)^2 + 2E[(\hat{\theta} - E[\hat{\theta}])(E[\hat{\theta}] - \theta)] = Var(\hat{\theta}) + (bias)^2 + 2E[(\hat{\theta} - E[\hat{\theta}])(E[\hat{\theta}] - \theta)] \\ &= Var(\hat{\theta}) + (bias)^2 + 2(E[\hat{\theta}] - E[\hat{\theta}])(E[\hat{\theta}] - \theta) = Var(\hat{\theta}) + (bias)^2 \end{aligned}$$

Sample Mean Variance and Bias

Theorem: Unbiased Sample Mean

Let X_1, \dots, X_n be a random sample of size n from a distribution with mean μ . Then the sample mean \bar{X} is an unbiased estimator for μ .

Since $\bar{X} = \frac{1}{n} \sum_{k=1}^n X_k$, then we see $\mu - E[\bar{X}] = \mu - \frac{1}{n}n\mu = 0$, thus unbiased.

Theorem: Sample Average Variance

Let \bar{X} be the sample of a random sample of size n from a distribution with mean μ and variance σ^2 . Then $Var \bar{X} = E[(\bar{X} - \mu)^2] = \frac{1}{n}\sigma^2$

Since $Var \bar{X} = Var(\frac{1}{n} \sum_{k=1}^n X_k) = \frac{1}{n^2} Var(\sum_{k=1}^n X_k)$, then check [ve401_note_4](#), we can see a fact that

$Var(X + Y) = Var X + Var Y + Cov(X, Y)$, where $Cov(X, Y) = E[(X - \mu_X)(Y - \mu_Y)]$, then we get $Var \bar{X} = \frac{1}{n^2}nVar X = \frac{1}{n}\sigma^2$

The standard deviation of \bar{X} is given by $\sqrt{Var \bar{X}} = \sigma/\sqrt{n}$ and is called standard error of mean.

Theorem: Sample Variance

Sample variance $S^2 = \frac{1}{n-1} \sum_{k=1}^n (X_k - \bar{X})^2$ is an unbiased estimator for σ^2 .

$$\begin{aligned} E\left[\frac{1}{n-1} \sum_{k=1}^n (X_k - \bar{X})^2\right] &= \frac{1}{n-1} E\left[\sum_{k=1}^n (X_k - \mu + \mu - \bar{X})^2\right] \\ &= \frac{1}{n-1} E\left[\sum_{k=1}^n (X_k - \mu)^2 - 2(\bar{X} - \mu)\left(\sum_{k=1}^n X_k - n\mu\right) + n(\mu - \bar{X})^2\right] \\ &= \frac{1}{n-1} E\left[\sum_{k=1}^n (X_k - \mu)^2 - n(\mu - \bar{X})^2\right] \\ &= \frac{1}{n-1} \left(\sum_{k=1}^n E[(X_k - \mu)^2] - nE[(\mu - \bar{X})^2]\right) = \frac{1}{n-1} \left(\sum_{k=1}^n \sigma^2 - n\frac{\sigma^2}{n}\right) = \sigma^2 \end{aligned}$$