

Introduction to Statistical Method

Hypothesis Testing

A second major statistical method for gaining information on a probability.

The goal is to reject or fail to reject statements (hypotheses) based on statistical data.

Hypothesis Definition

A statement about a population parameter θ . The hypothesis will compare θ to a null value denoted θ_0 .

Fisher's Null Hypothesis Test

This hypothesis will be denoted by H_0 and is null hypothesis.

Three forms: $H_0 : \theta = \theta_0$ or $H_0 : \theta \leq \theta_0$ or $H_0 : \theta \geq \theta_0$.

A hypothesis test is based on rejecting a hypothesis.

One-Tailed Test

The test of a hypothesis of the form $H_0 : \theta \leq \theta_0$ or $H_0 : \theta \geq \theta_0$ is said to be one-tailed tests.

P-Value for a One-Tailed Test

- Apply an example first

We want to find evidence that a new car design has a mean mileage greater than 26 mpg. Therefore, we set up the null hypothesis: $H_0 : \mu \leq 26$.

The goal is to reject the null hypothesis.

- Example explanation


We take a random sample and calculate \bar{X} , if it is much greater than 26, then there is reason to believe that H_0 is false.

Take a random sample of size n and find the value \bar{x} for the sample mean.

The probability of obtaining the measured value of \bar{x} or a larger result if H_0 is true is the **significance** or **P-Value** of the test.

(TM还是说中文吧：就是说，猜测 $\mu \leq 26$ ，然后我们得到了样本的平均的观测值 \bar{x} ，根据样本个数和标准差，我们可以对样本的平均值 \bar{X} 得到基于 $\mu \leq 26$ 的 test，然后可以放宽到 $\mu = 26$ ，然后可以用 $Z = \frac{\bar{X} - \mu_0}{\sigma/\sqrt{n}}$ 算出 P 的值来判断这个样本是否符合这个假设)

$$P[\bar{X} \geq \bar{x} | \mu \leq 26] \leq P[\bar{X} \geq \bar{x} | \mu = 26]$$



$H_0: \mu \leq \mu_0$ shows the case if $\mu = \mu_0$, the curve shift left if $\mu < \mu_0$

The shaded area shows the probability of obtaining $\bar{X} \geq \bar{x}$ if $\mu = \mu_0$.

- The P-value is therefore an upper bound of the probability of obtaining the data if H_0 is true.
- $P = P[D|H_0]$ if D represents the statistical data, we will reject H_0 if it is small.


So either:

- fail to reject H_0 at P level of significance
- reject the H_0 at P level of significance

The statistic on which the P is based is **test statistic**.

Two-Tailed Test

If we are testing a hypothesis of the form $H_0: \theta = \theta_0$, we say we are performing a two-tailed test.



$H_0: \mu = \mu_0$ The P is twice the value of one-tailed test.

Does a Small P-Value Provide Evidence that H_0 is False