

1. 6 marks Consider the problem [Vanderbei 5th edition, Exercise 1.1].
- (a) 3 marks Write this as a Linear Programming problem (in the standard form). You must explain your notation and variables.

Solution: Set the decision variables

$x_1 =$ the amount of Bands to be produced in tons

$x_2 =$ the amount of Coils to be produced in tons

We want to maximize the objective function

$$\text{Profit in \$} = 25x_1 + 30x_2$$

Before writing the constraints, we first translate the condition on production rates into the condition on time to produce x_1 amount of Bands, and x_2 amount of Coils.

producing x_1 tons will require $\frac{1}{200}x_1$ hours

producing x_2 tons will require $\frac{1}{140}x_2$ hours

Then we have the following constraints:

$$x_1 \leq 6000$$

$$x_2 \leq 4000$$

$$\frac{1}{200}x_1 + \frac{1}{140}x_2 \leq 40$$

$$x_1, x_2 \geq 0$$

The final LP problem is

$$\begin{aligned} & \max 25x_1 + 30x_2 \\ & \text{subject to. } x_1 \leq 6000 \\ & \quad x_2 \leq 4000 \\ & \quad \frac{1}{200}x_1 + \frac{1}{140}x_2 \leq 40 \\ & \quad x_1, x_2 \geq 0 \end{aligned}$$

- (b) 3 marks Solve the LP by writing down a code in the Python language using the Jupyter notebook; login to UBC syzygy website and the Jupyter notebook. Attach the screenshots.

Solution: See the file HW2-1b.ipynb.

2. 11 marks Consider the problem [Vanderbei. 5th edition, Exercise 1.2].
- (a) 5 marks Write this as a Linear Programming problem (in the standard inequality form). You must explain your notation and variables.

Solution: Notice that the flight is to fly from Ithaca to Newark, then to Boston. Set the decision variables. First name the routes:

R1 Ithaca to Newark

R2 Newark to Boston

R3 Ithaca to Boston

We use the letters Y, B, M, for the available classes. Then, the decision variables are, for each $i = 1, 2, 3$

x_{iY} = the number of Y class tickets for route i

x_{iB} = the number of B class tickets for route i

x_{iM} = the number of M class tickets for route i

Notice that there are 9 such variables.

We want to maximize the objective function (the revenue in \$)

$$300x_{1Y} + 220x_{1B} + 100x_{1M} + 160x_{2Y} + 130x_{2B} + 80x_{2M} + 360x_{3Y} + 280x_{3B} + 140x_{3M}$$

Now constraints. First, the obvious constraints: all variables are nonnegative and integer. Then, the upper bounds from the forecasted maximum demand.

$$x_{1Y} \leq 4, x_{1B} \leq 8, x_{1M} \leq 22, x_{2Y} \leq 8, x_{2B} \leq 13, x_{2M} \leq 20, x_{3Y} \leq 3, x_{3B} \leq 10, x_{3M} \leq 18.$$

Now the constraint from not allowing overbooking. The aircraft has 30 seats. So, we have

$$x_{1Y} + x_{1B} + x_{1M} + x_{3Y} + x_{3B} + x_{3M} \leq 30$$

$$x_{2Y} + x_{2B} + x_{2M} + x_{3Y} + x_{3B} + x_{3M} \leq 30$$

The first line is from the route between Ithaca and Newark, where there are route 1 (Ithaca to Newark) route 3 (Ithaca to Boston) passengers are to be onboard. The second line is from the route between Newark and Boston, where there are route 2 (Newark to Boston) route 3 (Ithaca to Boston) passengers are to be onboard.

The final LP problem is

max

$$300x_{1Y} + 220x_{1B} + 100x_{1M} + 160x_{2Y} + 130x_{1B} + 80x_{2M} + 360x_{3Y} + 280x_{3B} + 140x_{3M}$$

subject to

$$x_{1Y} \leq 4,$$

$$x_{1B} \leq 8,$$

$$x_{1M} \leq 22,$$

$$x_{2Y} \leq 8,$$

$$x_{2B} \leq 13,$$

$$x_{2M} \leq 20,$$

$$x_{3Y} \leq 3,$$

$$x_{3B} \leq 10,$$

$$x_{3M} \leq 18,$$

$$x_{1Y} + x_{1B} + x_{1M} + x_{3Y} + x_{3B} + x_{3M} \leq 30,$$

$$x_{2Y} + x_{2B} + x_{2M} + x_{3Y} + x_{3B} + x_{3M} \leq 30,$$

$$x_{1Y}, x_{1B}, x_{1M}, x_{2Y}, x_{1B}, x_{2M}, x_{3Y}, x_{3B}, x_{3M} \geq 0,$$

and $x_{1Y}, x_{1B}, x_{1M}, x_{2Y}, x_{1B}, x_{2M}, x_{3Y}, x_{3B}, x_{3M}$ are integers.

- (b) 6 marks Solve the LP by writing down a code in the Python language using the Jupyter notebook; login to UBC syzygy website and the Jupyter notebook.

Attach the screenshots.

[Hint1: You will need integer variables, those taking only integer values. For that you can add the command `cat='Integer'` like in the following example:

```
ticketvars = LpVariable.dicts("ticket", ticket, lowBound=0, cat='Integer')
```

Here the last part `cat='Integer'` restrict the variables to be integer variables.]

[Hint2: It is not necessary, but, to practice with 'for-loops' and 'dictionary', you can try to use dictionary variables as done in the blending (cat food) example.]

Solution: See the file HW2-2b.ipynb.

3. 3 marks For a nonempty set $S \subset \mathbf{R}^n$ and a positive real number $r \in \mathbf{R}$ (and $r > 0$), define the set rS as follows:

$$rS := \{z \in \mathbf{R}^n \mid z = rx, x \in S\}.$$

Here rx is the multiplication of the vector $x \in \mathbf{R}^n$ by the scalar $r \in \mathbf{R}$; in your more familiar notation, $r\vec{x}$. The set rS is the set of all points that are obtained by multiplying r with the vectors $x \in S$.

For a given nonempty set $S \subset \mathbf{R}^n$ and a given positive number $r > 0$, prove that if S is a convex set then rS is a convex set as well.

Solution:

- Assume S is convex.
- Suppose $x, y \in rS$, then $x = rx', y = ry'$ for some $x', y' \in S$.
- Now, consider for each $t \in [0, 1]$, $(1-t)x + ty = (1-t)rx' + t ry' = r[(1-t)x' + ty']$.
- Notice that since S is convex and $x', y' \in S$, we see that for that $t \in [0, 1]$, $(1-t)x' + ty' \in S$.
- Therefore by the definition of rS , $(1-t)x + ty = r[(1-t)x' + ty'] \in rS$.
- This verifies that rS is convex.

4. 5 marks For given two nonempty sets $S_1, S_2 \subset \mathbf{R}^n$, define the operation $S_1 + S_2$ as follows:

$$S_1 + S_2 := \{z \in \mathbf{R}^n \mid \text{there exist some } x \in S_1 \text{ and some } y \in S_2 \text{ such that } z = x + y\}$$

that is, each point $z \in S_1 + S_2$ is the one that can be expressed as the sum $x + y$ for some $x \in S_1$, and $y \in S_2$; here the sum $x + y$ is the vector sum between the two vectors. One does this for all $x \in S_1$ and $y \in S_2$ and get the set $S_1 + S_2$.

- (a) Consider $S_1 = \{(x_1, x_2) \in \mathbf{R}^2 \mid |x_1 - 1| \leq 1 \text{ \& } |x_2 - 2| \leq 1\}$ and $S_2 = \{(x_1, x_2) \in \mathbf{R}^2 \mid |x_1| \leq 2 \text{ \& } |x_2| \leq 1\}$. Sketch the set $S_1 + S_2$. You do not need to explain your solution for this question. But, your sketch should be neat and very clear, indicating all the relevant coordinate values.

Solution: The resulting set is

$$S_1 + S_2 = \{(x_1, x_2) \in \mathbf{R}^2 \mid |x_1 - 1| \leq 3 \text{ \& } |x_2 - 2| \leq 2\}.$$

We skip the sketch. Your sketch should clearly show that it is the square shape centered at $(1, 2)$ with horizontal side length 6 and vertical side length 4.

- (b) Is it true that $S_1 + S_2$ must be convex for **any** nonempty convex sets S_1 and S_2 in \mathbf{R}^n ? Justify your answer carefully. **[This problem is independent of part (a). The sets S_1, S_2 are arbitrary convex sets in this question, not the particular example given in part (a). If you do this problem only for the sets of part (a) or a particular example, you will get zero mark.]**

Solution: Such a sum $S_1 + S_2$ is called the Minkovski sum of S_1 and S_2 . It is convex if S_1 and S_2 are.

- To see this, let x, y be given two points in $S_1 + S_2$. It means there are points $x_1, y_1 \in S_1$ and $x_2, y_2 \in S_2$ such that $x = x_1 + x_2$, and $y = y_1 + y_2$.
- Now, consider for each $0 \leq t \leq 1$,

$$z_t = (1 - t)x + ty$$

- We claim that $z_t \in S_1 + S_2$ which will finish the proof.
- Notice that

$$\begin{aligned} z_t &= (1 - t)x + ty \\ &= (1 - t)(x_1 + x_2) + t(y_1 + y_2) \\ &= [(1 - t)x_1 + ty_1] + [(1 - t)x_2 + ty_2] \text{ by rearranging the terms.} \end{aligned}$$

- Notice that $[(1 - t)x_1 + ty_1] \in S_1$ from convexity of S_1 , since $x_1, y_1 \in S_1$ and $0 \leq t \leq 1$.

- Likewise, $[(1-t)x_2 + ty_2] \in S_2$ from convexity of S_2 , since $x_1, y_2 \in S_1$ and $0 \leq t \leq 1$.
- As z_t is given by their sum, we conclude that $z_t \in S_1 + S_2$ by the definition of the sum $S_1 + S_2$.
- Therefore we proved our claim, so this finishes the proof for convexity of $S_1 + S_2$ when S_1 and S_2 are convex.