

Differentiating Nonsmooth Solutions to Parametric Monotone Inclusion Problems

Jérôme Bolte, Tam Le, Edouard Pauwels, and Antonio Silveti-Falls



ISMP 2024 Montréal

- **Motivation**
- Conservative Gradients
- Results
- Applications
- Numerical Examples

Recall the LASSO problem:

$$\hat{x} \in \operatorname{argmin}_{x \in \mathbb{R}^p} \frac{1}{2} \|Ax - b\|_2^2 + e^\theta \|x\|_1.$$

Recall the LASSO problem:

$$\hat{x} \in \operatorname{argmin}_{x \in \mathbb{R}^p} \frac{1}{2} \|Ax - b\|_2^2 + e^\theta \|x\|_1.$$

Here, $A \in \mathbb{R}^{n \times p}$ is the design matrix for the n observations, $b \in \mathbb{R}^n$ is the associated response, and $\theta \in \mathbb{R}$ is a parameter.

Recall the LASSO problem:

$$\hat{x} \in \operatorname{argmin}_{x \in \mathbb{R}^p} \frac{1}{2} \|Ax - b\|_2^2 + e^\theta \|x\|_1.$$

Here, $A \in \mathbb{R}^{n \times p}$ is the design matrix for the n observations, $b \in \mathbb{R}^n$ is the associated response, and $\theta \in \mathbb{R}$ is a parameter.

Given some measure of task performance $C(\hat{x}(\theta))$, how to pick the "best" value of θ ?

The problem of choosing θ becomes a bilevel optimization problem:

$$\min_{\theta \in \mathbb{R}} C(\hat{x}(\theta)) \quad \text{such that} \quad \hat{x} \in \operatorname{argmin}_{x \in \mathbb{R}^p} \frac{1}{2} \|Ax - b\|_2^2 + e^\theta \|x\|_1.$$

The problem of choosing θ becomes a bilevel optimization problem:

$$\min_{\theta \in \mathbb{R}} C(\hat{x}(\theta)) \quad \text{such that} \quad \hat{x} \in \operatorname{argmin}_{x \in \mathbb{R}^p} \frac{1}{2} \|Ax - b\|_2^2 + e^\theta \|x\|_1.$$

If C and \hat{x} are smooth then we can use first-order optimization methods using the gradient:

$$\nabla C(\hat{x}(\theta)) = J_{\hat{x}}(\theta)^T \nabla_x C(\hat{x}(\theta)).$$

The problem of choosing θ becomes a bilevel optimization problem:

$$\min_{\theta \in \mathbb{R}} C(\hat{x}(\theta)) \quad \text{such that} \quad \hat{x} \in \operatorname{argmin}_{x \in \mathbb{R}^p} \frac{1}{2} \|Ax - b\|_2^2 + e^\theta \|x\|_1.$$

If C and \hat{x} are smooth then we can use first-order optimization methods using the gradient:

$$\nabla C(\hat{x}(\theta)) = J_{\hat{x}}(\theta)^T \nabla_x C(\hat{x}(\theta)).$$

However, $\hat{x}(\cdot)$ might not be smooth (often the case in machine learning settings). We need a method to derive functions like \hat{x} which are implicitly defined.


Classical Implicit Function Theorem

Consider the smooth function

$$F(x, y) = x^2 + y^2 - 1$$

and the equation

$$F(x, y) = 0.$$



Classical Implicit Function Theorem


Consider the smooth function

$$F(x, y) = x^2 + y^2 - 1$$

and the equation

$$F(x, y) = 0.$$

- Can we find a function $y = G(x)$ so that $F(x, G(x)) = 0$?



Classical Implicit Function Theorem


Consider the smooth function

$$F(x, y) = x^2 + y^2 - 1$$

and the equation

$$F(x, y) = 0.$$

- Can we find a function $y = G(x)$ so that $F(x, G(x)) = 0$?
- Can we compute the gradient of G ?



Classical Implicit Function Theorem


Consider the smooth function

$$F(x, y) = x^2 + y^2 - 1$$

and the equation

$$F(x, y) = 0.$$

- Can we find a function $y = G(x)$ so that $F(x, G(x)) = 0$?
- Can we compute the gradient of G ?



Classical Implicit Function Theorem

Consider the smooth function

$$F(x, y) = x^2 + y^2 - 1$$


and the equation

$$F(x, y) = 0.$$

- Can we find a function $y = G(x)$ so that $F(x, G(x)) = 0$?

Existence

- Can we compute the gradient of G ?



Classical Implicit Function Theorem

Consider the smooth function

$$F(x, y) = x^2 + y^2 - 1$$

and the equation


$$F(x, y) = 0.$$

- Can we find a function $y = G(x)$ so that $F(x, G(x)) = 0$?

Existence

- Can we compute the gradient of G ?

Calculus



In the nonsmooth world...

Existence: can we find implicit functions in the nonsmooth setting?

Calculus: can we compute the corresponding “gradient-like” objects in this setting?

In the nonsmooth world...

Existence: can we find implicit functions in the nonsmooth setting?

Yes! See (Clarke 1976, Hiriart-Urruty 1979, etc.) for locally Lipschitz functions.

Calculus: can we compute the corresponding “gradient-like” objects in this setting?

In the nonsmooth world...

Existence: can we find implicit functions in the nonsmooth setting?

Yes! See (Clarke 1976, Hiriart-Urruty 1979, etc.) for locally Lipschitz functions.

Calculus: can we compute the corresponding “gradient-like” objects in this setting?

Not with past theorems (Clarke, etc) - possible with **conservative Jacobians** [Bolte, Pauwels 2021] and **path differentiable functions**.

In the nonsmooth world...

Existence: can we find implicit functions in the nonsmooth setting?

Yes! See (Clarke 1976, Hiriart-Urruty 1979, etc.) for locally Lipschitz functions.

Calculus: can we compute the corresponding “gradient-like” objects in this setting?

Not with past theorems (Clarke, etc) - possible with **conservative Jacobians** [Bolte, Pauwels 2021] and **path differentiable functions**.

In practice one hopes for an algorithm of the form

$$x^+ = x - \gamma d(x)$$

where $d(x)$ is some descent direction or surrogate “gradient”.


Theorem (Clarke 1976, Hiriart-Urruty 1979)

Let $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be locally Lipschitz and $(\hat{x}, \hat{y}) \in \mathbb{R}^n \times \mathbb{R}^m$ such that

$$F(\hat{x}, \hat{y}) = 0.$$

If, $\forall [A \ B] \in \partial^c F(\hat{x}, \hat{y})$, B is invertible, then $\exists U \subset \mathbb{R}^n$ a neighborhood of \hat{x} and a locally Lipschitz function $G(x)$ so that

$$F(x, G(x)) = 0 \quad \forall x \in U.$$



Nonsmooth Implicit Function Theorem with Clarke Subdifferential


Theorem (Clarke 1976, Hiriart-Urruty 1979)

Let $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be locally Lipschitz and $(\hat{x}, \hat{y}) \in \mathbb{R}^n \times \mathbb{R}^m$ such that

$$F(\hat{x}, \hat{y}) = 0.$$

If, $\forall [A \ B] \in \partial^c F(\hat{x}, \hat{y})$, B is invertible, then $\exists U \subset \mathbb{R}^n$ a neighborhood of \hat{x} and a locally Lipschitz function $G(x)$ so that

$$F(x, G(x)) = 0 \quad \forall x \in U.$$



Nonsmooth Implicit Function Theorem with Clarke Subdifferential


Theorem (Clarke 1976, Hiriart-Urruty 1979)

Let $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be locally Lipschitz and $(\hat{x}, \hat{y}) \in \mathbb{R}^n \times \mathbb{R}^m$ such that

$$F(\hat{x}, \hat{y}) = 0.$$

If, $\forall [A \ B] \in \partial^c F(\hat{x}, \hat{y})$, B is invertible, then $\exists U \subset \mathbb{R}^n$ a neighborhood of \hat{x} and a locally Lipschitz function $G(x)$ so that

$$F(x, G(x)) = 0 \quad \forall x \in U.$$



Nonsmooth Implicit Function Theorem with Clarke Subdifferential


Theorem (Clarke 1976, Hiriart-Urruty 1979)

Let $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be locally Lipschitz and $(\hat{x}, \hat{y}) \in \mathbb{R}^n \times \mathbb{R}^m$ such that

$$F(\hat{x}, \hat{y}) = 0.$$

If, $\forall [A \ B] \in \partial^c F(\hat{x}, \hat{y})$, B is invertible, then $\exists U \subset \mathbb{R}^n$ a neighborhood of \hat{x} and a locally Lipschitz function $G(x)$ so that

$$F(x, G(x)) = 0 \quad \forall x \in U.$$



Let $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be locally Lipschitz and $(\hat{x}, \hat{y}) \in \mathbb{R}^n \times \mathbb{R}^m$ such that Clarke's IFT holds with implicit function $G(x)$.

Let $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be locally Lipschitz and $(\hat{x}, \hat{y}) \in \mathbb{R}^n \times \mathbb{R}^m$ such that Clarke's IFT holds with implicit function $G(x)$.

Recall from smooth IFT: $J_G(x) = -B^{-1}A$ $[A \ B] = J_F(x, G(x))$.

Let $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be locally Lipschitz and $(\hat{x}, \hat{y}) \in \mathbb{R}^n \times \mathbb{R}^m$ such that Clarke's IFT holds with implicit function $G(x)$.

Recall from smooth IFT: $J_G(x) = -B^{-1}A \quad [A \ B] = J_F(x, G(x))$.

Question

Can we have a **calculus** of the form:

$$\{-B^{-1}A : [A \ B] \in \partial^c F(\hat{x}, \hat{y})\} = \partial^c G(x)$$

for the Clarke Jacobian?

Let $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be locally Lipschitz and $(\hat{x}, \hat{y}) \in \mathbb{R}^n \times \mathbb{R}^m$ such that Clarke's IFT holds with implicit function $G(x)$.

Recall from smooth IFT: $J_G(x) = -B^{-1}A \quad [A \ B] = J_F(x, G(x))$.

Question

Can we have a **calculus** of the form:

$$\{-B^{-1}A : [A \ B] \in \partial^c F(\hat{x}, \hat{y})\} = \partial^c G(x)$$

for the Clarke Jacobian?

No - need something beyond ∂^c .

Let $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be locally Lipschitz and $(\hat{x}, \hat{y}) \in \mathbb{R}^n \times \mathbb{R}^m$ such that Clarke's IFT holds with implicit function $G(x)$.

Recall from smooth IFT: $J_G(x) = -B^{-1}A \quad [A \ B] = J_F(x, G(x))$.

Question

Can we have a **calculus** of the form:

$$\{-B^{-1}A : [A \ B] \in \partial^c F(\hat{x}, \hat{y})\} = \partial^c G(x)$$

for the Clarke Jacobian?


No - need something beyond ∂^c .

\exists piecewise linear $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ for which Clarke's inverse mapping theorem fails:

$$\exists M \in J_F^c(0,0) \quad \text{such that} \quad M^{-1} \notin J_{F^{-1}}^c(0,0)$$

Counterexample

$$F(x, y) = (|x| + y, 2x + |y|)$$



F^{-1} is linear on each region A, B, C, & D.

- Motivation
- **Conservative Gradients**
- Results
- Applications
- Numerical Examples

Definition (Conservative field (Bolte, Pauwels 2019))

A set valued mapping $D_F: \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ is a **conservative field** (or **conservative Jacobian**) for $F: \mathbb{R}^n \rightarrow \mathbb{R}$ locally Lipschitz if:

Definition (Conservative field (Bolte, Pauwels 2019))

A set valued mapping $D_F: \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ is a **conservative field** (or **conservative Jacobian**) for $F: \mathbb{R}^n \rightarrow \mathbb{R}$ locally Lipschitz if:

- 1 For all $x \in \mathbb{R}^n$, $D_F(x)$ is nonempty (ideally convex!).

Definition (Conservative field (Bolte, Pauwels 2019))

A set valued mapping $D_F: \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ is a **conservative field** (or **conservative Jacobian**) for $F: \mathbb{R}^n \rightarrow \mathbb{R}$ locally Lipschitz if:

- 1 For all $x \in \mathbb{R}^n$, $D_F(x)$ is nonempty (ideally convex!).
- 2 D_F has a closed graph and is locally bounded.

Definition (Conservative field (Bolte, Pauwels 2019))

A set valued mapping $D_F: \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ is a **conservative field** (or **conservative Jacobian**) for $F: \mathbb{R}^n \rightarrow \mathbb{R}$ locally Lipschitz if:

- ❶ For all $x \in \mathbb{R}^n$, $D_F(x)$ is nonempty (ideally convex!).
- ❷ D_F has a closed graph and is locally bounded.
- ❸ For any absolutely continuous curve $\gamma: [0, 1] \rightarrow \mathbb{R}^n$,

$$\frac{d}{dt} F(\gamma(t)) = \langle u, \dot{\gamma}(t) \rangle \quad \forall u \in D_F(\gamma(t))$$

for almost all $t \in [0, 1]$.

We call F **path differentiable**.

Definition (Conservative field (Bolte, Pauwels 2019))

A set valued mapping $D_F: \mathbb{R}^n \rightrightarrows \mathbb{R}^n$ is a **conservative field** (or **conservative Jacobian**) for $F: \mathbb{R}^n \rightarrow \mathbb{R}$ locally Lipschitz if:

- 1 For all $x \in \mathbb{R}^n$, $D_F(x)$ is nonempty (ideally convex!).
- 2 D_F has a closed graph and is locally bounded.
- 3 For any absolutely continuous curve $\gamma: [0, 1] \rightarrow \mathbb{R}^n$,

$$\frac{d}{dt} F(\gamma(t)) = \langle u, \dot{\gamma}(t) \rangle \quad \forall u \in D_F(\gamma(t))$$

for almost all $t \in [0, 1]$.

We call F **path differentiable**.


Take home message: **conservative fields/Jacobians** faithfully model what is computed by backpropagation in practice.

Visualizing Conservative Fields

For any absolutely continuous curve $\gamma : [0, 1] \rightarrow \mathbb{R}^n$,

$$\frac{d}{dt} F(\gamma(t)) = \langle u, \dot{\gamma}(t) \rangle$$

for all $u \in D_F(\gamma(t))$,
for almost all $t \in [0, 1]$.




Visualizing Conservative Fields

For any absolutely continuous curve $\gamma : [0, 1] \rightarrow \mathbb{R}^n$,

$$\frac{d}{dt} F(\gamma(t)) = \langle u, \dot{\gamma}(t) \rangle$$

for all $u \in D_F(\gamma(t))$,
for almost all $t \in [0, 1]$.




Visualizing Conservative Fields

For any absolutely continuous curve $\gamma : [0, 1] \rightarrow \mathbb{R}^n$,

$$\frac{d}{dt} F(\gamma(t)) = \langle u, \dot{\gamma}(t) \rangle$$

for all $u \in D_F(\gamma(t))$,
for almost all $t \in [0, 1]$.




Visualizing Conservative Fields

For any absolutely continuous curve $\gamma : [0, 1] \rightarrow \mathbb{R}^n$,

$$\frac{d}{dt} F(\gamma(t)) = \langle u, \dot{\gamma}(t) \rangle$$

for all $u \in D_F(\gamma(t))$,
for almost all $t \in [0, 1]$.




Visualizing Conservative Fields

For any absolutely continuous curve $\gamma : [0, 1] \rightarrow \mathbb{R}^n$,

$$\frac{d}{dt} F(\gamma(t)) = \langle u, \dot{\gamma}(t) \rangle$$

for all $u \in D_F(\gamma(t))$,
for almost all $t \in [0, 1]$.



- Motivation
- Conservative Gradients
- **Results**
- Applications
- Numerical Examples

Theorem (Bolte, Le, Pauwels, S.F. 2021)

Let $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be *path diff.* and $(\hat{x}, \hat{y}) \in \mathbb{R}^n \times \mathbb{R}^m$ be such that

$$F(\hat{x}, \hat{y}) = 0.$$


Assume $D_F(\hat{x}, \hat{y})$ is convex and $\forall [A \ B] \in D_F(\hat{x}, \hat{y})$, B is invertible. Then $\exists U \subset \mathbb{R}^n$ a neighborhood of \hat{x} and a *path diff.* function G such that

$$\forall x \in U \quad F(x, G(x)) = 0.$$

A conservative Jacobian of G is

$$D_G(x) = \{-B^{-1}A : [A \ B] \in D_F(x, G(x))\}$$

and is compatible with backprop.



Path Differentiable Nonsmooth Implicit Function Theorem

Theorem (Bolte, Le, Pauwels, S.F. 2021)

Let $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be *path diff.* and $(\hat{x}, \hat{y}) \in \mathbb{R}^n \times \mathbb{R}^m$ be such that

$$F(\hat{x}, \hat{y}) = 0.$$


Assume $D_F(\hat{x}, \hat{y})$ is convex and $\forall [A \ B] \in D_F(\hat{x}, \hat{y})$, B is invertible. Then $\exists U \subset \mathbb{R}^n$ a neighborhood of \hat{x} and a *path diff.* function G such that

$$\forall x \in U \quad F(x, G(x)) = 0.$$

A conservative Jacobian of G is

$$D_G(x) = \{-B^{-1}A : [A \ B] \in D_F(x, G(x))\}$$

and is compatible with backprop.



Path Differentiable Nonsmooth Implicit Function Theorem

Theorem (Bolte, Le, Pauwels, S.F. 2021)

Let $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be *path diff.* and $(\hat{x}, \hat{y}) \in \mathbb{R}^n \times \mathbb{R}^m$ be such that

$$F(\hat{x}, \hat{y}) = 0.$$


Assume $D_F(\hat{x}, \hat{y})$ is convex and $\forall [A \ B] \in D_F(\hat{x}, \hat{y})$, B is invertible. Then $\exists U \subset \mathbb{R}^n$ a neighborhood of \hat{x} and a *path diff.* function G such that

$$\forall x \in U \quad F(x, G(x)) = 0.$$

A conservative Jacobian of G is

$$D_G(x) = \{-B^{-1}A : [A \ B] \in D_F(x, G(x))\}$$

and is compatible with backprop.



Theorem (Bolte, Le, Pauwels, S.F. 2021)

Let $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be path diff. and $(\hat{x}, \hat{y}) \in \mathbb{R}^n \times \mathbb{R}^m$ be such that

$$F(\hat{x}, \hat{y}) = 0.$$


Assume $D_F(\hat{x}, \hat{y})$ is convex and $\forall [A \ B] \in D_F(\hat{x}, \hat{y})$, B is invertible. Then $\exists U \subset \mathbb{R}^n$ a neighborhood of \hat{x} and a path diff. function G such that

$$\forall x \in U \quad F(x, G(x)) = 0.$$

A conservative Jacobian of G is

$$D_G(x) = \{-B^{-1}A : [A \ B] \in D_F(x, G(x))\}$$

and is compatible with backprop.



Theorem (Bolte, Le, Pauwels, S.F. 2021)

Let $F : \mathbb{R}^n \times \mathbb{R}^m \rightarrow \mathbb{R}^m$ be *path diff.* and $(\hat{x}, \hat{y}) \in \mathbb{R}^n \times \mathbb{R}^m$ be such that

$$F(\hat{x}, \hat{y}) = 0.$$


Assume $D_F(\hat{x}, \hat{y})$ is convex and $\forall [A \ B] \in D_F(\hat{x}, \hat{y})$, B is invertible. Then $\exists U \subset \mathbb{R}^n$ a neighborhood of \hat{x} and a *path diff.* function G such that

$$\forall x \in U \quad F(x, G(x)) = 0.$$

A conservative Jacobian of G is

$$D_G(x) = \{-B^{-1}A : [A \ B] \in D_F(x, G(x))\}$$

and is compatible with backprop.



- Motivation
- Conservative Gradients
- Results
- **Applications**
- Numerical Examples

With our new theorem we can answer the question:

how to differentiate the solution to a nonsmooth convex optimization problem ?

$$\hat{x}(\theta) := \operatorname{argmin}_{x \in \mathbb{R}^n} f(x, \theta)$$

With our new theorem we can answer the question:

how to differentiate the solution to a nonsmooth convex optimization problem ?

$$\hat{x}(\theta) := \operatorname{argmin}_{x \in \mathbb{R}^n} f(x, \theta)$$

Solution: find a necessary and sufficient **path differentiable** optimality condition and apply our **implicit function theorem**.

With our new theorem we can answer the question:

how to differentiate the solution to a nonsmooth convex optimization problem ?

$$\hat{x}(\theta) := \operatorname{argmin}_{x \in \mathbb{R}^n} f(x, \theta)$$

Solution: find a necessary and sufficient **path differentiable** optimality condition and apply our **implicit function theorem**.

- Hyperparameter tuning of the LASSO [Bertrand, Klopfenstein, Blondel, Vaiter, Gramfort, Salmon 2020].

With our new theorem we can answer the question:

how to differentiate the solution to a nonsmooth convex optimization problem ?

$$\hat{x}(\theta) := \operatorname{argmin}_{x \in \mathbb{R}^n} f(x, \theta)$$

Solution: find a necessary and sufficient **path differentiable** optimality condition and apply our **implicit function theorem**.

- Hyperparameter tuning of the LASSO [[Bertrand, Klopfenstein, Blondel, Vaiter, Gramfort, Salmon 2020](#)].
- Differentiating monotone inclusions with protodifferentiability [[Adly, Rockafellar 2021](#)].

With our new theorem we can answer the question:

how to differentiate the solution to a nonsmooth convex optimization problem ?

$$\hat{x}(\theta) := \operatorname{argmin}_{x \in \mathbb{R}^n} f(x, \theta)$$

Solution: find a necessary and sufficient **path differentiable** optimality condition and apply our **implicit function theorem**.

- Hyperparameter tuning of the LASSO [[Bertrand, Klopfenstein, Blondel, Vaiter, Gramfort, Salmon 2020](#)].
- Differentiating monotone inclusions with protodifferentiability [[Adly, Rockafellar 2021](#)].
- Set-valued implicit function theorems + semismooth localizations [[Gferer, Outrata 2024](#)].
- etc.

We recall the LASSO problem:

$$\hat{x} \in \operatorname{argmin}_{x \in \mathbb{R}^p} \frac{1}{2} \|Ax - b\|_2^2 + e^\theta \|x\|_1 .$$

We recall the LASSO problem:

$$\hat{x} \in \operatorname{argmin}_{x \in \mathbb{R}^p} \frac{1}{2} \|Ax - b\|_2^2 + e^\theta \|x\|_1.$$

A fixed point condition for optimality:

$$\underbrace{\operatorname{prox}_{e^\theta \|\cdot\|_1}(\hat{x} - A^T(A\hat{x} - b)) - \hat{x}}_{F(\theta, x)} = 0.$$

the proximal mapping here is simply the “soft thresholding” operator, which is **path differentiable**. Thus, the function F is **path differentiable**.

It is necessary to verify the **invertibility** of the conservative Jacobian of F with respect to x evaluated at (θ, \hat{x}) .

It is necessary to verify the **invertibility** of the conservative Jacobian of F with respect to x evaluated at (θ, \hat{x}) .

We define the *equicorrelation* set:

$$\mathcal{E} := \{j \in \{1, \dots, p\} : |A_j^T(b - A\hat{x}(\theta))| = e^\theta\}.$$

It is necessary to verify the **invertibility** of the conservative Jacobian of F with respect to x evaluated at (θ, \hat{x}) .

We define the *equicorrelation* set:

$$\mathcal{E} := \{j \in \{1, \dots, p\} : |A_j^T(b - A\hat{x}(\theta))| = e^\theta\}.$$

- The set \mathcal{E} does NOT depend on the choice of solution $\hat{x}(\theta)$ [Tibshirani 2013].

It is necessary to verify the **invertibility** of the conservative Jacobian of F with respect to x evaluated at (θ, \hat{x}) .

We define the *equicorrelation* set:

$$\mathcal{E} := \{j \in \{1, \dots, p\} : |A_j^T(b - A\hat{x}(\theta))| = e^\theta\}.$$

- The set \mathcal{E} does NOT depend on the choice of solution $\hat{x}(\theta)$ [Tibshirani 2013].
- The set \mathcal{E} contains the *support* set $\mathcal{S} := \{i \in \{1, \dots, p\} : \hat{x}_i(\theta) \neq 0\}$.

Verification of Invertibility - Example With LASSO

It is necessary to verify the **invertibility** of the conservative Jacobian of F with respect to x evaluated at (θ, \hat{x}) .

We define the *equicorrelation* set:

$$\mathcal{E} := \{j \in \{1, \dots, p\} : |A_j^T(b - A\hat{x}(\theta))| = e^\theta\}.$$


- The set \mathcal{E} does NOT depend on the choice of solution $\hat{x}(\theta)$ [Tibshirani 2013].
- The set \mathcal{E} contains the *support* set $\mathcal{S} := \{i \in \{1, \dots, p\} : \hat{x}_i(\theta) \neq 0\}$.

Proposition (Prop. 5 [Bolte, Le, Pauwels, S.F. 21])


Define, $\forall \theta \in \mathbb{R}$, the matrix $A_\mathcal{E}$ by taking the columns of A indexed by \mathcal{E} . If, $\forall \theta \in \mathbb{R}$, **the matrix $A_\mathcal{E}^T A_\mathcal{E}$ is full rank**, then $\hat{x}(\cdot)$ is a **path differentiable** function with a conservative Jacobian given by

$$D_{\hat{x}} : \theta \mapsto \left\{ \left[-e^\theta (\text{Id}_p - \text{diag}(q)(\text{Id}_p - A^T A))^{-1} \text{diag}(q) \text{sign}(\hat{x} - A^T(A\hat{x} - b)) \right] : q \in \mathcal{M}(\theta) \right\}$$

$$\text{where } \mathcal{M}(\theta) = \left\{ q : q_i \in \begin{cases} \{1\} & \text{if } i \in \mathcal{S} \\ [0, 1] & \text{if } i \in \mathcal{E} \setminus \mathcal{S} \\ \{0\} & \text{if } i \notin \mathcal{E} \end{cases} \right\}.$$



- Deep equilibrium networks [Bai, Kolter, Koltun 2019].
- Implicit networks [El Ghaoui, Gu, Travacca, Askari, Tsai 2019].
- Declarative networks [Gould, Hartley, Campbell 2019].
- Monotone deep equilibrium networks [Winston, Kolter 2020].
- Optimization layers (OptNET) [Amos, Kolter 2017].
- General convex optimization layers [Agrawal, Amos, Barratt, Boyd, Diamond, Kolter 2019].



- Deep equilibrium networks [Bai, Kolter, Koltun 2019].
- Implicit networks [El Ghaoui, Gu, Travacca, Askari, Tsai 2019].
- Declarative networks [Gould, Hartley, Campbell 2019].
- Monotone deep equilibrium networks [Winston, Kolter 2020].
- Optimization layers (OptNET) [Amos, Kolter 2017].
- General convex optimization layers [Agrawal, Amos, Barratt, Boyd, Diamond, Kolter 2019].

conservative Jacobians + path differentiable implicit function theorem
 \implies convergence guarantees (every acc. point is a Clarke stationary point almost surely, objective values converge) for these network types.

Consider two parametric maximal monotone operators \mathcal{A}_θ and \mathcal{B}_θ and the inclusion

$$0 \in \mathcal{A}_\theta(x^*) + \mathcal{B}_\theta(x^*)$$

where \mathcal{A}_θ is set-valued but \mathcal{B}_θ is Lipschitz continuous. **Note:** We assume that $x^*(\theta)$ is unique for each θ .

Consider two parametric maximal monotone operators \mathcal{A}_θ and \mathcal{B}_θ and the inclusion

$$0 \in \mathcal{A}_\theta(x^\star) + \mathcal{B}_\theta(x^\star)$$

where \mathcal{A}_θ is set-valued but \mathcal{B}_θ is Lipschitz continuous. **Note:** We assume that $x^\star(\theta)$ is unique for each θ . Fixed point equation for the monotone inclusion

$$\underbrace{R_{\gamma\mathcal{A}_\theta}(x - \gamma\mathcal{B}_\theta x)}_{H(\theta, x)} = x$$

We call H the **Forward-Backward** mapping. Formally we denote $T(\theta, x) := R_{\gamma\mathcal{A}_\theta}(x)$ and $S(\theta, x) := x - \gamma\mathcal{B}_\theta(x)$ the **forward** and **backward** maps which gives an equation we can apply the IFT to:

$$F(\theta, x) := H(\theta, x) - x = 0.$$

Consider two parametric maximal monotone operators \mathcal{A}_θ and \mathcal{B}_θ and the inclusion

$$0 \in \mathcal{A}_\theta(x^\star) + \mathcal{B}_\theta(x^\star)$$

where \mathcal{A}_θ is set-valued but \mathcal{B}_θ is Lipschitz continuous. **Note:** We assume that $x^\star(\theta)$ is unique for each θ . Fixed point equation for the monotone inclusion

$$\underbrace{R_{\gamma\mathcal{A}_\theta}(x - \gamma\mathcal{B}_\theta x)}_{H(\theta, x)} = x$$

We call H the **Forward-Backward** mapping. Formally we denote $T(\theta, x) := R_{\gamma\mathcal{A}_\theta}(x)$ and $S(\theta, x) := x - \gamma\mathcal{B}_\theta(x)$ the **forward** and **backward** maps which gives an equation we can apply the IFT to:

$$F(\theta, x) := H(\theta, x) - x = 0.$$

We will assume that F is **path differentiable** jointly in (θ, x) .

Beware: conservative Jacobians are **not** unique and **not** defined pointwise!

Example: path differentiable $f : \mathbb{R} \rightarrow \mathbb{R}$,
$$\tilde{\mathcal{J}}_f(x) = \begin{cases} \mathcal{J}_f(x) \cup \{1\} & x \in \mathbb{N} \\ \mathcal{J}_f(x) & x \notin \mathbb{N} \end{cases}$$

Beware: conservative Jacobians are **not** unique and **not** defined pointwise!

Example: path differentiable $f : \mathbb{R} \rightarrow \mathbb{R}$,
$$\tilde{\mathcal{J}}_f(x) = \begin{cases} \mathcal{J}_f(x) \cup \{1\} & x \in \mathbb{N} \\ \mathcal{J}_f(x) & x \notin \mathbb{N} \end{cases}$$

We must make a choice for which **conservative Jacobian** to consider - it should reflect what is computed in practice and also be theoretically accessible.

Beware: conservative Jacobians are **not** unique and **not** defined pointwise!

$$\text{Example: path differentiable } f : \mathbb{R} \rightarrow \mathbb{R}, \quad \tilde{\mathcal{J}}_f(x) = \begin{cases} \mathcal{J}_f(x) \cup \{1\} & x \in \mathbb{N} \\ \mathcal{J}_f(x) & x \notin \mathbb{N} \end{cases}$$

We must make a choice for which conservative Jacobian to consider - it should reflect what is computed in practice and also be theoretically accessible.

We take the product of Clarke Jacobians of the forward and backward maps giving

$$\begin{aligned} \mathcal{J}_{H_\theta}(\theta, x) &= \text{Jac}_{\mathcal{T}}^c(S(\theta, x)) \times \text{Jac}_{\mathcal{S}}^c(\theta, x) \\ &= \{ [A \ B] \times \begin{bmatrix} \text{Id}_p & 0 \\ -C & \text{Id}_n - \gamma D \end{bmatrix} : [A \ B] \in \text{Jac}_{\mathcal{T}}^c(\theta, x - \gamma \mathcal{B}_\theta(x)), \\ &\quad [C \ D] \in \text{Jac}_{\mathcal{B}}^c(\theta, x) \} \\ &= \{ [A - BC \ B(\text{Id}_n - \gamma D)] : [A \ B] \in \text{Jac}_{\mathcal{T}}^c(\theta, x - \gamma \mathcal{B}_\theta(x)), [C \ D] \in \text{Jac}_{\mathcal{B}}^c(\theta, x) \} \end{aligned}$$

which is a conservative Jacobian for H .

Theorem (Bolte, Pauwels, S.F. 2024)

Assume that \mathcal{B}_θ is β -Lipschitz continuous and that either \mathcal{A}_θ or \mathcal{B}_θ is α -strongly monotone, for some $\alpha, \beta > 0$, uniformly in θ . For $\gamma \in (0, \frac{2\alpha}{(\alpha+\beta)^2})$, the invertibility condition holds and x^ is path differentiable with a conservative Jacobian whose formula is computable from $\mathcal{J}_H(x^*(\theta))$.*

Theorem (Bolte, Pauwels, S.F. 2024)

Assume that \mathcal{B}_θ is β -Lipschitz continuous and that either \mathcal{A}_θ or \mathcal{B}_θ is α -strongly monotone, for some $\alpha, \beta > 0$, uniformly in θ . For $\gamma \in (0, \frac{2\alpha}{(\alpha+\beta)^2})$, the invertibility condition holds and x^* is path differentiable with a conservative Jacobian whose formula is computable from $\mathcal{J}_H(x^*(\theta))$.

Proof.

If \mathcal{A}_θ or \mathcal{B}_θ is α -strongly monotone, then either \mathcal{T} or \mathcal{S} is a strict contraction, and we can choose γ to ensure that the composition H is a strict contraction. Then, the product of Clarke Jacobians will have norm < 1 . □

Theorem (Bolte, Pauwels, S.F. 2024)

Assume that \mathcal{B}_θ is β -Lipschitz continuous and that either \mathcal{A}_θ or \mathcal{B}_θ is α -strongly monotone, for some $\alpha, \beta > 0$, uniformly in θ . For $\gamma \in (0, \frac{2\alpha}{(\alpha+\beta)^2})$, the invertibility condition holds and x^* is path differentiable with a conservative Jacobian whose formula is computable from $\mathcal{J}_H(x^*(\theta))$.

Proof.

If \mathcal{A}_θ or \mathcal{B}_θ is α -strongly monotone, then either \mathcal{T} or \mathcal{S} is a strict contraction, and we can choose γ to ensure that the composition H is a strict contraction. Then, the product of Clarke Jacobians will have norm < 1 . \square

Corollary

The solution to the optimization problem

$$\min_{x \in \mathbb{R}^p} f_\theta(x) + g_\theta(x),$$

where f_θ is β -Lipschitz smooth and g_θ is nonsmooth, is path differentiable if either f_θ or g_θ is α -strongly convex.

- Motivation
- Conservative Gradients
- Results
- Applications
- **Numerical Examples**

Piecewise quadratic objective function posed as a bilevel problem:


$$\min_{x,y,s} (x - s_1)^2 + 4(y - s_2)^2 \quad \text{such that}$$

$$s \in \arg \max \left\{ (a + b)(-2x + y + 2) : a \in [0, 3], b \in [0, 5] \right\}$$

Piecewise quadratic objective function posed as a bilevel problem:

$$\min_{x,y,s} (x - s_1)^2 + 4(y - s_2)^2 \quad \text{such that}$$

$$s \in \arg \max \{ (a + b)(-2x + y + 2) : a \in [0, 3], b \in [0, 5] \}$$





Pathological Examples - Cyclic Trajectories

Piecewise quadratic objective function posed as a bilevel problem:

$$\min_{x,y,s} (x - s_1)^2 + 4(y - s_2)^2 \quad \text{such that}$$

$$s \in \arg \max \left\{ (a + b)(-2x + y + 2) : a \in [0, 3], b \in [0, 5] \right\}$$





Pathological Examples - Cyclic Trajectories

Piecewise quadratic objective function posed as a bilevel problem:

$$\min_{x,y,s} (x - s_1)^2 + 4(y - s_2)^2 \quad \text{such that}$$

$$s \in \arg \max \{ (a + b)(-2x + y + 2) : a \in [0, 3], b \in [0, 5] \}$$



$$\text{Let } L(u) = L(x, y, z) = \left(10(y - x), x(28 - z) - y, xy - \frac{8}{3}z\right)$$

Explicit formulation

$$\max_{u \in \mathbb{R}^3} u^T L(u)$$



Implicit formulation

$$\begin{aligned} \max_{u \in \mathbb{R}^3} u^T z \quad \text{such that} \\ z \in \operatorname{argmin}_{s \in \mathbb{R}^3} \|s - L(u)\|^4 \end{aligned}$$

Pathological Examples - Optimizing a Quadratic Two Ways

$$\text{Let } L(u) = L(x, y, z) = (10(y - x), x(28 - z) - y, xy - \frac{8}{3}z)$$


Explicit formulation

$$\max_{u \in \mathbb{R}^3} u^T L(u)$$



Implicit formulation

$$\begin{aligned} \max_{u \in \mathbb{R}^3} \quad & u^T z \quad \text{such that} \\ & z \in \operatorname{argmin}_{s \in \mathbb{R}^3} \|s - L(u)\|^4 \end{aligned}$$



Pathological Examples - Optimizing a Quadratic Two Ways

$$\text{Let } L(u) = L(x, y, z) = (10(y - x), x(28 - z) - y, xy - \frac{8}{3}z)$$



Explicit formulation

$$\max_{u \in \mathbb{R}^3} u^T L(u)$$



Implicit formulation

$$\max_{u \in \mathbb{R}^3} u^T z \quad \text{such that} \\ z \in \operatorname{argmin}_{s \in \mathbb{R}^3} \|s - L(u)\|^4$$



Nonsmooth implicit differentiation:

- Can we have a calculus for implicitly defined functions in the locally Lipschitz setting?

Nonsmooth implicit differentiation:

- Can we have a calculus for implicitly defined functions in the locally Lipschitz setting?
- Using the Clarke subdifferential: **No**.
 - ▶ The inverse of a Clarke Jacobian is not necessarily a Clarke Jacobian.

Nonsmooth implicit differentiation:

- Can we have a calculus for implicitly defined functions in the locally Lipschitz setting?
- Using the Clarke subdifferential: **No**.
 - ▶ The inverse of a Clarke Jacobian is not necessarily a Clarke Jacobian.
- Using conservative gradients **Yes**.
 - ▶ The inverse of a conservative Jacobian is again a conservative Jacobian.

Nonsmooth implicit differentiation:

- Can we have a calculus for implicitly defined functions in the locally Lipschitz setting?
- Using the Clarke subdifferential: **No**.
 - ▶ The inverse of a Clarke Jacobian is not necessarily a Clarke Jacobian.
- Using conservative gradients **Yes**.
 - ▶ The inverse of a conservative Jacobian is again a conservative Jacobian.

Practical implications:

- Method to compute the gradient of solutions to convex optimization problems.
- Applications in machine learning (bilevel hyperparameter tuning, implicit neural networks, ...).

**Nonsmooth Implicit Differentiation for Machine Learning
(NeurIPS, 2021)**

Jérôme Bolte, Tâm Lê, Edouard Pauwels, Antonio Silveti-Falls
<https://arxiv.org/abs/2106.04350>

**Differentiating Nonsmooth Solutions to Parametric Monotone Inclusion Problems
(SIAM Optimization, 2024)**

Jérôme Bolte, Edouard Pauwels, Antonio Silveti-Falls
<https://arxiv.org/abs/2212.07844>

N data points, L layers:

$$\min_{w \in \mathbb{R}^p} \ell(w) := \frac{1}{N} \sum_{i=1}^N \ell_i(w) \quad \text{with} \quad \ell_i := g_{i,L} \circ g_{i,L-1} \circ \dots \circ g_{i,1}$$

Each layer $g_{i,j}$ is semialgebraic (or definable) and path differentiable - can be explicit or implicit.

$N = 1, L = 2$ recovers bilevel optimization problem setting.

Define

$$w_{k+1} = w_k - s \alpha_k v_k \quad v_k \in J_{l_k}(w_k)$$

for $(\alpha_k)_{k \in \mathbb{N}} \in \ell^1 \setminus \ell^2$

For almost all w_0 , for almost all $s \in (s_{\min}, s_{\max})$, $\ell(w_k)$ converges and all acc. points of $(w_k)_{k \in \mathbb{N}}$ are clarke critical.