

The gradient's limit of a definable family of functions is a conservative field

Sholom Schechtman



IP PARIS

Introduction

Let $(f_a)_{a>0}$ be a family of smooth functions and $F : \mathbb{R}^d \rightarrow \mathbb{R}$ be smooth such that

$$\|f_a - F\| \xrightarrow{a \rightarrow 0} 0.$$

Does the following holds?

$$\|\nabla f_a - \nabla F\| \xrightarrow[a \rightarrow 0]{?} 0$$

Introduction

More formally,

$$\|f_a - F\| \xrightarrow{a \rightarrow 0} 0.$$

Introduction

More formally,

$$\|f_a - F\| \xrightarrow{a \rightarrow 0} 0.$$

Denote the gradients **limits**:

$$D_F(x) = \{v \in \mathbb{R}^d : x_n \rightarrow x, a_n \rightarrow 0, \nabla f_{a_n}(x_n) \rightarrow v\}.$$

Introduction

More formally,

$$\|f_a - F\| \xrightarrow{a \rightarrow 0} 0.$$

Denote the gradients **limits**:

$$D_F(x) = \{v \in \mathbb{R}^d : x_n \rightarrow x, a_n \rightarrow 0, \nabla f_{a_n}(x_n) \rightarrow v\}.$$

Does the following holds?

$$D_F(x) = \{\nabla F(x)\}.$$

Introduction

More generally

$f_a \xrightarrow{a \rightarrow 0} F$, with $f_a : \mathbb{R}^d \rightarrow \mathbb{R}$ locally Lipschitz.

Introduction

More generally

$f_a \xrightarrow{a \rightarrow 0} F$, with $f_a : \mathbb{R}^d \rightarrow \mathbb{R}$ locally Lipschitz.

$D_F(x) = \{v \in \mathbb{R}^d : \text{there is } (x_n, v_n, a_n) \rightarrow (x, v, 0) \text{ with } v_n \in \partial f_{a_n}(x_n)\}.$

→ $\partial f_a(x)$ some **subgradient** (convex, Frechet, Clarke) of f_a at x .

Introduction

More generally

$f_a \xrightarrow{a \rightarrow 0} F$, with $f_a : \mathbb{R}^d \rightarrow \mathbb{R}$ locally Lipschitz.

$D_F(x) = \{v \in \mathbb{R}^d : \text{there is } (x_n, v_n, a_n) \rightarrow (x, v, 0) \text{ with } v_n \in \partial f_{a_n}(x_n)\}.$

→ $\partial f_a(x)$ some **subgradient** (convex, Frechet, Clarke) of f_a at x .

Question. What is the link between D_F and ∂F ?

Introduction

More generally

$f_a \xrightarrow{a \rightarrow 0} F$, with $f_a : \mathbb{R}^d \rightarrow \mathbb{R}$ locally Lipschitz.

$D_F(x) = \{v \in \mathbb{R}^d : \text{there is } (x_n, v_n, a_n) \rightarrow (x, v, 0) \text{ with } v_n \in \partial f_{a_n}(x_n)\}.$

→ $\partial f_a(x)$ some **subgradient** (convex, Frechet, Clarke) of f_a at x .

Question. What is the link between D_F and ∂F ?

- One would wish that D_F reduces to some **common** first-order operators.
 - To the **gradient** if F is smooth.
 - To the **subgradient** if F is not.

Introduction

$$f_a \rightarrow F$$

$$D_F(x) = \{v \in \mathbb{R}^d : \text{there is } (x_n, v_n, a_n) \rightarrow (x, v, 0) \text{ with } x_n \in \partial f_a(x_n)\}.$$

Question. Link between D_F and ∂F ?

Motivation.

Introduction

$$f_a \rightarrow F$$

$$D_F(x) = \{v \in \mathbb{R}^d : \text{there is } (x_n, v_n, a_n) \rightarrow (x, v, 0) \text{ with } x_n \in \partial f_a(x_n)\}.$$

Question. Link between D_F and ∂F ?

Motivation.

- 1) Interesting in its own right.

Introduction

$$f_a \rightarrow F$$

$$D_F(x) = \{v \in \mathbb{R}^d : \text{there is } (x_n, v_n, a_n) \rightarrow (x, v, 0) \text{ with } x_n \in \partial f_a(x_n)\}.$$

Question. Link between D_F and ∂F ?

Motivation.

- 1) Interesting in its own right.
- 2) Consequences for smoothing methods.

Smoothing methods

Optimization problem: $\min_{x \in \mathbb{R}^d} F(x)$.

- Might be hard if F is nonsmooth and nonconvex.

Smoothing methods

Optimization problem: $\min_{x \in \mathbb{R}^d} F(x)$.

- ▶ Might be hard if F is nonsmooth and nonconvex.
- ▶ Construct $f_a \rightarrow F$, with f_a smooth.

Smoothing methods:

Smoothing methods

Optimization problem: $\min_{x \in \mathbb{R}^d} F(x)$.

- ▶ Might be hard if F is nonsmooth and nonconvex.
- ▶ Construct $f_a \rightarrow F$, with f_a smooth.

Smoothing methods:

- ▶ For $a_k, \varepsilon_k > 0$,

Solve $\min_{x \in \mathbb{R}^d} f_{a_k}(x)$

Smoothing methods

Optimization problem: $\min_{x \in \mathbb{R}^d} F(x)$.

- ▶ Might be hard if F is nonsmooth and nonconvex.
- ▶ Construct $f_a \rightarrow F$, with f_a smooth.

Smoothing methods:

- ▶ For $a_k, \varepsilon_k > 0$,

$$\text{Solve } \min_{x \in \mathbb{R}^d} f_{a_k}(x) \implies \text{find } \|\nabla f_{a_k}(x_k)\| \leq \varepsilon_k.$$

Smoothing methods

Optimization problem: $\min_{x \in \mathbb{R}^d} F(x)$.

- ▶ Might be hard if F is nonsmooth and nonconvex.
- ▶ Construct $f_a \rightarrow F$, with f_a smooth.

Smoothing methods:

- ▶ For $a_k, \varepsilon_k > 0$,

$$\text{Solve } \min_{x \in \mathbb{R}^d} f_{a_k}(x) \implies \text{find } \|\nabla f_{a_k}(x_k)\| \leq \varepsilon_k.$$

→ Easier because f_{a_k} is smooth.

Smoothing methods

Optimization problem: $\min_{x \in \mathbb{R}^d} F(x)$.

- ▶ Might be hard if F is nonsmooth and nonconvex.
- ▶ Construct $f_a \rightarrow F$, with f_a smooth.

Smoothing methods:

- ▶ For $a_k, \varepsilon_k > 0$,

$$\text{Solve } \min_{x \in \mathbb{R}^d} f_{a_k}(x) \implies \text{find } \|\nabla f_{a_k}(x_k)\| \leq \varepsilon_k.$$

- Easier because f_{a_k} is smooth.
- ▶ Decrease a_k and ε_k and repeat.

Smoothing methods

Optimization problem: $\min_{x \in \mathbb{R}^d} F(x)$.

- ▶ Might be hard if F is nonsmooth and nonconvex.
- ▶ Construct $f_a \rightarrow F$, with f_a smooth.

Smoothing methods:

- ▶ For $a_k, \varepsilon_k > 0$,

$$\text{Solve } \min_{x \in \mathbb{R}^d} f_{a_k}(x) \implies \text{find } \|\nabla f_{a_k}(x_k)\| \leq \varepsilon_k.$$

→ Easier because f_{a_k} is smooth.

- ▶ Decrease a_k and ε_k and repeat.
- ▶ When $a_k, \varepsilon_k \rightarrow 0$ and $x_k \rightarrow x^*$

$$\lim_{k \rightarrow +\infty} \nabla f_{a_k}(x_k) = 0$$

Smoothing methods

Optimization problem: $\min_{x \in \mathbb{R}^d} F(x)$.

- ▶ Might be hard if F is nonsmooth and nonconvex.
- ▶ Construct $f_a \rightarrow F$, with f_a smooth.

Smoothing methods:

- ▶ For $a_k, \varepsilon_k > 0$,

$$\text{Solve } \min_{x \in \mathbb{R}^d} f_{a_k}(x) \implies \text{find } \|\nabla f_{a_k}(x_k)\| \leq \varepsilon_k.$$

→ Easier because f_{a_k} is smooth.

- ▶ Decrease a_k and ε_k and repeat.
- ▶ When $a_k, \varepsilon_k \rightarrow 0$ and $x_k \rightarrow x^*$

$$\lim_{k \rightarrow +\infty} \nabla f_{a_k}(x_k) = 0 \in D_F(x^*).$$

Smoothing methods

Optimization problem: $\min_{x \in \mathbb{R}^d} F(x)$.

- ▶ Might be hard if F is nonsmooth and nonconvex.
- ▶ Construct $f_a \rightarrow F$, with f_a smooth.

Smoothing methods:

- ▶ For $a_k, \varepsilon_k > 0$,

$$\text{Solve } \min_{x \in \mathbb{R}^d} f_{a_k}(x) \implies \text{find } \|\nabla f_{a_k}(x_k)\| \leq \varepsilon_k.$$

- Easier because f_{a_k} is smooth.
- ▶ Decrease a_k and ε_k and repeat.
- ▶ When $a_k, \varepsilon_k \rightarrow 0$ and $x_k \rightarrow x^*$

$$\lim_{k \rightarrow +\infty} \nabla f_{a_k}(x_k) = 0 \in D_F(x^*).$$

- Necessary optimality condition if $D_F = \nabla F$ or ∂F .

Known results

Attouch's theorem

If for every $a > 0$, f_a is **convex**. Then $D_F(x) = \partial F(x)$.

Attouch, 1977

- ▶ With $\partial F(x)$ being **convex subgradient** of F at x .
- ▶ Several extensions to “**approximately convex**” case.
 - Poliquin, 1992; Levy, Poliquin, and Thibault, 1995; Zolezzi, 1985; Zolezzi, 1994; Czarnecki and Rifford, 2006 ...

Nonconvex case?

Can we do more?

Not really.

Nonconvex case?

Can we do more?

Not really.

$$\rightarrow f_a(x) = a \sin\left(\frac{x}{a}\right).$$

$$\rightarrow \sin\left(\frac{x}{a}\right) \rightarrow [-1, 1].$$




Figure: $f_a(x)$ and $f'_a(x)$

Nonconvex case?



Main result

- ▶ Assume that for each $a > 0$, f_a is locally Lipschitz continuous ¹
- ▶ $f_a \rightarrow F$ uniformly on compact sets.

Theorem

If the family (f_a) is **definable**, then D_F is a **conservative set-valued field** of F (in the sense of **Bolte and Pauwels**).

Schechtman, 2024

¹Can be relaxed.

Main result

- Assume that for each $a > 0$, f_a is locally Lipschitz continuous ¹
- $f_a \rightarrow F$ uniformly on compact sets.

Theorem

If the family (f_a) is **definable**, then D_F is a **conservative set-valued field** of F (in the sense of **Bolte and Pauwels**).

Schechtman, 2024

Some **consequences**.

- For almost every $x \in \mathbb{R}^d$, $D_F(x) = \{\nabla F(x)\}$.

¹Can be relaxed.

Main result

- ▶ Assume that for each $a > 0$, f_a is locally Lipschitz continuous ¹
- ▶ $f_a \rightarrow F$ uniformly on compact sets.

Theorem

If the family (f_a) is **definable**, then D_F is a **conservative set-valued field** of F (in the sense of **Bolte and Pauwels**).

Schechtman, 2024

Some **consequences**.

- ▶ For almost every $x \in \mathbb{R}^d$, $D_F(x) = \{\nabla F(x)\}$.
- ▶ $\partial F(x) \subset \text{conv } D_F(x)$
 - ➔ $0 \in \text{conv } D_F(x)$ is a **necessary** optimality condition.

¹Can be relaxed.

Main result

- ▶ Assume that for each $a > 0$, f_a is locally Lipschitz continuous¹
- ▶ $f_a \rightarrow F$ uniformly on compact sets.

Theorem

If the family (f_a) is **definable**, then D_F is a **conservative set-valued field** of F (in the sense of Bolte and Pauwels).

Schechtman, 2024

Some **consequences**.

- ▶ For almost every $x \in \mathbb{R}^d$, $D_F(x) = \{\nabla F(x)\}$.
- ▶ $\partial F(x) \subset \text{conv } D_F(x)$
 - $0 \in \text{conv } D_F(x)$ is a **necessary** optimality condition.
- ▶ The **geometric structure** of D_F is well understood.

¹Can be relaxed.

Outline

- I Definable/tame functions**
- II Conservative set-valued fields
- III Convergence of gradients

Outline

I Definable/tame functions

II Conservative set-valued fields

III Convergence of gradients

Outline

I Definable/tame functions

II Conservative set-valued fields


III Convergence of gradients

Definable functions


A continuous function in \mathbb{R}^d can be pathological.

Definable functions

A continuous function in \mathbb{R}^d can be pathological.




(a) $x \mapsto x \sin(1/x)$




(b) Weierstrass function

Definable functions

A continuous function in \mathbb{R}^d can be **pathological**.



(a) $x \mapsto x \sin(1/x)$



(b) Weierstrass function

There are functions $F : [0, 1] \rightarrow [0, 1]$ such that $\forall x, \partial F(x) = [0, 1]$

Definable functions

A **class** of function that includes:

- ▶ **Semialgebraic**: defined by a finite number of polynomial inequalities.

Definable functions

A **class** of function that includes:

- ▶ **Semialgebraic**: defined by a finite number of polynomial inequalities.
 - Piecewise polynomials.

Definable functions

A **class** of function that includes:

- ▶ **Semialgebraic**: defined by a finite number of polynomial inequalities.
 - Piecewise polynomials.
 - Piecewise rational functions.

Definable functions

A **class** of function that includes:

- ▶ **Semialgebraic**: defined by a finite number of polynomial inequalities.
 - Piecewise polynomials.
 - Piecewise rational functions.
 - $x \mapsto \|x\|_q$, with $q \geq 0$ rational.

Definable functions

A **class** of function that includes:

- ▶ **Semialgebraic**: defined by a finite number of polynomial inequalities.
 - Piecewise polynomials.
 - Piecewise rational functions.
 - $x \mapsto \|x\|_q$, with $q \geq 0$ rational.
 - \max, \min of semi-algebraic functions.

Definable functions

A **class** of function that includes:

- ▶ **Semialgebraic**: defined by a finite number of polynomial inequalities.
 - Piecewise polynomials.
 - Piecewise rational functions.
 - $x \mapsto \|x\|_q$, with $q \geq 0$ rational.
 - \max, \min of semi-algebraic functions.
- ▶ **Analytic** restricted to a (semialgebraic) compact.

Definable functions

A **class** of function that includes:

- ▶ **Semialgebraic**: defined by a finite number of polynomial inequalities.
 - Piecewise polynomials.
 - Piecewise rational functions.
 - $x \mapsto \|x\|_q$, with $q \geq 0$ rational.
 - \max, \min of semi-algebraic functions.
- ▶ **Analytic** restricted to a (semialgebraic) compact.
- ▶ **exp** and **log**

Definable functions

A **class** of function that includes:

- ▶ **Semialgebraic**: defined by a finite number of polynomial inequalities.
 - Piecewise polynomials.
 - Piecewise rational functions.
 - $x \mapsto \|x\|_q$, with $q \geq 0$ rational.
 - \max, \min of semi-algebraic functions.
- ▶ **Analytic** restricted to a (semialgebraic) compact.
- ▶ **exp** and **log**
- ▶ Stable by $\{+, -, \times, /, \circ^{-1}\}$ and **composition**.

Definable functions

A **class** of function that includes:

- ▶ **Semialgebraic**: defined by a finite number of polynomial inequalities.
 - Piecewise polynomials.
 - Piecewise rational functions.
 - $x \mapsto \|x\|_q$, with $q \geq 0$ rational.
 - \max, \min of semi-algebraic functions.
- ▶ **Analytic** restricted to a (semialgebraic) compact.
- ▶ **exp** and **log**
- ▶ Stable by $\{+, -, \times, /, \circ^{-1}\}$ and **composition**.

Definable functions are piecewise smooth 1/2




Figure: $F(y, z) = \frac{1}{2}y^2 + |z|$

- F is smooth on $X_1 = \{(y, z) : z > 0\}$ and $X_2 = \{(y, z) : z < 0\}$.

Definable functions are piecewise smooth 1/2




Figure: $F(y, z) = \frac{1}{2}y^2 + |z|$

- F is smooth on $X_1 = \{(y, z) : z > 0\}$ and $X_2 = \{(y, z) : z < 0\}$.
- F is smooth on $X_3 = \{(y, z) : z = 0\}$.

Definable functions are piecewise smooth 1/2




Figure: $F(y, z) = \frac{1}{2}y^2 + |z|$

- F is smooth on $X_1 = \{(y, z) : z > 0\}$ and $X_2 = \{(y, z) : z < 0\}$.
- F is smooth on $X_3 = \{(y, z) : z = 0\}$.

Definable functions are piecewise smooth 2/2

Theorem

Let $F : \mathbb{R}^d \rightarrow \mathbb{R}$ be **definable** and $p \in \mathbb{N}$. There exists (X_i) , a finite partition of \mathbb{R}^d into C^p manifolds, s.t. $F|_{X_i}$ is C^p .

Dries and Miller, 1996




Figure: $F(y, z) = \frac{1}{2}y^2 + |z|$ is smooth on $X_1, X_2, X_3 = z > 0, z < 0, z = 0$

Main result

Theorem

If the family (f_a) is definable in an o-minimal structure, then D_F is a conservative set-valued field of F .

Schechtman, 2024

► The family (f_a) is definable if

$(x, a) \mapsto f_a(x)$ is definable.

Conservative set-valued fields

Gradient

F smooth $\implies \nabla F$ conservative.

► For every a.c. curve $\mathbf{x} : [0, 1] \rightarrow \mathbb{R}^d$

Gradient

F smooth $\implies \nabla F$ conservative.

► For every a.c. curve $\mathbf{x} : [0, 1] \rightarrow \mathbb{R}^d$

$$F(\mathbf{x}_1) - F(\mathbf{x}_0) = \int_0^1 \langle \nabla F(\mathbf{x}_t), \dot{\mathbf{x}}_t \rangle dt$$

$$\Updownarrow$$

$$\frac{d}{dt} F(\mathbf{x}_t) = \langle \nabla F(\mathbf{x}_t), \dot{\mathbf{x}}_t \rangle .$$

Conservative set-valued field

$F : \mathbb{R}^d \rightarrow \mathbb{R}$, locally Lipschitz continuous.

$$D_F : \mathbb{R}^d \rightrightarrows \mathbb{R}^d \quad D(x) \subset \mathbb{R}^d.$$

Conservative set-valued field

$F : \mathbb{R}^d \rightarrow \mathbb{R}$, locally Lipschitz continuous.

$$D_F : \mathbb{R}^d \rightrightarrows \mathbb{R}^d \quad D(x) \subset \mathbb{R}^d.$$

D_F is a conservative set-valued field of F if (Bolte and Pauwels, 2021).

► For every a.c. curve $\mathbf{x} : [0, 1] \rightarrow \mathbb{R}^d$ and $\mathbf{v}_t \in D_F(\mathbf{x}_t)$

$$F(\mathbf{x}_1) - F(\mathbf{x}_0) = \int_0^1 \langle \mathbf{v}_t, \dot{\mathbf{x}}_t \rangle dt$$

$$\Updownarrow$$

$$\frac{d}{dt} F(\mathbf{x}_t) = \langle \mathbf{v}_t, \dot{\mathbf{x}}_t \rangle \quad \text{almost everywhere.}$$

Examples

$$F(\mathbf{x}_1) = F(\mathbf{x}_0) + \int_0^1 \langle \mathbf{v}_t, \dot{\mathbf{x}}_t \rangle dt .$$

- If F is C^1 , then $D_F(x) = \{\nabla F(x)\}$ is a conservative field.

Examples

$$F(x_1) = F(x_0) + \int_0^1 \langle v_t, \dot{x}_t \rangle dt .$$

- ▶ If F is C^1 , then $D_F(x) = \{\nabla F(x)\}$ is a conservative field.
- ▶ If F is definable, then ∂F is a conservative field. Bolte et al., 2007

Examples

$$F(\mathbf{x}_1) = F(\mathbf{x}_0) + \int_0^1 \langle \mathbf{v}_t, \dot{\mathbf{x}}_t \rangle dt .$$

- ▶ If F is C^1 , then $D_F(x) = \{\nabla F(x)\}$ is a conservative field.
- ▶ If F is definable, then ∂F is a conservative field. Bolte et al., 2007
- ▶ Automatic differentiation in TensorFlow or Pytorch outputs an element of a conservative field. Bolte and Pauwels, 2021

Examples

$$F(\mathbf{x}_1) = F(\mathbf{x}_0) + \int_0^1 \langle \mathbf{v}_t, \dot{\mathbf{x}}_t \rangle dt .$$


- ▶ If F is C^1 , then $D_F(x) = \{\nabla F(x)\}$ is a conservative field.
- ▶ If F is definable, then ∂F is a conservative field. Bolte et al., 2007
- ▶ Automatic differentiation in TensorFlow or Pytorch outputs an element of a conservative field. Bolte and Pauwels, 2021


Variational stratification

Let $F : \mathbb{R}^d \rightarrow \mathbb{R}$ and $D_F : \mathbb{R}^d \rightrightarrows \mathbb{R}^d$ be **definable**. D_F is a conservative field **if and only if** there is a partition of \mathbb{R}^d into manifolds (\mathcal{M}_i) such that for every $y \in \mathcal{M}_i$

$$D_F(y) \subset \nabla_{\mathcal{M}_i} F(y) + \mathcal{N}_{\mathcal{M}_i}(y).$$


Bolte and Pauwels, 2021; Lewis and Tian, 2021; Davis and Drusvyatskiy, 2022






$$D_F(x) \text{ and } \nabla_{\mathcal{M}} F(x)$$

► $D_F(x) = \{\nabla F(x)\}$ on a dense open set.



$$D_F(x) \text{ and } \nabla_{\mathcal{M}} F(x)$$


- $D_F(x) = \{\nabla F(x)\}$ on a dense open set.
- ∂F is a **conservative field** and $\partial F(x) \subset \text{conv } D_F(x)$.




$$D_F(x) \text{ and } \nabla_{\mathcal{M}} F(x)$$

- $D_F(x) = \{\nabla F(x)\}$ on a dense open set.
- ∂F is a **conservative field** and $\partial F(x) \subset \text{conv } D_F(x)$.
- x^* local minimum $\implies 0 \in \text{conv } D_F(x^*)$.


$\mathbf{x}(t)$ and $\dot{\mathbf{x}}(t)$



$$\frac{d}{dt}F(\mathbf{x}(t)) = \langle \mathbf{v}(t), \dot{\mathbf{x}}(t) \rangle = \langle \nabla_{\mathcal{M}} F(\mathbf{x}(t)), \dot{\mathbf{x}}(t) \rangle + \overbrace{\langle \mathcal{N}_{\mathcal{M}}(y), \dot{\mathbf{x}}(t) \rangle}^{=0}.$$




→ Conservative fields are **not** unique!



→ Conservative fields are **not** unique!

→ But $D_F(x) = \{\nabla F(x)\}$ on a dense open set.



- Conservative fields are **not** unique!
- But $D_F(x) = \{\nabla F(x)\}$ on a dense open set.
- ∂F is a **conservative field** and $\partial F(x) \subset \text{conv } D_F(x)$.
- x^* local minimum $\implies 0 \in \text{conv } D_F(x^*)$.

Theorem

If the family (f_a) is definable in an o-minimal structure, then D_F is a conservative set-valued field of F

Schechtman, 2024

$$D_F(x) = \{v \in \mathbb{R}^d : \text{there is } (x_n, v_n, a_n) \rightarrow (x, v, 0) \text{ with } v_n \in \partial f_{a_n}(x_n)\}.$$

→ ∂f_a is the Clarke's subgradient.

Convergence of gradients

Main result

- ▶ $f_a \rightarrow F$ uniformly on compact sets.

Main result

- ▶ $f_a \rightarrow F$ uniformly on compact sets.
- ▶ For each $a > 0$, D_a is a conservative field of f_a . (e.g. $D_a = \nabla f_a$ or ∂f_a).

Main result

- ▶ $f_a \rightarrow F$ uniformly on compact sets.
- ▶ For each $a > 0$, D_a is a **conservative field** of f_a . (e.g. $D_a = \nabla f_a$ or ∂f_a).
- ▶ The maps $(x, a) \mapsto f_a(x)$ and $(x, a) \mapsto D_a(x)$ are **definable** in the **same** o-minimal structure.

Main result

- ▶ $f_a \rightarrow F$ uniformly on compact sets.
- ▶ For each $a > 0$, D_a is a **conservative field** of f_a . (e.g. $D_a = \nabla f_a$ or ∂f_a).
- ▶ The maps $(x, a) \mapsto f_a(x)$ and $(x, a) \mapsto D_a(x)$ are **definable** in the **same** o-minimal structure.

Theorem

D_F is a **conservative set-valued field** of F .

$$D_F(x) = \{v \in \mathbb{R}^d : (x_n, v_n, a_n) \rightarrow (x, v, 0) \text{ with } v_n \in D_{a_n}(x_n)\}$$

Schechtman, 2024

Main result

- ▶ $f_a \rightarrow F$ uniformly on compact sets.
- ▶ For each $a > 0$, D_a is a **conservative field** of f_a . (e.g. $D_a = \nabla f_a$ or ∂f_a).
- ▶ The maps $(x, a) \mapsto f_a(x)$ and $(x, a) \mapsto D_a(x)$ are **definable** in the **same** o-minimal structure.

Theorem

D_F is a **conservative set-valued field** of F .

$$D_F(x) = \{v \in \mathbb{R}^d : (x_n, v_n, a_n) \rightarrow (x, v, 0) \text{ with } v_n \in D_{a_n}(x_n)\}$$


Schechtman, 2024

- ▶ D_F can be constructed from an **arbitrary** family of conservative fields of f_a .
- ▶ For almost every $x \in \mathbb{R}^d$, $D_F(x) = \{\nabla F(x)\}$.
- ▶ We can partition \mathbb{R}^d into (\mathcal{M}_i) such that for $y \in \mathcal{M}_i$,

$$D_F(y) \subset \nabla_{\mathcal{M}} F(y) + \mathcal{N}_{\mathcal{M}_i}(y).$$

Careful

$D_F \neq \partial F$ even if F and (f_a) are definable and smooth !



$f_a(x)$ and $f'_a(x)$

Careful

It is **not sufficient** to have f_a definable for each a .

We need the **whole** map $(x, a) \mapsto f_a(x)$ to be **definable!**

Careful

It is **not sufficient** to have f_a definable for each a .

We need the **whole** map $(x, a) \mapsto f_a(x)$ to be **definable!**

Recall $f_a(x) = a \sin(x/a)$.

- ▶ f_a restricted to $[-1, 1]$ is **subanalytic**.
- ▶ The map $(x, a) \mapsto f_a(x)$ **is not!**

non Lipschitz extensions?

- ▶ New proof, allowing non Lipschitz f_a .
 - Coming soon!
 - Using the **variational stratification** as a definition of a conservative field.

Summary

- ▶ The subgradients of a **definable family** of functions converge to a set-valued conservative field.
 - ▶ Similar result for conservative Jacobians.
- ▶ Smoothing methods converge to a D_F -critical point.
 - Meaningful first-order optimality condition.
- ▶ **Future works.** Extension to the non Lipschitz case?

Supplementary material

Clarke's subgradient

$F : \mathbb{R}^d \rightarrow \mathbb{R}$ locally Lipschitz continuous .

Clarke's subgradient

$F : \mathbb{R}^d \rightarrow \mathbb{R}$ locally Lipschitz continuous .

$$\partial F(x) := \overline{\text{conv}}\{v \in \mathbb{R}^d : x_n \rightarrow x, F \text{ differentiable at } x_n, \nabla F(x_n) \rightarrow v\}$$

Clarke's subgradient

$F : \mathbb{R}^d \rightarrow \mathbb{R}$ locally Lipschitz continuous.

$$\partial F(x) := \overline{\text{conv}}\{v \in \mathbb{R}^d : x_n \rightarrow x, F \text{ differentiable at } x_n, \nabla F(x_n) \rightarrow v\}$$

Properties

► Clarke's subgradient: $\partial F : \mathbb{R}^d \rightrightarrows \mathbb{R}^d$, i.e. $\partial F(x) \subset \mathbb{R}^d$.

Clarke's subgradient

$F : \mathbb{R}^d \rightarrow \mathbb{R}$ locally Lipschitz continuous .

$$\partial F(x) := \overline{\text{conv}}\{v \in \mathbb{R}^d : x_n \rightarrow x, F \text{ differentiable at } x_n, \nabla F(x_n) \rightarrow v\}$$

Properties

- ▶ Clarke's subgradient: $\partial F : \mathbb{R}^d \rightrightarrows \mathbb{R}^d$, i.e. $\partial F(x) \subset \mathbb{R}^d$.
- ▶ If F is C^1 around x , then $\partial F(x) = \{\nabla F(x)\}$.

Clarke's subgradient

$F : \mathbb{R}^d \rightarrow \mathbb{R}$ locally Lipschitz continuous.

$$\partial F(x) := \overline{\text{conv}}\{v \in \mathbb{R}^d : x_n \rightarrow x, F \text{ differentiable at } x_n, \nabla F(x_n) \rightarrow v\}$$

Properties

- ▶ Clarke's subgradient: $\partial F : \mathbb{R}^d \rightrightarrows \mathbb{R}^d$, i.e. $\partial F(x) \subset \mathbb{R}^d$.
- ▶ If F is C^1 around x , then $\partial F(x) = \{\nabla F(x)\}$.
- ▶ If F is convex, then ∂F is the convex subgradient.

Clarke's subgradient

$F : \mathbb{R}^d \rightarrow \mathbb{R}$ locally Lipschitz continuous.

$$\partial F(x) := \overline{\text{conv}}\{v \in \mathbb{R}^d : x_n \rightarrow x, F \text{ differentiable at } x_n, \nabla F(x_n) \rightarrow v\}$$

Properties

- ▶ Clarke's subgradient: $\partial F : \mathbb{R}^d \rightrightarrows \mathbb{R}^d$, i.e. $\partial F(x) \subset \mathbb{R}^d$.
- ▶ If F is C^1 around x , then $\partial F(x) = \{\nabla F(x)\}$.
- ▶ If F is convex, then ∂F is the convex subgradient.
- ▶ **Optimality condition.** If x^* a **local minimum**, then

$$0 \in \partial F(x^*).$$

Sketch of proof

Goal.

$$F(\mathbf{x}(1)) - F(\mathbf{x}(0)) = \int_0^1 \langle \mathbf{v}(t), \dot{\mathbf{x}}(t) \rangle dt \quad \text{with } \mathbf{v}(t) \in D_F(\mathbf{x}(t)).$$

Sketch of proof

Goal.

$$F(\mathbf{x}(1)) - F(\mathbf{x}(0)) = \int_0^1 \langle \mathbf{v}(t), \dot{\mathbf{x}}(t) \rangle dt \quad \text{with } \mathbf{v}(t) \in D_F(\mathbf{x}(t)).$$

- 1) **One dimensional version.** If $\mathbf{x}_a(t) \rightarrow \mathbf{x}(t)$, then $\dot{\mathbf{x}}_a(t) \rightarrow \dot{\mathbf{x}}(t)$ almost everywhere.

Sketch of proof

Goal.

$$F(x(1)) - F(x(0)) = \int_0^1 \langle v(t), \dot{x}(t) \rangle dt \quad \text{with } v(t) \in D_F(x(t)).$$

- 1) **One dimensional version.** If $x_a(t) \rightarrow x(t)$, then $\dot{x}_a(t) \rightarrow \dot{x}(t)$ almost everywhere.
- 2) **Approximating family.** Construct $x_a(t), v_a(t) \rightarrow x(t), v(t)$ such that $v_a(t) \in D_a(x_a(t))$ **a.e.**

Sketch of proof

Goal.

$$F(x(1)) - F(x(0)) = \int_0^1 \langle v(t), \dot{x}(t) \rangle dt \quad \text{with } v(t) \in D_F(x(t)).$$

- 1) **One dimensional version.** If $x_a(t) \rightarrow x(t)$, then $\dot{x}_a(t) \rightarrow \dot{x}(t)$ almost everywhere.
- 2) **Approximating family.** Construct $x_a(t), v_a(t) \rightarrow x(t), v(t)$ such that $v_a(t) \in D_a(x_a(t))$ **a.e.**
- 3) **Thus.**

$$f_a(x_a(1)) - f_a(x_a(0)) = \int_0^1 \langle \dot{x}_a(t), v_a(t) \rangle dt$$

Sketch of proof

Goal.

$$F(\mathbf{x}(1)) - F(\mathbf{x}(0)) = \int_0^1 \langle \mathbf{v}(t), \dot{\mathbf{x}}(t) \rangle dt \quad \text{with } \mathbf{v}(t) \in D_F(\mathbf{x}(t)).$$

- 1) **One dimensional version.** If $\mathbf{x}_a(t) \rightarrow \mathbf{x}(t)$, then $\dot{\mathbf{x}}_a(t) \rightarrow \dot{\mathbf{x}}(t)$ almost everywhere.
- 2) **Approximating family.** Construct $\mathbf{x}_a(t), \mathbf{v}_a(t) \rightarrow \mathbf{x}(t), \mathbf{v}(t)$ such that $\mathbf{v}_a(t) \in D_a(\mathbf{x}_a(t))$ **a.e.**
- 3) **Thus.**

$$f_a(\mathbf{x}_a(1)) - f_a(\mathbf{x}_a(0)) = \int_0^1 \langle \dot{\mathbf{x}}_a(t), \mathbf{v}_a(t) \rangle dt$$

passing to the limit

$$F(\mathbf{x}(1)) - F(\mathbf{x}(0)) = \int_0^1 \langle \dot{\mathbf{x}}(t), \mathbf{v}(t) \rangle dt.$$