

ARIMA

Data 624
March 5, 2024

Our team

Miya Josephs
Mohamed Hassan

Nick Climaco
Umer Farooq
Mohammed Rahman

Table of contents

01

Introduction

02

Components
of ARIMA

03

Parameter
Selection

Table of contents

04

Model
Estimation

05

Model
Diagnostic

06

Forecasting
with ARIMA
Models

Table of contents

07

Seasonal
ARIMA

08

Conclusion

01


Introduction

+


.....



+



Introduction




- ARIMA is an acronym for AutoRegressive Integrated Moving Average
- ARIMA models are another way to forecast a time series. It is one of the most widely used approaches.
- ARIMA models aim to describe the autocorrelations in the data.
- There are two main concepts to discuss before we discuss Arima further:
 - Stationary time series
 - Differencing time series

02

Components of ARIMA Models

+

.....




.....

Stationary

- Time Series whose statistical properties do not depend on the time at which the series was observed
- Does not have seasonality
- For example, a white noise time series is stationary.

(b) Change in google price




Differencing


- Differencing allows us to take non stationary time series to stationary.
- Transformations such as logarithms can help to stabilise the variance of a time series.
- Differencing can help stabilise the mean of a time series and therefore eliminating (or reducing) trend and seasonality.

.....

(a) Google closing price



(b) Change in google price



ARIMA

Autoregressive

- Autoregressive model is regression of a time series against the lagged values of the series.
- The last p observations are used as predictors in the regression equation.
- Autoregressive models are remarkably flexible at handling a wide range of different time series patterns.
- We refer to this as an AR(p) model, an autoregressive model of order p.
- Changing the phi of the formula results in different patterns and the variance in the error terms changes the scale.

$$y_t = c + \phi_1 y_{t-1} + \phi_2 y_{t-2} + \cdots + \phi_p y_{t-p} + \varepsilon_t,$$

.....

ARIMA

Moving Average

- Moving Average can also be considered a regression but it regresses against lagged errors.
- The last q errors are used as predictors.
- We refer to this as an MA(q) model, a moving average model of order q .
- Changing the theta changes the time series pattern.

$$y_t = c + \varepsilon_t + \theta_1 \varepsilon_{t-1} + \theta_2 \varepsilon_{t-2} + \cdots + \theta_q \varepsilon_{t-q},$$

.....

ARMA


- Putting these together you get ARMA (autoregressive moving average model)
- Multiple regression with lagged observations and errors as predictors.
- This can only work with stationary data, so we have to difference the data.

.....

+

ARIMA Integrated

- The I in Arima stands for integrated - it combines ARMA model with differencing
 - If the time series needs to be differenced a number of times to make it stationary it results in an **ARIMA(p,d,q)** model.




03

Parameter Selection

+

.....



+

Parameter Selection – Nonseasonal

- Three parameters: p, d, q
- AR - p , I - d , MA - q
- p : Number of lag observations in the model; order of the autoregressive part; measures how much the current values depends on the previous values.
- d : Degrees of differencing; value effects prediction interval - higher the value, the faster it increases in size.
- q : Order of Moving Average part; number of lag errors in the model; measures how much current value depends on past errors.
- If time series is non-stationary, use differencing d .
- p can be determined by the PACF plot - last significant spike.
- q can be determined by the ACF plot - last significant spike.

Google Stock Price (Daily Open)

```
fit <- google_stock |>  
  filter(Symbol == "GOOG") |>  
  model(ARIMA(Open))  
report(fit)
```

Series: Open

Model: ARIMA(2,1,2)


Coefficients:

	ar1	ar2	ma1	ma2
	-0.2537	-0.8339	0.170	0.8555
s.e.	0.0549	0.0581	0.049	0.0578


sigma^2 estimated as 163.3: log likelihood=-4984.33

AIC=9978.67 AICC=9978.71 BIC=10004.35


Google Stock Price (Daily Open)



Differenced



Differenced Twice



Parameter Selection – Seasonal


- Incorporates both seasonal and nonseasonal factors in a multiplicative model
- ARIMA(p,d,q) \times (P,D,Q) S
- p = non-seasonal AR order, d = non-seasonal differencing, q = non-seasonal MA order, P = seasonal AR order, D = seasonal differencing, Q = seasonal MA order, and S = time span of repeating seasonal pattern.

* Source: <https://online.stat.psu.edu/stat510/lesson/4/4.1>

04

Model Estimation

.....



+

.....

Maximum Likelihood Estimation (MLE)


- Maximum likelihood estimation (MLE) is a commonly used method of estimating the parameters of a statistical model given a set of observations.
- It is based on the premise that the best choice of the parameter values should maximize the likelihood of making the observations given these parameters.
- Estimates coefficients

* Source:

https://mfe.baruch.cuny.edu/wp-content/uploads/2014/12/TS_Lecture1_2019.pdf

05

Model Diagnostic and Evaluation




+

.....

.....

Google Stock Price - Model Diagnostic and Evaluation



Model Diagnostic and Evaluation

	coef	std err	z	P> z	[0.025	0.975]
ar.L1	-0.2798	0.065	-4.289	0.000	-0.408	-0.152
ar.L2	-0.8129	0.062	-13.121	0.000	-0.934	-0.691
ma.L1	0.2091	0.064	3.273	0.001	0.084	0.334
ma.L2	0.8357	0.058	14.377	0.000	0.722	0.950
sigma2	0.0002	3.86e-06	62.616	0.000	0.000	0.000


Ljung-Box (L1) (Q):	0.06	Jarque-Bera (JB):	6463.73
Prob(Q):	0.81	Prob(JB):	0.00
Heteroskedasticity (H):	0.94	Skew:	0.27
Prob(H) (two-sided):	0.55	Kurtosis:	14.10

06

Forecasting with ARIMA models

+


.....



Forecasting with ARIMA Models



Forecasting with ARIMA Models




07

Seasonal ARIMA


+

.....




+


Australian Beer Production




Beer Production in Australia (Monthly)



After Log Transformation



STL Decomposition



```
```{r warning=FALSE}
beer_ts |>
 gg_tsdisplay(difference(log(production), 12) |> difference(),
 plot_type='partial', lag=36) +
 labs(title="Seasonally differenced", y="")
```

```


```
```{r}
beer_ts |>
 mutate(log_prod = difference(log(production), 12)) |>
 features(log_prod, unitroot_ndiffs)
```



```

A tibble: 1 × 1

| ndiffs |
|--------|
| <int> |
| 1 |

1 row





```
```{r}
fit <- beer_ts |>
 model(arima0111013 = ARIMA(log(production) ~ 0 + pdq(0,1,11)+PDQ(0,1,3)),
 arima1110210 = ARIMA(log(production) ~ 0 + pdq(11,1,0)+PDQ(2,1,0)),
 arima314111 = ARIMA(log(production) ~ 0 + pdq(3,1,4)+PDQ(1,1,1)),
 arima310110 = ARIMA(log(production) ~ 0 + pdq(3,1,0)+PDQ(1,1,0)),
 arima313112 = ARIMA(log(production) ~ 0 + pdq(3,1,3)+PDQ(1,1,2)),
 auto = ARIMA(log(production), stepwise = FALSE, approximation = FALSE))
```
```

```
```{r}
glance(fit) |> arrange(AICc) |> select(.model:BIC)
```

```

A tibble: 6 × 6

| .model | sigma2 | logLik | AIC | AICc | BIC |
|--------------|-------------|----------|-----------|------------|------------|
| <chr> | <dbl> | <dbl> | <dbl> | <dbl> | <dbl> |
| arima314111 | 0.003823228 | 564.2942 | -1108.588 | -1108.0492 | -1068.2097 |
| arima0111013 | 0.004370602 | 542.1744 | -1054.349 | -1053.1578 | -993.7808 |
| auto | 0.004494746 | 532.8018 | -1051.604 | -1051.3310 | -1023.3384 |
| arima313112 | 0.004549141 | 531.4799 | -1042.960 | -1042.4205 | -1002.5810 |
| arimal110210 | 0.004844871 | 524.4654 | -1020.931 | -1019.8912 | -964.4006 |
| arima310110 | 0.007452891 | 432.2530 | -854.506 | -854.3607 | -834.3166 |

6 rows

```
```{r}
fit |>
accuracy() |> arrange(RMSE)
```


```

A tibble: 6 × 10

| .model | .type | ME | RMSE | MAE | MPE | MAPE | MASE | RMSSE | ACF1 |
|--------------|----------|-------------|-----------|----------|------------|----------|-----------|-----------|--------------|
| <chr> | <chr> | <dbl> | <dbl> | <dbl> | <dbl> | <dbl> | <dbl> | <dbl> | <dbl> |
| arima314111 | Training | -0.13755617 | 8.652973 | 6.346916 | -0.2215620 | 4.620051 | 0.6769872 | 0.6855461 | 0.055546810 |
| arima0111013 | Training | -0.18539222 | 9.171343 | 6.678233 | -0.2791862 | 4.890190 | 0.7123267 | 0.7266149 | -0.006852976 |
| arima313112 | Training | -0.21795962 | 9.397983 | 6.811001 | -0.2921734 | 4.987327 | 0.7264883 | 0.7445708 | 0.017259665 |
| auto | Training | -0.22563239 | 9.399475 | 6.804963 | -0.2888263 | 4.978114 | 0.7258443 | 0.7446890 | 0.005041232 |
| arimal110210 | Training | -0.07435153 | 9.648806 | 6.959582 | -0.2517469 | 5.101980 | 0.7423365 | 0.7644427 | 0.024689270 |
| arima310110 | Training | -0.02959673 | 12.319454 | 9.023652 | -0.3454046 | 6.489179 | 0.9624984 | 0.9760292 | -0.031233961 |

6 rows

```
{r}  
fit |> select(arima314111) |> gg_tsresiduals()
```




```
```{r}
augment(fit) |>
 filter(.model == "arima314111") |>
 features(.innov, ljung_box, lag=36)
~~~
```

A tibble: 1 × 3

.model	lb_stat	lb_pvalue
arima314111	42.29477	0.2176564

1 row

Australia Beer Production (Monthly)



08


# Conclusion

+

.....



.....



# Conclusion


## Advantage of ARIMA model:


- ARIMA models offer the flexibility to capture various types of patterns and behaviors in the data, such as seasonality, cycles, or trends.
- ARIMA models are very easy to implement, as they have only three parameters and some statistical assumptions.
- ARIMA models are widely used and supported by many software packages and libraries, such as R.
- ARIMA models provide confidence interval and error measure in forecasting, such as, standard errors or root mean squared error.

.....

## Disadvantages of ARIMA model:

- ARIMA models cannot handle non linear relationships or complex dynamics, as they are linear models.
- ARIMA models are not suitable for very short or very long time series as they may not have enough information or become unstable over time.
- Outliers and missing values can affect the model estimation and forecasting performance in ARIMA model.





# Thanks!



**CREDITS:** This presentation template was created by [Slidesgo](#), and includes icons by [Flaticon](#), and infographics & images by [Freepik](#)

Please keep this slide for attribution

+

# Questions?

# Resources

- <https://online.stat.psu.edu/stat510/lesson/4/4.1>
- [https://mfe.baruch.cuny.edu/wp-content/uploads/2014/12/TS\\_Lecture1\\_2019.pdf](https://mfe.baruch.cuny.edu/wp-content/uploads/2014/12/TS_Lecture1_2019.pdf)
- <https://otexts.com/fpp3/>
- <https://towardsdatascience.com/time-series-forecasting-arima-models-7f221e9eee06>
- <https://medium.com/@ooemma83/how-to-interpret-acf-and-pacf-plots-for-identifying-ar-ma-arma-or-arima-models-498717e815b6>
- <https://myweb.uiowa.edu/cavaugh/doc/pub/aicaicc.pdf>
-