

COMS 4771 Lecture 26

1. Lagrangian duality

LAGRANGIAN DUALITY

LAGRANGIAN DUALITY

Uses of Lagrangian duality:

- ▶ Analysis: get insight about convex optimization problems.
- ▶ Algorithms: simple algorithms for constrained convex optimization.

CONSTRAINED CONVEX OPTIMIZATION PROBLEM

For convex functions $f_0, f_1, f_2, \dots, f_n: \mathbb{R}^d \rightarrow \mathbb{R}$:

$$\begin{array}{ll} \min_{\mathbf{x} \in \mathbb{R}^d} & f_0(\mathbf{x}) \\ \text{s.t.} & f_i(\mathbf{x}) \leq 0 \quad i = 1, 2, \dots, n. \end{array}$$

CONSTRAINED CONVEX OPTIMIZATION PROBLEM

For convex functions $f_0, f_1, f_2, \dots, f_n: \mathbb{R}^d \rightarrow \mathbb{R}$:

$$\begin{array}{ll} \min_{\mathbf{x} \in \mathbb{R}^d} & f_0(\mathbf{x}) \\ \text{s.t.} & f_i(\mathbf{x}) \leq 0 \quad i = 1, 2, \dots, n. \end{array}$$

Always think of optimization as a game against an adversary who is trying to prevent you from solving the problem.

CONSTRAINED CONVEX OPTIMIZATION PROBLEM

For convex functions $f_0, f_1, f_2, \dots, f_n: \mathbb{R}^d \rightarrow \mathbb{R}$:

$$\begin{array}{ll} \min_{\mathbf{x} \in \mathbb{R}^d} & f_0(\mathbf{x}) \\ \text{s.t.} & f_i(\mathbf{x}) \leq 0 \quad i = 1, 2, \dots, n. \end{array}$$

Always think of optimization as a game against an adversary who is trying to prevent you from solving the problem.

- You MUST find \mathbf{x} satisfying $f_i(\mathbf{x}) \leq 0$ for each $i = 1, 2, \dots, n$ (i.e., a *feasible* \mathbf{x}).

CONSTRAINED CONVEX OPTIMIZATION PROBLEM

For convex functions $f_0, f_1, f_2, \dots, f_n: \mathbb{R}^d \rightarrow \mathbb{R}$:

$$\begin{array}{ll} \min_{\mathbf{x} \in \mathbb{R}^d} & f_0(\mathbf{x}) \\ \text{s.t.} & f_i(\mathbf{x}) \leq 0 \quad i = 1, 2, \dots, n. \end{array}$$

Always think of optimization as a game against an adversary who is trying to prevent you from solving the problem.

- ▶ You **MUST** find \mathbf{x} satisfying $f_i(\mathbf{x}) \leq 0$ for each $i = 1, 2, \dots, n$ (i.e., a *feasible* \mathbf{x}).
- ▶ Therefore, **you are playing the following game:**

CONSTRAINED CONVEX OPTIMIZATION PROBLEM

For convex functions $f_0, f_1, f_2, \dots, f_n: \mathbb{R}^d \rightarrow \mathbb{R}$:

$$\begin{array}{ll} \min_{\mathbf{x} \in \mathbb{R}^d} & f_0(\mathbf{x}) \\ \text{s.t.} & f_i(\mathbf{x}) \leq 0 \quad i = 1, 2, \dots, n. \end{array}$$

Always think of optimization as a game against an adversary who is trying to prevent you from solving the problem.

- ▶ You **MUST** find \mathbf{x} satisfying $f_i(\mathbf{x}) \leq 0$ for each $i = 1, 2, \dots, n$ (i.e., a *feasible* \mathbf{x}).
- ▶ Therefore, **you are playing the following game:**
 - ▶ If you pick a feasible \mathbf{x} , then you are charged $f_0(\mathbf{x})$.

CONSTRAINED CONVEX OPTIMIZATION PROBLEM

For convex functions $f_0, f_1, f_2, \dots, f_n: \mathbb{R}^d \rightarrow \mathbb{R}$:

$$\begin{array}{ll} \min_{\mathbf{x} \in \mathbb{R}^d} & f_0(\mathbf{x}) \\ \text{s.t.} & f_i(\mathbf{x}) \leq 0 \quad i = 1, 2, \dots, n. \end{array}$$

Always think of optimization as a game against an adversary who is trying to prevent you from solving the problem.

- ▶ You **MUST** find \mathbf{x} satisfying $f_i(\mathbf{x}) \leq 0$ for each $i = 1, 2, \dots, n$ (i.e., a *feasible* \mathbf{x}).
- ▶ Therefore, **you are playing the following game:**
 - ▶ If you pick a feasible \mathbf{x} , then you are charged $f_0(\mathbf{x})$.
 - ▶ If you pick an infeasible \mathbf{x} , then you are charged $f_0(\mathbf{x})$, plus $\lambda_i \cdot f_i(\mathbf{x})$ for each constraint $i \in [n]$ that you violate.

CONSTRAINED CONVEX OPTIMIZATION PROBLEM

For convex functions $f_0, f_1, f_2, \dots, f_n: \mathbb{R}^d \rightarrow \mathbb{R}$:

$$\begin{array}{ll} \min_{\mathbf{x} \in \mathbb{R}^d} & f_0(\mathbf{x}) \\ \text{s.t.} & f_i(\mathbf{x}) \leq 0 \quad i = 1, 2, \dots, n. \end{array}$$

Always think of optimization as a game against an adversary who is trying to prevent you from solving the problem.

- ▶ You **MUST** find \mathbf{x} satisfying $f_i(\mathbf{x}) \leq 0$ for each $i = 1, 2, \dots, n$ (i.e., a *feasible* \mathbf{x}).
- ▶ Therefore, **you are playing the following game:**
 - ▶ If you pick a feasible \mathbf{x} , then you are charged $f_0(\mathbf{x})$.
 - ▶ If you pick an infeasible \mathbf{x} , then you are charged $f_0(\mathbf{x})$, plus $\lambda_i \cdot f_i(\mathbf{x})$ for each constraint $i \in [n]$ that you violate.
 - ▶ The adversary gets to pick any non-negative $\lambda_i \geq 0$, and he gets to pick them **after you pick your \mathbf{x} !!!!**

THE GAME AGAINST THE ADVERSARY

The game you are playing: you pick $\mathbf{x} \in \mathbb{R}^d$, then adversary picks $\boldsymbol{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_n) \geq \mathbf{0}$, then you incur cost

$$f_0(\mathbf{x}) + \sum_{i=1}^n \mathbb{1}\{f_i(\mathbf{x}) > 0\} \cdot \lambda_i \cdot f_i(\mathbf{x}).$$

THE GAME AGAINST THE ADVERSARY

The game you are playing: you pick $\mathbf{x} \in \mathbb{R}^d$, then adversary picks $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n) \geq \mathbf{0}$, then you incur cost

$$f_0(\mathbf{x}) + \sum_{i=1}^n \mathbb{1}\{f_i(\mathbf{x}) > 0\} \cdot \lambda_i \cdot f_i(\mathbf{x}).$$

- **Observation:** If $f_i(\mathbf{x}) \leq 0$, adversary might as well pick $\lambda_i = 0$.

THE GAME AGAINST THE ADVERSARY

The game you are playing: you pick $\mathbf{x} \in \mathbb{R}^d$, then adversary picks $\boldsymbol{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_n) \geq \mathbf{0}$, then you incur cost

$$f_0(\mathbf{x}) + \sum_{i=1}^n \mathbb{1}\{f_i(\mathbf{x}) > 0\} \cdot \lambda_i \cdot f_i(\mathbf{x}).$$

► **Observation:** If $f_i(\mathbf{x}) \leq 0$, adversary might as well pick $\lambda_i = 0$.

Equivalent game: you pick $\mathbf{x} \in \mathbb{R}^d$, then adversary picks $\boldsymbol{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_n) \geq \mathbf{0}$, then you incur cost

$$\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) := f_0(\mathbf{x}) + \sum_{i=1}^n \lambda_i \cdot f_i(\mathbf{x}).$$

THE GAME AGAINST THE ADVERSARY

The game you are playing: you pick $\mathbf{x} \in \mathbb{R}^d$, then adversary picks $\boldsymbol{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_n) \geq \mathbf{0}$, then you incur cost

$$f_0(\mathbf{x}) + \sum_{i=1}^n \mathbb{1}\{f_i(\mathbf{x}) > 0\} \cdot \lambda_i \cdot f_i(\mathbf{x}).$$

► **Observation:** If $f_i(\mathbf{x}) \leq 0$, adversary might as well pick $\lambda_i = 0$.

Equivalent game: you pick $\mathbf{x} \in \mathbb{R}^d$, then adversary picks $\boldsymbol{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_n) \geq \mathbf{0}$, then you incur cost

$$\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) := f_0(\mathbf{x}) + \sum_{i=1}^n \lambda_i \cdot f_i(\mathbf{x}).$$

- If you manage to pick a feasible \mathbf{x} , you incur cost $f_0(\mathbf{x})$.
(Adversary would not want to set $\lambda_i > 0$ for any i in this case.)

THE GAME AGAINST THE ADVERSARY

The game you are playing: you pick $\mathbf{x} \in \mathbb{R}^d$, then adversary picks $\boldsymbol{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_n) \geq \mathbf{0}$, then you incur cost

$$f_0(\mathbf{x}) + \sum_{i=1}^n \mathbb{1}\{f_i(\mathbf{x}) > 0\} \cdot \lambda_i \cdot f_i(\mathbf{x}).$$

- **Observation:** If $f_i(\mathbf{x}) \leq 0$, adversary might as well pick $\lambda_i = 0$.

Equivalent game: you pick $\mathbf{x} \in \mathbb{R}^d$, then adversary picks $\boldsymbol{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_n) \geq \mathbf{0}$, then you incur cost

$$\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) := f_0(\mathbf{x}) + \sum_{i=1}^n \lambda_i \cdot f_i(\mathbf{x}).$$

- If you manage to pick a feasible \mathbf{x} , you incur cost $f_0(\mathbf{x})$.
(Adversary would not want to set $\lambda_i > 0$ for any i in this case.)
- If you pick an infeasible \mathbf{x} , **adversary punishes you by sending $\lambda_i \rightarrow +\infty$ for the constraint $i \in [n]$ you violate.**

THE GAME AGAINST THE ADVERSARY

The game you are playing: you pick $\mathbf{x} \in \mathbb{R}^d$, then adversary picks $\boldsymbol{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_n) \geq \mathbf{0}$, then you incur cost

$$f_0(\mathbf{x}) + \sum_{i=1}^n \mathbb{1}\{f_i(\mathbf{x}) > 0\} \cdot \lambda_i \cdot f_i(\mathbf{x}).$$

- **Observation:** If $f_i(\mathbf{x}) \leq 0$, adversary might as well pick $\lambda_i = 0$.

Equivalent game: you pick $\mathbf{x} \in \mathbb{R}^d$, then adversary picks $\boldsymbol{\lambda} = (\lambda_1, \lambda_2, \dots, \lambda_n) \geq \mathbf{0}$, then you incur cost

$$\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) := f_0(\mathbf{x}) + \sum_{i=1}^n \lambda_i \cdot f_i(\mathbf{x}).$$

- If you manage to pick a feasible \mathbf{x} , you incur cost $f_0(\mathbf{x})$.
(Adversary would not want to set $\lambda_i > 0$ for any i in this case.)
- If you pick an infeasible \mathbf{x} , **adversary punishes you by sending $\lambda_i \rightarrow +\infty$ for the constraint $i \in [n]$ you violate.**

In this case, you incur $+\infty$ cost.

THE BEST YOU CAN DO AGAINST THE ADVERSARY

Best cost for you:

$$\min_{\mathbf{x} \in \mathbb{R}^d} \max_{\lambda_1, \lambda_2, \dots, \lambda_n \geq 0} \underbrace{f_0(\mathbf{x}) + \sum_{i=1}^n \lambda_i \cdot f_i(\mathbf{x})}_{\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda})}.$$

THE BEST YOU CAN DO AGAINST THE ADVERSARY

Best cost for you:

$$\min_{\mathbf{x} \in \mathbb{R}^d} \max_{\lambda_1, \lambda_2, \dots, \lambda_n \geq 0} \underbrace{f_0(\mathbf{x}) + \sum_{i=1}^n \lambda_i \cdot f_i(\mathbf{x})}_{\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda})}.$$

$\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda})$ is called the **Lagrangian function**.

THE BEST YOU CAN DO AGAINST THE ADVERSARY

Best cost for you:

$$\min_{\mathbf{x} \in \mathbb{R}^d} \max_{\lambda_1, \lambda_2, \dots, \lambda_n \geq 0} \underbrace{f_0(\mathbf{x}) + \sum_{i=1}^n \lambda_i \cdot f_i(\mathbf{x})}_{\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda})}.$$

$\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda})$ is called the **Lagrangian function**.

Summary of game:

- ▶ You pick $\mathbf{x} \in \mathbb{R}^d$.
- ▶ Adversary picks $\boldsymbol{\lambda} \geq \mathbf{0}$.
- ▶ You incur cost $\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda})$.

THE BEST YOU CAN DO AGAINST THE ADVERSARY

Best cost for you:

$$\min_{\mathbf{x} \in \mathbb{R}^d} \max_{\lambda_1, \lambda_2, \dots, \lambda_n \geq 0} \underbrace{f_0(\mathbf{x}) + \sum_{i=1}^n \lambda_i \cdot f_i(\mathbf{x})}_{\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda})}.$$

$\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda})$ is called the **Lagrangian function**.

Summary of game:

- ▶ You pick $\mathbf{x} \in \mathbb{R}^d$.
- ▶ Adversary picks $\boldsymbol{\lambda} \geq \mathbf{0}$.
- ▶ You incur cost $\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda})$.

Only way for you not to incur $+\infty$ cost is to pick a feasible $\mathbf{x} \in \mathbb{R}^d$.

WEAK DUALITY

$$\min_{\mathbf{x} \in \mathbb{R}^d} \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) \geq \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}).$$

WEAK DUALITY

$$\min_{\mathbf{x} \in \mathbb{R}^d} \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) \geq \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}).$$

- **LHS**: best you can do in game where you must pick \mathbf{x} first, and then adversary gets to pick $\boldsymbol{\lambda}$ after seeing what you picked.

WEAK DUALITY

$$\min_{\mathbf{x} \in \mathbb{R}^d} \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) \geq \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}).$$

- ▶ **LHS**: best you can do in game where **you must pick \mathbf{x} first**, and then **adversary gets to pick $\boldsymbol{\lambda}$ after seeing what you picked**.
- ▶ **RHS**: worst you can do in game where **adversary must pick $\boldsymbol{\lambda}$ first**, and then **you get to pick \mathbf{x} after seeing what adversary picked**.

WEAK DUALITY

$$\min_{\mathbf{x} \in \mathbb{R}^d} \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) \geq \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}).$$

- ▶ **LHS**: best you can do in game where **you must pick \mathbf{x} first**, and then **adversary gets to pick $\boldsymbol{\lambda}$ after seeing what you picked**.
- ▶ **RHS**: worst you can do in game where **adversary must pick $\boldsymbol{\lambda}$ first**, and then **you get to pick \mathbf{x} after seeing what adversary picked**.
- ▶ RHS is better for you in terms of guaranteeing smaller cost.

WEAK DUALITY

lower bound, the right part always convex, have solution as the lower bound for the original problem

$$\min_{\mathbf{x} \in \mathbb{R}^d} \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) \geq \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}).$$

- ▶ **LHS**: best you can do in game where **you must pick \mathbf{x} first**, and then **adversary gets to pick $\boldsymbol{\lambda}$ after seeing what you picked**.
- ▶ **RHS**: worst you can do in game where **adversary must pick $\boldsymbol{\lambda}$ first**, and then **you get to pick \mathbf{x} after seeing what adversary picked**.
- ▶ RHS is better for you in terms of guaranteeing smaller cost.

In fact, for all $\tilde{\mathbf{x}} \in \mathbb{R}^d$ and all $\tilde{\boldsymbol{\lambda}} \geq \mathbf{0}$,

$$\max_{\boldsymbol{\lambda} \geq \mathbf{0}} \mathcal{L}(\tilde{\mathbf{x}}, \boldsymbol{\lambda}) \geq \min_{\mathbf{x} \in \mathbb{R}^d} \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) \geq \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) \geq \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \tilde{\boldsymbol{\lambda}})$$

STRONG DUALITY

Under some conditions (e.g., some $\mathbf{x} \in \mathbb{R}^d$ strictly satisfies all constraints),

$$\min_{\mathbf{x} \in \mathbb{R}^d} \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) = \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}).$$

STRONG DUALITY

Under some conditions (e.g., some $x \in \mathbb{R}^d$ strictly satisfies all constraints),

$$\min_{x \in \mathbb{R}^d} \max_{\lambda \geq 0} \mathcal{L}(x, \lambda) = \max_{\lambda \geq 0} \min_{x \in \mathbb{R}^d} \mathcal{L}(x, \lambda).$$

Doesn't matter if you go first or if adversary goes first.

STRONG DUALITY

Under some conditions (e.g., some $x \in \mathbb{R}^d$ strictly satisfies all constraints),

$$\min_{x \in \mathbb{R}^d} \max_{\lambda \geq 0} \mathcal{L}(x, \lambda) = \max_{\lambda \geq 0} \min_{x \in \mathbb{R}^d} \mathcal{L}(x, \lambda).$$

Doesn't matter if you go first or if adversary goes first.

- **LHS**: the primal (i.e., original) problem.

STRONG DUALITY

Under some conditions (e.g., some $\mathbf{x} \in \mathbb{R}^d$ strictly satisfies all constraints),

$$\min_{\mathbf{x} \in \mathbb{R}^d} \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) = \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}).$$

Doesn't matter if you go first or if adversary goes first.

- ▶ **LHS:** the primal (i.e., original) problem.
- ▶ **RHS:**

$$\max_{\boldsymbol{\lambda} \geq \mathbf{0}} g(\boldsymbol{\lambda})$$

where

$$g(\boldsymbol{\lambda}) := \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}).$$

This maximization problem is called the **Lagrange dual problem**.

STRONG DUALITY

Under some conditions (e.g., some $\mathbf{x} \in \mathbb{R}^d$ strictly satisfies all constraints),

$$\min_{\mathbf{x} \in \mathbb{R}^d} \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) = \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}).$$

Doesn't matter if you go first or if adversary goes first.

- ▶ **LHS:** the primal (i.e., original) problem.
- ▶ **RHS:**

$$\max_{\boldsymbol{\lambda} \geq \mathbf{0}} g(\boldsymbol{\lambda})$$

where

$$g(\boldsymbol{\lambda}) := \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}).$$

This maximization problem is called the **Lagrange dual problem**.

- ▶ $\boldsymbol{\lambda} \geq \mathbf{0}$: **Lagrange multipliers** (dual variables).

STRONG DUALITY

Under some conditions (e.g., some $\mathbf{x} \in \mathbb{R}^d$ strictly satisfies all constraints),

$$\min_{\mathbf{x} \in \mathbb{R}^d} \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) = \max_{\boldsymbol{\lambda} \geq \mathbf{0}} \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}).$$

Doesn't matter if you go first or if adversary goes first.

► **LHS:** the primal (i.e., original) problem.

$\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda})$ is concave

► **RHS:**

$\min \{\text{concave functions}\}$ is
also a concave function

$$\max_{\boldsymbol{\lambda} \geq \mathbf{0}} g(\boldsymbol{\lambda})$$

where

$$g(\boldsymbol{\lambda}) := \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}).$$

This maximization problem is called the **Lagrange dual problem**.

► $\boldsymbol{\lambda} \geq \mathbf{0}$: **Lagrange multipliers** (dual variables).

► $g: \mathbb{R}^n \rightarrow \mathbb{R}$: **Lagrange dual function**, which is always **concave**.

EQUALITY CONSTRAINTS: SHORTCUT

For convex functions $f_0, f_1, f_2, \dots, f_n: \mathbb{R}^d \rightarrow \mathbb{R}$, $\mathbf{A} \in \mathbb{R}^{k \times d}$, and $\mathbf{b} \in \mathbb{R}^k$:

$$\begin{array}{ll} \min_{\mathbf{x} \in \mathbb{R}^d} & f_0(\mathbf{x}) \\ \text{s.t.} & f_i(\mathbf{x}) \leq 0 \quad i = 1, 2, \dots, n; \\ & \mathbf{Ax} - \mathbf{b} = \mathbf{0}. \end{array}$$

EQUALITY CONSTRAINTS: SHORTCUT

For convex functions $f_0, f_1, f_2, \dots, f_n: \mathbb{R}^d \rightarrow \mathbb{R}$, $\mathbf{A} \in \mathbb{R}^{k \times d}$, and $\mathbf{b} \in \mathbb{R}^k$:

$$\begin{array}{ll} \min_{\mathbf{x} \in \mathbb{R}^d} & f_0(\mathbf{x}) \\ \text{s.t.} & f_i(\mathbf{x}) \leq 0 \quad i = 1, 2, \dots, n; \\ & \mathbf{Ax} - \mathbf{b} = \mathbf{0}. \end{array}$$

Equality constraint $\mathbf{Ax} - \mathbf{b} = \mathbf{0}$ is same as conjunction of two inequality constraints:

$$\mathbf{Ax} - \mathbf{b} \leq \mathbf{0} \quad \text{and} \quad \mathbf{b} - \mathbf{Ax} \leq \mathbf{0}.$$

EQUALITY CONSTRAINTS: SHORTCUT

For convex functions $f_0, f_1, f_2, \dots, f_n: \mathbb{R}^d \rightarrow \mathbb{R}$, $\mathbf{A} \in \mathbb{R}^{k \times d}$, and $\mathbf{b} \in \mathbb{R}^k$:

$$\begin{array}{ll} \min_{\mathbf{x} \in \mathbb{R}^d} & f_0(\mathbf{x}) \\ \text{s.t.} & f_i(\mathbf{x}) \leq 0 \quad i = 1, 2, \dots, n; \\ & \mathbf{Ax} - \mathbf{b} = \mathbf{0}. \end{array}$$

Equality constraint $\mathbf{Ax} - \mathbf{b} = \mathbf{0}$ is same as conjunction of two inequality constraints:

$$\mathbf{Ax} - \mathbf{b} \leq \mathbf{0} \quad \text{and} \quad \mathbf{b} - \mathbf{Ax} \leq \mathbf{0}.$$

Shortcut: Lagrange multipliers associated with equality constraints do not have to be non-negative. (Easy to derive by reasoning about “game against adversary”.)

STRONG DUALITY

Under some conditions (e.g., some $\mathbf{x} \in \mathbb{R}^d$ strictly satisfies all constraints),

$$\min_{\mathbf{x} \in \mathbb{R}^d} \max_{\boldsymbol{\lambda} \geq \mathbf{0}, \boldsymbol{\nu} \in \mathbb{R}^k} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) = \max_{\boldsymbol{\lambda} \geq \mathbf{0}, \boldsymbol{\nu} \in \mathbb{R}^k} \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}).$$

STRONG DUALITY

Under some conditions (e.g., some $\mathbf{x} \in \mathbb{R}^d$ strictly satisfies all constraints),

$$\min_{\mathbf{x} \in \mathbb{R}^d} \max_{\boldsymbol{\lambda} \geq \mathbf{0}, \boldsymbol{\nu} \in \mathbb{R}^k} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}) = \max_{\boldsymbol{\lambda} \geq \mathbf{0}, \boldsymbol{\nu} \in \mathbb{R}^k} \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}).$$

- ▶ **LHS:** the primal (i.e., original) problem.
- ▶ **RHS:**

$$\max_{\boldsymbol{\lambda} \geq \mathbf{0}, \boldsymbol{\nu} \in \mathbb{R}^k} g(\boldsymbol{\lambda}, \boldsymbol{\nu})$$

where

$$g(\boldsymbol{\lambda}, \boldsymbol{\nu}) := \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}, \boldsymbol{\nu}).$$

MAXIMUM ENTROPY

EXAMPLE: MAXIMUM ENTROPY

The maximum entropy optimization problem (over domain $[d]$):

$$\begin{aligned} \min_{\mathbf{w} \in \mathbb{R}^d} \quad & \sum_{i=1}^d w_i \ln \frac{w_i}{\pi_i} \\ \text{s.t.} \quad & \mu_j - \sum_{i=1}^d t_{i,j} w_i = 0 \quad j = 1, 2, \dots, k; \\ & 1 - \sum_{i=1}^d w_i = 0. \end{aligned}$$

(We drop the constraint $\mathbf{w} \geq 0$, since it'll be satisfied w/o explicitly imposing it.)

EXAMPLE: MAXIMUM ENTROPY

The maximum entropy optimization problem (over domain $[d]$):

$$\begin{aligned} \min_{\mathbf{w} \in \mathbb{R}^d} \quad & \sum_{i=1}^d w_i \ln \frac{w_i}{\pi_i} \\ \text{s.t.} \quad & \mu_j - \sum_{i=1}^d t_{i,j} w_i = 0 \quad j = 1, 2, \dots, k; \\ & 1 - \sum_{i=1}^d w_i = 0. \end{aligned}$$

(We drop the constraint $\mathbf{w} \geq 0$, since it'll be satisfied w/o explicitly imposing it.)

Lagrange multipliers: $\boldsymbol{\eta} \in \mathbb{R}^k$, $\nu \in \mathbb{R}$.

EXAMPLE: MAXIMUM ENTROPY

The maximum entropy optimization problem (over domain $[d]$):

$$\begin{aligned} \min_{\mathbf{w} \in \mathbb{R}^d} \quad & \sum_{i=1}^d w_i \ln \frac{w_i}{\pi_i} \\ \text{s.t.} \quad & \mu_j - \sum_{i=1}^d t_{i,j} w_i = 0 \quad j = 1, 2, \dots, k; \\ & 1 - \sum_{i=1}^d w_i = 0. \end{aligned}$$

(We drop the constraint $\mathbf{w} \geq \mathbf{0}$, since it'll be satisfied w/o explicitly imposing it.)

Lagrange multipliers: $\boldsymbol{\eta} \in \mathbb{R}^k$, $\nu \in \mathbb{R}$.

Lagrangian:

$$\mathcal{L}(\mathbf{w}, \boldsymbol{\eta}, \nu) = \sum_{i=1}^d w_i \ln \frac{w_i}{\pi_i} + \sum_{j=1}^k \eta_j \left(\mu_j - \sum_{i=1}^d t_{i,j} w_i \right) + \nu \left(1 - \sum_{i=1}^d w_i \right).$$

EXAMPLE: MAXIMUM ENTROPY

$$\mathcal{L}(\mathbf{w}, \boldsymbol{\eta}, \nu) = \sum_{i=1}^d w_i \ln \frac{w_i}{\pi_i} + \sum_{j=1}^k \eta_j \left(\mu_j - \sum_{i=1}^d t_{i,j} w_i \right) + \nu \left(1 - \sum_{i=1}^d w_i \right).$$

We want to minimize this w.r.t. \mathbf{w} .

EXAMPLE: MAXIMUM ENTROPY

$$\mathcal{L}(\mathbf{w}, \boldsymbol{\eta}, \nu) = \sum_{i=1}^d w_i \ln \frac{w_i}{\pi_i} + \sum_{j=1}^k \eta_j \left(\mu_j - \sum_{i=1}^d t_{i,j} w_i \right) + \nu \left(1 - \sum_{i=1}^d w_i \right).$$

We want to minimize this w.r.t. \mathbf{w} .

Partial derivative w.r.t. w_i is

$$1 + \ln(w_i) - \ln(\pi_i) - \sum_{j=1}^k \eta_j t_{i,j} - \nu,$$

which is zero when

$$w_i = \pi_i \exp \left(\sum_{j=1}^k \eta_j t_{i,j} - (1 - \nu) \right).$$

(This is always non-negative!)

EXAMPLE: MAXIMUM ENTROPY

Plugging-in the choice of w_i for each $i = 1, 2, \dots, n$ that minimizes $\mathcal{L}(w, \boldsymbol{\eta}, \nu)$ yields the dual problem

$$\max_{\boldsymbol{\eta} \in \mathbb{R}^k, \nu \in \mathbb{R}} g(\boldsymbol{\eta}, \nu),$$

where

$$g(\boldsymbol{\eta}, \nu) = - \sum_{i=1}^d \pi_i \exp \left(\sum_{j=1}^k \eta_j t_{i,j} - (1 - \nu) \right) + \sum_{j=1}^k \eta_j \mu_j + \nu.$$

EXAMPLE: MAXIMUM ENTROPY

Maximum entropy dual optimization problem:

$$\max_{\boldsymbol{\eta} \in \mathbb{R}^k, \nu \in \mathbb{R}} \quad - \sum_{i=1}^d \pi_i \exp \left(\sum_{j=1}^k \eta_j t_{i,j} - (1 - \nu) \right) + \sum_{j=1}^k \eta_j \mu_j + \nu.$$

EXAMPLE: MAXIMUM ENTROPY

Maximum entropy dual optimization problem:

$$\max_{\boldsymbol{\eta} \in \mathbb{R}^k, \nu \in \mathbb{R}} \quad - \sum_{i=1}^d \pi_i \exp \left(\sum_{j=1}^k \eta_j t_{i,j} - (1 - \nu) \right) + \sum_{j=1}^k \eta_j \mu_j + \nu.$$

This is concave in $\boldsymbol{\eta}$ and ν .

EXAMPLE: MAXIMUM ENTROPY

Maximum entropy dual optimization problem:

$$\max_{\boldsymbol{\eta} \in \mathbb{R}^k, \nu \in \mathbb{R}} - \sum_{i=1}^d \pi_i \exp \left(\sum_{j=1}^k \eta_j t_{i,j} - (1 - \nu) \right) + \sum_{j=1}^k \eta_j \mu_j + \nu.$$

This is concave in $\boldsymbol{\eta}$ and ν .

Derivative w.r.t. ν is

$$- \sum_{i=1}^d \pi_i \exp \left(\sum_{j=1}^k \eta_j t_{i,j} - (1 - \nu) \right) + 1,$$

which is zero when

$$1 - \nu = \ln \left\{ \sum_{i=1}^d \pi_i \exp \left(\sum_{j=1}^k \eta_j t_{i,j} \right) \right\}.$$

EXAMPLE: MAXIMUM ENTROPY

Plugging-in maximizing choice of ν leaves us with

$$\max_{\boldsymbol{\eta} \in \mathbb{R}^k} -\ln \left\{ \sum_{i=1}^d \pi_i \exp \left(\sum_{j=1}^k \eta_j t_{i,j} \right) \right\} + \sum_{j=1}^k \eta_j \mu_j.$$

EXAMPLE: MAXIMUM ENTROPY

Plugging-in maximizing choice of ν leaves us with

$$\max_{\boldsymbol{\eta} \in \mathbb{R}^k} -\ln \left\{ \sum_{i=1}^d \pi_i \exp \left(\sum_{j=1}^k \eta_j t_{i,j} \right) \right\} + \sum_{j=1}^k \eta_j \mu_j.$$

Derivative w.r.t. η_j is

$$-\frac{\sum_{i=1}^d \pi_i \exp \left(\sum_{j'=1}^k \eta_{j'} t_{i,j'} \right) t_{i,j}}{\sum_{i=1}^d \pi_i \exp \left(\sum_{j'=1}^k \eta_{j'} t_{i,j'} \right)} + \mu_j,$$

which is zero when

$$\sum_{i=1}^d \hat{w}_i t_{i,j} = \mu_j$$

where

$$\hat{w}_i = \frac{\pi_i \exp \left(\sum_{j'=1}^k \eta_{j'} t_{i,j'} \right)}{\sum_{i=1}^d \pi_i \exp \left(\sum_{j'=1}^k \eta_{j'} t_{i,j'} \right)} \quad i = 1, 2, \dots, n.$$

LAGRANGIAN RELAXATION METHOD

LAGRANGIAN RELAXATION METHOD

Equality constrained convex optimization problem for $\mathbf{A} \in \mathbb{R}^{k \times d}$ and $\mathbf{b} \in \mathbb{R}^k$:

$$\begin{array}{ll} \min_{\mathbf{x} \in \mathbb{R}^d} & f_0(\mathbf{x}) \\ \text{s.t.} & \mathbf{Ax} - \mathbf{b} = \mathbf{0}. \end{array}$$

LAGRANGIAN RELAXATION METHOD

Equality constrained convex optimization problem for $\mathbf{A} \in \mathbb{R}^{k \times d}$ and $\mathbf{b} \in \mathbb{R}^k$:

$$\begin{array}{ll} \min_{\mathbf{x} \in \mathbb{R}^d} & f_0(\mathbf{x}) \\ \text{s.t.} & \mathbf{Ax} - \mathbf{b} = \mathbf{0}. \end{array}$$

Lagrangian: $\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) = f_0(\mathbf{x}) + \langle \boldsymbol{\lambda}, \mathbf{Ax} - \mathbf{b} \rangle$.

LAGRANGIAN RELAXATION METHOD

Equality constrained convex optimization problem for $A \in \mathbb{R}^{k \times d}$ and $b \in \mathbb{R}^k$:

$$\begin{array}{ll} \min_{x \in \mathbb{R}^d} & f_0(x) \\ \text{s.t.} & Ax - b = 0. \end{array}$$

Lagrangian: $\mathcal{L}(x, \lambda) = f_0(x) + \langle \lambda, Ax - b \rangle$.

Lagrangian relaxation algorithm

Initialize $\lambda^{(1)} \in \mathbb{R}^k$.

For $t = 1, 2, \dots$:

- ▶ Let $x^{(t)}$ be minimizer of $\mathcal{L}(x, \lambda^{(t)})$ w.r.t. x (e.g., via gradient descent).
- ▶ If $Ax^{(t)} = b$, then done; return current $x^{(t)}$.
- ▶ If $Ax^{(t)} \neq b$, then update $\lambda^{(t)}$ (via **gradient ascent step** of size $\eta_t > 0$):

$$\lambda^{(t+1)} := \lambda^{(t)} + \eta_t (Ax^{(t)} - b). \quad \text{gradient of } \lambda \text{ w.r.t. } b$$

LAGRANGIAN RELAXATION METHOD

Since $\mathbf{x}^{(t)}$ is the minimizer of $\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}^{(t)})$,

$$\mathcal{L}(\mathbf{x}^{(t)}, \boldsymbol{\lambda}^{(t)}) = \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}^{(t)}) \leq \max_{\boldsymbol{\lambda} \in \mathbb{R}^k} \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) \leq \min_{\mathbf{x} \in \mathbb{R}^d} \max_{\boldsymbol{\lambda} \in \mathbb{R}^k} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda})$$

where last step uses weak duality.

LAGRANGIAN RELAXATION METHOD

Since $\mathbf{x}^{(t)}$ is the minimizer of $\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}^{(t)})$,

$$\mathcal{L}(\mathbf{x}^{(t)}, \boldsymbol{\lambda}^{(t)}) = \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}^{(t)}) \leq \max_{\boldsymbol{\lambda} \in \mathbb{R}^k} \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) \leq \min_{\mathbf{x} \in \mathbb{R}^d} \max_{\boldsymbol{\lambda} \in \mathbb{R}^k} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda})$$

where last step uses weak duality.

- $\mathcal{L}(\mathbf{x}^{(t)}, \boldsymbol{\lambda}^{(t)})$ is always a lower-bound on optimal value.

LAGRANGIAN RELAXATION METHOD

Since $\mathbf{x}^{(t)}$ is the minimizer of $\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}^{(t)})$,

$$\mathcal{L}(\mathbf{x}^{(t)}, \boldsymbol{\lambda}^{(t)}) = \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}^{(t)}) \leq \max_{\boldsymbol{\lambda} \in \mathbb{R}^k} \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) \leq \min_{\mathbf{x} \in \mathbb{R}^d} \max_{\boldsymbol{\lambda} \in \mathbb{R}^k} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda})$$

where last step uses weak duality.

- ▶ $\mathcal{L}(\mathbf{x}^{(t)}, \boldsymbol{\lambda}^{(t)})$ is always a lower-bound on optimal value.
- ▶ If $A\mathbf{x}^{(t)} = \mathbf{b}$, then $\mathbf{x}^{(t)}$ achieves optimal value.

LAGRANGIAN RELAXATION METHOD

Since $\mathbf{x}^{(t)}$ is the minimizer of $\mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}^{(t)})$,

$$\mathcal{L}(\mathbf{x}^{(t)}, \boldsymbol{\lambda}^{(t)}) = \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}^{(t)}) \leq \max_{\boldsymbol{\lambda} \in \mathbb{R}^k} \min_{\mathbf{x} \in \mathbb{R}^d} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda}) \leq \min_{\mathbf{x} \in \mathbb{R}^d} \max_{\boldsymbol{\lambda} \in \mathbb{R}^k} \mathcal{L}(\mathbf{x}, \boldsymbol{\lambda})$$

where last step uses weak duality.

- ▶ $\mathcal{L}(\mathbf{x}^{(t)}, \boldsymbol{\lambda}^{(t)})$ is always a lower-bound on optimal value.
- ▶ If $A\mathbf{x}^{(t)} = \mathbf{b}$, then $\mathbf{x}^{(t)}$ achieves optimal value.

Can also apply to inequality constrained problems via slack variables.

- ▶ **Lagrangian duality:** every convex optimization problem has an associated dual problem.
- ▶ **Lagrange dual problem:** maximization of concave objective function subject to convex constraints.

Actually, we can apply this approach to **non-convex** optimization problems: **the dual problem always has a concave objective function and convex constraints!**

- ▶ In MaxEnt, dual problem reveals form of solution.
- ▶ **Lagrangian relaxation method:** exploit Lagrangian duality to get a simple algorithm for constrained optimization.