

Location is a Dynamic Variable Quantum Supercomputing and the Frequency–Location Hypothesis

Christopher Woodyard
Vers3Dynamics
Washington, D.C., USA

Abstract—This paper develops a mathematical framework that reinterprets spatial localization as an emergent property of coupled matter–scalar-field configurations. We propose that quantum computational resources approaching million-qubit fault-tolerance may enable the investigation of non-equilibrium field dynamics relevant to such models. The framework introduces a frequency-parameterized location operator whose eigenstates represent stable matter configurations, with re-localization interpreted as resonance-driven transitions between eigenstates. We explicitly derive the relationship to standard quantum mechanics, demonstrate recovery of classical limits, identify parameter regimes where predictions deviate measurably from standard theory, and propose falsifiable experimental tests. While macroscopic applications remain computationally and energetically prohibitive, the framework offers a mathematically consistent exploration of position as a dynamical field variable.

Index Terms—Quantum Field Theory, Scalar Fields, Position Operators, Emergent Spacetime, Quantum Computing, Falsifiable Physics

I. INTRODUCTION

Physical position in quantum mechanics is typically represented by the Hermitian position operator $\hat{x}$ with continuous eigenvalue spectrum. Recent developments in emergent spacetime theories and analog gravity suggest position may arise from deeper field-theoretic structures. We explore a complementary formalism where localization is parameterized by resonance frequencies of coupled matter–scalar-field systems.

Scope and Limitations: This work is explicitly *speculative theoretical physics*. We do not claim standard quantum mechanics is incorrect, nor that macroscopic matter teleportation is achievable. Rather, we investigate whether a frequency-based formalism can: (1) recover standard QM in appropriate limits, (2) generate testable deviations in extreme parameter regimes, and (3) provide alternative mathematical intuition for certain quantum phenomena.

II. COMPUTATIONAL PREREQUISITES

Modeling driven quantum fields in curved spacetime requires computational resources exceeding classical capabilities. We outline a quantum computing roadmap enabling such simulations.

A. Quantum Architecture Scaling

Phase I–II (100–10,000 Qubits): Development of logical qubits via surface-code error correction with physical error rates below 10^{-3} .

Phase III–IV (100,000–1,000,000 Qubits): Modular fault-tolerant arrays enabling real-time simulation of $(3 + 1)$ -dimensional quantum field dynamics on discrete spacetime lattices.

Such systems could numerically solve the coupled field equations derived in Section IV with sufficient precision to identify observable deviations from standard predictions.




Fig. 1. **Computational Roadmap.** A layered architecture scaling from physical qubits to the “Resonance Engine” required for non-equilibrium scalar field simulation.

III. RELATIONSHIP TO STANDARD QUANTUM MECHANICS

A. Standard Position Operator

In non-relativistic quantum mechanics, position is represented by the self-adjoint operator $\hat{x}$ satisfying:

$$\hat{x}|\psi\rangle = x|\psi\rangle, \quad [\hat{x}, \hat{p}] = i\hbar. \quad (1)$$

Position measurement probabilities follow Born’s rule:

$$P_{\text{std}}(x, t) = |\langle x|\psi(t)\rangle|^2 = |\psi(x, t)|^2. \quad (2)$$

B. Proposed Frequency-Location Operator

We introduce a modified operator $\mathcal{L}$ acting on an extended Hilbert space $\mathcal{H} \otimes \mathcal{H}_\phi$ (matter tensor scalar field):

$$\mathcal{L}|\Psi\rangle = \omega_{\text{loc}}|\Psi\rangle, \quad (3)$$

where $|\Psi\rangle = |\psi\rangle_{\text{matter}} \otimes |\phi\rangle_{\text{field}}$ and ω_{loc} is the dominant resonance frequency of the coupled system.

C. Mapping Between Formalisms

The relationship between $\mathcal{L}$ and $\hat{x}$ is given by the functional:

$$\mathcal{L} = \int d^3x \omega(x) |\phi(x)\rangle\langle\phi(x)| \otimes \hat{P}_x, \quad (4)$$

where $\hat{P}_x = |x\rangle\langle x|$ projects onto position x , and $\omega(x)$ is the local scalar field resonance frequency.

Key Insight: When the scalar field is in its ground state ($|\phi\rangle = |\phi_0\rangle$, uniform), we recover:

$$\langle\mathcal{L}\rangle|_{\phi=\phi_0} = \omega_0\langle\hat{x}\rangle, \quad (5)$$

where ω_0 is a constant proportionality factor. Standard position measurement is the limiting case of our framework when field coupling is negligible.

D. Correspondence Principle

For weak scalar coupling ($\xi \ll 1$) and low field amplitudes ($|\phi| \approx \phi_0$), position probabilities differ from standard QM by:

$$P_{\text{model}}(x, t) = P_{\text{std}}(x, t) [1 + \epsilon(x, t) + \mathcal{O}(\epsilon^2)], \quad (6)$$

where the correction term is:

$$\epsilon(x, t) = \xi^2 \frac{|\phi(x, t) - \phi_0|^2}{\phi_0^2} \cdot f\left(\frac{\omega_{\text{loc}}(x)}{\omega_w}\right). \quad (7)$$

For current laboratory conditions ($\xi \sim 10^{-10}$, $|\phi - \phi_0|/\phi_0 \sim 10^{-6}$), we have $\epsilon \sim 10^{-26}$, well below experimental resolution.

IV. FIELD-THEORETIC FRAMEWORK

A. Effective Action

We propose an effective action incorporating curvature-coupled scalar dynamics:

$$S = \int d^4x \sqrt{-g} \left[\frac{R}{16\pi G} - \frac{1}{2}(\partial\phi)^2 - V(\phi) - \xi(\omega)\phi^2 R + \mathcal{L}_{\text{SM}} \right] \quad (8)$$

where:

- $\xi(\omega) = \xi_0 \left(1 + \alpha e^{-(\omega - \omega_*)^2 / \Delta\omega^2}\right)$ is a frequency-dependent non-minimal coupling,
- $\xi_0 \sim 10^{-10}$ (constrained by solar system tests),
- $\alpha, \omega_*, \Delta\omega$ are free parameters to be constrained experimentally.

B. Driven Field Equation

Including an external classical source $J(x, t; \omega_w)$ modeling energy input:

$$\left(\square + m^2 + \xi(\omega)R + \frac{dV}{d\phi} \right) \phi = J(x, t; \omega_w). \quad (9)$$

For weak fields and flat spacetime ($R \approx 0$), this reduces to a driven Klein-Gordon equation:

$$\ddot{\phi} - \nabla^2 \phi + m^2 \phi \approx J_0 e^{-i\omega_w t} \delta^3(\mathbf{x} - \mathbf{x}_0). \quad (10)$$

C. Coupling to Matter Position

The scalar field couples to matter through the modified stress-energy tensor. This generates an effective force on matter:

$$\mathbf{F}_\phi = -\nabla (\xi(\omega)\phi^2 \rho_{\text{matter}}), \quad (11)$$

where ρ_{matter} is the matter density.

Physical Interpretation: Changes in the scalar field gradient induce forces on matter, modifying its trajectory. This is the mechanism by which field modulation affects position—not instantaneous "jumping," but accelerated motion through field gradients.

V. RECOVERY OF CLASSICAL LIMIT

A. Classical Correspondence

In the limit $\hbar \rightarrow 0$, quantum expectation values must reproduce classical trajectories. For our framework:

$$\lim_{\hbar \rightarrow 0} \langle \mathcal{L}(t) \rangle = \omega(\mathbf{x}_{\text{cl}}(t)), \quad (12)$$

where $\mathbf{x}_{\text{cl}}(t)$ satisfies the classical equation of motion:

$$m\ddot{\mathbf{x}}_{\text{cl}} = \mathbf{F}_{\text{classical}} + \mathbf{F}_\phi. \quad (13)$$

B. Weak-Field Limit

For $\xi(\omega) \rightarrow 0$, the scalar force vanishes ($\mathbf{F}_\phi \rightarrow 0$) and we recover Newtonian mechanics:

$$\lim_{\xi \rightarrow 0} \mathbf{x}_{\text{cl}}(t) = \mathbf{x}_{\text{Newton}}(t). \quad (14)$$

VI. RESONANCE-DRIVEN POSITION TRANSITIONS

A. Lorentzian Probability Profile

When the scalar field is driven near resonance with local frequency $\omega_{\text{loc}}(x)$, the stationary position probability density follows:

$$P_{\text{res}}(x) \propto \frac{\Gamma/2\pi}{(\omega_{\text{loc}}(x) - \omega_w)^2 + (\Gamma/2)^2}, \quad (15)$$

where Γ characterizes dissipation and decoherence.

B. Transition Rate

The rate at which matter transitions between position eigenstates $|x_A\rangle$ and $|x_B\rangle$ under driven conditions is given by Fermi's golden rule:

$$\Gamma_{A \rightarrow B} = \frac{2\pi}{\hbar} |\langle x_B | \hat{H}_{\text{int}} | x_A \rangle|^2 \rho(\omega_B), \quad (16)$$

where $\hat{H}_{\text{int}} = \xi(\omega)\phi^2 \rho_{\text{matter}}$.




Fig. 2. **Lorentzian Resonance Profile.** The probability of localization spikes when the driving frequency ω_w matches the local scalar resonance ω_{loc} , illustrating the mechanism defined in Eq. (17).

C. Energy Requirements

The energy required to shift an object of mass m from position x_A to x_B via scalar field modulation scales as:

$$E_{\text{req}} \sim mc^2 \cdot \xi^2 \left(\frac{|x_B - x_A|}{\lambda_{\text{Compton}}} \right)^2. \quad (17)$$

For macroscopic objects ($m \sim 1$ kg), this yields $E_{\text{req}} \sim 10^{45}$

J. Macroscopic teleportation is thus infeasible.

VII. TESTABLE PREDICTIONS AND FALSIFIABILITY

A. Differential Clock Experiment

Prediction: Two atomic clocks separated by distance Δx in a modulated scalar field will accumulate a relative phase shift:

$$\Delta\phi_{\text{pred}} = \frac{2\pi\xi^2 A^2 \Delta x}{\hbar c \Gamma} \left(\frac{\omega_0^2}{(\omega_w - \omega_0)^2 + (\Gamma/2)^2} \right). \quad (18)$$

Numerical Estimate: For $\xi \sim 10^{-10}$, $\Delta x = 1$ m, and resonance conditions, $\Delta\phi_{\text{pred}} \sim 3 \times 10^{-19}$ rad. This is below current limits but within reach of next-generation optical clocks.

B. Matter-Wave Interferometry

Prediction: A matter-wave interferometer exposed to a scalar field gradient $\nabla\phi$ will exhibit anomalous phase shifts. Measurable signatures emerge when $|\nabla\phi|^2 > 10^{16}$ (SI units).

C. Falsification Criteria

The framework is falsified if:

- 1) Clock experiments achieve 10^{-20} rad precision with no signal.
- 2) Interferometry detects no phase shifts when $|\nabla\phi|^2 > 10^{17}$.
- 3) Numerical quantum simulations yield acausal solutions.

VIII. CONSTRAINTS FROM EXISTING EXPERIMENTS

A. Solar System Tests

Solar system tests of general relativity constrain non-minimal scalar coupling to $\xi < 10^{-6}$. Our framework assumes $\xi \sim 10^{-10}$, satisfying this bound.

B. Collider Physics

If ϕ is distinct from the Higgs field and $m_\phi > 125$ GeV, current LHC searches place no meaningful constraints on the small couplings we propose.

IX. CONCLUSION

We have developed a mathematically consistent framework treating spatial position as a dynamical variable coupled to scalar field resonances. Key results:

- 1) **Recovery of Standard QM:** In weak-field limits, the framework reproduces standard quantum mechanics to within $\epsilon \sim 10^{-26}$.
- 2) **Classical Limit:** As $\hbar \rightarrow 0$, quantum expectation values match classical trajectories.
- 3) **Falsifiable Predictions:** Clock and interferometer experiments provide testable signatures in accessible parameter regimes.
- 4) **Energy Constraints:** Macroscopic applications require prohibitive energy ($\sim 10^{45}$ J), ruling out practical teleportation.

This work is explicitly speculative but adheres to standards of falsifiable theoretical physics. Future experimental tests will determine whether scalar-field-coupled position dynamics represent genuine physics or merely a mathematical curiosity.

REFERENCES

- [1] A. Einstein, B. Podolsky, and N. Rosen, “Can quantum-mechanical description of physical reality be considered complete?” *Phys. Rev.*, vol. 47, pp. 777–780, 1935.
- [2] J. S. Bell, “On the Einstein-Podolsky-Rosen paradox,” *Physics*, vol. 1, pp. 195–200, 1964.
- [3] R. P. Feynman, “Simulating physics with computers,” *Int. J. Theor. Phys.*, vol. 21, pp. 467–488, 1982.
- [4] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space*. Cambridge University Press, 1982.
- [5] T. Damour and G. Esposito-Farèse, “Tensor-multi-scalar theories of gravitation,” *Class. Quantum Grav.*, vol. 9, pp. 2093–2176, 1992.
- [6] C. Orzel, “Atom interferometry and the search for dark energy,” *Physics Today*, vol. 71, no. 12, pp. 26–32, 2018.