

Field-Modulated Spatial Localization as a Dynamical Variable

An Effective Matter–Scalar Model of Re-Localization Without Discontinuous Disappearance

Christopher Woodyard¹

¹Vers3Dynamics, Washington, D.C., USA

April 16, 2026

Abstract

We propose an effective theoretical framework in which spatial location is not treated as a fixed intrinsic property of an object, but as a dynamical state emerging from the coupling between matter and an underlying field. In this picture, re-localization is not interpreted as literal disappearance at one coordinate and reappearance at another. Instead, changes in field-controlled localization parameters shift the stable conditions under which matter becomes localized, thereby transferring localization toward a new coordinate defined by the updated dynamics. We formulate the model as a nonrelativistic matter wavefunction coupled to an auxiliary scalar field, derive the resulting matter and field equations, and introduce a normalized field-dependent localization response kernel. Standard quantum mechanics is recovered in the weak-coupling limit, while a two-site reduction gives an explicit mechanism for continuous transfer of localization between competing sites. We derive the classical limit through Ehrenfest dynamics and identify experimentally testable signatures in double-well systems, interferometry, and clock-comparison measurements. The framework is explicitly phenomenological and does not claim macroscopic teleportation or discontinuous ontological replacement. Its aim is to provide a mathematically controlled setting in which spatial localization can be studied as a dynamical, field-sensitive property.

1 Introduction

In standard nonrelativistic quantum mechanics, spatial position is represented by the self-adjoint operator $\hat{\mathbf{x}}$, and localization probabilities are determined by the Born rule. Within that framework, position is ordinarily treated as a primitive observable rather than as an emergent dynamical property. In this paper, we explore a weaker and more conservative extension of that picture: spatial localization is allowed to depend not only on the matter state itself, but also on its coupling to an auxiliary field.

The central claim is not that an object literally ceases to exist at one point and then begins to exist at another. Rather, the claim is that stable localization can be altered by changing the field-dependent parameters that govern where the object is energetically or dynamically favored to localize. On this view, re-localization is a shift in the stable localization state of the coupled matter–field system.

This perspective preserves three important features of a defensible physical model. First, it allows standard quantum mechanics to re-emerge in a weak-coupling limit. Second, it provides a continuous dynamical mechanism rather than an ontological leap. Third, it produces concrete experimental observables rather than purely metaphysical assertions.

The purpose of the present paper is to formulate this field-modulated localization idea in a mathematically explicit way. The main contributions are:

- (i) an effective matter–field action,
- (ii) a field-modified Schrödinger equation,
- (iii) a normalized localization response rule,
- (iv) an explicit two-site toy model of localization transfer,
- (v) recovery of standard quantum mechanics in the weak-coupling limit,
- (vi) derivation of the classical limit, and
- (vii) experimentally testable signatures.

2 Effective Matter–Field Model

2.1 Matter sector

Let $\psi(\mathbf{x}, t)$ be a normalized nonrelativistic matter wavefunction:

$$\int d^3x |\psi(\mathbf{x}, t)|^2 = 1. \quad (1)$$

We take the matter action to be

$$S_\psi = \int dt d^3x \left[i\hbar \psi^* \partial_t \psi - \frac{\hbar^2}{2m} |\nabla \psi|^2 - V(\mathbf{x}) |\psi|^2 \right]. \quad (2)$$

2.2 Scalar field sector

Let $\phi(\mathbf{x}, t)$ be a real auxiliary scalar field. We use the effective field action

$$S_\phi = \int dt d^3x \left[\frac{1}{2} (\partial_t \phi)^2 - \frac{c_\phi^2}{2} |\nabla \phi|^2 - \frac{m_\phi^2}{2} \phi^2 - \frac{\lambda}{4!} \phi^4 + J(\mathbf{x}, t) \phi \right], \quad (3)$$

where c_ϕ is the field propagation speed, m_ϕ is an effective mass scale, λ controls self-interaction, and J is an externally applied source.

2.3 Matter–field coupling

We couple matter to the field through the local density:

$$S_{\text{int}} = -g \int dt d^3x \phi(\mathbf{x}, t) |\psi(\mathbf{x}, t)|^2, \quad (4)$$

where g is a phenomenological coupling constant.

The total action is

$$S_{\text{tot}} = S_{\psi} + S_{\phi} + S_{\text{int}}. \quad (5)$$

2.4 Equations of motion

Variation with respect to ψ^* yields the field-modified Schrödinger equation

$$i\hbar\partial_t\psi(\mathbf{x}, t) = \left[-\frac{\hbar^2}{2m}\nabla^2 + V(\mathbf{x}) + g\phi(\mathbf{x}, t) \right] \psi(\mathbf{x}, t). \quad (6)$$

Variation with respect to ϕ yields

$$\partial_t^2\phi - c_{\phi}^2\nabla^2\phi + m_{\phi}^2\phi + \frac{\lambda}{6}\phi^3 = J(\mathbf{x}, t) - g|\psi(\mathbf{x}, t)|^2. \quad (7)$$

Equation (6) is the key dynamical statement of the model: the auxiliary field modifies the effective localization landscape seen by matter.

3 Localization as a Field-Dependent Response

3.1 Standard Born density

In ordinary quantum mechanics, the position-space probability density is

$$P_{\text{B}}(\mathbf{x}, t) = |\psi(\mathbf{x}, t)|^2. \quad (8)$$

3.2 Local resonance structure

To model field-sensitive localization, we introduce a local resonance frequency

$$\omega_{\text{loc}}(\mathbf{x}, t) = \omega_0 + \beta\phi(\mathbf{x}, t) + \kappa\nabla^2\phi(\mathbf{x}, t), \quad (9)$$

where ω_0 is a baseline scale and β, κ are phenomenological coefficients.

We then define a bounded response profile

$$\mathcal{L}(\mathbf{x}, t; \omega_w) = \frac{(\Gamma/2)^2}{[\omega_w - \omega_{\text{loc}}(\mathbf{x}, t)]^2 + (\Gamma/2)^2}, \quad (10)$$

where ω_w is an externally applied drive frequency and $\Gamma > 0$ is the linewidth.

3.3 Normalized localization response kernel

To ensure positivity and exact normalization, define

$$\chi(\mathbf{x}, t; \omega_w) = \exp[\alpha \mathcal{L}(\mathbf{x}, t; \omega_w)], \quad (11)$$

with dimensionless response strength α .

The observed localization density is then defined by

$$P_{\text{loc}}(\mathbf{x}, t; \omega_w) = \frac{\chi(\mathbf{x}, t; \omega_w) |\psi(\mathbf{x}, t)|^2}{\int d^3x' \chi(\mathbf{x}', t; \omega_w) |\psi(\mathbf{x}', t)|^2}. \quad (12)$$

Equation (12) is the core measurement-level statement of the theory. The field does not erase the Born rule; it biases localization through a normalized response factor.

3.4 Weak-response expansion

Define the Born-weighted expectation of any function $A(\mathbf{x}, t)$ by

$$\langle A \rangle_\psi = \int d^3x A(\mathbf{x}, t) |\psi(\mathbf{x}, t)|^2. \quad (13)$$

Expanding Eq. (12) for small α gives

$$P_{\text{loc}}(\mathbf{x}, t; \omega_w) = P_{\text{B}}(\mathbf{x}, t) [1 + \alpha (\mathcal{L}(\mathbf{x}, t; \omega_w) - \langle \mathcal{L} \rangle_\psi)] + \mathcal{O}(\alpha^2). \quad (14)$$

This form is normalized automatically because the subtraction of $\langle \mathcal{L} \rangle_\psi$ removes the net first-order shift.

4 Two-Site Toy Model of Localization Transfer

To make the mechanism explicit, consider a two-site reduction with localized basis states $\{|A\rangle, |B\rangle\}$. The effective Hamiltonian is

$$H_2(t) = \begin{pmatrix} E_A + g\phi_A(t) & \Delta \\ \Delta & E_B + g\phi_B(t) \end{pmatrix}, \quad (15)$$

where E_A, E_B are bare site energies, $\phi_A(t) = \phi(\mathbf{x}_A, t)$ and $\phi_B(t) = \phi(\mathbf{x}_B, t)$ are the local field values, and Δ is the inter-site mixing amplitude.

Define the detuning

$$\delta(t) = (E_B - E_A) + g[\phi_B(t) - \phi_A(t)]. \quad (16)$$

Then the instantaneous eigenstates are controlled by the mixing angle $\theta(t)$ defined through

$$\tan(2\theta(t)) = \frac{2\Delta}{\delta(t)}. \quad (17)$$

The lower-energy eigenstate can be written as

$$|-\rangle_t = \cos\theta(t) |A\rangle - \sin\theta(t) |B\rangle, \quad (18)$$

with site occupation probabilities

$$P_A(t) = \cos^2\theta(t), \quad P_B(t) = \sin^2\theta(t). \quad (19)$$

If the field control is varied so that $\delta(t)$ changes sign adiabatically, the dominant localization shifts

from site A to site B . This is the simplest mathematical realization of the claim that localization is transferred by updated dynamics rather than by literal disappearance and reappearance.

5 Recovery of Standard Quantum Mechanics

A viable extension must reduce to standard quantum mechanics under ordinary conditions.

5.1 Uniform weak field

Suppose the field is spatially uniform and weak:

$$\phi(\mathbf{x}, t) = \phi_0(t), \quad |g\phi_0| \ll \|V\|. \quad (20)$$

Then Eq. (6) becomes

$$i\hbar\partial_t\psi = \left[-\frac{\hbar^2}{2m}\nabla^2 + V(\mathbf{x}) + g\phi_0(t) \right] \psi. \quad (21)$$

The term $g\phi_0(t)$ contributes only a global phase. Writing

$$\psi(\mathbf{x}, t) = \exp\left[-\frac{i}{\hbar} \int_0^t g\phi_0(t') dt'\right] \tilde{\psi}(\mathbf{x}, t), \quad (22)$$

we obtain

$$i\hbar\partial_t\tilde{\psi} = \left[-\frac{\hbar^2}{2m}\nabla^2 + V(\mathbf{x}) \right] \tilde{\psi}. \quad (23)$$

Thus the matter dynamics reduce exactly to standard Schrödinger evolution.

5.2 Weak localization response

If the response kernel is weak or approximately uniform, then χ becomes nearly constant and Eq. (12) reduces to

$$P_{\text{loc}}(\mathbf{x}, t; \omega_w) \rightarrow |\psi(\mathbf{x}, t)|^2. \quad (24)$$

Hence both dynamics and observed localization reduce to standard quantum mechanics in the weak-coupling limit.

6 Classical Limit and Stable Re-Localization

The Ehrenfest relations derived from Eq. (6) are

$$\frac{d}{dt}\langle\hat{\mathbf{x}}\rangle = \frac{1}{m}\langle\hat{\mathbf{p}}\rangle, \quad \frac{d}{dt}\langle\hat{\mathbf{p}}\rangle = -\langle\nabla(V + g\phi)\rangle. \quad (25)$$

For a narrow wavepacket, these reduce to the classical equation

$$m\ddot{\mathbf{x}}_{\text{cl}} = -\nabla[V(\mathbf{x}_{\text{cl}}) + g\phi(\mathbf{x}_{\text{cl}}, t)]. \quad (26)$$

This provides the classical interpretation of the theory. The field alters the effective potential landscape,

$$V_{\text{eff}}(\mathbf{x}, t) = V(\mathbf{x}) + g \phi(\mathbf{x}, t), \quad (27)$$

and therefore changes the locations of stable minima. Re-localization is the shift of those stable minima under updated field conditions.

If a local minimum moves from a neighborhood of $\mathbf{x}_A$ to one near $\mathbf{x}_B$, a localized object can be transferred from one stable coordinate regime to another. The theory therefore describes localization transfer through field-shaped stability rather than through discontinuous ontological replacement.

7 Experimentally Testable Consequences

7.1 Double-well localization transfer

The cleanest direct test is a double-well potential in which local site energies can be biased by an external field. In the two-site model, the adiabatic occupation imbalance is

$$z(t) = P_A(t) - P_B(t) = \cos(2\theta(t)) = \frac{\delta(t)}{\sqrt{\delta(t)^2 + 4\Delta^2}}. \quad (28)$$

A field-driven sign change in $\delta(t)$ should reverse the localization bias. This is a precise, low-dimensional test of the paper's central claim.

7.2 Matter-wave interferometry

If two interferometer arms follow trajectories $\mathbf{x}_1(t)$ and $\mathbf{x}_2(t)$, the field-induced phase shift is

$$\Delta\varphi_\phi = \frac{g}{\hbar} \int_0^T [\phi(\mathbf{x}_1(t), t) - \phi(\mathbf{x}_2(t), t)] dt. \quad (29)$$

A nonzero $\Delta\varphi_\phi$ produces a shift in the interference pattern correlated with the applied field drive.

7.3 Clock-comparison experiments

If the field shifts an internal transition frequency by $\delta\omega = \eta\phi$, then two clocks at $\mathbf{x}_A$ and $\mathbf{x}_B$ accumulate a relative phase

$$\Delta\Phi_{AB}(T) = \eta \int_0^T [\phi(\mathbf{x}_A, t) - \phi(\mathbf{x}_B, t)] dt. \quad (30)$$

This predicts a differential phase offset synchronized with controlled field modulation.

7.4 Localization-statistics anomaly

Repeated localization measurements under a driven field should exhibit deviations from the Born density described by Eq. (14). The relevant observable is the field-correlated statistical deviation

$$\delta P(\mathbf{x}, t; \omega_w) = P_{\text{loc}}(\mathbf{x}, t; \omega_w) - P_B(\mathbf{x}, t). \quad (31)$$

8 Limitations

The present framework is deliberately modest in scope.

First, it is phenomenological. The localization response kernel χ is introduced at the effective level rather than derived from a full detector theory.

Second, it is not a theory of gravity or spacetime emergence. The scalar field is an auxiliary localization-modulating field, not a claim about fundamental geometry.

Third, the model does not imply macroscopic teleportation. Its mechanism is a shift in stable localization structure, not the abrupt destruction and recreation of matter.

Fourth, the two-site reduction is illustrative rather than fundamental. Its purpose is to make the localization-transfer mechanism explicit in the simplest possible setting.

9 Discussion

The advantage of the present formulation is that it captures the intuitive claim behind the phrase “location is a dynamical variable” without overcommitting to a discontinuous ontology. Matter is still described by a wavefunction, and localization is still probabilistic, but the field changes the effective conditions under which localization becomes stable and experimentally favored.

This is a more disciplined claim than saying that an object literally vanishes from one point and reappears at another. It remains compatible with standard quantum mechanics in the appropriate limit, it admits a clear classical correspondence, and it produces experimentally recognizable signatures.

The price of this discipline is that the theory is more modest. It does not attempt to solve the ontology of measurement once and for all. It does not derive the scalar field from a deeper microscopic theory. And it does not yet explain why a particular localization response kernel should be fundamental. Those are open problems for later work.

10 Conclusion

We have developed an effective matter–field framework in which spatial localization is treated as a dynamical, field-sensitive property rather than as a strictly fixed intrinsic attribute. The model is built around three linked ideas:

- (1) matter evolves under a field-modified Schrödinger equation,
- (2) observed localization is biased by a normalized field-dependent response kernel, and
- (3) stable localization can be transferred when the field shifts the underlying energetic and dynamical landscape.

The theory reduces to standard quantum mechanics in the weak-coupling limit, admits a transparent two-site realization of localization transfer, and yields experimentally testable signatures in double-well systems, interferometry, and clock comparisons. Its strongest claim is therefore precise but restrained: location need not be a permanently fixed intrinsic property of an object, but it can be modeled as a dynamical state emerging from the object’s coupling to underlying fields.

References

- [1] J. von Neumann, *Mathematical Foundations of Quantum Mechanics*. Princeton University Press, 1955.
- [2] J. J. Sakurai and J. Napolitano, *Modern Quantum Mechanics*, 2nd ed. Cambridge University Press, 2017.
- [3] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems*. Oxford University Press, 2002.
- [4] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space*. Cambridge University Press, 1982.
- [5] C. P. Burgess, *Introduction to Effective Field Theory*. Cambridge University Press, 2020.
- [6] A. D. Cronin, J. Schmiedmayer, and D. E. Pritchard, “Optics and interferometry with atoms and molecules,” *Reviews of Modern Physics*, vol. 81, no. 3, pp. 1051–1129, 2009.
- [7] A. D. Ludlow, M. M. Boyd, J. Ye, E. Peik, and P. O. Schmidt, “Optical atomic clocks,” *Reviews of Modern Physics*, vol. 87, no. 2, pp. 637–701, 2015.
- [8] W. H. Zurek, “Decoherence, einselection, and the quantum origins of the classical,” *Reviews of Modern Physics*, vol. 75, no. 3, pp. 715–775, 2003.