

# Tensor Products of Pointed Sets

The Clowder Project Authors

May 3, 2024

**00C3** In this chapter we introduce, construct, and study tensor products of pointed sets. The most well-known among these is the *smash product of pointed sets*

$$\wedge: \mathbf{Sets}_* \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*,$$

introduced in [Section 5.1](#), defined via a universal property as inducing a bijection between the following data:

- Pointed maps  $f: X \wedge Y \rightarrow Z$ .
- Maps of sets  $f: X \times Y \rightarrow Z$  satisfying

$$\begin{aligned} f(x_0, y) &= z_0, \\ f(x, y_0) &= z_0 \end{aligned}$$

for each  $x \in X$  and each  $y \in Y$ .

As it turns out, however, dropping either of the *bilinearity* conditions

$$\begin{aligned} f(x_0, y) &= z_0, \\ f(x, y_0) &= z_0 \end{aligned}$$

while retaining the other leads to two other tensor products of pointed sets,

$$\begin{aligned} \triangleleft: \mathbf{Sets}_* \times \mathbf{Sets}_* &\rightarrow \mathbf{Sets}_*, \\ \triangleright: \mathbf{Sets}_* \times \mathbf{Sets}_* &\rightarrow \mathbf{Sets}_*, \end{aligned}$$

called the *left* and *right tensor products of pointed sets*. In contrast to  $\wedge$ , which turns out to endow  $\mathbf{Sets}_*$  with a monoidal category structure ([Proposition 5.9.1.1](#)), these do not admit invertible associators and unitors, but do endow  $\mathbf{Sets}_*$  with the structure of a skew monoidal category, however ([Propositions 3.8.1.1](#) and [4.8.1.1](#)).

Finally, in addition to the tensor products  $\triangleleft$ ,  $\triangleright$ , and  $\wedge$ , we also have a “tensor product” of the form

$$\odot: \mathbf{Sets} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*,$$

called the *tensor* of sets with pointed sets. All in all, these tensor products assemble into a family of functors of the form

$$\begin{aligned} \otimes_{k,\ell}: \mathbf{Mon}_{\mathbb{E}_k}(\mathbf{Sets}) \times \mathbf{Mon}_{\mathbb{E}_\ell}(\mathbf{Sets}) &\rightarrow \mathbf{Mon}_{\mathbb{E}_{k+\ell}}(\mathbf{Sets}), \\ \triangleleft_{i,k}: \mathbf{Mon}_{\mathbb{E}_k}(\mathbf{Sets}) \times \mathbf{Mon}_{\mathbb{E}_k}(\mathbf{Sets}) &\rightarrow \mathbf{Mon}_{\mathbb{E}_k}(\mathbf{Sets}), \\ \triangleright_{i,k}: \mathbf{Mon}_{\mathbb{E}_k}(\mathbf{Sets}) \times \mathbf{Mon}_{\mathbb{E}_k}(\mathbf{Sets}) &\rightarrow \mathbf{Mon}_{\mathbb{E}_k}(\mathbf{Sets}), \end{aligned}$$

where  $k, \ell, i \in \mathbb{N}$  with  $i \leq k - 1$ . Together with the Cartesian product  $\times$  of  $\mathbf{Sets}$ , the tensor products studied in this chapter form the cases:

- $(k, \ell) = (-1, -1)$  for the Cartesian product of  $\mathbf{Sets}$ ;
- $(k, \ell) = (0, -1)$  and  $(-1, 0)$  for the tensor of sets with pointed sets of [Definition 2.1.1.1](#);
- $(i, k) = (-1, 0)$  for the left and right tensor products of pointed sets of [Sections 3](#) and [4](#);
- $(k, \ell) = (-1, -1)$  for the smash product of pointed sets of [Section 5](#).

In this chapter, we will carefully define and study bilinearity for pointed sets, as well as all the tensor products described above. Then, in [??](#), we will extend these to tensor products involving also monoids and commutative monoids, which will end up covering all cases up to  $k, \ell \leq 2$ , and hence *all* cases since  $\mathbb{E}_k$ -monoids on  $\mathbf{Sets}$  are the same as  $\mathbb{E}_2$ -monoids on  $\mathbf{Sets}$  when  $k \geq 2$ .

## Contents

<b>1</b>	<b>Bilinear Morphisms of Pointed Sets.....</b>	<b>4</b>
1.1	Left Bilinear Morphisms of Pointed Sets.....	4
1.2	Right Bilinear Morphisms of Pointed Sets.....	5
1.3	Bilinear Morphisms of Pointed Sets.....	5
<b>2</b>	<b>Tensors and Cotensors of Pointed Sets by Sets.....</b>	<b>7</b>
2.1	Tensors of Pointed Sets by Sets.....	7
2.2	Cotensors of Pointed Sets by Sets.....	15

<b>3</b>	<b>The Left Tensor Product of Pointed Sets.....</b>	<b>22</b>
3.1	Foundations.....	22
3.2	The Left Internal Hom of Pointed Sets.....	26
3.3	The Left Skew Unit.....	28
3.4	The Left Skew Associator.....	28
3.5	The Left Skew Left Unitor.....	31
3.6	The Left Skew Right Unitor.....	33
3.7	The Diagonal.....	35
3.8	The Left Skew Monoidal Structure on Pointed Sets Associated	
to $\triangleleft$ .....		36
3.9	Monoids With Respect to the Left Tensor Product of Pointed	
Sets.....		41
<b>4</b>	<b>The Right Tensor Product of Pointed Sets.....</b>	<b>45</b>
4.1	Foundations.....	45
4.2	The Right Internal Hom of Pointed Sets.....	49
4.3	The Right Skew Unit.....	51
4.4	The Right Skew Associator.....	51
4.5	The Right Skew Left Unitor.....	54
4.6	The Right Skew Right Unitor.....	55
4.7	The Diagonal.....	58
4.8	The Right Skew Monoidal Structure on Pointed Sets Associated	
to $\triangleright$ .....		59
4.9	Monoids With Respect to the Right Tensor Product of Pointed	
Sets.....		63
<b>5</b>	<b>The Smash Product of Pointed Sets.....</b>	<b>67</b>
5.1	Foundations.....	67
5.2	The Internal Hom of Pointed Sets.....	77
5.3	The Monoidal Unit.....	80
5.4	The Associator.....	80
5.5	The Left Unitor.....	82
5.6	The Right Unitor.....	84
5.7	The Symmetry.....	87
5.8	The Diagonal.....	89
5.9	The Monoidal Structure on Pointed Sets Associated to $\wedge$ .....	92
5.10	Universal Properties of the Smash Product of Pointed Sets I...	97
5.11	Universal Properties of the Smash Product of Pointed Sets II..	97
5.12	Monoids With Respect to the Smash Product of Pointed Sets..	98
5.13	Comonoids With Respect to the Smash Product of Pointed Sets	98

<b>6</b>	<b>Miscellany.....</b>	<b>98</b>
6.1	The Smash Product of a Family of Pointed Sets.....	98
<b>A</b>	<b>Other Chapters.....</b>	<b>99</b>

## 00C4 1 Bilinear Morphisms of Pointed Sets

### 00C5 1.1 Left Bilinear Morphisms of Pointed Sets

Let  $(X, x_0)$ ,  $(Y, y_0)$ , and  $(Z, z_0)$  be pointed sets.

**00C6 Definition 1.1.1.1.** A **left bilinear morphism of pointed sets from  $(X \times Y, (x_0, y_0))$  to  $(Z, z_0)$**  is a map of sets

$$f: X \times Y \rightarrow Z$$

satisfying the following condition:<sup>1,2</sup>

( $\star$ ) *Left Unital Bilinearity.* The diagram

$$\begin{array}{ccccc}
 & & \text{pt} \times \text{pt} & & \\
 & \text{id}_{\text{pt}} \times \epsilon_Y \nearrow & & \searrow \sim & \\
 \text{pt} \times Y & & & & \text{pt} \\
 & \downarrow [x_0] \times \text{id}_Y & & & \downarrow [z_0] \\
 X \times Y & \xrightarrow{f} & Z & & 
 \end{array}$$

commutes, i.e. for each  $y \in Y$ , we have

$$f(x_0, y) = z_0.$$

**00C7 Definition 1.1.1.2.** The **set of left bilinear morphisms of pointed sets from  $(X \times Y, (x_0, y_0))$  to  $(Z, z_0)$**  is the set  $\text{Hom}_{\text{Sets}_*}^{\otimes, \text{L}}(X \times Y, Z)$  defined by

$$\text{Hom}_{\text{Sets}_*}^{\otimes, \text{L}}(X \times Y, Z) \stackrel{\text{def}}{=} \{f \in \text{Hom}_{\text{Sets}}(X \times Y, Z) \mid f \text{ is left bilinear}\}.$$

<sup>1</sup>*Slogan:* The map  $f$  is left bilinear if it preserves basepoints in its first argument.

<sup>2</sup>Succinctly,  $f$  is bilinear if we have

$$f(x_0, y) = z_0$$

**00C8 1.2 Right Bilinear Morphisms of Pointed Sets**

Let  $(X, x_0)$ ,  $(Y, y_0)$ , and  $(Z, z_0)$  be pointed sets.

**00C9 Definition 1.2.1.1.** A **right bilinear morphism of pointed sets** from  $(X \times Y, (x_0, y_0))$  **to**  $(Z, z_0)$  is a map of sets

$$f: X \times Y \rightarrow Z$$

satisfying the following condition:<sup>3,4</sup>

( $\star$ ) *Right Unital Bilinearity.* The diagram

$$\begin{array}{ccccc}
 & & \text{pt} \times \text{pt} & & \\
 & \nearrow \epsilon_X \times \text{id}_{\text{pt}} & & \searrow \sim & \\
 X \times \text{pt} & & & & \text{pt} \\
 \downarrow \text{id}_X \times [y_0] & & & & \downarrow [z_0] \\
 X \times Y & \xrightarrow{f} & Z & & 
 \end{array}$$

commutes, i.e. for each  $x \in X$ , we have

$$f(x, y_0) = z_0.$$

**00CA Definition 1.2.1.2.** The **set of right bilinear morphisms of pointed sets from**  $(X \times Y, (x_0, y_0))$  **to**  $(Z, z_0)$  is the set  $\text{Hom}_{\text{Sets}_*}^{\otimes, \text{R}}(X \times Y, Z)$  defined by

$$\text{Hom}_{\text{Sets}_*}^{\otimes, \text{R}}(X \times Y, Z) \stackrel{\text{def}}{=} \{f \in \text{Hom}_{\text{Sets}}(X \times Y, Z) \mid f \text{ is right bilinear}\}.$$

**00CB 1.3 Bilinear Morphisms of Pointed Sets**

Let  $(X, x_0)$ ,  $(Y, y_0)$ , and  $(Z, z_0)$  be pointed sets.

---

for each  $y \in Y$ .

<sup>3</sup>*Slogan:* The map  $f$  is right bilinear if it preserves basepoints in its second argument.

<sup>4</sup>Succinctly,  $f$  is bilinear if we have

$$f(x, y_0) = z_0$$

for each  $x \in X$ .

**00CC Definition 1.3.1.1.** A **bilinear morphism of pointed sets** from  $(X \times Y, (x_0, y_0))$  to  $(Z, z_0)$  is a map of sets

$$f: X \times Y \rightarrow Z$$

that is both left bilinear and right bilinear.

**00CD Remark 1.3.1.2.** In detail, a **bilinear morphism of pointed sets** from  $(X \times Y, (x_0, y_0))$  to  $(Z, z_0)$  is a map of sets

$$f: (X \times Y, (x_0, y_0)) \rightarrow (Z, z_0)$$

satisfying the following conditions:<sup>5,6</sup>

1. *Left Unital Bilinearity.* The diagram

$$\begin{array}{ccc}
 & \text{pt} \times \text{pt} & \\
 \text{id}_{\text{pt}} \times \epsilon_Y \nearrow & & \searrow \sim \\
 \text{pt} \times Y & & \text{pt} \\
 [\text{x}_0] \times \text{id}_Y \searrow & & \nearrow [\text{z}_0] \\
 X \times Y & \xrightarrow{f} & Z
 \end{array}$$

commutes, i.e. for each  $y \in Y$ , we have

$$f(x_0, y) = z_0.$$

2. *Right Unital Bilinearity.* The diagram

$$\begin{array}{ccc}
 & \text{pt} \times \text{pt} & \\
 \epsilon_X \times \text{id}_{\text{pt}} \nearrow & & \searrow \sim \\
 X \times \text{pt} & & \text{pt} \\
 \text{id}_X \times [\text{y}_0] \searrow & & \nearrow [\text{z}_0] \\
 X \times Y & \xrightarrow{f} & Z
 \end{array}$$

<sup>5</sup>*Slogan:* The map  $f$  is bilinear if it preserves basepoints in each argument.

<sup>6</sup>Succinctly,  $f$  is bilinear if we have

$$\begin{aligned}
 f(x_0, y) &= z_0, \\
 f(x, y_0) &= z_0
 \end{aligned}$$

commutes, i.e. for each  $x \in X$ , we have

$$f(x, y_0) = z_0.$$

**00CE Definition 1.3.1.3.** The set of bilinear morphisms of pointed sets from  $(X \times Y, (x_0, y_0))$  to  $(Z, z_0)$  is the set  $\text{Hom}_{\text{Sets}_*}^{\otimes}(X \times Y, Z)$  defined by

$$\text{Hom}_{\text{Sets}_*}^{\otimes}(X \times Y, Z) \stackrel{\text{def}}{=} \{f \in \text{Hom}_{\text{Sets}}(X \times Y, Z) \mid f \text{ is bilinear}\}.$$

## 00CF 2 Tensors and Cotensors of Pointed Sets by Sets

### 00CG 2.1 Tensors of Pointed Sets by Sets

Let  $(X, x_0)$  be a pointed set and let  $A$  be a set.

**00CH Definition 2.1.1.1.** The tensor of  $(X, x_0)$  by  $A$ <sup>7</sup> is the pointed set<sup>8</sup>  $A \odot (X, x_0)$  satisfying the following universal property:

(UP) We have a bijection

$$\text{Sets}_*(A \odot X, K) \cong \text{Sets}(A, \text{Sets}_*(X, K)),$$

natural in  $(K, k_0) \in \text{Obj}(\text{Sets}_*)$ .

**00CJ Remark 2.1.1.2.** The universal property in Definition 2.1.1.1 is equivalent to the following one:

(UP) We have a bijection

$$\text{Sets}_*(A \odot X, K) \cong \text{Sets}_{\mathbb{E}_0}^{\otimes}(A \times X, K),$$

natural in  $(K, k_0) \in \text{Obj}(\text{Sets}_*)$ , where  $\text{Sets}_{\mathbb{E}_0}^{\otimes}(A \times X, K)$  is the set defined by

$$\text{Sets}_{\mathbb{E}_0}^{\otimes}(A \times X, K) \stackrel{\text{def}}{=} \left\{ f \in \text{Sets}(A \times X, K) \mid \begin{array}{l} \text{for each } a \in A, \text{ we} \\ \text{have } f(a, x_0) = k_0 \end{array} \right\}.$$

for each  $x \in X$  and each  $y \in Y$ .

<sup>7</sup>Further Terminology: Also called the **copower of  $(X, x_0)$  by  $A$** .

<sup>8</sup>Further Notation: Often written  $A \odot X$  for simplicity.

*Proof.* We claim we have a bijection

$$\mathbf{Sets}(A, \mathbf{Sets}_*(X, K)) \cong \mathbf{Sets}_{\mathbb{E}_0}^{\otimes}(A \times X, K)$$

natural in  $(K, k_0) \in \mathbf{Obj}(\mathbf{Sets}_*)$ . Indeed, this bijection is a restriction of the bijection

$$\mathbf{Sets}(A, \mathbf{Sets}(X, K)) \cong \mathbf{Sets}(A \times X, K)$$

of **Constructions With Sets, Item 2** of **Proposition 1.3.1.2**:

- A map

$$\begin{aligned} \xi: A &\longrightarrow \mathbf{Sets}_*(X, K), \\ a &\mapsto (\xi_a: X \rightarrow K), \end{aligned}$$

in  $\mathbf{Sets}(A, \mathbf{Sets}_*(X, K))$  gets sent to the map

$$\xi^\dagger: A \times X \rightarrow K$$

defined by

$$\xi^\dagger(a, x) \stackrel{\text{def}}{=} \xi_a(x)$$

for each  $(a, x) \in A \times X$ , which indeed lies in  $\mathbf{Sets}_{\mathbb{E}_0}^{\otimes}(A \times X, K)$ , as we have

$$\begin{aligned} \xi^\dagger(a, x_0) &\stackrel{\text{def}}{=} \xi_a(x_0) \\ &\stackrel{\text{def}}{=} k_0 \end{aligned}$$

for each  $a \in A$ , where we have used that  $\xi_a \in \mathbf{Sets}_*(X, K)$  is a morphism of pointed sets.

- Conversely, a map

$$\xi: A \times X \rightarrow K$$

in  $\mathbf{Sets}_{\mathbb{E}_0}^{\otimes}(A \times X, K)$  gets sent to the map

$$\begin{aligned} \xi^\dagger: A &\longrightarrow \mathbf{Sets}_*(X, K), \\ a &\mapsto (\xi_a^\dagger: X \rightarrow K), \end{aligned}$$

where

$$\xi_a^\dagger: X \rightarrow K$$

is the map defined by

$$\xi_a^\dagger(x) \stackrel{\text{def}}{=} \xi(a, x)$$



for each  $x \in X$ , and indeed lies in  $\mathbf{Sets}_*(X, K)$ , as we have

$$\begin{aligned}\xi_a^\dagger(x_0) &\stackrel{\text{def}}{=} \xi(a, x_0) \\ &\stackrel{\text{def}}{=} k_0.\end{aligned}$$

This finishes the proof.  $\square$

**00CK Construction 2.1.1.3.** Concretely, the **tensor of  $(X, x_0)$  by  $A$**  is the pointed set  $A \odot (X, x_0)$  consisting of:

- *The Underlying Set.* The set  $A \odot X$  given by

$$A \odot X \cong \bigvee_{a \in A} (X, x_0),$$

where  $\bigvee_{a \in A} (X, x_0)$  is the wedge product of the  $A$ -indexed family  $((X, x_0))_{a \in A}$  of **Pointed Sets, Definition 3.2.1.1**.

- *The Basepoint.* The point  $[(a, x_0)] = [(a', x_0)]$  of  $\bigvee_{a \in A} (X, x_0)$ .

*Proof.* (Proven below in a bit.)  $\square$

**00CL Notation 2.1.1.4.** We write  $a \odot x$  for the element  $[(a, x)]$  of

$$\begin{aligned}A \odot X &\cong \bigvee_{a \in A} (X, x_0) \\ &\stackrel{\text{def}}{=} \left( \prod_{i \in I} X_i \right) / \sim.\end{aligned}$$

**00CM Remark 2.1.1.5.** Taking the tensor of any element of  $A$  with the basepoint  $x_0$  of  $X$  leads to the same element in  $A \odot X$ , i.e. we have

$$a \odot x_0 = a' \odot x_0,$$

for each  $a, a' \in A$ . This is due to the equivalence relation  $\sim$  on

$$\bigvee_{a \in A} (X, x_0) \stackrel{\text{def}}{=} \prod_{a \in A} X / \sim$$

identifying  $(a, x_0)$  with  $(a', x_0)$ , so that the equivalence class  $a \odot x_0$  is independent from the choice of  $a \in A$ .

*Proof.* We claim we have a bijection

$$\mathbf{Sets}_*(A \odot X, K) \cong \mathbf{Sets}(A, \mathbf{Sets}_*(X, K))$$

natural in  $(K, k_0) \in \mathbf{Obj}(\mathbf{Sets}_*)$ .

- *Map I.* We define a map

$$\Phi_K: \mathbf{Sets}_*(A \odot X, K) \rightarrow \mathbf{Sets}(A, \mathbf{Sets}_*(X, K))$$

by sending a morphism of pointed sets

$$\xi: (A \odot X, a \odot x_0) \rightarrow (K, k_0)$$

to the map of sets

$$\begin{aligned} \xi^\dagger: A &\longrightarrow \mathbf{Sets}_*(X, K), \\ a &\mapsto (\xi_a: X \rightarrow K), \end{aligned}$$

where

$$\xi_a: (X, x_0) \rightarrow (K, k_0)$$

is the morphism of pointed sets defined by

$$\xi_a(x) \stackrel{\text{def}}{=} \xi(a \odot x)$$

for each  $x \in X$ . Note that we have

$$\begin{aligned} \xi_a(x_0) &\stackrel{\text{def}}{=} \xi(a \odot x_0) \\ &= k_0, \end{aligned}$$

so that  $\xi_a$  is indeed a morphism of pointed sets, where we have used that  $\xi$  is a morphism of pointed sets.

- *Map II.* We define a map

$$\Psi_K: \mathbf{Sets}(A, \mathbf{Sets}_*(X, K)) \rightarrow \mathbf{Sets}_*(A \odot X, K)$$

given by sending a map

$$\begin{aligned} \xi: A &\longrightarrow \mathbf{Sets}_*(X, K), \\ a &\mapsto (\xi_a: X \rightarrow K), \end{aligned}$$

to the morphism of pointed sets

$$\xi^\dagger: (A \odot X, a \odot x_0) \rightarrow (K, k_0)$$

defined by

$$\xi^\dagger(a \odot x) \stackrel{\text{def}}{=} \xi_a(x)$$

for each  $a \odot x \in A \odot X$ . Note that  $\xi^\dagger$  is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \xi^\dagger(a \odot x_0) &\stackrel{\text{def}}{=} \xi_a(x_0) \\ &= k_0, \end{aligned}$$

where we have used that  $\xi(a) \in \mathbf{Sets}_*(X, K)$  is a morphism of pointed sets.

- *Invertibility I.* We claim that

$$\Psi_K \circ \Phi_K = \text{id}_{\mathbf{Sets}_*(A \odot X, K)}.$$

Indeed, given a morphism of pointed sets

$$\xi: (A \odot X, a \odot x_0) \rightarrow (K, k_0),$$

we have

$$\begin{aligned} [\Psi_K \circ \Phi_K](\xi) &= \Psi_K(\Phi_K(\xi)) \\ &= \Psi_K(\llbracket a \mapsto \llbracket x \mapsto \xi(a \odot x) \rrbracket \rrbracket) \\ &= \Psi_K(\llbracket a' \mapsto \llbracket x' \mapsto \xi(a' \odot x') \rrbracket \rrbracket) \\ &= \llbracket a \odot x \mapsto \text{ev}_x(\text{ev}_a(\llbracket a' \mapsto \llbracket x' \mapsto \xi(a' \odot x') \rrbracket \rrbracket)) \rrbracket \\ &= \llbracket a \odot x \mapsto \text{ev}_x(\llbracket x' \mapsto \xi(a \odot x') \rrbracket) \rrbracket \\ &= \llbracket a \odot x \mapsto \xi(a \odot x) \rrbracket \\ &= \xi. \end{aligned}$$

- *Invertibility II.* We claim that

$$\Phi_K \circ \Psi_K = \text{id}_{\mathbf{Sets}(A, \mathbf{Sets}_*(X, K))}.$$

Indeed, given a morphism  $\xi: A \rightarrow \mathbf{Sets}_*(X, K)$ , we have

$$\begin{aligned} [\Phi_K \circ \Psi_K](\xi) &= \Phi_K(\Psi_K(\xi)) \\ &= \Phi_K(\llbracket a \odot x \mapsto \xi_a(x) \rrbracket) \\ &= \llbracket a \mapsto \llbracket x \mapsto \xi_a(x) \rrbracket \rrbracket \\ &= \llbracket a \mapsto \xi(a) \rrbracket \\ &= \xi. \end{aligned}$$

- *Naturality of  $\Phi$ .* We need to show that, given a morphism of pointed sets

$$\phi: (K, k_0) \rightarrow (K', k'_0),$$

the diagram

$$\begin{array}{ccc} \mathbf{Sets}_*(A \odot X, K) & \xrightarrow{\Phi_K} & \mathbf{Sets}(A, \mathbf{Sets}_*(X, K)) \\ \phi_* \downarrow & & \downarrow (\phi_*)_* \\ \mathbf{Sets}_*(A \odot X, K') & \xrightarrow{\Phi_{K'}} & \mathbf{Sets}(A, \mathbf{Sets}_*(X, K')) \end{array}$$

commutes. Indeed, given a morphism of pointed sets

$$\xi: (A \odot X, a \odot x_0) \rightarrow (K, k_0),$$

we have

$$\begin{aligned} [\Phi_{K'} \circ \phi_*](\xi) &= \Phi_{K'}(\phi_*(\xi)) \\ &= \Phi_{K'}(\phi \circ \xi) \\ &= (\phi \circ \xi)^\dagger \\ &= \llbracket a \mapsto \phi \circ \xi(a \odot -) \rrbracket \\ &= \llbracket a \mapsto \phi_*(\xi(a \odot -)) \rrbracket \\ &= (\phi_*)_* (\llbracket a \mapsto \xi(a \odot -) \rrbracket) \\ &= (\phi_*)_*(\Phi_K(\xi)) \\ &= [(\phi_*)_* \circ \Phi_K](\xi). \end{aligned}$$

- *Naturality of  $\Psi$ .* Since  $\Phi$  is natural and  $\Phi$  is a componentwise inverse to  $\Psi$ , it follows from **Categories, Item 2** of **Proposition 8.6.1.2** that  $\Psi$  is also natural.

This finishes the proof.  $\square$

**00CN Proposition 2.1.1.6.** Let  $(X, x_0)$  be a pointed set and let  $A$  be a set.

- 00CP** 1. *Functoriality.* The assignments  $A, (X, x_0), (A, (X, x_0))$  define functors

$$\begin{aligned} A \odot -: \mathbf{Sets}_* &\rightarrow \mathbf{Sets}_*, \\ - \odot X: \mathbf{Sets} &\rightarrow \mathbf{Sets}_*, \\ -_1 \odot -_2: \mathbf{Sets} \times \mathbf{Sets}_* &\rightarrow \mathbf{Sets}_*. \end{aligned}$$

In particular, given:

- A map of sets  $f: A \rightarrow B$ ;
- A pointed map  $\phi: (X, x_0) \rightarrow (Y, y_0)$ ;

the induced map

$$f \odot \phi: A \odot X \rightarrow B \odot Y$$

is given by

$$[f \odot \phi](a \odot x) \stackrel{\text{def}}{=} f(a) \odot \phi(x)$$

for each  $a \odot x \in A \odot X$ .

**00CQ** 2. *Adjointness I.* We have an adjunction

$$(- \odot X \dashv \text{Sets}_*(X, -)): \text{Sets} \begin{array}{c} \xrightarrow{- \odot X} \\ \perp \\ \xleftarrow{\text{Sets}_*(X, -)} \end{array} \text{Sets}_*,$$

witnessed by a bijection

$$\text{Sets}_*(A \odot X, K) \cong \text{Sets}(A, \text{Sets}_*(X, K)),$$

natural in  $A \in \text{Obj}(\text{Sets})$  and  $X, Y \in \text{Obj}(\text{Sets}_*)$ .

**00CR** 3. *Adjointness II.* We have an adjunctions

$$(A \odot - \dashv A \pitchfork -): \text{Sets}_* \begin{array}{c} \xrightarrow{A \odot -} \\ \perp \\ \xleftarrow{A \pitchfork -} \end{array} \text{Sets}_*,$$

witnessed by a bijection

$$\text{Hom}_{\text{Sets}_*}(A \odot X, Y) \cong \text{Hom}_{\text{Sets}_*}(X, A \pitchfork Y),$$

natural in  $A \in \text{Obj}(\text{Sets})$  and  $X, Y \in \text{Obj}(\text{Sets}_*)$ .

**00CS** 4. *As a Weighted Colimit.* We have

$$A \odot X \cong \text{colim}^{[A]}(X),$$

where in the right hand side we write:

- $A$  for the functor  $A: \text{pt} \rightarrow \text{Sets}$  picking  $A \in \text{Obj}(\text{Sets})$ ;
- $X$  for the functor  $X: \text{pt} \rightarrow \text{Sets}_*$  picking  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ .

- 00CT** 5. *Iterated Tensors.* We have an isomorphism of pointed sets

$$A \odot (B \odot X) \cong (A \times B) \odot X,$$

natural in  $A, B \in \text{Obj}(\text{Sets})$  and  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ .

- 00CU** 6. *Interaction With Homs.* We have a natural isomorphism

$$\text{Sets}_*(A \odot X, -) \cong A \bowtie \text{Sets}_*(X, -).$$

- 00CV** 7. *The Tensor Evaluation Map.* For each  $X, Y \in \text{Obj}(\text{Sets}_*)$ , we have a map

$$\text{ev}_{X,Y}^\odot : \text{Sets}_*(X, Y) \odot X \rightarrow Y,$$

natural in  $X, Y \in \text{Obj}(\text{Sets}_*)$ , and given by

$$\text{ev}_{X,Y}^\odot(f \odot x) \stackrel{\text{def}}{=} f(x)$$

for each  $f \odot x \in \text{Sets}_*(X, Y) \odot X$ .

- 00CW** 8. *The Tensor Coevaluation Map.* For each  $A \in \text{Obj}(\text{Sets})$  and each  $X \in \text{Obj}(\text{Sets}_*)$ , we have a map

$$\text{coev}_{A,X}^\odot : A \rightarrow \text{Sets}_*(X, A \odot X),$$

natural in  $A \in \text{Obj}(\text{Sets})$  and  $X \in \text{Obj}(\text{Sets}_*)$ , and given by

$$\text{coev}_{A,X}^\odot(a) \stackrel{\text{def}}{=} \llbracket x \mapsto a \odot x \rrbracket$$

for each  $a \in A$ .

*Proof. Item 1, Functoriality:* This is the special case of ??, ?? of ?? for when  $C = \text{Sets}_*$ .

*Item 2, Adjointness I:* This is simply a rephrasing of **Definition 2.1.1.1**.

*Item 3, : Adjointness II:* This is the special case of ??, ?? of ?? for when  $C = \text{Sets}_*$ .

*Item 4, As a Weighted Colimit:* This is the special case of ??, ?? of ?? for when  $C = \text{Sets}_*$ .

*Item 5, Iterated Tensors:* This is the special case of ??, ?? of ?? for when  $C = \text{Sets}_*$ .

*Item 6, Interaction With Homs:* This is the special case of ??, ?? of ?? for when  $C = \text{Sets}_*$ .

*Item 7, The Tensor Evaluation Map:* This is the special case of ??, ?? of ?? for when  $C = \text{Sets}_*$ .

*Item 8, The Tensor Coevaluation Map:* This is the special case of ??, ?? of ?? for when  $C = \text{Sets}_*$ .  $\square$

**00CX 2.2 Cotensors of Pointed Sets by Sets**

Let  $(X, x_0)$  be a pointed set and let  $A$  be a set.

**00CY Definition 2.2.1.1.** The **cotensor of  $(X, x_0)$  by  $A$** <sup>9</sup> is the pointed set<sup>10</sup>  $A \pitchfork (X, x_0)$  satisfying the following universal property:

(UP) We have a bijection

$$\mathbf{Sets}_*(K, A \pitchfork X) \cong \mathbf{Sets}(A, \mathbf{Sets}_*(K, X)),$$

natural in  $(K, k_0) \in \mathbf{Obj}(\mathbf{Sets}_*)$ .

**00CZ Remark 2.2.1.2.** The universal property of **Definition 2.2.1.1** is equivalent to the following one:

(UP) We have a bijection

$$\mathbf{Sets}_*(K, A \pitchfork X) \cong \mathbf{Sets}_{\mathbb{E}_0}^{\otimes}(A \times K, X),$$

natural in  $(K, k_0) \in \mathbf{Obj}(\mathbf{Sets}_*)$ , where  $\mathbf{Sets}_{\mathbb{E}_0}^{\otimes}(A \times K, X)$  is the set defined by

$$\mathbf{Sets}_{\mathbb{E}_0}^{\otimes}(A \times K, X) \stackrel{\text{def}}{=} \left\{ f \in \mathbf{Sets}(A \times K, X) \left| \begin{array}{l} \text{for each } a \in A, \text{ we} \\ \text{have } f(a, k_0) = x_0 \end{array} \right. \right\}.$$

*Proof.* This follows from the bijection

$$\mathbf{Sets}(A, \mathbf{Sets}_*(K, X)) \cong \mathbf{Sets}_{\mathbb{E}_0}^{\otimes}(A \times K, X),$$

natural in  $(K, k_0) \in \mathbf{Obj}(\mathbf{Sets}_*)$  constructed in the proof of **Remark 2.1.1.2**.  $\square$

**00D0 Construction 2.2.1.3.** Concretely, the **cotensor of  $(X, x_0)$  by  $A$**  is the pointed set  $A \pitchfork (X, x_0)$  consisting of:

- *The Underlying Set.* The set  $A \pitchfork X$  given by

$$A \pitchfork X \cong \bigwedge_{a \in A} (X, x_0),$$

where  $\bigwedge_{a \in A} (X, x_0)$  is the smash product of the  $A$ -indexed family  $((X, x_0))_{a \in A}$  of **Definition 6.1.1.1**.

<sup>9</sup> *Further Terminology:* Also called the **power of  $(X, x_0)$  by  $A$** .

<sup>10</sup> *Further Notation:* Often written  $A \pitchfork X$  for simplicity.

- *The Basepoint.* The point  $[(x_0)_{a \in A}] = [(x_0, x_0, x_0, \dots)]$  of  $\bigwedge_{a \in A}(X, x_0)$ .

*Proof.* We claim we have a bijection

$$\mathbf{Sets}_*(K, A \pitchfork X) \cong \mathbf{Sets}(A, \mathbf{Sets}_*(K, X)),$$

natural in  $(K, k_0) \in \mathbf{Obj}(\mathbf{Sets}_*)$ .

- *Map I.* We define a map

$$\Phi_K: \mathbf{Sets}_*(K, A \pitchfork X) \rightarrow \mathbf{Sets}(A, \mathbf{Sets}_*(K, X)),$$

by sending a morphism of pointed sets

$$\xi: (K, k_0) \rightarrow (A \pitchfork X, [(x_0)_{a \in A}])$$

to the map of sets

$$\begin{aligned} \xi^\dagger: A &\rightarrow \mathbf{Sets}_*(K, X), \\ a &\mapsto (\xi_a: K \rightarrow X), \end{aligned}$$

where

$$\xi_a: (K, k_0) \rightarrow (X, x_0)$$

is the morphism of pointed sets defined by

$$\xi_a(k) = \begin{cases} x_a^k & \text{if } \xi(k) \neq [(x_0)_{a \in A}], \\ x_0 & \text{if } \xi(k) = [(x_0)_{a \in A}] \end{cases}$$

for each  $k \in K$ , where  $x_a^k$  is the  $a$ th component of  $\xi(k) = \left[ \left( x_a^k \right)_{a \in A} \right]$ .  
Note that:

1. The definition of  $\xi_a(k)$  is independent of the choice of equivalence class. Indeed, suppose we have

$$\begin{aligned} \xi(k) &= \left[ \left( x_a^k \right)_{a \in A} \right] \\ &= \left[ \left( y_a^k \right)_{a \in A} \right] \end{aligned}$$

with  $x_a^k \neq y_a^k$  for some  $a \in A$ . Then there exist  $a_x, a_y \in A$  such



that  $x_{a_x}^k = y_{a_y}^k = x_0$ . The equivalence relation  $\sim$  on  $\prod_{a \in A} X$  then forces

$$\begin{aligned} \left[ \left( x_a^k \right)_{a \in A} \right] &= [(x_0)_{a \in A}], \\ \left[ \left( y_a^k \right)_{a \in A} \right] &= [(x_0)_{a \in A}], \end{aligned}$$

however, and  $\xi_a(k)$  is defined to be  $x_0$  in this case.

2. The map  $\xi_a$  is indeed a morphism of pointed sets, as we have

$$\xi_a(k_0) = x_0$$

since  $\xi(k_0) = [(x_0)_{a \in A}]$  as  $\xi$  is a morphism of pointed sets and  $\xi_a(k_0)$ , defined to be the  $a$ th component of  $[(x_0)_{a \in A}]$ , is equal to  $x_0$ .

- *Map II.* We define a map

$$\Psi_K : \mathbf{Sets}(A, \mathbf{Sets}_*(K, X)) \rightarrow \mathbf{Sets}_*(K, A \pitchfork X),$$

given by sending a map

$$\begin{aligned} \xi : A &\rightarrow \mathbf{Sets}_*(K, X), \\ a &\mapsto (\xi_a : K \rightarrow X), \end{aligned}$$

to the morphism of pointed sets

$$\xi^\dagger : (K, k_0) \rightarrow (A \pitchfork X, [(x_0)_{a \in A}])$$

defined by

$$\xi^\dagger(k) \stackrel{\text{def}}{=} [(\xi_a(k))_{a \in A}]$$

for each  $k \in K$ . Note that  $\xi^\dagger$  is indeed a morphism of pointed sets, as we have

$$\begin{aligned} \xi^\dagger(k_0) &\stackrel{\text{def}}{=} [(\xi_a(k_0))_{a \in A}] \\ &= x_0, \end{aligned}$$

where we have used that  $\xi_a \in \mathbf{Sets}_*(K, X)$  is a morphism of pointed sets for each  $a \in A$ .

- *Naturality of  $\Psi$ .* We need to show that, given a morphism of pointed sets

$$\phi: (K, k_0) \rightarrow (K', k'_0),$$

the diagram

$$\begin{array}{ccc} \mathbf{Sets}(A, \mathbf{Sets}_*(K', X)) & \xrightarrow{\Psi_{K'}} & \mathbf{Sets}_*(K', A \pitchfork X) \\ (\phi^*)_* \downarrow & & \downarrow \phi^* \\ \mathbf{Sets}(A, \mathbf{Sets}_*(K, X)) & \xrightarrow{\Psi_K} & \mathbf{Sets}_*(K, A \pitchfork X) \end{array}$$

commutes. Indeed, given a map of sets

$$\begin{aligned} \xi: A &\rightarrow \mathbf{Sets}_*(K', X), \\ a &\mapsto (\xi_a: K' \rightarrow X), \end{aligned}$$

we have

$$\begin{aligned} [\Psi_K \circ (\phi^*)_*](\xi) &= \Psi_K((\phi^*)_*(\xi)) \\ &= \Psi_K((\phi^*)_*([a \mapsto \xi_a])) \\ &= \Psi_K([a \mapsto \phi^*(\xi_a)]) \\ &= \Psi_K([a \mapsto [k \mapsto \xi_a(\phi(k))]]) \\ &= [k \mapsto [(\xi_a(\phi(k)))_{a \in A}]] \\ &= \phi^*([k' \mapsto [(\xi_a(k'))_{a \in A}]] \\ &= \phi^*(\Psi_{K'}(\xi)) \\ &= [\phi^* \circ \Psi_{K'}](\xi). \end{aligned}$$

- *Naturality of  $\Phi$ .* Since  $\Psi$  is natural and  $\Psi$  is a componentwise inverse to  $\Phi$ , it follows from [Categories, Item 2](#) of [Proposition 8.6.1.2](#) that  $\Phi$  is also natural.
- *Invertibility I.* We claim that

$$\Psi_K \circ \Phi_K = \text{id}_{\mathbf{Sets}_*(K, A \pitchfork X)}.$$

Indeed, given a morphism of pointed sets

$$\xi: (K, k_0) \rightarrow (A \pitchfork X, [(x_0)_{a \in A}])$$

we have

$$\begin{aligned}
 [\Psi_K \circ \Phi_K](\xi) &= \Psi_K(\Phi_K(\xi)) \\
 &= \Psi_K(\llbracket a \mapsto \xi_a \rrbracket) \\
 &= \Psi_K(\llbracket a' \mapsto \xi_{a'} \rrbracket) \\
 &= \llbracket k \mapsto \left[ (\text{ev}_a(\llbracket a' \mapsto \xi_{a'}(k) \rrbracket))_{a \in A} \right] \rrbracket \\
 &= \llbracket k \mapsto [(\xi_a(k))_{a \in A}] \rrbracket.
 \end{aligned}$$

Now, we have two cases:

1. If  $\xi(k) = [(x_0)_{a \in A}]$ , we have

$$\begin{aligned}
 [\Psi_K \circ \Phi_K](\xi) &= \dots \\
 &= \llbracket k \mapsto [(\xi_a(k))_{a \in A}] \rrbracket \\
 &= \llbracket k \mapsto [(x_0)_{a \in A}] \rrbracket \\
 &= \llbracket k \mapsto \xi(k) \rrbracket \\
 &= \xi.
 \end{aligned}$$

2. If  $\xi(k) \neq [(x_0)_{a \in A}]$  and  $\xi(k) = \left[ \left( x_a^k \right)_{a \in A} \right]$  instead, we have

$$\begin{aligned}
 [\Psi_K \circ \Phi_K](\xi) &= \dots \\
 &= \llbracket k \mapsto [(\xi_a(k))_{a \in A}] \rrbracket \\
 &= \llbracket k \mapsto \left[ \left( x_a^k \right)_{a \in A} \right] \rrbracket \\
 &= \llbracket k \mapsto \xi(k) \rrbracket \\
 &= \xi.
 \end{aligned}$$

In both cases, we have  $[\Psi_K \circ \Phi_K](\xi) = \xi$ , and thus we are done.

- *Invertibility II.* We claim that

$$\Phi_K \circ \Psi_K = \text{id}_{\mathbf{Sets}(A, \mathbf{Sets}_*(K, X))}.$$

Indeed, given a morphism  $\xi: A \rightarrow \mathbf{Sets}_*(K, X)$ , we have

$$\begin{aligned}
 [\Phi_K \circ \Psi_K](\xi) &= \Phi_K(\Psi_K(\xi)) \\
 &= \Phi_K(\llbracket k \mapsto [(\xi_a(k))_{a \in A}] \rrbracket) \\
 &= \llbracket a \mapsto \llbracket k \mapsto \xi_a(k) \rrbracket \rrbracket \\
 &= \xi
 \end{aligned}$$

This finishes the proof.  $\square$

**00D1 Proposition 2.2.1.4.** Let  $(X, x_0)$  be a pointed set and let  $A$  be a set.

**00D2** 1. *Functoriality.* The assignments  $A, (X, x_0), (A, (X, x_0))$  define functors

$$\begin{aligned} A \pitchfork - &: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*, \\ - \pitchfork X &: \mathbf{Sets}^{\text{op}} \rightarrow \mathbf{Sets}_*, \\ -_1 \pitchfork -_2 &: \mathbf{Sets}^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*. \end{aligned}$$

In particular, given:

- A map of sets  $f: A \rightarrow B$ ;
- A pointed map  $\phi: (X, x_0) \rightarrow (Y, y_0)$ ;

the induced map

$$f \odot \phi: A \pitchfork X \rightarrow B \pitchfork Y$$

is given by

$$[f \odot \phi]([(x_a)_{a \in A}]) \stackrel{\text{def}}{=} \left[ \left( \phi(x_{f(a)}) \right)_{a \in A} \right]$$

for each  $[(x_a)_{a \in A}] \in A \pitchfork X$ .

**00D3** 2. *Adjointness I.* We have an adjunction

$$(- \pitchfork X \dashv \mathbf{Sets}_*(-, X)) : \mathbf{Sets}^{\text{op}} \begin{array}{c} \xrightarrow{- \pitchfork X} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(-, X)} \end{array} \mathbf{Sets}_*,$$

witnessed by a bijection

$$\mathbf{Sets}_*^{\text{op}}(A \pitchfork X, K) \cong \mathbf{Sets}(A, \mathbf{Sets}_*(K, X)),$$

i.e. by a bijection

$$\mathbf{Sets}_*(K, A \pitchfork X) \cong \mathbf{Sets}(A, \mathbf{Sets}_*(K, X)),$$

natural in  $A \in \text{Obj}(\mathbf{Sets})$  and  $X, Y \in \text{Obj}(\mathbf{Sets}_*)$ .

00D4 3. *Adjointness II.* We have an adjunctions

$$(A \odot - \dashv A \pitchfork -): \text{Sets}_* \begin{array}{c} \xrightarrow{A \odot -} \\ \perp \\ \xleftarrow{A \pitchfork -} \end{array} \text{Sets}_*,$$

witnessed by a bijection

$$\text{Hom}_{\text{Sets}_*}(A \odot X, Y) \cong \text{Hom}_{\text{Sets}_*}(X, A \pitchfork Y),$$

natural in  $A \in \text{Obj}(\text{Sets})$  and  $X, Y \in \text{Obj}(\text{Sets}_*)$ .

00D5 4. *As a Weighted Limit.* We have

$$A \pitchfork X \cong \lim^{[A]}(X),$$

where in the right hand side we write:

- $A$  for the functor  $A: \text{pt} \rightarrow \text{Sets}$  picking  $A \in \text{Obj}(\text{Sets})$ ;
- $X$  for the functor  $X: \text{pt} \rightarrow \text{Sets}_*$  picking  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ .

00D6 5. *Iterated Cotensors.* We have an isomorphism of pointed sets

$$A \pitchfork (B \pitchfork X) \cong (A \times B) \pitchfork X,$$

natural in  $A, B \in \text{Obj}(\text{Sets})$  and  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ .

00D7 6. *Commutativity With Homs.* We have natural isomorphisms

$$\begin{aligned} A \pitchfork \text{Sets}_*(X, -) &\cong \text{Sets}_*(A \odot X, -), \\ A \pitchfork \text{Sets}_*(-, Y) &\cong \text{Sets}_*(-, A \pitchfork Y). \end{aligned}$$

00D8 7. *The Cotensor Evaluation Map.* For each  $X, Y \in \text{Obj}(\text{Sets}_*)$ , we have a map

$$\text{ev}_{X,Y}^{\pitchfork}: X \rightarrow \text{Sets}_*(X, Y) \pitchfork Y,$$

natural in  $X, Y \in \text{Obj}(\text{Sets}_*)$ , and given by

$$\text{ev}_{X,Y}^{\pitchfork}(x) \stackrel{\text{def}}{=} \left[ (f(x))_{f \in \text{Sets}_*(X, Y)} \right]$$

for each  $x \in X$ .

00D9 8. *The Cotensor Coevaluation Map.* For each  $X \in \text{Obj}(\text{Sets}_*)$  and each

$A \in \text{Obj}(\text{Sets})$ , we have a map

$$\text{coev}_{A,X}^{\dashv} : A \rightarrow \text{Sets}_*(A \dashv X, X),$$

natural in  $X \in \text{Obj}(\text{Sets}_*)$  and  $A \in \text{Obj}(\text{Sets})$ , and given by

$$\text{coev}_{A,X}^{\dashv}(a) \stackrel{\text{def}}{=} \llbracket (x_b)_{b \in A} \rrbracket \mapsto x_a$$

for each  $a \in A$ .

*Proof. Item 1, Functoriality:* This is the special case of ??, ?? of ?? for when  $C = \text{Sets}_*$ .

*Item 2, Adjointness I:* This is simply a rephrasing of [Definition 2.2.1.1](#).

*Item 3, : Adjointness II:* This is the special case of ??, ?? of ?? for when  $C = \text{Sets}_*$ .

*Item 4, As a Weighted Limit:* This is the special case of ??, ?? of ?? for when  $C = \text{Sets}_*$ .

*Item 5, Iterated Cotensors:* This is the special case of ??, ?? of ?? for when  $C = \text{Sets}_*$ .

*Item 6, Commutativity With Homs:* This is the special case of ??, ?? of ?? for when  $C = \text{Sets}_*$ .

*Item 7, The Cotensor Evaluation Map:* This is the special case of ??, ?? of ?? for when  $C = \text{Sets}_*$ .

*Item 8, The Cotensor Coevaluation Map:* This is the special case of ??, ?? of ?? for when  $C = \text{Sets}_*$ .  $\square$

## 00DA 3 The Left Tensor Product of Pointed Sets

### 00DB 3.1 Foundations

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

**00DC Definition 3.1.1.1.** The **left tensor product of pointed sets** is the functor<sup>11</sup>

$$\triangleleft : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

defined as the composition

$$\text{Sets}_* \times \text{Sets}_* \xrightarrow{\text{id} \times \text{忘}} \text{Sets}_* \times \text{Sets} \xrightarrow{\beta_{\text{Sets}_*, \text{Sets}}^{\text{Cats}_2}} \text{Sets} \times \text{Sets}_* \xrightarrow{\odot} \text{Sets}_*,$$

where:

<sup>11</sup> *Further Notation:* Also written  $\triangleleft_{\text{Sets}_*}$ .

- $\omega: \mathbf{Sets}_* \rightarrow \mathbf{Sets}$  is the forgetful functor from pointed sets to sets.
- $\beta_{\mathbf{Sets}_*, \mathbf{Sets}}^{\mathbf{Cats}_2}: \mathbf{Sets}_* \times \mathbf{Sets} \xrightarrow{\cong} \mathbf{Sets} \times \mathbf{Sets}_*$  is the braiding of  $\mathbf{Cats}_2$ , i.e. the functor witnessing the isomorphism

$$\mathbf{Sets}_* \times \mathbf{Sets} \cong \mathbf{Sets} \times \mathbf{Sets}_*.$$

- $\odot: \mathbf{Sets} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$  is the tensor functor of **Item 1** of **Proposition 2.1.1.6**.

**00DD Remark 3.1.1.2.** The left tensor product of pointed sets satisfies the following natural bijection:

$$\mathbf{Sets}_*(X \triangleleft Y, Z) \cong \mathrm{Hom}_{\mathbf{Sets}_*}^{\otimes, L}(X \times Y, Z).$$

That is to say, the following data are in natural bijection:

1. Pointed maps  $f: X \triangleleft Y \rightarrow Z$ .
2. Maps of sets  $f: X \times Y \rightarrow Z$  satisfying  $f(x_0, y) = z_0$  for each  $y \in Y$ .

**00DE Remark 3.1.1.3.** The left tensor product of pointed sets may be described as follows:

- The left tensor product of  $(X, x_0)$  and  $(Y, y_0)$  is the pair  $((X \triangleleft Y, x_0 \triangleleft y_0), \iota)$  consisting of
  - A pointed set  $(X \triangleleft Y, x_0 \triangleleft y_0)$ ;
  - A left bilinear morphism of pointed sets  $\iota: (X \times Y, (x_0, y_0)) \rightarrow X \triangleleft Y$ ;

satisfying the following universal property:

(UP) Given another such pair  $((Z, z_0), f)$  consisting of

- \* A pointed set  $(Z, z_0)$ ;
- \* A left bilinear morphism of pointed sets  $f: (X \times Y, (x_0, y_0)) \rightarrow Z$ ;

there exists a unique morphism of pointed sets  $X \triangleleft Y \xrightarrow{\exists!} Z$  making the diagram

$$\begin{array}{ccc} & & X \triangleleft Y \\ & \nearrow \iota & \downarrow \exists! \\ X \times Y & \xrightarrow{f} & Z \end{array}$$

commute.

**00DF Construction 3.1.1.4.** In detail, the **left tensor product of**  $(X, x_0)$  and  $(Y, y_0)$  is the pointed set  $(X \triangleleft Y, [x_0])$  consisting of

- *The Underlying Set.* The set  $X \triangleleft Y$  defined by

$$\begin{aligned} X \triangleleft Y &\stackrel{\text{def}}{=} |Y| \odot X \\ &\cong \bigvee_{y \in Y} (X, x_0), \end{aligned}$$

where  $|Y|$  denotes the underlying set of  $(Y, y_0)$ ;

- *The Underlying Basepoint.* The point  $[(y_0, x_0)]$  of  $\bigvee_{y \in Y} (X, x_0)$ , which is equal to  $[(y, x_0)]$  for any  $y \in Y$ .

**00DG Notation 3.1.1.5.** We write<sup>12</sup>  $x \triangleleft y$  for the element  $[(y, x)]$  of

$$X \triangleleft Y \cong |Y| \odot X.$$

**00DH Remark 3.1.1.6.** Employing the notation introduced in **Notation 3.1.1.5**, we have

$$x_0 \triangleleft y_0 = x_0 \triangleleft y$$

for each  $y \in Y$ , and

$$x_0 \triangleleft y = x_0 \triangleleft y'$$

for each  $y, y' \in Y$ .

**00DJ Proposition 3.1.1.7.** Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

**00DK** 1. *Functoriality.* The assignments  $X, Y, (X, Y) \mapsto X \triangleleft Y$  define functors

$$\begin{aligned} X \triangleleft - &: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*, \\ - \triangleleft Y &: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*, \\ -_1 \triangleleft -_2 &: \mathbf{Sets}_* \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*. \end{aligned}$$

In particular, given pointed maps

$$\begin{aligned} f &: (X, x_0) \rightarrow (A, a_0), \\ g &: (Y, y_0) \rightarrow (B, b_0), \end{aligned}$$

the induced map

$$f \triangleleft g: X \triangleleft Y \rightarrow A \triangleleft B$$

---

<sup>12</sup> *Further Notation:* Also written  $x \triangleleft_{\mathbf{Sets}_*} y$ .



is given by

$$[f \triangleleft g](x \triangleleft y) \stackrel{\text{def}}{=} f(x) \triangleleft g(y)$$

for each  $x \triangleleft y \in X \triangleleft Y$ .

**00DL** 2. *Adjointness I.* We have an adjunction

$$\left( - \triangleleft Y \dashv [Y, -]_{\mathbf{Sets}_*}^{\triangleleft} \right): \quad \mathbf{Sets}_* \begin{array}{c} \xrightarrow{- \triangleleft Y} \\ \perp \\ \xleftarrow{[Y, -]_{\mathbf{Sets}_*}^{\triangleleft}} \end{array} \mathbf{Sets}_*,$$

witnessed by a bijection of sets

$$\text{Hom}_{\mathbf{Sets}_*}(X \triangleleft Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}\left(X, [Y, Z]_{\mathbf{Sets}_*}^{\triangleleft}\right)$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$ , where  $[X, Y]_{\mathbf{Sets}_*}^{\triangleleft}$  is the pointed set of **Definition 3.2.1.1**.

**00DM** 3. *Adjointness II.* The functor

$$X \triangleleft - : \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

does not admit a right adjoint.

**00DN** 4. *Adjointness III.* We have a bijection of sets

$$\text{Hom}_{\mathbf{Sets}_*}(X \triangleleft Y, Z) \cong \text{Hom}_{\mathbf{Sets}}(|Y|, \mathbf{Sets}_*(X, Z))$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$ .

*Proof. Item 1, Functoriality:* Clear.

*Item 2, Adjointness I:* This follows from **Item 3** of **Proposition 2.1.1.6**.

*Item 3, Adjointness II:* For  $X \triangleleft -$  to admit a right adjoint would require it to preserve colimits by **??**, **??** of **??**. However, we have

$$\begin{aligned} X \triangleleft \text{pt} &\stackrel{\text{def}}{=} |\text{pt}| \odot X \\ &\cong X \\ &\not\cong \text{pt}, \end{aligned}$$

and thus we see that  $X \triangleleft -$  does not have a right adjoint.

*Item 4, Adjointness III:* This follows from **Item 2** of **Proposition 2.1.1.6**.  $\square$

**00DP Remark 3.1.1.8.** Here is some intuition on why  $X \triangleleft -$  fails to be a left adjoint. **Item 4** of **Proposition 3.1.1.7** states that we have a natural bijection

$$\mathrm{Hom}_{\mathbf{Sets}_*}(X \triangleleft Y, Z) \cong \mathrm{Hom}_{\mathbf{Sets}}(|Y|, \mathbf{Sets}_*(X, Z)),$$

so it would be reasonable to wonder whether a natural bijection of the form

$$\mathrm{Hom}_{\mathbf{Sets}_*}(X \triangleleft Y, Z) \cong \mathrm{Hom}_{\mathbf{Sets}_*}(Y, \mathbf{Sets}_*(X, Z)),$$

also holds, which would give  $X \triangleleft - \dashv \mathbf{Sets}_*(X, -)$ . However, such a bijection would require every map

$$f: X \triangleleft Y \rightarrow Z$$

to satisfy

$$f(x \triangleleft y_0) = z_0$$

for each  $x \in X$ , whereas we are imposing such a basepoint preservation condition only for elements of the form  $x_0 \triangleleft y$ . Thus  $\mathbf{Sets}_*(X, -)$  can't be a right adjoint for  $X \triangleleft -$ , and as shown by **Item 3** of **Proposition 3.1.1.7**, no functor can.<sup>13</sup>

## 00DQ 3.2 The Left Internal Hom of Pointed Sets

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

**00DR Definition 3.2.1.1.** The left internal Hom of pointed sets is the functor

$$[-, -]_{\mathbf{Sets}_*}^{\triangleleft} : \mathbf{Sets}_*^{\mathrm{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

defined as the composition

$$\mathbf{Sets}_*^{\mathrm{op}} \times \mathbf{Sets}_* \xrightarrow{\omega \times \mathrm{id}} \mathbf{Sets}^{\mathrm{op}} \times \mathbf{Sets}_* \xrightarrow{\pitchfork} \mathbf{Sets}_*,$$

where:

- $\omega: \mathbf{Sets}_* \rightarrow \mathbf{Sets}$  is the forgetful functor from pointed sets to sets.
- $\pitchfork: \mathbf{Sets}^{\mathrm{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$  is the cotensor functor of **Item 1** of **Proposition 2.2.1.4**.

*Proof.* For a proof that  $[-, -]_{\mathbf{Sets}_*}^{\triangleleft}$  is indeed the left internal Hom of  $\mathbf{Sets}_*$  with respect to the left tensor product of pointed sets, see **Item 2** of **Proposition 3.1.1.7**.  $\square$

<sup>13</sup>The functor  $\mathbf{Sets}_*(X, -)$  is instead right adjoint to  $X \wedge -$ , the smash product of

**00DS Remark 3.2.1.2.** The left internal Hom of pointed sets satisfies the following universal property:

$$\mathbf{Sets}_*(X \triangleleft Y, Z) \cong \mathbf{Sets}_*\left(X, [Y, Z]_{\mathbf{Sets}_*}^{\triangleleft}\right)$$

That is to say, the following data are in bijection:

1. Pointed maps  $f: X \triangleleft Y \rightarrow Z$ .
2. Pointed maps  $f: X \rightarrow [Y, Z]_{\mathbf{Sets}_*}^{\triangleleft}$ .

**00DT Remark 3.2.1.3.** In detail, the **left internal Hom of**  $(X, x_0)$  **and**  $(Y, y_0)$  is the pointed set  $\left([X, Y]_{\mathbf{Sets}_*}^{\triangleleft}, [(y_0)_{x \in X}]\right)$  consisting of

- *The Underlying Set.* The set  $[X, Y]_{\mathbf{Sets}_*}^{\triangleleft}$  defined by

$$\begin{aligned} [X, Y]_{\mathbf{Sets}_*}^{\triangleleft} &\stackrel{\text{def}}{=} |X| \curvearrowright Y \\ &\cong \bigwedge_{x \in X} (Y, y_0), \end{aligned}$$

where  $|X|$  denotes the underlying set of  $(X, x_0)$ ;

- *The Underlying Basepoint.* The point  $[(y_0)_{x \in X}]$  of  $\bigwedge_{x \in X} (Y, y_0)$ .

**00DU Proposition 3.2.1.4.** Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

**00DV** 1. *Functoriality.* The assignments  $X, Y, (X, Y) \mapsto [X, Y]_{\mathbf{Sets}_*}^{\triangleleft}$  define functors

$$\begin{aligned} [X, -]_{\mathbf{Sets}_*}^{\triangleleft} &: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*, \\ [-, Y]_{\mathbf{Sets}_*}^{\triangleleft} &: \mathbf{Sets}_*^{\text{op}} \rightarrow \mathbf{Sets}_*, \\ [-1, -2]_{\mathbf{Sets}_*}^{\triangleleft} &: \mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*. \end{aligned}$$

In particular, given pointed maps

$$\begin{aligned} f &: (X, x_0) \rightarrow (A, a_0), \\ g &: (Y, y_0) \rightarrow (B, b_0), \end{aligned}$$

the induced map

$$[f, g]_{\mathbf{Sets}_*}^{\triangleleft} : [A, Y]_{\mathbf{Sets}_*}^{\triangleleft} \rightarrow [X, B]_{\mathbf{Sets}_*}^{\triangleleft}$$

is given by

$$[f, g]_{\mathbf{Sets}_*}^{\triangleleft}([(y_a)_{a \in A}]) \stackrel{\text{def}}{=} \left[ \left( g(y_{f(x)}) \right)_{x \in X} \right]$$

for each  $[(y_a)_{a \in A}] \in [A, Y]_{\mathbf{Sets}_*}^{\triangleleft}$ .

**00DW** 2. *Adjointness I.* We have an adjunction

$$\left( - \triangleleft Y \dashv [Y, -]_{\mathbf{Sets}_*}^{\triangleleft} \right): \mathbf{Sets}_* \begin{array}{c} \xrightarrow{- \triangleleft Y} \\ \perp \\ \xleftarrow{[Y, -]_{\mathbf{Sets}_*}^{\triangleleft}} \end{array} \mathbf{Sets}_*,$$

witnessed by a bijection of sets

$$\mathrm{Hom}_{\mathbf{Sets}_*}(X \triangleleft Y, Z) \cong \mathrm{Hom}_{\mathbf{Sets}_*}\left(X, [Y, Z]_{\mathbf{Sets}_*}^{\triangleleft}\right)$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \mathrm{Obj}(\mathbf{Sets}_*)$

**00DX** 3. *Adjointness II.* The functor

$$X \triangleleft - : \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

does not admit a right adjoint.

*Proof.* **Item 1, Functoriality:** Clear.

**Item 2, Adjointness I:** This is a repetition of **Item 2** of **Proposition 3.1.1.7**, and is proved there.

**Item 3, Adjointness II:** This is a repetition of **Item 3** of **Proposition 3.1.1.7**, and is proved there.  $\square$

### **00DY** 3.3 The Left Skew Unit

**00DZ** **Definition 3.3.1.1.** The left skew unit of the left tensor product of pointed sets is the functor

$$\mathbb{1}^{\mathbf{Sets}_*, \triangleleft} : \mathbf{pt} \rightarrow \mathbf{Sets}_*$$

defined by

$$\mathbb{1}_{\mathbf{Sets}_*}^{\triangleleft} \stackrel{\mathrm{def}}{=} S^0.$$

### **00E0** 3.4 The Left Skew Associator

**00E1** **Definition 3.4.1.1.** The skew associator of the left tensor product of pointed sets is the natural transformation

$$\alpha^{\mathbf{Sets}_*, \triangleleft} : \triangleleft \circ (\triangleleft \times \mathrm{id}_{\mathbf{Sets}_*}) \Longrightarrow \triangleleft \circ (\mathrm{id}_{\mathbf{Sets}_*} \times \triangleleft) \circ \alpha_{\mathbf{Sets}_*, \mathbf{Sets}_*, \mathbf{Sets}_*}^{\mathbf{Cats}}$$

pointed sets of **Definition 5.1.1.1**. See **Item 2** of **Proposition 5.1.1.9**.

as in the diagram

$$\begin{array}{ccc}
 & \text{Sets}_* \times (\text{Sets}_* \times \text{Sets}_*) & \\
 \alpha_{\text{Sets}_*, \text{Sets}_*, \text{Sets}_*}^{\text{Cats}} \swarrow & \nearrow \text{id} \times \triangleleft & \\
 (\text{Sets}_* \times \text{Sets}_*) \times \text{Sets}_* & & \text{Sets}_* \times \text{Sets}_* \\
 \triangleleft \times \text{id} \searrow & \alpha_{\text{Sets}_*, \triangleleft} \nearrow & \searrow \triangleleft \\
 \text{Sets}_* \times \text{Sets}_* & \xrightarrow{\triangleleft} & \text{Sets}_*
 \end{array}$$

whose component

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft} : (X \triangleleft Y) \triangleleft Z \rightarrow X \triangleleft (Y \triangleleft Z)$$

at  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$  is given by

$$\begin{aligned}
 (X \triangleleft Y) \triangleleft Z &\stackrel{\text{def}}{=} |Z| \odot (X \triangleleft Y) \\
 &\stackrel{\text{def}}{=} |Z| \odot (|Y| \odot X) \\
 &\cong \bigvee_{z \in Z} |Y| \odot X \\
 &\cong \bigvee_{z \in Z} \left( \bigvee_{y \in Y} X \right) \\
 &\rightarrow \bigvee_{[(z,y)] \in \bigvee_{z \in Z} Y} X \\
 &\cong \bigvee_{[(z,y)] \in |Z| \odot Y} X \\
 &\cong ||Z| \odot Y| \odot X \\
 &\stackrel{\text{def}}{=} |Y \triangleleft Z| \odot X \\
 &\stackrel{\text{def}}{=} X \triangleleft (Y \triangleleft Z),
 \end{aligned}$$

where the map

$$\bigvee_{z \in Z} \left( \bigvee_{y \in Y} X \right) \rightarrow \bigvee_{(z,y) \in \bigvee_{z \in Z} Y} X$$

is given by  $[(z, [(y, x)])] \mapsto [[(z, y)], x]$ .

*Proof.* (Proven below in a bit.)

□

**00E2 Remark 3.4.1.2.** Unwinding the notation for elements, we have

$$\begin{aligned} [(z, [(y, x)])] &\stackrel{\text{def}}{=} [(z, x \triangleleft y)] \\ &\stackrel{\text{def}}{=} (x \triangleleft y) \triangleleft z \end{aligned}$$

and

$$\begin{aligned} [([(z, y)], x)] &\stackrel{\text{def}}{=} [(y \triangleleft z, x)] \\ &\stackrel{\text{def}}{=} x \triangleleft (y \triangleleft z). \end{aligned}$$

So, in other words,  $\alpha_{X,Y,Z}^{\mathbf{Sets}_*, \triangleleft}$  acts on elements via

$$\alpha_{X,Y,Z}^{\mathbf{Sets}_*, \triangleleft}((x \triangleleft y) \triangleleft z) \stackrel{\text{def}}{=} x \triangleleft (y \triangleleft z)$$

for each  $(x \triangleleft y) \triangleleft z \in (X \triangleleft Y) \triangleleft Z$ .

**00E3 Remark 3.4.1.3.** Taking  $y = y_0$ , we see that the morphism  $\alpha_{X,Y,Z}^{\mathbf{Sets}_*, \triangleleft}$  acts on elements as

$$\alpha_{X,Y,Z}^{\mathbf{Sets}_*, \triangleleft}((x \triangleleft y_0) \triangleleft z) \stackrel{\text{def}}{=} x \triangleleft (y_0 \triangleleft z).$$

However, by the definition of  $\triangleleft$ , we have  $y_0 \triangleleft z = y_0 \triangleleft z'$  for all  $z, z' \in Z$ , preventing  $\alpha_{X,Y,Z}^{\mathbf{Sets}_*, \triangleleft}$  from being non-invertible.

*Proof.* Firstly, note that, given  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$ , the map

$$\alpha_{X,Y,Z}^{\mathbf{Sets}_*, \triangleleft}: (X \triangleleft Y) \triangleleft Z \rightarrow X \triangleleft (Y \triangleleft Z)$$

is indeed a morphism of pointed sets, as we have

$$\alpha_{X,Y,Z}^{\mathbf{Sets}_*, \triangleleft}((x_0 \triangleleft y_0) \triangleleft z_0) = x_0 \triangleleft (y_0 \triangleleft z_0).$$

Next, we claim that  $\alpha^{\mathbf{Sets}_*, \triangleleft}$  is a natural transformation. We need to show that, given morphisms of pointed sets

$$\begin{aligned} f: (X, x_0) &\rightarrow (X', x'_0), \\ g: (Y, y_0) &\rightarrow (Y', y'_0), \\ h: (Z, z_0) &\rightarrow (Z', z'_0) \end{aligned}$$

the diagram

$$\begin{array}{ccc} (X \triangleleft Y) \triangleleft Z & \xrightarrow{(f \triangleleft g) \triangleleft h} & (X' \triangleleft Y') \triangleleft Z' \\ \alpha_{X,Y,Z}^{\mathbf{Sets}_*, \triangleleft} \downarrow & & \downarrow \alpha_{X',Y',Z'}^{\mathbf{Sets}_*, \triangleleft} \\ X \triangleleft (Y \triangleleft Z) & \xrightarrow{f \triangleleft (g \triangleleft h)} & X' \triangleleft (Y' \triangleleft Z') \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} (x \triangleleft y) \triangleleft z & \longmapsto & (f(x) \triangleleft g(y)) \triangleleft h(z) \\ \downarrow & & \downarrow \\ x \triangleleft (y \triangleleft z) & \longmapsto & f(x) \triangleleft (g(y) \triangleleft h(z)) \end{array}$$

and hence indeed commutes, showing  $\alpha^{\mathbf{Sets}_*, \triangleleft}$  to be a natural transformation. This finishes the proof.  $\square$

### 00E4 3.5 The Left Skew Left Unitor

00E5 **Definition 3.5.1.1.** The **skew left unitor of the left tensor product of pointed sets** is the natural transformation

$$\lambda^{\mathbf{Sets}_*, \triangleleft} : \triangleleft \circ (\mathbb{1}^{\mathbf{Sets}_*} \times \text{id}_{\mathbf{Sets}_*}) \xrightarrow{\sim} \lambda_{\mathbf{Sets}_*}^{\mathbf{Cats}_2}$$

whose component

$$\lambda_X^{\mathbf{Sets}_*, \triangleleft} : S^0 \triangleleft X \rightarrow X$$

at  $(X, x_0) \in \text{Obj}(\mathbf{Sets}_*)$  is given by the composition

$$\begin{aligned} S^0 \triangleleft X &\cong |X| \odot S^0 \\ &\cong \bigvee_{x \in X} S^0 \\ &\rightarrow X, \end{aligned}$$

where  $\bigvee_{x \in X} S^0 \rightarrow X$  is the map given by

$$\begin{aligned} [(x, 0)] &\mapsto x_0, \\ [(x, 1)] &\mapsto x. \end{aligned}$$

*Proof.* (Proven below in a bit.)  $\square$

**00E6 Remark 3.5.1.2.** In other words,  $\lambda_X^{\text{Sets}_*, \triangleleft}$  acts on elements as

$$\begin{aligned}\lambda_X^{\text{Sets}_*, \triangleleft}(0 \triangleleft x) &\stackrel{\text{def}}{=} x_0, \\ \lambda_X^{\text{Sets}_*, \triangleleft}(1 \triangleleft x) &\stackrel{\text{def}}{=} x\end{aligned}$$

for each  $1 \triangleleft x \in S^0 \triangleleft X$ .

**00E7 Remark 3.5.1.3.** The morphism  $\lambda_X^{\text{Sets}_*, \triangleleft}$  is almost invertible, with its would-be-inverse

$$\phi_X: X \rightarrow S^0 \triangleleft X$$

given by

$$\phi_X(x) \stackrel{\text{def}}{=} 1 \triangleleft x$$

for each  $x \in X$ . Indeed, we have

$$\begin{aligned}\left[\lambda_X^{\text{Sets}_*, \triangleleft} \circ \phi\right](x) &= \lambda_X^{\text{Sets}_*, \triangleleft}(\phi(x)) \\ &= \lambda_X^{\text{Sets}_*, \triangleleft}(1 \triangleleft x) \\ &= x \\ &= [\text{id}_X](x)\end{aligned}$$

so that

$$\lambda_X^{\text{Sets}_*, \triangleleft} \circ \phi = \text{id}_X$$

and

$$\begin{aligned}\left[\phi \circ \lambda_X^{\text{Sets}_*, \triangleleft}\right](1 \triangleleft x) &= \phi\left(\lambda_X^{\text{Sets}_*, \triangleleft}(1 \triangleleft x)\right) \\ &= \phi(x) \\ &= 1 \triangleleft x \\ &= [\text{id}_{S^0 \triangleleft X}](1 \triangleleft x),\end{aligned}$$

but

$$\begin{aligned}\left[\phi \circ \lambda_X^{\text{Sets}_*, \triangleleft}\right](0 \triangleleft x) &= \phi\left(\lambda_X^{\text{Sets}_*, \triangleleft}(0 \triangleleft x)\right) \\ &= \phi(x_0) \\ &= 1 \triangleleft x_0,\end{aligned}$$

where  $0 \triangleleft x \neq 1 \triangleleft x_0$ . Thus

$$\phi \circ \lambda_X^{\text{Sets}_*, \triangleleft} \stackrel{?}{=} \text{id}_{S^0 \triangleleft X}$$

holds for all elements in  $S^0 \triangleleft X$  except one.



*Proof.* Firstly, note that, given  $(X, x_0) \in \text{Obj}(\mathbf{Sets}_*)$ , the map

$$\lambda_X^{\mathbf{Sets}_*, \triangleleft} : S^0 \triangleleft X \rightarrow X$$

is indeed a morphism of pointed sets, as we have

$$\lambda_X^{\mathbf{Sets}_*, \triangleleft}(0 \triangleleft x_0) = x_0.$$

Next, we claim that  $\lambda^{\mathbf{Sets}_*, \triangleleft}$  is a natural transformation. We need to show that, given a morphism of pointed sets

$$f : (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} S^0 \triangleleft X & \xrightarrow{\text{id}_{S^0} \triangleleft f} & S^0 \triangleleft Y \\ \lambda_X^{\mathbf{Sets}_*, \triangleleft} \downarrow & & \downarrow \lambda_Y^{\mathbf{Sets}_*, \triangleleft} \\ X & \xrightarrow{f} & Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} 0 \triangleleft x & & 0 \triangleleft x \mapsto 0 \triangleleft f(x) \\ \downarrow & & \downarrow \\ x_0 & \mapsto & f(x_0) \end{array} \quad \begin{array}{ccc} & & y_0 \end{array}$$

and

$$\begin{array}{ccc} 1 \triangleleft x & \mapsto & 1 \triangleleft f(x) \\ \downarrow & & \downarrow \\ x & \mapsto & f(x) \end{array}$$

and hence indeed commutes, showing  $\lambda^{\mathbf{Sets}_*, \triangleleft}$  to be a natural transformation. This finishes the proof.  $\square$

### 00E8 3.6 The Left Skew Right Unitor

**00E9 Definition 3.6.1.1.** The **skew right unitor of the left tensor product of pointed sets** is the natural transformation

$$\rho^{\text{Sets}_*, \triangleleft} : \rho_{\text{Sets}_*}^{\text{Cats}_2} \xrightarrow{\sim} \triangleleft \circ (\text{id} \times \mathbb{1}^{\text{Sets}_*}),$$

whose component

$$\rho_X^{\text{Sets}_*, \triangleleft} : X \rightarrow X \triangleleft S^0$$

at  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$  is given by the composition

$$\begin{aligned} X &\rightarrow X \vee X \\ &\cong |S^0| \odot X \\ &\cong X \triangleleft S^0, \end{aligned}$$

where  $X \rightarrow X \vee X$  is the map sending  $X$  to the second factor of  $X$  in  $X \vee X$ .

*Proof.* (Proven below in a bit.) □

**00EA Remark 3.6.1.2.** In other words,  $\rho_X^{\text{Sets}_*, \triangleleft}$  acts on elements as

$$\rho_X^{\text{Sets}_*, \triangleleft}(x) \stackrel{\text{def}}{=} [(1, x)]$$

i.e. by

$$\rho_X^{\text{Sets}_*, \triangleleft}(x) \stackrel{\text{def}}{=} x \triangleleft 1$$

for each  $x \in X$ .

**00EB Remark 3.6.1.3.** The morphism  $\rho_X^{\text{Sets}_*, \triangleleft}$  is non-invertible, as it is non-surjective when viewed as a map of sets, since the elements  $x \triangleleft 0$  of  $X \triangleleft S^0$  with  $x \neq x_0$  are outside the image of  $\rho_X^{\text{Sets}_*, \triangleleft}$ , which sends  $x$  to  $x \triangleleft 1$ .

*Proof.* Firstly, note that, given  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ , the map

$$\rho_X^{\text{Sets}_*, \triangleleft} : X \rightarrow X \triangleleft S^0$$

is indeed a morphism of pointed sets as we have

$$\begin{aligned}\rho_X^{\mathbf{Sets}_*, \triangleleft}(x_0) &= x_0 \triangleleft 1 \\ &= x_0 \triangleleft 0.\end{aligned}$$

Next, we claim that  $\rho^{\mathbf{Sets}_*, \triangleleft}$  is a natural transformation. We need to show that, given a morphism of pointed sets

$$f: (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \rho_X^{\mathbf{Sets}_*, \triangleleft} \downarrow & & \downarrow \rho_Y^{\mathbf{Sets}_*, \triangleleft} \\ X \triangleleft S^0 & \xrightarrow{f \triangleleft \text{id}_{S^0}} & Y \triangleleft S^0 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x & \longmapsto & f(x) \\ \downarrow & & \downarrow \\ x \triangleleft 0 & \longmapsto & f(x) \triangleleft 0 \end{array}$$

and hence indeed commutes, showing  $\rho^{\mathbf{Sets}_*, \triangleleft}$  to be a natural transformation. This finishes the proof.  $\square$

### 00EC 3.7 The Diagonal

00ED **Definition 3.7.1.1.** The **diagonal of the left tensor product of pointed sets** is the natural transformation

$$\Delta^{\triangleleft}: \text{id}_{\mathbf{Sets}_*} \Rightarrow \triangleleft \circ \Delta_{\mathbf{Sets}_*}^{\mathbf{Cats}_2},$$

whose component

$$\Delta_X^{\triangleleft}: (X, x_0) \rightarrow (X \triangleleft X, x_0 \triangleleft x_0)$$

at  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$  is given by

$$\Delta_X^\triangleleft(x) \stackrel{\text{def}}{=} x \triangleleft x$$

for each  $x \in X$ .

*Proof. Being a Morphism of Pointed Sets:* We have

$$\Delta_X^\triangleleft(x_0) \stackrel{\text{def}}{=} x_0 \triangleleft x_0,$$

and thus  $\Delta_X^\triangleleft$  is a morphism of pointed sets.

*Naturality:* We need to show that, given a morphism of pointed sets

$$f: (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \Delta_X^\triangleleft \downarrow & & \downarrow \Delta_Y^\triangleleft \\ X \triangleleft X & \xrightarrow{f \triangleleft f} & Y \triangleleft Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x & \xrightarrow{\quad} & f(x) \\ \downarrow & & \downarrow \\ x \triangleleft x & \xrightarrow{\quad} & f(x) \triangleleft f(x) \end{array}$$

and hence indeed commutes, showing  $\Delta^\triangleleft$  to be natural.  $\square$

### 3.8 The Left Skew Monoidal Structure on Pointed Sets Associated to $\triangleleft$

00EE

**00EF Proposition 3.8.1.1.** The category  $\text{Sets}_*$  admits a left-closed left skew monoidal category structure consisting of

- *The Underlying Category.* The category  $\text{Sets}_*$  of pointed sets;
- *The Left Skew Monoidal Product.* The left tensor product functor

$$\triangleleft: \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

of [Definition 3.1.1.1](#);

- *The Left Internal Skew Hom.* The left internal Hom functor

$$[-, -]_{\mathbf{Sets}_*}^{\triangleleft} : \mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

of [Definition 3.2.1.1](#);

- *The Left Skew Monoidal Unit.* The functor

$$\mathbb{1}^{\mathbf{Sets}_*, \triangleleft} : \mathbf{pt} \rightarrow \mathbf{Sets}_*$$

of [Definition 3.3.1.1](#);

- *The Left Skew Associators.* The natural transformation

$$\alpha^{\mathbf{Sets}_*, \triangleleft} : \triangleleft \circ (\triangleleft \times \text{id}_{\mathbf{Sets}_*}) \Longrightarrow \triangleleft \circ (\text{id}_{\mathbf{Sets}_*} \times \triangleleft) \circ \alpha_{\mathbf{Sets}_*, \mathbf{Sets}_*, \mathbf{Sets}_*}^{\mathbf{Cats}}$$

of [Definition 3.4.1.1](#);

- *The Left Skew Left Unitors.* The natural transformation

$$\lambda^{\mathbf{Sets}_*, \triangleleft} : \triangleleft \circ (\mathbb{1}^{\mathbf{Sets}_*} \times \text{id}_{\mathbf{Sets}_*}) \xrightarrow{\sim} \lambda_{\mathbf{Sets}_*}^{\mathbf{Cats}_2}$$

of [Definition 3.5.1.1](#);

- *The Left Skew Right Unitors.* The natural transformation

$$\rho^{\mathbf{Sets}_*, \triangleleft} : \rho_{\mathbf{Sets}_*}^{\mathbf{Cats}_2} \xrightarrow{\sim} \triangleleft \circ (\text{id} \times \mathbb{1}^{\mathbf{Sets}_*})$$

of [Definition 3.6.1.1](#).

*Proof. The Pentagon Identity:* Let  $(W, w_0)$ ,  $(X, x_0)$ ,  $(Y, y_0)$  and  $(Z, z_0)$  be

pointed sets. We have to show that the diagram

$$\begin{array}{ccc}
 & (W \triangleleft (X \triangleleft Y)) \triangleleft Z & \\
 \alpha_{W,X,Y}^{\text{Sets}_*, \triangleleft} \triangleleft \text{id}_Z \nearrow & & \searrow \alpha_{W,X \triangleleft Y,Z}^{\text{Sets}_*, \triangleleft} \\
 ((W \triangleleft X) \triangleleft Y) \triangleleft Z & & W \triangleleft ((X \triangleleft Y) \triangleleft Z) \\
 \alpha_{W \triangleleft X,Y,Z}^{\text{Sets}_*, \triangleleft} \searrow & & \nearrow \text{id}_W \triangleleft \alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft} \\
 (W \triangleleft X) \triangleleft (Y \triangleleft Z) & \xrightarrow{\alpha_{W,X,Y \triangleleft Z}^{\text{Sets}_*, \triangleleft}} & W \triangleleft (X \triangleleft (Y \triangleleft Z))
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 & (w \triangleleft (x \triangleleft y)) \triangleleft z & \\
 \nwarrow & & \nearrow \\
 ((w \triangleleft x) \triangleleft y) \triangleleft z & & w \triangleleft ((x \triangleleft y) \triangleleft z) \\
 \nwarrow & & \nearrow \\
 (w \triangleleft x) \triangleleft (y \triangleleft z) & \longmapsto & w \triangleleft (x \triangleleft (y \triangleleft z))
 \end{array}$$

and thus we see that the pentagon identity is satisfied.

*The Left Skew Left Triangle Identity:* Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

We have to show that the diagram

$$\begin{array}{ccc}
 (S^0 \triangleleft X) \triangleleft Y & \xrightarrow{\alpha_{S^0, X, Y}^{\text{Sets}_*, \triangleleft}} & S^0 \triangleleft (X \triangleleft Y) \\
 \searrow \lambda_X^{\text{Sets}_*, \triangleleft} \triangleleft \text{id}_Y & & \downarrow \lambda_{X \triangleleft Y}^{\text{Sets}_*, \triangleleft} \\
 & & X \triangleleft Y
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 (0 \triangleleft x) \triangleleft y & \mapsto & 0 \triangleleft (x \triangleleft y) \\
 \searrow & & \downarrow \\
 & & x_0 \triangleleft y = x_0 \triangleleft y_0
 \end{array}$$

and

$$\begin{array}{ccc}
 (1 \triangleleft x) \triangleleft y & \mapsto & 1 \triangleleft (x \triangleleft y) \\
 \searrow & & \downarrow \\
 & & x \triangleleft y
 \end{array}$$

and hence indeed commutes. Thus the left skew triangle identity is satisfied.

*The Left Skew Right Triangle Identity:* Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccc}
 X \triangleleft Y & & \\
 \downarrow \rho_{X \triangleleft Y}^{\text{Sets}_*, \triangleleft} & \searrow \text{id}_X \triangleleft \rho_Y^{\text{Sets}_*, \triangleleft} & \\
 (X \triangleleft Y) \triangleleft S^0 & \xrightarrow{\alpha_{X, Y, S^0}^{\text{Sets}_*, \triangleleft}} & X \triangleleft (Y \triangleleft S^0)
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 x \triangleleft y & & \\
 \downarrow & \searrow & \\
 (x \triangleleft y) \triangleleft 1 & \mapsto & x \triangleleft (y \triangleleft 1)
 \end{array}$$

and hence indeed commutes. Thus the right skew triangle identity is satisfied.

### 3.8 The Left Skew Monoidal Structure on Pointed Sets Associated to $\triangleleft$ 40

*The Left Skew Middle Triangle Identity:* Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccc}
 X \triangleleft Y & \xlongequal{\quad} & X \triangleleft Y \\
 \downarrow \rho_X^{\text{Sets}_*, \triangleleft} \triangleleft \text{id}_Y & & \uparrow \text{id}_A \triangleleft \lambda_Y^{\text{Sets}_*, \triangleleft} \\
 (X \triangleleft S^0) \triangleleft Y & \xrightarrow[\alpha_{X, S^0, Y}^{\text{Sets}_*, \triangleleft}]{} & X \triangleleft (S^0 \triangleleft Y)
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 x \triangleleft y & \xrightarrow{\quad} & x \triangleleft y \\
 \downarrow & & \uparrow \\
 (x \triangleleft 1) \triangleleft y & \xrightarrow{\quad} & x \triangleleft (1 \triangleleft y)
 \end{array}$$

and hence indeed commutes. Thus the right skew triangle identity is satisfied.

*The Zig-Zag Identity:* We have to show that the diagram

$$\begin{array}{ccc}
 S^0 & \xrightarrow[\rho_{S^0}^{\text{Sets}_*, \triangleleft}]{} & S^0 \triangleleft S^0 \\
 & \searrow & \downarrow \lambda_{S^0}^{\text{Sets}_*, \triangleleft} \\
 & & S^0
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 0 & \xrightarrow{\quad} & 0 \triangleleft 1 \\
 & \searrow & \downarrow \\
 & & 0
 \end{array}$$

and

$$\begin{array}{ccc}
 1 & \xrightarrow{\quad} & 1 \triangleleft 1 \\
 & \searrow & \downarrow \\
 & & 1
 \end{array}$$

and hence indeed commutes. Thus the zig-zag identity is satisfied.

*Left Skew Monoidal Left-Closedness:* This follows from [Item 2](#) of [Proposition 3.1.1.7](#).  $\square$



### 3.9 Monoids With Respect to the Left Tensor Product of Pointed Sets

00EG

**00EH Proposition 3.9.1.1.** The category of monoids on  $(\mathbf{Sets}_*, \triangleleft, S^0)$  is isomorphic to the category of “monoids with left zero”<sup>14</sup> and morphisms between them.

*Proof.* Monoids on  $(\mathbf{Sets}_*, \triangleleft, S^0)$ : A monoid on  $(\mathbf{Sets}_*, \triangleleft, S^0)$  consists of:

- *The Underlying Object.* A pointed set  $(A, 0_A)$ .
- *The Multiplication Morphism.* A morphism of pointed sets

$$\mu_A: A \triangleleft A \rightarrow A,$$

determining a left bilinear morphism of pointed sets

$$\begin{aligned} A \times A &\longrightarrow A \\ (a, b) &\longmapsto ab. \end{aligned}$$

- *The Unit Morphism.* A morphism of pointed sets

$$\eta_A: S^0 \rightarrow A$$

picking an element  $1_A$  of  $A$ .

satisfying the following conditions:

1. *Associativity.* The diagram

$$\begin{array}{ccc} & A \triangleleft (A \triangleleft A) & \\ \alpha_{A,A,A}^{\mathbf{Sets}_*, \triangleleft} \nearrow & & \searrow \text{id}_A \triangleleft \mu_A \\ (A \triangleleft A) \triangleleft A & & A \triangleleft A \\ \mu_A \triangleleft \text{id}_A \searrow & & \swarrow \mu_A \\ A \triangleleft A & \xrightarrow{\mu_A} & A \end{array}$$

<sup>14</sup>A monoid with left zero is defined similarly as the monoids with zero of ???. Succinctly, they are monoids  $(A, \mu_A, \eta_A)$  with a special element  $0_A$  satisfying

$$0_A a = 0_A$$

for each  $a \in A$ .

2. *Left Unitality.* The diagram

$$\begin{array}{ccc}
 S^0 \triangleleft A & \xrightarrow{\eta_A \times \text{id}_A} & A \triangleleft A \\
 & \searrow \lambda_A^{\text{Sets}^*, \triangleleft} & \downarrow \mu_A \\
 & & A
 \end{array}$$

commutes.

3. *Right Unitality.* The diagram

$$\begin{array}{ccc}
 A & \xrightarrow{\rho_A^{\text{Sets}^*, \triangleleft}} & A \triangleleft S^0 \\
 \parallel & & \downarrow \text{id}_A \times \eta_A \\
 A & \xleftarrow{\mu_A} & A \triangleleft A
 \end{array}$$

commutes.

Being a left-bilinear morphism of pointed sets, the multiplication map satisfies

$$0_A a = 0_A$$

for each  $a \in A$ . Now, the associativity, left unitality, and right unitality conditions act on elements as follows:

1. *Associativity.* The associativity condition acts as

$$\begin{array}{ccc}
 & & a \triangleleft (b \triangleleft c) \\
 & \swarrow & \searrow \\
 (a \triangleleft b) \triangleleft c & & (a \triangleleft b) \triangleleft c \quad a \triangleleft bc \\
 \searrow & & \swarrow \\
 ab \triangleleft c & \xrightarrow{\quad} & (ab)c \\
 & & \downarrow \\
 & & a(bc)
 \end{array}$$

This gives

$$(ab)c = a(bc)$$

for each  $a, b, c \in A$ .

2. *Left Unitality.* The left unitality condition acts:

(a) On  $0 \triangleleft a$  as

$$\begin{array}{ccc} 0 \triangleleft a & & 0 \triangleleft a \mapsto 0_A \triangleleft a \\ & \searrow & \searrow \downarrow \\ & 0_A & 0_A a. \end{array}$$

(b) On  $1 \triangleleft a$  as

$$\begin{array}{ccc} 1 \triangleleft a & & 1 \triangleleft a \mapsto 1_A \triangleleft a \\ & \searrow & \searrow \downarrow \\ & a & 1_A a. \end{array}$$

This gives

$$\begin{aligned} 1_A a &= a, \\ 0_A a &= 0_A \end{aligned}$$

for each  $a \in A$ .

3. *Right Unitality.* The right unitality condition acts as

$$\begin{array}{ccc} a & \xrightarrow{\quad} & a \triangleleft 1 \\ \downarrow & & \downarrow \\ a & \xleftarrow{\quad} & a \triangleleft 1_A \end{array}$$

This gives

$$a 1_A = a$$

for each  $a \in A$ .

Thus we see that monoids with respect to  $\triangleleft$  are exactly monoids with left zero.

*Morphisms of Monoids on  $(\mathbf{Sets}_*, \triangleleft, S^0)$ :* A morphism of monoids on  $(\mathbf{Sets}_*, \triangleleft, S^0)$  from  $(A, \mu_A, \eta_A, 0_A)$  to  $(B, \mu_B, \eta_B, 0_B)$  is a morphism of pointed sets

$$f: (A, 0_A) \rightarrow (B, 0_B)$$

satisfying the following conditions:

1. *Compatibility With the Multiplication Morphisms.* The diagram

$$\begin{array}{ccc}
 A \triangleleft A & \xrightarrow{f \triangleleft f} & B \triangleleft B \\
 \mu_A \downarrow & & \downarrow \mu_B \\
 A & \xrightarrow{f} & B
 \end{array}$$

commutes.

2. *Compatibility With the Unit Morphisms.* The diagram

$$\begin{array}{ccc}
 S^0 & \xrightarrow{\eta_A} & A \\
 & \searrow \eta_B & \downarrow f \\
 & & B
 \end{array}$$

commutes.

These act on elements as

$$\begin{array}{ccc}
 a \triangleleft b & & a \triangleleft b \mapsto f(a) \triangleleft f(b) \\
 \downarrow & & \downarrow \\
 ab \mapsto f(ab) & & f(a)f(b)
 \end{array}$$

and

$$\begin{array}{ccc}
 0 & \xrightarrow{\quad} & 0_A \\
 & \searrow & \downarrow \\
 & & f(0_A)
 \end{array}$$

and

$$\begin{array}{ccc}
 1 & \xrightarrow{\quad} & 1_A \\
 & \searrow & \downarrow \\
 & & f(1_A)
 \end{array}$$

giving

$$\begin{aligned} f(ab) &= f(a)f(b), \\ f(0_A) &= 0_B, \\ f(1_A) &= 1_B, \end{aligned}$$

for each  $a, b \in A$ , which is exactly a morphism of monoids with left zero.

*Identities and Composition:* Similarly, the identities and composition of  $\text{Mon}(\text{Sets}_*, \triangleleft, S^0)$  can be easily seen to agree with those of monoids with left zero, which finishes the proof.  $\square$

## 00EJ 4 The Right Tensor Product of Pointed Sets

### 00EK 4.1 Foundations

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

00EL **Definition 4.1.1.1.** The **right tensor product of pointed sets** is the functor<sup>15</sup>

$$\triangleright : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

defined as the composition

$$\text{Sets}_* \times \text{Sets}_* \xrightarrow{\text{忘} \times \text{id}} \text{Sets} \times \text{Sets}_* \xrightarrow{\odot} \text{Sets}_*,$$

where:

- $\text{忘} : \text{Sets}_* \rightarrow \text{Sets}$  is the forgetful functor from pointed sets to sets.
- $\odot : \text{Sets} \times \text{Sets}_* \rightarrow \text{Sets}_*$  is the tensor functor of **Item 1** of **Proposition 2.1.1.6**.

00EM **Remark 4.1.1.2.** The right tensor product of pointed sets satisfies the following natural bijection:

$$\text{Sets}_*(X \triangleright Y, Z) \cong \text{Hom}_{\text{Sets}_*}^{\otimes, \text{R}}(X \times Y, Z).$$

That is to say, the following data are in natural bijection:

1. Pointed maps  $f : X \triangleright Y \rightarrow Z$ .
2. Maps of sets  $f : X \times Y \rightarrow Z$  satisfying  $f(x, y_0) = z_0$  for each  $x \in X$ .

<sup>15</sup> *Further Notation:* Also written  $\triangleright_{\text{Sets}_*}$ .

**00EN Remark 4.1.1.3.** The right tensor product of pointed sets may be described as follows:

- The right tensor product of  $(X, x_0)$  and  $(Y, y_0)$  is the pair  $((X \triangleright Y, x_0 \triangleright y_0), \iota)$  consisting of
  - A pointed set  $(X \triangleright Y, x_0 \triangleright y_0)$ ;
  - A right bilinear morphism of pointed sets  $\iota: (X \times Y, (x_0, y_0)) \rightarrow X \triangleright Y$ ;

satisfying the following universal property:

(UP) Given another such pair  $((Z, z_0), f)$  consisting of

- \* A pointed set  $(Z, z_0)$ ;
- \* A right bilinear morphism of pointed sets  $f: (X \times Y, (x_0, y_0)) \rightarrow Z$ ;

there exists a unique morphism of pointed sets  $X \triangleright Y \xrightarrow{\exists!} Z$  making the diagram

$$\begin{array}{ccc} & & X \triangleright Y \\ & \nearrow \iota & \downarrow \exists! \\ X \times Y & \xrightarrow{f} & Z \end{array}$$

commute.

**00EP Construction 4.1.1.4.** In detail, the **right tensor product of**  $(X, x_0)$  **and**  $(Y, y_0)$  is the pointed set  $(X \triangleright Y, [y_0])$  consisting of:

- *The Underlying Set.* The set  $X \triangleright Y$  defined by

$$\begin{aligned} X \triangleright Y &\stackrel{\text{def}}{=} |X| \odot Y \\ &\cong \bigvee_{x \in X} (Y, y_0), \end{aligned}$$

where  $|X|$  denotes the underlying set of  $(X, x_0)$ .

- *The Underlying Basepoint.* The point  $[(x_0, y_0)]$  of  $\bigvee_{x \in X} (Y, y_0)$ , which is equal to  $[(x, y_0)]$  for any  $x \in X$ .

**00EQ** **Notation 4.1.1.5.** We write<sup>16</sup>  $x \triangleright y$  for the element  $[(x, y)]$  of

$$X \triangleright Y \cong |X| \odot Y.$$

**00ER** **Remark 4.1.1.6.** Employing the notation introduced in **Notation 4.1.1.5**, we have

$$x_0 \triangleright y_0 = x \triangleright y_0$$

for each  $x \in X$ , and

$$x \triangleright y_0 = x' \triangleright y_0$$

for each  $x, x' \in X$ .

**00ES** **Proposition 4.1.1.7.** Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

**00ET** 1. *Functoriality.* The assignments  $X, Y, (X, Y) \mapsto X \triangleright Y$  define functors

$$\begin{aligned} X \triangleright - &: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*, \\ - \triangleright Y &: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*, \\ -_1 \triangleright -_2 &: \mathbf{Sets}_* \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*. \end{aligned}$$

In particular, given pointed maps

$$\begin{aligned} f &: (X, x_0) \rightarrow (A, a_0), \\ g &: (Y, y_0) \rightarrow (B, b_0), \end{aligned}$$

the induced map

$$f \triangleright g: X \triangleright Y \rightarrow A \triangleright B$$

is given by

$$[f \triangleright g](x \triangleright y) \stackrel{\text{def}}{=} f(x) \triangleright g(y)$$

for each  $x \triangleright y \in X \triangleright Y$ .

**00EU** 2. *Adjointness I.* We have an adjunction

$$\left( X \triangleright - \dashv [X, -]_{\mathbf{Sets}_*}^{\triangleright} \right): \mathbf{Sets}_* \begin{array}{c} \xrightarrow{X \triangleright -} \\ \perp \\ \xleftarrow{[X, -]_{\mathbf{Sets}_*}^{\triangleright}} \end{array} \mathbf{Sets}_*,$$

witnessed by a bijection of sets

$$\text{Hom}_{\mathbf{Sets}_*}(X \triangleright Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}\left(Y, [X, Z]_{\mathbf{Sets}_*}^{\triangleright}\right)$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$ , where  $[X, Y]_{\mathbf{Sets}_*}^{\triangleright}$  is the pointed set of **Definition 4.2.1.1**.

---

<sup>16</sup> *Further Notation:* Also written  $x \triangleright_{\mathbf{Sets}_*} y$ .

**00EV** 3. *Adjointness II.* The functor

$$- \triangleright Y : \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

does not admit a right adjoint.

**00EW** 4. *Adjointness III.* We have a bijection of sets

$$\mathrm{Hom}_{\mathbf{Sets}_*}(X \triangleright Y, Z) \cong \mathrm{Hom}_{\mathbf{Sets}}(|X|, \mathbf{Sets}_*(Y, Z))$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \mathrm{Obj}(\mathbf{Sets}_*)$ .

*Proof. Item 1, Functoriality:* Clear.

*Item 2, Adjointness I:* This follows from **Item 3** of **Proposition 2.1.1.6**.

*Item 3, Adjointness II:* For  $- \triangleright Y$  to admit a right adjoint would require it to preserve colimits by ??, ?? of ??. However, we have

$$\begin{aligned} \mathrm{pt} \triangleright X &\stackrel{\mathrm{def}}{=} |\mathrm{pt}| \odot X \\ &\cong X \\ &\not\cong \mathrm{pt}, \end{aligned}$$

and thus we see that  $- \triangleright Y$  does not have a right adjoint.

*Item 4, Adjointness III:* This follows from **Item 2** of **Proposition 2.1.1.6**.  $\square$

**00EX Remark 4.1.1.8.** Here is some intuition on why  $- \triangleright Y$  fails to be a left adjoint. **Item 4** of **Proposition 3.1.1.7** states that we have a natural bijection

$$\mathrm{Hom}_{\mathbf{Sets}_*}(X \triangleright Y, Z) \cong \mathrm{Hom}_{\mathbf{Sets}}(|X|, \mathbf{Sets}_*(Y, Z)),$$

so it would be reasonable to wonder whether a natural bijection of the form

$$\mathrm{Hom}_{\mathbf{Sets}_*}(X \triangleright Y, Z) \cong \mathrm{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(Y, Z)),$$

also holds, which would give  $- \triangleright Y \dashv \mathbf{Sets}_*(Y, -)$ . However, such a bijection would require every map

$$f : X \triangleright Y \rightarrow Z$$

to satisfy

$$f(x_0 \triangleright y) = z_0$$

for each  $x \in X$ , whereas we are imposing such a basepoint preservation condition only for elements of the form  $x \triangleright y_0$ . Thus  $\mathbf{Sets}_*(Y, -)$  can't be a right adjoint for  $- \triangleright Y$ , and as shown by **Item 3** of **Proposition 4.1.1.7**, no functor can.<sup>17</sup>

<sup>17</sup>The functor  $\mathbf{Sets}_*(Y, -)$  is instead right adjoint to  $- \wedge Y$ , the smash product of pointed



**00EY 4.2 The Right Internal Hom of Pointed Sets**

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

**00EZ Definition 4.2.1.1.** The **right internal Hom of pointed sets** is the functor

$$[-, -]_{\mathbf{Sets}_*}^{\triangleright} : \mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

defined as the composition

$$\mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \xrightarrow{\omega \times \text{id}} \mathbf{Sets}^{\text{op}} \times \mathbf{Sets}_* \xrightarrow{\pitchfork} \mathbf{Sets}_*,$$

where:

- $\omega : \mathbf{Sets}_* \rightarrow \mathbf{Sets}$  is the forgetful functor from pointed sets to sets.
- $\pitchfork : \mathbf{Sets}^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$  is the cotensor functor of [Item 1 of Proposition 2.2.1.4](#).

*Proof.* For a proof that  $[-, -]_{\mathbf{Sets}_*}^{\triangleright}$  is indeed the right internal Hom of  $\mathbf{Sets}_*$  with respect to the right tensor product of pointed sets, see [Item 2 of Proposition 4.1.1.7](#).  $\square$

**00F0 Remark 4.2.1.2.** We have

$$[-, -]_{\mathbf{Sets}_*}^{\triangleleft} = [-, -]_{\mathbf{Sets}_*}^{\triangleright}.$$

**00F1 Remark 4.2.1.3.** The right internal Hom of pointed sets satisfies the following universal property:

$$\mathbf{Sets}_*(X \triangleright Y, Z) \cong \mathbf{Sets}_*\left(Y, [X, Z]_{\mathbf{Sets}_*}^{\triangleright}\right)$$

That is to say, the following data are in bijection:

1. Pointed maps  $f : X \triangleright Y \rightarrow Z$ .
2. Pointed maps  $f : Y \rightarrow [X, Z]_{\mathbf{Sets}_*}^{\triangleright}$ .

**00F2 Remark 4.2.1.4.** In detail, the **right internal Hom of**  $(X, x_0)$  and  $(Y, y_0)$  is the pointed set  $\left([X, Y]_{\mathbf{Sets}_*}^{\triangleright}, [(y_0)_{x \in X}]\right)$  consisting of

sets of [Definition 5.1.1.1](#). See [Item 2 of Proposition 5.1.1.9](#).

- *The Underlying Set.* The set  $[X, Y]_{\mathbf{Sets}_*}^\triangleright$  defined by

$$\begin{aligned} [X, Y]_{\mathbf{Sets}_*}^\triangleright &\stackrel{\text{def}}{=} |X| \curvearrowright Y \\ &\cong \bigwedge_{x \in X} (Y, y_0), \end{aligned}$$

where  $|X|$  denotes the underlying set of  $(X, x_0)$ ;

- *The Underlying Basepoint.* The point  $[(y_0)_{x \in X}]$  of  $\bigwedge_{x \in X} (Y, y_0)$ .

**00F3 Proposition 4.2.1.5.** Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

- 00F4** 1. *Functoriality.* The assignments  $X, Y, (X, Y) \mapsto [X, Y]_{\mathbf{Sets}_*}^\triangleright$  define functors

$$\begin{aligned} [X, -]_{\mathbf{Sets}_*}^\triangleright &: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*, \\ [-, Y]_{\mathbf{Sets}_*}^\triangleright &: \mathbf{Sets}_*^{\text{op}} \rightarrow \mathbf{Sets}_*, \\ [-1, -2]_{\mathbf{Sets}_*}^\triangleright &: \mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*. \end{aligned}$$

In particular, given pointed maps

$$\begin{aligned} f &: (X, x_0) \rightarrow (A, a_0), \\ g &: (Y, y_0) \rightarrow (B, b_0), \end{aligned}$$

the induced map

$$[f, g]_{\mathbf{Sets}_*}^\triangleright : [A, Y]_{\mathbf{Sets}_*}^\triangleright \rightarrow [X, B]_{\mathbf{Sets}_*}^\triangleright$$

is given by

$$[f, g]_{\mathbf{Sets}_*}^\triangleright ((y_a)_{a \in A}) \stackrel{\text{def}}{=} \left[ \left( g(y_{f(x)}) \right)_{x \in X} \right]$$

for each  $[(y_a)_{a \in A}] \in [A, Y]_{\mathbf{Sets}_*}^\triangleright$ .

- 00F5** 2. *Adjointness I.* We have an adjunction

$$\left( X \triangleright - \dashv [X, -]_{\mathbf{Sets}_*}^\triangleright \right) : \mathbf{Sets}_* \begin{array}{c} \xrightarrow{X \triangleright -} \\ \perp \\ \xleftarrow{[X, -]_{\mathbf{Sets}_*}^\triangleright} \end{array} \mathbf{Sets}_*,$$

witnessed by a bijection of sets

$$\text{Hom}_{\mathbf{Sets}_*}(X \triangleright Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}(Y, [X, Z]_{\mathbf{Sets}_*}^\triangleright)$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$ , where  $[X, Y]_{\mathbf{Sets}_*}^\triangleright$  is the pointed set of **Definition 4.2.1.1**.

00F6 3. *Adjointness II*. The functor

$$- \triangleright Y : \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

does not admit a right adjoint.

*Proof. Item 1, Functoriality:* Clear.

*Item 2, Adjointness I:* This is a repetition of *Item 2* of *Proposition 4.1.1.7*, and is proved there.

*Item 3, Adjointness II:* This is a repetition of *Item 3* of *Proposition 4.1.1.7*, and is proved there.  $\square$

### 00F7 4.3 The Right Skew Unit

00F8 **Definition 4.3.1.1.** The **right skew unit of the right tensor product of pointed sets** is the functor

$$\mathbb{1}^{\mathbf{Sets}_*, \triangleright} : \mathbf{pt} \rightarrow \mathbf{Sets}_*$$

defined by

$$\mathbb{1}_{\mathbf{Sets}_*}^{\triangleright} \stackrel{\text{def}}{=} S^0.$$

### 00F9 4.4 The Right Skew Associator

00FA **Definition 4.4.1.1.** The **skew associator of the right tensor product of pointed sets** is the natural transformation

$$\alpha^{\mathbf{Sets}_*, \triangleright} : \triangleright \circ (\text{id}_{\mathbf{Sets}_*} \times \triangleright) \Longrightarrow \triangleright \circ (\triangleright \times \text{id}_{\mathbf{Sets}_*}) \circ \alpha_{\mathbf{Sets}_*, \mathbf{Sets}_*, \mathbf{Sets}_*}^{\mathbf{Cats}, -1}$$

as in the diagram

$$\begin{array}{ccc}
 & (\mathbf{Sets}_* \times \mathbf{Sets}_*) \times \mathbf{Sets}_* & \\
 \alpha_{\mathbf{Sets}_*, \mathbf{Sets}_*, \mathbf{Sets}_*}^{\mathbf{Cats}, -1} \swarrow & & \searrow \triangleright \times \text{id} \\
 \mathbf{Sets}_* \times (\mathbf{Sets}_* \times \mathbf{Sets}_*) & & \mathbf{Sets}_* \times \mathbf{Sets}_* \\
 \text{id} \times \triangleright \searrow & \alpha_{\mathbf{Sets}_*, \triangleright} \nearrow & \searrow \triangleright \\
 \mathbf{Sets}_* \times \mathbf{Sets}_* & \xrightarrow{\triangleright} & \mathbf{Sets}_*
 \end{array}$$

whose component

$$\alpha_{X, Y, Z}^{\mathbf{Sets}_*, \triangleright} : X \triangleright (Y \triangleright Z) \rightarrow (X \triangleright Y) \triangleright Z$$

at  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$  is given by

$$\begin{aligned}
 X \triangleright (Y \triangleright Z) &\stackrel{\text{def}}{=} |X| \odot (Y \triangleright Z) \\
 &\stackrel{\text{def}}{=} |X| \odot (|Y| \odot Z) \\
 &\cong \bigvee_{x \in X} (|Y| \odot Z) \\
 &\cong \bigvee_{x \in X} \left( \bigvee_{y \in Y} Z \right) \\
 &\rightarrow \bigvee_{[(x,y)] \in \bigvee_{x \in X} Y} Z \\
 &\cong \bigvee_{[(x,y)] \in |X| \odot Y} Z \\
 &\cong ||X| \odot Y| \odot Z \\
 &\stackrel{\text{def}}{=} |X \triangleright Y| \odot Z \\
 &\stackrel{\text{def}}{=} (X \triangleright Y) \triangleright Z,
 \end{aligned}$$

where the map

$$\bigvee_{x \in X} \left( \bigvee_{y \in Y} Z \right) \rightarrow \bigvee_{[(x,y)] \in \bigvee_{x \in X} Y} Z$$

is given by  $[(x, [(y, z)])] \mapsto [([(x, y)], z)]$ .

*Proof.* (Proven below in a bit.) □

**00FB Remark 4.4.1.2.** Unwinding the notation for elements, we have

$$\begin{aligned}
 [(x, [(y, z)])] &\stackrel{\text{def}}{=} [(x, y \triangleright z)] \\
 &\stackrel{\text{def}}{=} x \triangleright (y \triangleright z)
 \end{aligned}$$

and

$$\begin{aligned}
 [([(x, y)], z)] &\stackrel{\text{def}}{=} [(x \triangleright y, z)] \\
 &\stackrel{\text{def}}{=} (x \triangleright y) \triangleright z.
 \end{aligned}$$

So, in other words,  $\alpha_{X,Y,Z}^{\mathbf{Sets}_*, \triangleright}$  acts on elements via

$$\alpha_{X,Y,Z}^{\mathbf{Sets}_*, \triangleright} (x \triangleright (y \triangleright z)) \stackrel{\text{def}}{=} (x \triangleright y) \triangleright z$$

for each  $x \triangleright (y \triangleright z) \in X \triangleright (Y \triangleright Z)$ .

**00FC Remark 4.4.1.3.** Taking  $y = y_0$ , we see that the morphism  $\alpha_{X,Y,Z}^{\mathbf{Sets}_*, \triangleright}$  acts on elements as

$$\alpha_{X,Y,Z}^{\mathbf{Sets}_*, \triangleright} (x \triangleright (y_0 \triangleright z)) \stackrel{\text{def}}{=} (x \triangleright y_0) \triangleright z.$$

However, by the definition of  $\triangleright$ , we have  $x \triangleright y_0 = x' \triangleright y_0$  for all  $x, x' \in X$ , preventing  $\alpha_{X,Y,Z}^{\mathbf{Sets}_*, \triangleright}$  from being non-invertible.

*Proof.* Firstly, note that, given  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$ , the map

$$\alpha_{X,Y,Z}^{\mathbf{Sets}_*, \triangleright} : X \triangleright (Y \triangleright Z) \rightarrow (X \triangleright Y) \triangleright Z$$

is indeed a morphism of pointed sets, as we have

$$\alpha_{X,Y,Z}^{\mathbf{Sets}_*, \triangleright} (x_0 \triangleright (y_0 \triangleright z_0)) = (x_0 \triangleright y_0) \triangleright z_0.$$

Next, we claim that  $\alpha^{\mathbf{Sets}_*, \triangleright}$  is a natural transformation. We need to show that, given morphisms of pointed sets

$$\begin{aligned} f &: (X, x_0) \rightarrow (X', x'_0), \\ g &: (Y, y_0) \rightarrow (Y', y'_0), \\ h &: (Z, z_0) \rightarrow (Z', z'_0) \end{aligned}$$

the diagram

$$\begin{array}{ccc} X \triangleright (Y \triangleright Z) & \xrightarrow{f \triangleright (g \triangleright h)} & X' \triangleright (Y' \triangleright Z') \\ \alpha_{X,Y,Z}^{\mathbf{Sets}_*, \triangleright} \downarrow & & \downarrow \alpha_{X',Y',Z'}^{\mathbf{Sets}_*, \triangleright} \\ (X \triangleright Y) \triangleright Z & \xrightarrow{(f \triangleright g) \triangleright h} & (X' \triangleright Y') \triangleright Z' \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x \triangleright (y \triangleright z) & \longmapsto & f(x) \triangleright (g(y) \triangleright h(z)) \\ \downarrow & & \downarrow \\ (x \triangleright y) \triangleright z & \longmapsto & (f(x) \triangleright g(y)) \triangleright h(z) \end{array}$$

and hence indeed commutes, showing  $\alpha^{\mathbf{Sets}_*, \triangleright}$  to be a natural transformation. This finishes the proof.  $\square$

**00FD 4.5 The Right Skew Left Unitor**

**00FE Definition 4.5.1.1.** The **skew left unitor of the right tensor product of pointed sets** is the natural transformation

$$\lambda^{\mathbf{Sets}_*, \triangleright} : \lambda_{\mathbf{Sets}_*}^{\mathbf{Cats}_2} \xRightarrow{\sim} \triangleright \circ (\mathbb{1}_{\mathbf{Sets}_*} \times \text{id}_{\mathbf{Sets}_*})$$

whose component

$$\lambda_X^{\mathbf{Sets}_*, \triangleright} : X \rightarrow S^0 \triangleright X$$

at  $(X, x_0) \in \text{Obj}(\mathbf{Sets}_*)$  is given by the composition

$$\begin{aligned} X &\rightarrow X \vee X \\ &\cong |S^0| \odot X \\ &\cong S^0 \triangleright X, \end{aligned}$$

where  $X \rightarrow X \vee X$  is the map sending  $X$  to the second factor of  $X$  in  $X \vee X$ .

*Proof.* (Proven below in a bit.) □

**00FF Remark 4.5.1.2.** In other words,  $\lambda_X^{\mathbf{Sets}_*, \triangleright}$  acts on elements as

$$\lambda_X^{\mathbf{Sets}_*, \triangleright}(x) \stackrel{\text{def}}{=} [(1, x)]$$

i.e. by

$$\lambda_X^{\mathbf{Sets}_*, \triangleright}(x) \stackrel{\text{def}}{=} 1 \triangleright x$$

for each  $x \in X$ .

**00FG Remark 4.5.1.3.** The morphism  $\lambda_X^{\mathbf{Sets}_*, \triangleright}$  is non-invertible, as it is non-surjective when viewed as a map of sets, since the elements  $0 \triangleright x$  of  $S^0 \triangleright X$  with  $x \neq x_0$  are outside the image of  $\lambda_X^{\mathbf{Sets}_*, \triangleright}$ , which sends  $x$  to  $1 \triangleright x$ .

*Proof.* Firstly, note that, given  $(X, x_0) \in \text{Obj}(\mathbf{Sets}_*)$ , the map

$$\lambda_X^{\mathbf{Sets}_*, \triangleright} : X \rightarrow S^0 \triangleright X$$

is indeed a morphism of pointed sets, as we have

$$\begin{aligned}\lambda_X^{\mathbf{Sets}_*, \triangleright}(x_0) &= 1 \triangleright x_0 \\ &= 0 \triangleright x_0.\end{aligned}$$

Next, we claim that  $\lambda^{\mathbf{Sets}_*, \triangleright}$  is a natural transformation. We need to show that, given a morphism of pointed sets

$$f: (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \lambda_X^{\mathbf{Sets}_*, \triangleright} \downarrow & & \downarrow \lambda_Y^{\mathbf{Sets}_*, \triangleright} \\ S^0 \triangleright X & \xrightarrow{\text{id}_{S^0} \triangleright f} & S^0 \triangleright Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x & \xrightarrow{\quad} & f(x) \\ \downarrow & & \downarrow \\ 1 \triangleright x & \xrightarrow{\quad} & 1 \triangleright f(x) \end{array}$$

and hence indeed commutes, showing  $\lambda^{\mathbf{Sets}_*, \triangleright}$  to be a natural transformation. This finishes the proof.  $\square$

## 00FH 4.6 The Right Skew Right Unitor

00FJ **Definition 4.6.1.1.** The **skew right unitor of the right tensor product of pointed sets** is the natural transformation

$$\rho^{\mathbf{Sets}_*, \triangleright} : \triangleright \circ (\text{id} \times \mathbb{1}^{\mathbf{Sets}_*}) \xRightarrow{\sim} \rho_{\mathbf{Sets}_*}^{\mathbf{Cats}_2},$$

The diagram illustrates the naturality of the skew right unitor. It shows a commutative square with a curved arrow. The top-left node is  $\mathbf{Sets}_* \times \text{pt}$ , the top-right node is  $\mathbf{Sets}_* \times \mathbf{Sets}_*$ , and the bottom-right node is  $\mathbf{Sets}_*$ . A vertical arrow labeled  $\triangleright$  (representing  $\lambda$ ) points from  $\mathbf{Sets}_* \times \mathbf{Sets}_*$  to  $\mathbf{Sets}_*$ . A curved arrow labeled  $\rho_{\mathbf{Sets}_*}^{\mathbf{Cats}_2}$  points from  $\mathbf{Sets}_* \times \text{pt}$  to  $\mathbf{Sets}_*$ . A double arrow labeled  $\rho^{\mathbf{Sets}_*, \triangleright}$  points from  $\mathbf{Sets}_* \times \text{pt}$  to  $\mathbf{Sets}_* \times \mathbf{Sets}_*$ . The horizontal arrow from  $\mathbf{Sets}_* \times \text{pt}$  to  $\mathbf{Sets}_* \times \mathbf{Sets}_*$  is labeled  $\text{id} \times \mathbb{1}^{\mathbf{Sets}_*}$ .

whose component

$$\rho_X^{\mathbf{Sets}_*, \triangleright} : X \triangleright S^0 \rightarrow X$$

at  $(X, x_0) \in \text{Obj}(\mathbf{Sets}_*)$  is given by the composition

$$\begin{aligned} X \triangleright S^0 &\cong |X| \odot S^0 \\ &\cong \bigvee_{x \in X} S^0 \\ &\rightarrow X, \end{aligned}$$

where  $\bigvee_{x \in X} S^0 \rightarrow X$  is the map given by

$$\begin{aligned} [(x, 0)] &\mapsto x_0, \\ [(x, 1)] &\mapsto x. \end{aligned}$$

*Proof.* (Proven below in a bit.) □

**00FK Remark 4.6.1.2.** In other words,  $\rho_X^{\mathbf{Sets}_*, \triangleright}$  acts on elements as

$$\begin{aligned} \rho_X^{\mathbf{Sets}_*, \triangleright}(x \triangleright 0) &\stackrel{\text{def}}{=} x_0, \\ \rho_X^{\mathbf{Sets}_*, \triangleright}(x \triangleright 1) &\stackrel{\text{def}}{=} x \end{aligned}$$

for each  $x \triangleright 1 \in X \triangleright S^0$ .

**00FL Remark 4.6.1.3.** The morphism  $\rho_X^{\mathbf{Sets}_*, \triangleright}$  is almost invertible, with its would-be-inverse

$$\phi_X : X \rightarrow X \triangleright S^0$$

given by

$$\phi_X(x) \stackrel{\text{def}}{=} x \triangleright 1$$

for each  $x \in X$ . Indeed, we have

$$\begin{aligned} [\rho_X^{\mathbf{Sets}_*, \triangleright} \circ \phi](x) &= \rho_X^{\mathbf{Sets}_*, \triangleright}(\phi(x)) \\ &= \rho_X^{\mathbf{Sets}_*, \triangleright}(x \triangleright 1) \\ &= x \\ &= [\text{id}_X](x) \end{aligned}$$

so that

$$\rho_X^{\mathbf{Sets}_*, \triangleright} \circ \phi = \text{id}_X$$



and

$$\begin{aligned} [\phi \circ \rho_X^{\mathbf{Sets}_*, \triangleright}](x \triangleright 1) &= \phi(\rho_X^{\mathbf{Sets}_*, \triangleright}(x \triangleright 1)) \\ &= \phi(x) \\ &= x \triangleright 1 \\ &= [\mathrm{id}_{X \triangleright S^0}](x \triangleright 1), \end{aligned}$$

but

$$\begin{aligned} [\phi \circ \rho_X^{\mathbf{Sets}_*, \triangleright}](x \triangleright 0) &= \phi(\rho_X^{\mathbf{Sets}_*, \triangleright}(x \triangleright 0)) \\ &= \phi(x_0) \\ &= 1 \triangleright x_0, \end{aligned}$$

where  $x \triangleright 0 \neq 1 \triangleright x_0$ . Thus

$$\phi \circ \rho_X^{\mathbf{Sets}_*, \triangleright} \stackrel{?}{=} \mathrm{id}_{X \triangleright S^0}$$

holds for all elements in  $X \triangleright S^0$  except one.

*Proof.* Firstly, note that, given  $(X, x_0) \in \mathrm{Obj}(\mathbf{Sets}_*)$ , the map

$$\rho_X^{\mathbf{Sets}_*, \triangleright} : X \triangleright S^0 \rightarrow X$$

is indeed a morphism of pointed sets as we have

$$\rho_X^{\mathbf{Sets}_*, \triangleright}(x_0 \triangleright 0) = x_0.$$

Next, we claim that  $\rho^{\mathbf{Sets}_*, \triangleright}$  is a natural transformation. We need to show that, given a morphism of pointed sets

$$f : (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X \triangleright S^0 & \xrightarrow{f \triangleright \mathrm{id}_{S^0}} & Y \triangleright S^0 \\ \rho_X^{\mathbf{Sets}_*, \triangleright} \downarrow & & \downarrow \rho_Y^{\mathbf{Sets}_*, \triangleright} \\ X & \xrightarrow{f} & Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x \triangleright 0 & & x \triangleright 0 \mapsto f(x) \triangleright 0 \\ \downarrow & & \downarrow \\ x_0 & \mapsto & f(x_0) \end{array} \qquad \begin{array}{ccc} x \triangleright 0 & \mapsto & f(x) \triangleright 0 \\ \downarrow & & \downarrow \\ y_0 & & \end{array}$$

and

$$\begin{array}{ccc} x \triangleright 1 & \longmapsto & f(x) \triangleright 1 \\ \downarrow & & \downarrow \\ x & \longmapsto & f(x) \end{array}$$

and hence indeed commutes, showing  $\rho^{\mathbf{Sets}_*, \triangleright}$  to be a natural transformation. This finishes the proof.  $\square$

## 00FM 4.7 The Diagonal

**00FN Definition 4.7.1.1.** The **diagonal of the right tensor product of pointed sets** is the natural transformation

$$\Delta^{\triangleright} : \text{id}_{\mathbf{Sets}_*} \Rightarrow \triangleright \circ \Delta_{\mathbf{Sets}_*}^{\mathbf{Cats}_2},$$

whose component

$$\Delta_X^{\triangleright} : (X, x_0) \rightarrow (X \triangleright X, x_0 \triangleright x_0)$$

at  $(X, x_0) \in \text{Obj}(\mathbf{Sets}_*)$  is given by

$$\Delta_X^{\triangleright}(x) \stackrel{\text{def}}{=} x \triangleright x$$

for each  $x \in X$ .

*Proof. Being a Morphism of Pointed Sets:* We have

$$\Delta_X^{\triangleright}(x_0) \stackrel{\text{def}}{=} x_0 \triangleright x_0,$$

and thus  $\Delta_X^{\triangleright}$  is a morphism of pointed sets.

*Naturality:* We need to show that, given a morphism of pointed sets

$$f : (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \Delta_X^{\triangleright} \downarrow & & \downarrow \Delta_Y^{\triangleright} \\ X \triangleright X & \xrightarrow{f \triangleright f} & Y \triangleright Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x & \xrightarrow{\quad} & f(x) \\ \downarrow & & \downarrow \\ x \triangleright x & \xrightarrow{\quad} & f(x) \triangleright f(x) \end{array}$$

and hence indeed commutes, showing  $\Delta^\triangleright$  to be natural.  $\square$

#### 4.8 The Right Skew Monoidal Structure on Pointed Sets Associated to $\triangleright$

00FP

**00FQ Proposition 4.8.1.1.** The category  $\mathbf{Sets}_*$  admits a right-closed right skew monoidal category structure consisting of

- *The Underlying Category.* The category  $\mathbf{Sets}_*$  of pointed sets;
- *The Right Skew Monoidal Product.* The right tensor product functor

$$\triangleright : \mathbf{Sets}_* \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

of Definition 4.1.1.1;

- *The Right Internal Skew Hom.* The right internal Hom functor

$$[-, -]_{\mathbf{Sets}_*}^\triangleright : \mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

of Definition 4.2.1.1;

- *The Right Skew Monoidal Unit.* The functor

$$\mathbb{1}^{\mathbf{Sets}_*, \triangleright} : \mathbf{pt} \rightarrow \mathbf{Sets}_*$$

of Definition 4.3.1.1;

- *The Right Skew Associators.* The natural transformation

$$\alpha^{\mathbf{Sets}_*, \triangleright} : \triangleright \circ (\text{id}_{\mathbf{Sets}_*} \times \triangleright) \Longrightarrow \triangleright \circ (\triangleright \times \text{id}_{\mathbf{Sets}_*}) \circ \alpha_{\mathbf{Sets}_*, \mathbf{Sets}_*, \mathbf{Sets}_*}^{\mathbf{Cats}, -1}$$

of Definition 4.4.1.1;

- *The Right Skew Left Unitors.* The natural transformation

$$\lambda^{\mathbf{Sets}_*, \triangleright} : \lambda_{\mathbf{Sets}_*}^{\mathbf{Cats}_2} \xrightarrow{\sim} \triangleright \circ (\mathbb{1}^{\mathbf{Sets}_*} \times \text{id}_{\mathbf{Sets}_*})$$

of Definition 4.5.1.1;

- *The Right Skew Right Unitors.* The natural transformation

$$\rho^{\text{Sets}_*, \triangleright} : \triangleright \circ (\text{id} \times \mathbb{1}^{\text{Sets}_*}) \xRightarrow{\sim} \rho_{\text{Sets}_*}^{\text{Cats}_2}$$

of [Definition 4.6.1.1](#).

*Proof. The Pentagon Identity:* Let  $(W, w_0)$ ,  $(X, x_0)$ ,  $(Y, y_0)$  and  $(Z, z_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccccc}
 & & W \triangleright ((X \triangleright Y) \triangleright Z) & & \\
 & \nearrow \alpha_{W, X, Y \triangleright Z}^{\text{Sets}_*, \triangleright} & & \nwarrow \alpha_{W, X \triangleright Y, Z}^{\text{Sets}_*, \triangleright} & \\
 W \triangleright (X \triangleright (Y \triangleright Z)) & & & & (W \triangleright (X \triangleright Y)) \triangleright Z \\
 \searrow \alpha_{W \triangleright X, Y, Z}^{\text{Sets}_*, \triangleright} & & & & \swarrow \text{id}_W \triangleright \alpha_{X, Y, Z}^{\text{Sets}_*, \triangleright} \\
 (W \triangleright X) \triangleright (Y \triangleright Z) & \xrightarrow{\alpha_{W, X, Y \triangleright Z}^{\text{Sets}_*, \triangleright}} & & & ((W \triangleright X) \triangleright Y) \triangleright Z
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 & w \triangleright ((x \triangleright y) \triangleright z) & \\
 \swarrow & & \searrow \\
 w \triangleright (x \triangleright (y \triangleright z)) & & (w \triangleright (x \triangleright y)) \triangleright z \\
 \searrow & & \swarrow \\
 (w \triangleright x) \triangleright (y \triangleright z) & \longmapsto & ((w \triangleright x) \triangleright y) \triangleright z
 \end{array}$$

and thus we see that the pentagon identity is satisfied.

*The Right Skew Left Triangle Identity:* Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccc}
 X \triangleright Y & & \\
 \lambda_{X \triangleright Y}^{\text{Sets}_*, \triangleright} \downarrow & \searrow \lambda_X^{\text{Sets}_*, \triangleright} \triangleright \text{id}_Y & \\
 S^0 \triangleright (X \triangleright Y) & \xrightarrow{\alpha_{S^0, X, Y}^{\text{Sets}_*, \triangleright}} & (S^0 \triangleright X) \triangleright Y
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 x \triangleright y & & \\
 \downarrow & \searrow & \\
 1 \triangleright (x \triangleright y) & \longmapsto & (1 \triangleright x) \triangleright y
 \end{array}$$

and hence indeed commutes. Thus the left skew triangle identity is satisfied.

*The Right Skew Right Triangle Identity:* Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed

sets. We have to show that the diagram

$$\begin{array}{ccc}
 X \triangleright (Y \triangleright S^0) & \xrightarrow{\text{id}_X \triangleright \rho_Y^{\text{Sets}_*, \triangleright}} & (X \triangleright Y) \triangleright S^0 \\
 & \searrow \alpha_{S^0, X, Y}^{\text{Sets}_*, \triangleright} & \downarrow \rho_{X \triangleright Y}^{\text{Sets}_*, \triangleright} \\
 & & X \triangleright Y
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 x \triangleright (y \triangleright 0) & \mapsto & (x \triangleright y) \triangleright 0 \\
 & \searrow & \downarrow \\
 & & x \triangleright y_0 = x_0 \triangleright y_0
 \end{array}$$

and

$$\begin{array}{ccc}
 x \triangleright (y \triangleright 1) & \mapsto & (x \triangleright y) \triangleright 1 \\
 & \searrow & \downarrow \\
 & & x \triangleright y
 \end{array}$$

and hence indeed commutes. Thus the right skew triangle identity is satisfied.

*The Right Skew Middle Triangle Identity:* Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets. We have to show that the diagram

$$\begin{array}{ccc}
 X \triangleright Y & \xlongequal{\quad} & X \triangleright Y \\
 \text{id}_X \triangleright \lambda_Y^{\text{Sets}_*, \triangleright} \downarrow & & \uparrow \rho_X^{\text{Sets}_*, \triangleright} \triangleright \text{id}_Y \\
 X \triangleright (S^0 \triangleright Y) & \xrightarrow{\alpha_{X, S^0, Y}^{\text{Sets}_*, \triangleright}} & (X \triangleright S^0) \triangleright Y
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 x \triangleright y & \xrightarrow{\quad} & x \triangleright y \\
 \downarrow & & \uparrow \\
 x \triangleright (1 \triangleright y) & \xrightarrow{\quad} & (x \triangleright 1) \triangleright y
 \end{array}$$

and hence indeed commutes. Thus the right skew triangle identity is satisfied.

*The Zig-Zag Identity:* We have to show that the diagram

$$\begin{array}{ccc}
 S^0 & \xrightarrow{\lambda_{S^0}^{\mathbf{Sets}_*, \triangleright}} & S^0 \triangleright S^0 \\
 & \searrow & \downarrow \rho_{S^0}^{\mathbf{Sets}_*, \triangleright} \\
 & & S^0
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 0 & \mapsto & 1 \triangleright 0 \\
 & \searrow & \downarrow \\
 & & 0
 \end{array}$$

and

$$\begin{array}{ccc}
 1 & \mapsto & 1 \triangleright 1 \\
 & \searrow & \downarrow \\
 & & 1
 \end{array}$$

and hence indeed commutes. Thus the zig-zag identity is satisfied.

*Right Skew Monoidal Right-Closedness:* This follows from **Item 2** of **Proposition 4.1.1.7**.  $\square$

## 4.9 Monoids With Respect to the Right Tensor Product of Pointed Sets

00FR

**00FS Proposition 4.9.1.1.** The category of monoids on  $(\mathbf{Sets}_*, \triangleright, S^0)$  is isomorphic to the category of “monoids with right zero”<sup>18</sup> and morphisms between them.

*Proof. Monoids on  $(\mathbf{Sets}_*, \triangleright, S^0)$ :* A monoid on  $(\mathbf{Sets}_*, \triangleright, S^0)$  consists of:

- *The Underlying Object.* A pointed set  $(A, 0_A)$ .

---

<sup>18</sup>A monoid with right zero is defined similarly as the monoids with zero of ???. Succinctly, they are monoids  $(A, \mu_A, \eta_A)$  with a special element  $0_A$  satisfying

$$0_A a = 0_A$$

for each  $a \in A$ .

- *The Multiplication Morphism.* A morphism of pointed sets

$$\mu_A: A \triangleright A \rightarrow A,$$

determining a right bilinear morphism of pointed sets

$$\begin{aligned} A \times A &\longrightarrow A \\ (a, b) &\longmapsto ab. \end{aligned}$$

- *The Unit Morphism.* A morphism of pointed sets

$$\eta_A: S^0 \rightarrow A$$

picking an element  $1_A$  of  $A$ .

satisfying the following conditions:

1. *Associativity.* The diagram

$$\begin{array}{ccc} & A \triangleright (A \triangleright A) & \\ \alpha_{A,A,A}^{\text{Sets}_*, \triangleright} \nearrow & & \searrow \text{id}_A \triangleright \mu_A \\ (A \triangleright A) \triangleright A & & A \triangleright A \\ \mu_A \triangleright \text{id}_A \searrow & & \swarrow \mu_A \\ & A \triangleright A & \xrightarrow{\mu_A} A \end{array}$$

2. *Left Unitality.* The diagram

$$\begin{array}{ccc} A & \xrightarrow{\lambda_A^{\text{Sets}_*, \triangleright}} & S^0 \triangleright A \\ \parallel & & \downarrow \eta_A \times \text{id}_A \\ A & \xleftarrow{\mu_A} & A \triangleright A \end{array}$$

commutes.

3. *Right Unitality.* The diagram

$$\begin{array}{ccc} A \triangleright S^0 & \xrightarrow{\text{id}_A \times \eta_A} & A \triangleright A \\ \searrow \rho_A^{\text{Sets}_*, \triangleright} & & \downarrow \mu_A \\ & & A \end{array}$$

commutes.




Being a right-bilinear morphism of pointed sets, the multiplication map satisfies

$$0_A a = 0_A$$

for each  $a \in A$ . Now, the associativity, left unitality, and right unitality conditions act on elements as follows:

1. *Associativity.* The associativity condition acts as




This gives

$$(ab)c = a(bc)$$

for each  $a, b, c \in A$ .

2. *Left Unitality.* The left unitality condition acts as




This gives

$$1_A a = a$$

for each  $a \in A$ .

3. *Right Unitality.* The right unitality condition acts:

(a) On  $1 \triangleright 0$  as



(b) On  $a \triangleright 1$  as

$$\begin{array}{ccc} a \triangleright 1 & & a \triangleright 1 \mapsto a \triangleright 1_A \\ & \searrow & \searrow \downarrow \\ & a & a1_A. \end{array}$$

This gives

$$\begin{aligned} a1_A &= a, \\ a0_A &= 0_A \end{aligned}$$

for each  $a \in A$ .

Thus we see that monoids with respect to  $\triangleright$  are exactly monoids with right zero.

*Morphisms of Monoids on  $(\mathbf{Sets}_*, \triangleright, S^0)$ :* A morphism of monoids on  $(\mathbf{Sets}_*, \triangleright, S^0)$  from  $(A, \mu_A, \eta_A, 0_A)$  to  $(B, \mu_B, \eta_B, 0_B)$  is a morphism of pointed sets

$$f: (A, 0_A) \rightarrow (B, 0_B)$$

satisfying the following conditions:

1. *Compatibility With the Multiplication Morphisms.* The diagram

$$\begin{array}{ccc} A \triangleright A & \xrightarrow{f \triangleright f} & B \triangleright B \\ \mu_A \downarrow & & \downarrow \mu_B \\ A & \xrightarrow{f} & B \end{array}$$

commutes.

2. *Compatibility With the Unit Morphisms.* The diagram

$$\begin{array}{ccc} S^0 & \xrightarrow{\eta_A} & A \\ & \searrow \eta_B & \downarrow f \\ & & B \end{array}$$

commutes.

These act on elements as

$$\begin{array}{ccc}
 a \triangleright b & & a \triangleright b \mapsto f(a) \triangleright f(b) \\
 \downarrow & & \downarrow \\
 ab \mapsto f(ab) & & f(a)f(b)
 \end{array}$$

and

$$\begin{array}{ccc}
 0 & & 0 \mapsto 0_A \\
 \searrow & & \downarrow \\
 & & f(0_A) \\
 & & \downarrow \\
 & & 0_B
 \end{array}$$

and

$$\begin{array}{ccc}
 1 & & 1 \mapsto 1_A \\
 \searrow & & \downarrow \\
 & & f(1_A) \\
 & & \downarrow \\
 & & 1_B
 \end{array}$$

giving

$$\begin{aligned}
 f(ab) &= f(a)f(b), \\
 f(0_A) &= 0_B, \\
 f(1_A) &= 1_B,
 \end{aligned}$$

for each  $a, b \in A$ , which is exactly a morphism of monoids with right zero.

*Identities and Composition:* Similarly, the identities and composition of  $\text{Mon}(\text{Sets}_*, \triangleright, S^0)$  can be easily seen to agree with those of monoids with right zero, which finishes the proof.  $\square$

## 00FT 5 The Smash Product of Pointed Sets

### 00FU 5.1 Foundations

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

**00FV Definition 5.1.1.1.** The **smash product of**  $(X, x_0)$  **and**  $(Y, y_0)$ <sup>19</sup> is the pointed set  $X \wedge Y$ <sup>20</sup> satisfying the bijection

$$\mathbf{Sets}_*(X \wedge Y, Z) \cong \mathbf{Hom}_{\mathbf{Sets}_*}^{\otimes}(X \times Y, Z),$$

naturally in  $(X, x_0), (Y, y_0), (Z, z_0) \in \mathbf{Obj}(\mathbf{Sets}_*)$ .

**00FW Remark 5.1.1.2.** That is to say, the smash product of pointed sets is defined so as to induce a bijection between the following data:

- Pointed maps  $f: X \wedge Y \rightarrow Z$ .
- Maps of sets  $f: X \times Y \rightarrow Z$  satisfying

$$\begin{aligned} f(x_0, y) &= z_0, \\ f(x, y_0) &= z_0 \end{aligned}$$

for each  $x \in X$  and each  $y \in Y$ .

**00FX Remark 5.1.1.3.** The smash product of pointed sets may be described as follows:

- The smash product of  $(X, x_0)$  and  $(Y, y_0)$  is the pair  $((X \wedge Y, x_0 \wedge y_0), \iota)$  consisting of
  - A pointed set  $(X \wedge Y, x_0 \wedge y_0)$ ;
  - A bilinear morphism of pointed sets  $\iota: (X \times Y, (x_0, y_0)) \rightarrow X \wedge Y$ ;

satisfying the following universal property:

(UP) Given another such pair  $((Z, z_0), f)$  consisting of

- \* A pointed set  $(Z, z_0)$ ;
- \* A bilinear morphism of pointed sets  $f: (X \times Y, (x_0, y_0)) \rightarrow X \wedge Y$ ;

there exists a unique morphism of pointed sets  $X \wedge Y \xrightarrow{\exists!} Z$

<sup>19</sup>*Further Terminology:* In the context of monoids with zero as models for  $\mathbb{F}_1$ -algebras, the smash product  $X \wedge Y$  is also called the **tensor product of  $\mathbb{F}_1$ -modules of**  $(X, x_0)$  **and**  $(Y, y_0)$  or the **tensor product of**  $(X, x_0)$  **and**  $(Y, y_0)$  **over**  $\mathbb{F}_1$ .

<sup>20</sup>*Further Notation:* In the context of monoids with zero as models for  $\mathbb{F}_1$ -algebras, the smash product  $X \wedge Y$  is also denoted  $X \otimes_{\mathbb{F}_1} Y$ .

making the diagram

$$\begin{array}{ccc} & & X \wedge Y \\ & \nearrow \iota & \downarrow \exists! \\ X \times Y & \xrightarrow{f} & Z \end{array}$$

commute.

**00FY Construction 5.1.1.4.** Concretely, the **smash product** of  $(X, x_0)$  and  $(Y, y_0)$  is the pointed set  $(X \wedge Y, x_0 \wedge y_0)$  consisting of

- *The Underlying Set.* The set  $X \wedge Y$  defined by

$$X \wedge Y \cong (X \times Y) / \sim_R,$$

where  $\sim_R$  is the equivalence relation on  $X \times Y$  obtained by declaring

$$\begin{aligned} (x_0, y) &\sim_R (x_0, y'), \\ (x, y_0) &\sim_R (x', y_0) \end{aligned}$$

for each  $x, x' \in X$  and each  $y, y' \in Y$ ;

- *The Basepoint.* The element  $[(x_0, y_0)]$  of  $X \wedge Y$  given by the equivalence class of  $(x_0, y_0)$  under the equivalence relation  $\sim$  on  $X \times Y$ .

*Proof.* By **Equivalence Relations and Apartness Relations, Item 6** of **Proposition 5.2.1.3**, we have a natural bijection

$$\mathbf{Sets}_*(X \wedge Y, Z) \cong \mathbf{Hom}_{\mathbf{Sets}}^R(X \times Y, Z).$$

Now, by definition,  $\mathbf{Hom}_{\mathbf{Sets}}^R(X \times Y, Z)$  is the set

$$\mathbf{Hom}_{\mathbf{Sets}}^R(X \times Y, Z) \stackrel{\text{def}}{=} \left\{ f \in \mathbf{Hom}_{\mathbf{Sets}}(X \times Y, Z) \left| \begin{array}{l} \text{for each } x, y \in X, \text{ if} \\ (x, y) \sim_R (x', y'), \text{ then} \\ f(x, y) = f(x', y') \end{array} \right. \right\}.$$

However, the condition  $(x, y) \sim_R (x', y')$  only holds when:

1. We have  $x = x'$  and  $y = y'$ .
2. The following conditions are satisfied:

- (a) We have  $x = x_0$  or  $y = y_0$ .
- (b) We have  $x' = x_0$  or  $y' = y_0$ .

So, given  $f \in \text{Hom}_{\text{Sets}}(X \times Y, Z)$  with a corresponding  $\bar{f}: X \wedge Y \rightarrow Z$ , the latter case above implies

$$\begin{aligned} f(x_0, y) &= f(x, y_0) \\ &= f(x_0, y_0), \end{aligned}$$

and since  $\bar{f}: X \wedge Y \rightarrow Z$  is a pointed map, we have

$$\begin{aligned} f(x_0, y_0) &= \bar{f}(x_0, y_0) \\ &= z_0. \end{aligned}$$

Thus the elements  $f$  in  $\text{Hom}_{\text{Sets}}(X \times Y, Z)$  are precisely those functions  $f: X \times Y \rightarrow Z$  satisfying the equalities

$$\begin{aligned} f(x_0, y) &= z_0, \\ f(x, y_0) &= z_0 \end{aligned}$$

for each  $x \in X$  and each  $y \in Y$ , giving an equality

$$\text{Hom}_{\text{Sets}}^R(X \times Y, Z) = \text{Hom}_{\text{Sets}_*}^{\otimes}(X \times Y, Z)$$

of sets, which when composed with our earlier isomorphism

$$\text{Sets}_*(X \wedge Y, Z) \cong \text{Hom}_{\text{Sets}}^R(X \times Y, Z)$$

gives our desired natural bijection, finishing the proof.  $\square$

**00FZ Remark 5.1.1.5.** It is also somewhat common to write

$$X \wedge Y \stackrel{\text{def}}{=} \frac{X \times Y}{X \vee Y},$$

identifying  $X \vee Y$  with the subspace  $(\{x_0\} \times Y) \cup (X \times \{y_0\})$  of  $X \times Y$ , and having the quotient be defined by declaring  $(x, y) \sim (x', y')$  iff we have  $(x, y), (x', y') \in X \vee Y$ .

**00G0 Notation 5.1.1.6.** We write  $x \wedge y$  for the element  $[(x, y)]$  of

$$X \wedge Y \cong X \times Y / \sim.$$

**00G1 Remark 5.1.1.7.** Employing the notation introduced in [Notation 5.1.1.6](#), we have

$$\begin{aligned} x_0 \wedge y_0 &= x \wedge y_0, \\ &= x_0 \wedge y \end{aligned}$$

for each  $x \in X$  and each  $y \in Y$ , and

$$\begin{aligned} x \wedge y_0 &= x' \wedge y_0, \\ x_0 \wedge y &= x_0 \wedge y' \end{aligned}$$

for each  $x, x' \in X$  and each  $y, y' \in Y$ .

**00G2 Example 5.1.1.8.** Here are some examples of smash products of pointed sets.

**00G3** 1. *Smashing With pt.* For any pointed set  $X$ , we have isomorphisms of pointed sets

$$\begin{aligned} \text{pt} \wedge X &\cong \text{pt}, \\ X \wedge \text{pt} &\cong \text{pt}. \end{aligned}$$

**00G4** 2. *Smashing With  $S^0$ .* For any pointed set  $X$ , we have isomorphisms of pointed sets

$$\begin{aligned} S^0 \wedge X &\cong X, \\ X \wedge S^0 &\cong X. \end{aligned}$$

**00G5 Proposition 5.1.1.9.** Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

**00G6** 1. *Functoriality.* The assignments  $X, Y, (X, Y) \mapsto X \wedge Y$  define functors

$$\begin{aligned} X \wedge - &: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*, \\ - \wedge Y &: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*, \\ -_1 \wedge -_2 &: \mathbf{Sets}_* \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*. \end{aligned}$$

In particular, given pointed maps

$$\begin{aligned} f &: (X, x_0) \rightarrow (A, a_0), \\ g &: (Y, y_0) \rightarrow (B, b_0), \end{aligned}$$

the induced map

$$f \wedge g: X \wedge Y \rightarrow A \wedge B$$

is given by

$$[f \wedge g](x \wedge y) \stackrel{\text{def}}{=} f(x) \wedge g(y)$$

for each  $x \wedge y \in X \wedge Y$ .

**00G7** 2. *Adjointness.* We have adjunctions

$$\begin{aligned} (X \wedge - \dashv \mathbf{Sets}_*(X, -)) &: \mathbf{Sets}_* \begin{array}{c} \xrightarrow{X \wedge -} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(X, -)} \end{array} \mathbf{Sets}_*, \\ (- \wedge Y \dashv \mathbf{Sets}_*(Y, -)) &: \mathbf{Sets}_* \begin{array}{c} \xrightarrow{- \wedge Y} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(Y, -)} \end{array} \mathbf{Sets}_*, \end{aligned}$$

witnessed by bijections

$$\begin{aligned} \text{Hom}_{\mathbf{Sets}_*}(X \wedge Y, Z) &\cong \text{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(Y, Z)), \\ \text{Hom}_{\mathbf{Sets}_*}(X \wedge Y, Z) &\cong \text{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(A, Z)), \end{aligned}$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$ .

**00G8** 3. *Enriched Adjointness.* We have  $\mathbf{Sets}_*$ -enriched adjunctions

$$\begin{aligned} (X \wedge - \dashv \mathbf{Sets}_*(X, -)) &: \mathbf{Sets}_* \begin{array}{c} \xrightarrow{X \wedge -} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(X, -)} \end{array} \mathbf{Sets}_*, \\ (- \wedge Y \dashv \mathbf{Sets}_*(Y, -)) &: \mathbf{Sets}_* \begin{array}{c} \xrightarrow{- \wedge Y} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(Y, -)} \end{array} \mathbf{Sets}_*, \end{aligned}$$

witnessed by isomorphisms of pointed sets

$$\begin{aligned} \mathbf{Sets}_*(X \wedge Y, Z) &\cong \mathbf{Sets}_*(X, \mathbf{Sets}_*(Y, Z)), \\ \mathbf{Sets}_*(X \wedge Y, Z) &\cong \mathbf{Sets}_*(X, \mathbf{Sets}_*(A, Z)), \end{aligned}$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$ .



**00G9** 4. *As a Pushout.* We have an isomorphism

$$X \wedge Y \cong \text{pt} \coprod_{X \vee Y} (X \times Y), \quad \begin{array}{ccc} X \wedge Y & \leftarrow & X \times Y \\ \uparrow \ulcorner & & \uparrow \iota \\ \text{pt} & \xleftarrow{\quad} & X \vee Y \end{array}$$

natural in  $X, Y \in \text{Obj}(\mathbf{Sets}_*)$ , where the pushout is taken in  $\mathbf{Sets}$ , and the embedding  $\iota: X \vee Y \hookrightarrow X \times Y$  is defined following [Remark 5.1.1.5](#).

**00GA** 5. *Distributivity Over Wedge Sums.* We have isomorphisms of pointed sets

$$\begin{aligned} X \wedge (Y \vee Z) &\cong (X \wedge Y) \vee (X \wedge Z), \\ (X \vee Y) \wedge Z &\cong (X \wedge Z) \vee (Y \wedge Z), \end{aligned}$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$ .

*Proof. Item 1, Functoriality:* The map  $f \wedge g$  comes from [Equivalence Relations and Apartness Relations](#), [Item 4](#) of [Proposition 5.2.1.3](#) via the map

$$f \wedge g: X \times Y \rightarrow A \wedge B$$

sending  $(x, y)$  to  $f(x) \wedge g(y)$ , which we need to show satisfies

$$[f \wedge g](x, y) = [f \wedge g](x', y')$$

for each  $(x, y), (x', y') \in X \times Y$  with  $(x, y) \sim_R (x', y')$ , where  $\sim_R$  is the relation constructing  $X \wedge Y$  as

$$X \wedge Y \cong (X \times Y) / \sim_R$$

in [Construction 5.1.1.4](#). The condition defining  $\sim$  is that at least one of the following conditions is satisfied:

1. We have  $x = x'$  and  $y = y'$ ;
2. Both of the following conditions are satisfied:
  - (a) We have  $x = x_0$  or  $y = y_0$ .
  - (b) We have  $x' = x_0$  or  $y' = y_0$ .

We have five cases:

1. In the first case, we clearly have

$$[f \wedge g](x, y) = [f \wedge g](x', y')$$

since  $x = x'$  and  $y = y'$ .

2. If  $x = x_0$  and  $x' = x_0$ , we have

$$\begin{aligned} [f \wedge g](x_0, y) &\stackrel{\text{def}}{=} f(x_0) \wedge g(y) \\ &= a_0 \wedge g(y) \\ &= a_0 \wedge g(y') \\ &= f(x_0) \wedge g(y') \\ &\stackrel{\text{def}}{=} [f \wedge g](x_0, y'). \end{aligned}$$

3. If  $x = x_0$  and  $y' = y_0$ , we have

$$\begin{aligned} [f \wedge g](x_0, y) &\stackrel{\text{def}}{=} f(x_0) \wedge g(y) \\ &= a_0 \wedge g(y) \\ &= a_0 \wedge b_0 \\ &= f(x') \wedge b_0 \\ &= f(x') \wedge g(y_0) \\ &\stackrel{\text{def}}{=} [f \wedge g](x', y_0). \end{aligned}$$

4. If  $y = y_0$  and  $x' = x_0$ , we have

$$\begin{aligned} [f \wedge g](x, y_0) &\stackrel{\text{def}}{=} f(x) \wedge g(y_0) \\ &= f(x) \wedge b_0 \\ &= a_0 \wedge b_0 \\ &= a_0 \wedge g(y') \\ &= f(x_0) \wedge g(y') \\ &\stackrel{\text{def}}{=} [f \wedge g](x_0, y'). \end{aligned}$$

5. If  $y = y_0$  and  $y' = y_0$ , we have

$$\begin{aligned} [f \wedge g](x, y_0) &\stackrel{\text{def}}{=} f(x) \wedge g(y_0) \\ &= f(x) \wedge b_0 \\ &= f(x') \wedge b_0 \\ &= f(x) \wedge g(y_0) \\ &\stackrel{\text{def}}{=} [f \wedge g](x', y_0). \end{aligned}$$

Thus  $f \wedge g$  is well-defined. Next, we claim that  $\wedge$  preserves identities and composition:

- *Preservation of Identities.* We have

$$\begin{aligned} [\text{id}_X \wedge \text{id}_Y](x \wedge y) &\stackrel{\text{def}}{=} \text{id}_X(x) \wedge \text{id}_Y(y) \\ &= x \wedge y \\ &= [\text{id}_{X \wedge Y}](x \wedge y) \end{aligned}$$

for each  $x \wedge y \in X \wedge Y$ , and thus

$$\text{id}_X \wedge \text{id}_Y = \text{id}_{X \wedge Y}.$$

- *Preservation of Composition.* Given pointed maps

$$\begin{aligned} f &: (X, x_0) \rightarrow (X', x'_0), \\ h &: (X', x'_0) \rightarrow (X'', x''_0), \\ g &: (Y, y_0) \rightarrow (Y', y'_0), \\ k &: (Y', y'_0) \rightarrow (Y'', y''_0), \end{aligned}$$

we have

$$\begin{aligned} [(h \circ f) \wedge (k \circ g)](x \wedge y) &\stackrel{\text{def}}{=} h(f(x)) \wedge k(g(y)) \\ &\stackrel{\text{def}}{=} [h \wedge k](f(x) \wedge g(y)) \\ &\stackrel{\text{def}}{=} [h \wedge k]([f \wedge g](x \wedge y)) \\ &\stackrel{\text{def}}{=} [(h \wedge k) \circ (f \wedge g)](x \wedge y) \end{aligned}$$

for each  $x \wedge y \in X \wedge Y$ , and thus

$$(h \circ f) \wedge (k \circ g) = (h \wedge k) \circ (f \wedge g).$$

This finishes the proof.

*Item 2, Adjointness:* We prove only the adjunction  $- \wedge Y \dashv \mathbf{Sets}_*(Y, -)$ , witnessed by a natural bijection

$$\text{Hom}_{\mathbf{Sets}_*}(X \wedge Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(Y, Z)),$$

as the proof of the adjunction  $X \wedge - \dashv \mathbf{Sets}_*(X, -)$  is similar. We claim we have a bijection

$$\text{Hom}_{\mathbf{Sets}_*}^{\otimes}(X \times Y, Z) \cong \text{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(Y, Z))$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$ , implying the desired adjunction. Indeed, this bijection is a restriction of the bijection

$$\mathbf{Sets}(X \times Y, Z) \cong \mathbf{Sets}(X, \mathbf{Sets}(Y, Z))$$

of **Constructions With Sets, Item 2** of **Proposition 1.3.1.2**:

- A map

$$\xi: X \times Y \rightarrow Z$$

in  $\text{Hom}_{\mathbf{Sets}_*}^{\otimes}(X \times Y, Z)$  gets sent to the pointed map

$$\begin{aligned} \xi^\dagger: (X, x_0) &\rightarrow (\mathbf{Sets}_*(Y, Z), \Delta_{z_0}), \\ x &\longmapsto (\xi_x^\dagger: Y \rightarrow Z), \end{aligned}$$

where  $\xi_x^\dagger: Y \rightarrow Z$  is the map defined by

$$\xi_x^\dagger(y) \stackrel{\text{def}}{=} \xi(x, y)$$

for each  $y \in Y$ , where:

- The map  $\xi^\dagger$  is indeed pointed, as we have

$$\begin{aligned} \xi_{x_0}^\dagger(y) &\stackrel{\text{def}}{=} \xi(x_0, y) \\ &\stackrel{\text{def}}{=} z_0 \end{aligned}$$

for each  $y \in Y$ . Thus  $\xi_{x_0}^\dagger = \Delta_{z_0}$  and  $\xi^\dagger$  is pointed.

- The map  $\xi_x^\dagger$  indeed lies in  $\mathbf{Sets}_*(Y, Z)$ , as we have

$$\begin{aligned} \xi_x^\dagger(y_0) &\stackrel{\text{def}}{=} \xi(x, y_0) \\ &\stackrel{\text{def}}{=} z_0. \end{aligned}$$

- Conversely, a map

$$\begin{aligned} \xi: (X, x_0) &\rightarrow (\mathbf{Sets}_*(Y, Z), \Delta_{z_0}), \\ x &\longmapsto (\xi_x: Y \rightarrow Z), \end{aligned}$$

in  $\text{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(Y, Z))$  gets sent to the map

$$\xi^\dagger: X \times Y \rightarrow Z$$

defined by

$$\xi^\dagger(x, y) \stackrel{\text{def}}{=} \xi_x(y)$$

for each  $(x, y) \in X \times Y$ , which indeed lies in  $\text{Hom}_{\mathbf{Sets}_*}^{\otimes}(X \times Y, Z)$ , as:

– *Left Bilinearity.* We have

$$\begin{aligned}\xi^\dagger(x_0, y) &\stackrel{\text{def}}{=} \xi_{x_0}(y) \\ &\stackrel{\text{def}}{=} \Delta_{z_0}(y) \\ &\stackrel{\text{def}}{=} z_0\end{aligned}$$

for each  $y \in Y$ , since  $\xi_{x_0} = \Delta_{z_0}$  as  $\xi$  is assumed to be a pointed map.

– *Right Bilinearity.* We have

$$\begin{aligned}\xi^\dagger(x, y_0) &\stackrel{\text{def}}{=} \xi_x(y_0) \\ &\stackrel{\text{def}}{=} z_0\end{aligned}$$

for each  $x \in X$ , since  $\xi_x \in \mathbf{Sets}_*(Y, Z)$  is a morphism of pointed sets.

This finishes the proof.

*Item 3, Enriched Adjointness:* This follows from [Item 2](#) and [??](#), [??](#) of [??](#).

*Item 4, As a Pushout:* Following the description of [Constructions With Sets](#), [Remark 2.4.1.2](#), we have

$$\text{pt} \coprod_{X \vee Y} (X \times Y) \cong (\text{pt} \times (X \times Y)) / \sim,$$

where  $\sim$  identifies the element  $\star$  in  $\text{pt}$  with all elements of the form  $(x_0, y)$  and  $(x, y_0)$  in  $X \times Y$ . Thus [Equivalence Relations and Apartness Relations](#), [Item 4](#) of [Proposition 5.2.1.3](#) coupled with [Remark 5.1.1.7](#) then gives us a well-defined map

$$\text{pt} \coprod_{X \vee Y} (X \times Y) \rightarrow X \wedge Y$$

via  $[(\star, (x, y))] \mapsto x \wedge y$ , with inverse

$$X \wedge Y \rightarrow \text{pt} \coprod_{X \vee Y} (X \times Y)$$

given by  $x \wedge y \mapsto [(\star, (x, y))]$ .

*Item 5, Distributivity Over Wedge Sums:* This follows from [Proposition 5.9.1.1](#), [??](#), [??](#) of [??](#), and the fact that  $\vee$  is the coproduct in  $\mathbf{Sets}_*$  ([Pointed Sets](#), [Definition 3.3.1.1](#)).  $\square$

## **00GB** 5.2 The Internal Hom of Pointed Sets

Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

**00GC Definition 5.2.1.1.** The **internal Hom**<sup>21</sup> of pointed sets from  $(X, x_0)$  to  $(Y, y_0)$  is the pointed set  $\mathbf{Sets}_*((X, x_0), (Y, y_0))$ <sup>22</sup> consisting of:

- *The Underlying Set.* The set  $\mathbf{Sets}_*((X, x_0), (Y, y_0))$  of morphisms of pointed sets from  $(X, x_0)$  to  $(Y, y_0)$ .
- *The Basepoint.* The element

$$\Delta_{y_0}: (X, x_0) \rightarrow (Y, y_0)$$

of  $\mathbf{Sets}_*((X, x_0), (Y, y_0))$  given by

$$\Delta_{y_0}(x) \stackrel{\text{def}}{=} y_0$$

for each  $x \in X$ .

*Proof.* For a proof that  $\mathbf{Sets}_*$  is indeed the internal Hom of  $\mathbf{Sets}_*$  with respect to the smash product of pointed sets, see **Item 2** of **Proposition 5.1.1.9**.  $\square$

**00GD Proposition 5.2.1.2.** Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets.

**00GE** 1. *Functoriality.* The assignments  $X, Y, (X, Y) \mapsto \mathbf{Sets}_*(X, Y)$  define functors

$$\begin{aligned} \mathbf{Sets}_*(X, -) &: \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*, \\ \mathbf{Sets}_*(-, Y) &: \mathbf{Sets}_*^{\text{op}} \rightarrow \mathbf{Sets}_*, \\ \mathbf{Sets}_*(-, -) &: \mathbf{Sets}_*^{\text{op}} \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*. \end{aligned}$$

In particular, given pointed maps

$$\begin{aligned} f &: (X, x_0) \rightarrow (A, a_0), \\ g &: (Y, y_0) \rightarrow (B, b_0), \end{aligned}$$

the induced map

$$\mathbf{Sets}_*(f, g): \mathbf{Sets}_*(A, Y) \rightarrow \mathbf{Sets}_*(X, B)$$

is given by

$$[\mathbf{Sets}_*(f, g)](\phi) \stackrel{\text{def}}{=} g \circ \phi \circ f$$

for each  $\phi \in \mathbf{Sets}_*(A, Y)$ .

<sup>21</sup>The pointed set  $\mathbf{Sets}_*(X, Y)$  is the internal **Hom** of  $\mathbf{Sets}_*$  with respect to the smash product of **Tensor Products of Pointed Sets**, **Definition 5.1.1.1**; see **Tensor Products of Pointed Sets**, **Item 2** of **Proposition 5.1.1.9**.

<sup>22</sup>*Further Notation:* Also written  $\mathbf{Hom}_{\mathbf{Sets}_*}(X, Y)$ .

**00GF** 2. *Adjointness.* We have adjunctions

$$\begin{aligned} (X \wedge - \dashv \mathbf{Sets}_*(X, -)) : \quad & \mathbf{Sets}_* \begin{array}{c} \xrightarrow{X \wedge -} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(X, -)} \end{array} \mathbf{Sets}_*, \\ (- \wedge Y \dashv \mathbf{Sets}_*(Y, -)) : \quad & \mathbf{Sets}_* \begin{array}{c} \xrightarrow{- \wedge Y} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(Y, -)} \end{array} \mathbf{Sets}_*, \end{aligned}$$

witnessed by bijections

$$\begin{aligned} \mathrm{Hom}_{\mathbf{Sets}_*}(X \wedge Y, Z) &\cong \mathrm{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(Y, Z)), \\ \mathrm{Hom}_{\mathbf{Sets}_*}(X \wedge Y, Z) &\cong \mathrm{Hom}_{\mathbf{Sets}_*}(X, \mathbf{Sets}_*(A, Z)), \end{aligned}$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \mathrm{Obj}(\mathbf{Sets}_*)$ .

**00GG** 3. *Enriched Adjointness.* We have  $\mathbf{Sets}_*$ -enriched adjunctions

$$\begin{aligned} (X \wedge - \dashv \mathbf{Sets}_*(X, -)) : \quad & \mathbf{Sets}_* \begin{array}{c} \xrightarrow{X \wedge -} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(X, -)} \end{array} \mathbf{Sets}_*, \\ (- \wedge Y \dashv \mathbf{Sets}_*(Y, -)) : \quad & \mathbf{Sets}_* \begin{array}{c} \xrightarrow{- \wedge Y} \\ \perp \\ \xleftarrow{\mathbf{Sets}_*(Y, -)} \end{array} \mathbf{Sets}_*, \end{aligned}$$

witnessed by isomorphisms of pointed sets

$$\begin{aligned} \mathbf{Sets}_*(X \wedge Y, Z) &\cong \mathbf{Sets}_*(X, \mathbf{Sets}_*(Y, Z)), \\ \mathbf{Sets}_*(X \wedge Y, Z) &\cong \mathbf{Sets}_*(X, \mathbf{Sets}_*(A, Z)), \end{aligned}$$

natural in  $(X, x_0), (Y, y_0), (Z, z_0) \in \mathrm{Obj}(\mathbf{Sets}_*)$ .

*Proof. Item 1, Functoriality:* This follows from **Constructions With Sets**, **Item 1** of **Proposition 3.5.1.2** and from the equalities

$$\begin{aligned} g \circ \Delta_{y_0} &= \Delta_{z_0}, \\ \Delta_{y_0} \circ f &= \Delta_{y_0} \end{aligned}$$

for morphisms  $f: (K, k_0) \rightarrow (X, x_0)$  and  $g: (Y, y_0) \rightarrow (Z, z_0)$ , which guarantee pre- and postcomposition by morphisms of pointed sets to also be morphisms of pointed sets.

*Item 2, Adjointness:* This is a repetition of **Item 2** of **Proposition 5.1.1.9**, and is proved there.

*Item 3, Enriched Adjointness:* This is a repetition of **Item 3** of **Proposition 5.1.1.9**, and is proved there.  $\square$

### 00GH 5.3 The Monoidal Unit

00GJ **Definition 5.3.1.1.** The monoidal unit of the smash product of pointed sets is the functor

$$\mathbb{1}^{\mathbf{Sets}_*} : \mathbf{pt} \rightarrow \mathbf{Sets}_*$$

defined by

$$\mathbb{1}_{\mathbf{Sets}_*} \stackrel{\text{def}}{=} S^0.$$

### 00GK 5.4 The Associator

00GL **Definition 5.4.1.1.** The associator of the smash product of pointed sets is the natural isomorphism

$$\alpha^{\mathbf{Sets}_*} : \wedge \circ (\wedge \times \text{id}_{\mathbf{Sets}_*}) \xrightarrow{\sim} \wedge \circ (\text{id}_{\mathbf{Sets}_*} \times \wedge) \circ \alpha_{\mathbf{Sets}_*, \mathbf{Sets}_*, \mathbf{Sets}_*}^{\mathbf{Cats}},$$

as in the diagram

$$\begin{array}{ccc}
 & \mathbf{Sets}_* \times (\mathbf{Sets}_* \times \mathbf{Sets}_*) & \\
 \alpha_{\mathbf{Sets}_*, \mathbf{Sets}_*, \mathbf{Sets}_*}^{\mathbf{Cats}} \swarrow & \nearrow \text{id} \times \wedge & \\
 (\mathbf{Sets}_* \times \mathbf{Sets}_*) \times \mathbf{Sets}_* & & \mathbf{Sets}_* \times \mathbf{Sets}_* \\
 \wedge \times \text{id} \searrow & \nearrow \alpha^{\mathbf{Sets}_*} & \searrow \wedge \\
 \mathbf{Sets}_* \times \mathbf{Sets}_* & \xrightarrow{\wedge} & \mathbf{Sets}_*
 \end{array}$$

whose component

$$\alpha_{X,Y,Z}^{\mathbf{Sets}_*} : (X \wedge Y) \wedge Z \xrightarrow{\cong} X \wedge (Y \wedge Z)$$

at  $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\mathbf{Sets}_*)$  is given by

$$\alpha_{X,Y,Z}^{\mathbf{Sets}_*}((x \wedge y) \wedge z) \stackrel{\text{def}}{=} x \wedge (y \wedge z)$$

for each  $(x \wedge y) \wedge z \in (X \wedge Y) \wedge Z$ .

*Proof. Well-Definedness:* Let  $[(x, y), z] = [(x', y'), z']$  be an element in  $(X \wedge Y) \wedge Z$ . Then either:

1. We have  $x = x'$ ,  $y = y'$ , and  $z = z'$ .



2. Both of the following conditions are satisfied:

- (a) We have  $x = x_0$  or  $y = y_0$  or  $z = z_0$ .
- (b) We have  $x' = x_0$  or  $y' = y_0$  or  $z' = z_0$ .

In the first case,  $\alpha_{X,Y,Z}^{\text{Sets}_*}$  clearly sends both elements to the same element in  $X \wedge (Y \wedge Z)$ . Meanwhile, in the latter case both elements are equal to the basepoint  $(x_0 \wedge y_0) \wedge z_0$  of  $(X \wedge Y) \wedge Z$ , which gets sent to the basepoint  $x_0 \wedge (y_0 \wedge z_0)$  of  $X \wedge (Y \wedge Z)$ .

*Being a Morphism of Pointed Sets:* As just mentioned, we have

$$\alpha_{X,Y,Z}^{\text{Sets}_*}((x_0 \wedge y_0) \wedge z_0) \stackrel{\text{def}}{=} x_0 \wedge (y_0 \wedge z_0),$$

and thus  $\alpha_{X,Y,Z}^{\text{Sets}_*}$  is a morphism of pointed sets.

*Invertibility:* Clearly, the inverse of  $\alpha_{X,Y,Z}^{\text{Sets}_*}$  is given by the morphism

$$\alpha_{X,Y,Z}^{\text{Sets}_*, -1}: X \wedge (Y \wedge Z) \xrightarrow{\cong} (X \wedge Y) \wedge Z$$

defined by

$$\alpha_{X,Y,Z}^{\text{Sets}_*, -1}(x \wedge (y \wedge z)) \stackrel{\text{def}}{=} (x \wedge y) \wedge z$$

for each  $x \wedge (y \wedge z) \in X \wedge (Y \wedge Z)$ .

*Naturality:* We need to show that, given morphisms of pointed sets

$$\begin{aligned} f &: (X, x_0) \rightarrow (X', x'_0), \\ g &: (Y, y_0) \rightarrow (Y', y'_0), \\ h &: (Z, z_0) \rightarrow (Z', z'_0) \end{aligned}$$

the diagram

$$\begin{array}{ccc} (X \wedge Y) \wedge Z & \xrightarrow{(f \wedge g) \wedge h} & (X' \wedge Y') \wedge Z' \\ \alpha_{X,Y,Z}^{\text{Sets}_*} \downarrow & & \downarrow \alpha_{X',Y',Z'}^{\text{Sets}_*} \\ X \wedge (Y \wedge Z) & \xrightarrow{f \wedge (g \wedge h)} & X' \wedge (Y' \wedge Z') \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} (x \wedge y) \wedge z & \longmapsto & (f(x) \wedge g(y)) \wedge h(z) \\ \downarrow & & \downarrow \\ x \wedge (y \wedge z) & \longmapsto & f(x) \wedge (g(y) \wedge h(z)) \end{array}$$

and hence indeed commutes, showing  $\alpha^{\mathbf{Sets}_*}$  to be a natural transformation. *Being a Natural Isomorphism:* Since  $\alpha^{\mathbf{Sets}_*}$  is natural and  $\alpha^{\mathbf{Sets}_*, -1}$  is a componentwise inverse to  $\alpha^{\mathbf{Sets}_*}$ , it follows from **Categories, Item 2 of Proposition 8.6.1.2** that  $\alpha^{\mathbf{Sets}_*, -1}$  is also natural. Thus  $\alpha^{\mathbf{Sets}_*}$  is a natural isomorphism.  $\square$

## 00GM 5.5 The Left Unitor

00GN **Definition 5.5.1.1.** The left unitor of the smash product of pointed sets is the natural isomorphism

$$\lambda^{\mathbf{Sets}_*} : \wedge \circ (\mathbb{1}^{\mathbf{Sets}_*} \times \text{id}_{\mathbf{Sets}_*}) \xrightarrow{\sim} \lambda_{\mathbf{Sets}_*}^{\mathbf{Cats}_2}$$

whose component

$$\lambda_X^{\mathbf{Sets}_*} : S^0 \wedge X \xrightarrow{\cong} X$$

at  $X \in \text{Obj}(\mathbf{Sets}_*)$  is given by

$$\begin{aligned} 0 \wedge x &\mapsto x_0, \\ 1 \wedge x &\mapsto x. \end{aligned}$$

*Proof. Well-Definedness:* Let  $[(x, y)] = [(x', y')]$  be an element in  $S^0 \wedge X$ . Then either:

1. We have  $x = x'$  and  $y = y'$ .
2. Both of the following conditions are satisfied:
  - (a) We have  $x = 0$  or  $y = x_0$ .
  - (b) We have  $x' = 0$  or  $y' = x_0$ .

In the first case,  $\lambda_X^{\mathbf{Sets}_*}$  clearly sends both elements to the same element in  $X$ . Meanwhile, in the latter case both elements are equal to the basepoint  $0 \wedge x_0$  of  $S^0 \wedge X$ , which gets sent to the basepoint  $x_0$  of  $X$ .

*Being a Morphism of Pointed Sets:* As just mentioned, we have

$$\lambda_X^{\mathbf{Sets}_*}(0 \wedge x_0) \stackrel{\text{def}}{=} x_0,$$

and thus  $\lambda_X^{\mathbf{Sets}_*}$  is a morphism of pointed sets.

*Invertibility:* The inverse of  $\lambda_X^{\mathbf{Sets}_*}$  is the morphism

$$\lambda_X^{\mathbf{Sets}_*, -1} : X \xrightarrow{\cong} S^0 \wedge X$$

defined by

$$\lambda_X^{\mathbf{Sets}_*, -1}(x) \stackrel{\text{def}}{=} 1 \wedge x$$

for each  $x \in X$ . Indeed:

- *Invertibility I.* We have

$$\begin{aligned} \left[ \lambda_X^{\mathbf{Sets}_*, -1} \circ \lambda_X^{\mathbf{Sets}_*} \right] (0 \wedge x) &= \lambda_X^{\mathbf{Sets}_*, -1} \left( \lambda_X^{\mathbf{Sets}_*} (0 \wedge x) \right) \\ &= \lambda_X^{\mathbf{Sets}_*, -1} (x_0) \\ &= 1 \wedge x_0 \\ &= 0 \wedge x, \end{aligned}$$

and

$$\begin{aligned} \left[ \lambda_X^{\mathbf{Sets}_*, -1} \circ \lambda_X^{\mathbf{Sets}_*} \right] (1 \wedge x) &= \lambda_X^{\mathbf{Sets}_*, -1} \left( \lambda_X^{\mathbf{Sets}_*} (1 \wedge x) \right) \\ &= \lambda_X^{\mathbf{Sets}_*, -1} (x) \\ &= 1 \wedge x \end{aligned}$$

for each  $x \in X$ , and thus we have

$$\lambda_X^{\mathbf{Sets}_*, -1} \circ \lambda_X^{\mathbf{Sets}_*} = \text{id}_{S^0 \wedge X}.$$

- *Invertibility II.* We have

$$\begin{aligned} \left[ \lambda_X^{\mathbf{Sets}_*} \circ \lambda_X^{\mathbf{Sets}_*, -1} \right] (x) &= \lambda_X^{\mathbf{Sets}_*} \left( \lambda_X^{\mathbf{Sets}_*, -1} (x) \right) \\ &= \lambda_X^{\mathbf{Sets}_*, -1} (1 \wedge x) \\ &= x \end{aligned}$$

for each  $x \in X$ , and thus we have

$$\lambda_X^{\mathbf{Sets}_*} \circ \lambda_X^{\mathbf{Sets}_*, -1} = \text{id}_X.$$

This shows  $\lambda_X^{\mathbf{Sets}_*}$  to be invertible.

*Naturality:* We need to show that, given a morphism of pointed sets

$$f: (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} S^0 \wedge X & \xrightarrow{\text{id}_{S^0} \wedge f} & S^0 \wedge Y \\ \lambda_X^{\mathbf{Sets}_*} \downarrow & & \downarrow \lambda_Y^{\mathbf{Sets}_*} \\ X & \xrightarrow{f} & Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} 0 \wedge x & & 0 \wedge x \mapsto 0 \wedge f(x) \\ \downarrow & & \downarrow \\ x_0 \mapsto & f(x_0) & y_0 \end{array}$$

and

$$\begin{array}{ccc} 1 \wedge x \mapsto & 1 \wedge f(x) & \\ \downarrow & \downarrow & \\ x \mapsto & f(x) & \end{array}$$

and hence indeed commutes, showing  $\lambda^{\mathbf{Sets}_*}$  to be a natural transformation.

*Being a Natural Isomorphism:* Since  $\lambda^{\mathbf{Sets}_*}$  is natural and  $\lambda^{\mathbf{Sets}_*, -1}$  is a componentwise inverse to  $\lambda^{\mathbf{Sets}_*}$ , it follows from **Categories, Item 2** of **Proposition 8.6.1.2** that  $\lambda^{\mathbf{Sets}_*, -1}$  is also natural. Thus  $\lambda^{\mathbf{Sets}_*}$  is a natural isomorphism.  $\square$

## 00GP 5.6 The Right Unitor

**00GQ Definition 5.6.1.1.** The **right unitor of the smash product of pointed sets** is the natural isomorphism

$$\rho^{\mathbf{Sets}_*} : \wedge \circ (\mathrm{id} \times \mathbb{1}^{\mathbf{Sets}_*}) \xrightarrow{\sim} \rho_{\mathbf{Sets}_*}^{\mathbf{Cats}_2},$$

whose component

$$\rho_X^{\mathbf{Sets}_*} : X \wedge S^0 \xrightarrow{\cong} X$$

at  $X \in \mathrm{Obj}(\mathbf{Sets}_*)$  is given by

$$\begin{aligned} x \wedge 0 &\mapsto x_0, \\ x \wedge 1 &\mapsto x. \end{aligned}$$

*Proof. Well-Definedness:* Let  $[(x, y)] = [(x', y')]$  be an element in  $X \wedge S^0$ . Then either:

1. We have  $x = x'$  and  $y = y'$ .
2. Both of the following conditions are satisfied:
  - (a) We have  $x = x_0$  or  $y = 0$ .
  - (b) We have  $x' = x_0$  or  $y' = 0$ .

In the first case,  $\rho_X^{\mathbf{Sets}_*}$  clearly sends both elements to the same element in  $X$ . Meanwhile, in the latter case both elements are equal to the basepoint  $x_0 \wedge 0$  of  $X \wedge S^0$ , which gets sent to the basepoint  $x_0$  of  $X$ .

*Being a Morphism of Pointed Sets:* As just mentioned, we have

$$\rho_X^{\mathbf{Sets}_*}(x_0 \wedge 0) \stackrel{\mathrm{def}}{=} x_0,$$

and thus  $\rho_X^{\mathbf{Sets}_*}$  is a morphism of pointed sets.

*Invertibility:* The inverse of  $\rho_X^{\mathbf{Sets}_*}$  is the morphism

$$\rho_X^{\mathbf{Sets}_*, -1} : X \xrightarrow{\cong} X \wedge S^0$$

defined by

$$\rho_X^{\mathbf{Sets}_*, -1}(x) \stackrel{\mathrm{def}}{=} x \wedge 1$$

for each  $x \in X$ . Indeed:

- *Invertibility I.* We have

$$\begin{aligned}
 [\rho_X^{\mathbf{Sets}_*, -1} \circ \rho_X^{\mathbf{Sets}_*}](x \wedge 0) &= \rho_X^{\mathbf{Sets}_*, -1}(\rho_X^{\mathbf{Sets}_*}(x \wedge 0)) \\
 &= \rho_X^{\mathbf{Sets}_*, -1}(x_0) \\
 &= x_0 \wedge 1 \\
 &= x \wedge 0,
 \end{aligned}$$

and

$$\begin{aligned}
 [\rho_X^{\mathbf{Sets}_*, -1} \circ \rho_X^{\mathbf{Sets}_*}](x \wedge 1) &= \rho_X^{\mathbf{Sets}_*, -1}(\rho_X^{\mathbf{Sets}_*}(x \wedge 1)) \\
 &= \rho_X^{\mathbf{Sets}_*, -1}(x) \\
 &= x \wedge 1
 \end{aligned}$$

for each  $x \in X$ , and thus we have

$$\rho_X^{\mathbf{Sets}_*, -1} \circ \rho_X^{\mathbf{Sets}_*} = \text{id}_{X \wedge S^0}.$$

- *Invertibility II.* We have

$$\begin{aligned}
 [\rho_X^{\mathbf{Sets}_*} \circ \rho_X^{\mathbf{Sets}_*, -1}](x) &= \rho_X^{\mathbf{Sets}_*}(\rho_X^{\mathbf{Sets}_*, -1}(x)) \\
 &= \rho_X^{\mathbf{Sets}_*, -1}(x \wedge 1) \\
 &= x
 \end{aligned}$$

for each  $x \in X$ , and thus we have

$$\rho_X^{\mathbf{Sets}_*} \circ \rho_X^{\mathbf{Sets}_*, -1} = \text{id}_X.$$

This shows  $\rho_X^{\mathbf{Sets}_*}$  to be invertible.

*Naturality:* We need to show that, given a morphism of pointed sets

$$f: (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc}
 X \wedge S^0 & \xrightarrow{f \wedge \text{id}_{S^0}} & Y \wedge S^0 \\
 \rho_X^{\mathbf{Sets}_*} \downarrow & & \downarrow \rho_Y^{\mathbf{Sets}_*} \\
 X & \xrightarrow{f} & Y
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x \wedge 0 & & x \wedge 0 \mapsto f(x) \wedge 0 \\ \downarrow & & \downarrow \\ x_0 \mapsto f(x_0) & & y_0 \end{array}$$

and

$$\begin{array}{ccc} x \wedge 1 \mapsto f(x) \wedge 1 & & \\ \downarrow & & \downarrow \\ x \mapsto f(x) & & \end{array}$$

and hence indeed commutes, showing  $\rho^{\mathbf{Sets}_*}$  to be a natural transformation. *Being a Natural Isomorphism:* Since  $\rho^{\mathbf{Sets}_*}$  is natural and  $\rho^{\mathbf{Sets}_*, -1}$  is a componentwise inverse to  $\rho^{\mathbf{Sets}_*}$ , it follows from **Categories, Item 2 of Proposition 8.6.1.2** that  $\rho^{\mathbf{Sets}_*, -1}$  is also natural. Thus  $\rho^{\mathbf{Sets}_*}$  is a natural isomorphism.  $\square$

## 00GR 5.7 The Symmetry

**00GS Definition 5.7.1.1.** The **symmetry of the smash product of pointed sets** is the natural isomorphism

$$\sigma^{\mathbf{Sets}_*} : \wedge \xrightarrow{\sim} \wedge \circ \sigma_{\mathbf{Sets}_*, \mathbf{Sets}_*}^{\mathbf{Cats}_2}, \quad \begin{array}{ccc} \mathbf{Sets}_* \times \mathbf{Sets}_* & \xrightarrow{\wedge} & \mathbf{Sets}_* \\ \sigma_{\mathbf{Sets}_*, \mathbf{Sets}_*}^{\mathbf{Cats}_2} \searrow & \sigma_{\mathbf{Sets}_*}^{\parallel} \downarrow & \nearrow \wedge \\ & \mathbf{Sets}_* \times \mathbf{Sets}_* & \end{array}$$

whose component

$$\sigma_{X,Y}^{\mathbf{Sets}_*} : X \wedge Y \xrightarrow{\cong} Y \wedge X$$

at  $X, Y \in \mathbf{Obj}(\mathbf{Sets}_*)$  is defined by

$$\sigma_{X,Y}^{\mathbf{Sets}_*}(x \wedge y) \stackrel{\text{def}}{=} y \wedge x$$

for each  $x \wedge y \in X \wedge Y$ .

*Proof. Well-Definedness:* Let  $[(x, y)] = [(x', y')]$  be an element in  $X \wedge Y$ . Then either:

1. We have  $x = x'$  and  $y = y'$ .
2. Both of the following conditions are satisfied:
  - (a) We have  $x = x_0$  or  $y = y_0$ .
  - (b) We have  $x' = x_0$  or  $y' = y_0$ .

In the first case,  $\sigma_X^{\mathbf{Sets}_*}$  clearly sends both elements to the same element in  $X$ . Meanwhile, in the latter case both elements are equal to the basepoint  $x_0 \wedge y_0$  of  $X \wedge Y$ , which gets sent to the basepoint  $y_0 \wedge x_0$  of  $Y \wedge X$ .

*Being a Morphism of Pointed Sets:* As just mentioned, we have

$$\sigma_X^{\mathbf{Sets}_*}(x_0 \wedge y_0) \stackrel{\text{def}}{=} y_0 \wedge x_0,$$

and thus  $\sigma_X^{\mathbf{Sets}_*}$  is a morphism of pointed sets.

*Invertibility:* Clearly, the inverse of  $\sigma_{X,Y}^{\mathbf{Sets}_*}$  is given by the morphism

$$\sigma_{X,Y}^{\mathbf{Sets}_*, -1}: Y \wedge X \xrightarrow{\cong} X \wedge Y$$

defined by

$$\sigma_{X,Y}^{\mathbf{Sets}_*, -1}(y \wedge x) \stackrel{\text{def}}{=} x \wedge y$$

for each  $y \wedge x \in Y \wedge X$ .

*Naturality:* We need to show that, given morphisms of pointed sets

$$\begin{aligned} f &: (X, x_0) \rightarrow (A, a_0), \\ g &: (Y, y_0) \rightarrow (B, b_0) \end{aligned}$$

the diagram

$$\begin{array}{ccc} X \wedge Y & \xrightarrow{f \wedge g} & A \wedge B \\ \sigma_{X,Y}^{\mathbf{Sets}_*} \downarrow & & \downarrow \sigma_{A,B}^{\mathbf{Sets}_*} \\ Y \wedge X & \xrightarrow{g \wedge f} & B \wedge A \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x \wedge y & \longmapsto & f(x) \wedge g(y) \\ \downarrow & & \downarrow \\ y \wedge x & \longmapsto & g(y) \wedge f(x) \end{array}$$



and hence indeed commutes, showing  $\sigma^{\mathbf{Sets}_*}$  to be a natural transformation. *Being a Natural Isomorphism:* Since  $\sigma^{\mathbf{Sets}_*}$  is natural and  $\sigma^{\mathbf{Sets}_*, -1}$  is a componentwise inverse to  $\sigma^{\mathbf{Sets}_*}$ , it follows from **Categories, Item 2 of Proposition 8.6.1.2** that  $\sigma^{\mathbf{Sets}_*, -1}$  is also natural. Thus  $\sigma^{\mathbf{Sets}_*}$  is a natural isomorphism.  $\square$

## 00GT 5.8 The Diagonal

00GU **Definition 5.8.1.1.** The **diagonal of the smash product of pointed sets** is the natural transformation

$$\Delta^\wedge : \mathrm{id}_{\mathbf{Sets}_*} \Longrightarrow \wedge \circ \Delta_{\mathbf{Sets}_*}^{\mathbf{Cats}_2},$$

whose component

$$\Delta_X^\wedge : (X, x_0) \rightarrow (X \wedge X, x_0 \wedge x_0)$$

at  $(X, x_0) \in \mathrm{Obj}(\mathbf{Sets}_*)$  is given by the composition

$$\begin{aligned} (X, x_0) &\xrightarrow{\Delta_X^\wedge} (X \times X, (x_0, x_0)) \\ &\longrightarrow ((X \times X)/\sim, [(x_0, x_0)]) \\ &\stackrel{\text{def}}{=} (X \wedge X, x_0 \wedge x_0) \end{aligned}$$

in  $\mathbf{Sets}_*$ , and thus by

$$\Delta_X^\wedge(x) \stackrel{\text{def}}{=} x \wedge x$$

for each  $x \in X$ .

*Proof. Being a Morphism of Pointed Sets:* We have

$$\Delta_X^\wedge(x_0) \stackrel{\text{def}}{=} x_0 \wedge x_0,$$

and thus  $\Delta_X^\wedge$  is a morphism of pointed sets.

*Naturality:* We need to show that, given a morphism of pointed sets

$$f : (X, x_0) \rightarrow (Y, y_0),$$

the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \Delta_X^\wedge \downarrow & & \downarrow \Delta_Y^\wedge \\ X \wedge X & \xrightarrow{f \wedge f} & Y \wedge Y \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x & \mapsto & f(x) \\ \downarrow & & \downarrow \\ x \wedge x & \mapsto & f(x) \wedge f(x) \end{array}$$

and hence indeed commutes, showing  $\Delta^\wedge$  to be natural.  $\square$

**00GV Proposition 5.8.1.2.** Let  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ .

**00GW** 1. *Monoidality.* The diagonal

$$\Delta^\wedge : \text{id}_{\text{Sets}_*} \Longrightarrow \wedge \circ \Delta_{\text{Sets}_*}^{\text{Cats}_2},$$

of the smash product of pointed sets is a monoidal natural transformation:

**00GX** (a) *Compatibility With Strong Monoidality Constraints.* For each  $(X, x_0), (Y, y_0) \in \text{Obj}(\text{Sets}_*)$ , the diagram

$$\begin{array}{ccc} X \wedge Y & \xrightarrow{\Delta_X^\wedge \wedge \Delta_Y^\wedge} & (X \wedge X) \wedge (Y \wedge Y) \\ & \searrow \Delta_{X \wedge Y}^\wedge & \downarrow \wr \\ & & (X \wedge Y) \wedge (X \wedge Y) \end{array}$$

commutes.

**00GY** (b) *Compatibility With Strong Unitality Constraints.* The diagrams

$$\begin{array}{ccc} S^0 & \xrightarrow{\Delta_{S^0}^\wedge} & S^0 \wedge S^0 \\ \parallel & \downarrow \lambda_{S^0}^{\text{Sets}_*} & \\ & S^0 & \end{array} \quad \begin{array}{ccc} S^0 & \xrightarrow{\Delta_{S^0}^\wedge} & S^0 \wedge S^0 \\ \parallel & \downarrow \rho_{S^0}^{\text{Sets}_*} & \\ & S^0 & \end{array}$$

commute, i.e. we have

$$\begin{aligned}\Delta_{S^0}^\wedge &= \lambda_{S^0}^{\mathbf{Sets}_*, -1} \\ &= \rho_{S^0}^{\mathbf{Sets}_*, -1},\end{aligned}$$

where we recall that the equalities

$$\begin{aligned}\lambda_{S^0}^{\mathbf{Sets}_*} &= \rho_{S^0}^{\mathbf{Sets}_*}, \\ \lambda_{S^0}^{\mathbf{Sets}_*, -1} &= \rho_{S^0}^{\mathbf{Sets}_*, -1}\end{aligned}$$

are always true in any monoidal category by ??, ?? of ??.

**00GZ** 2. *The Diagonal of the Unit.* The component

$$\Delta_{S^0}^\wedge : S^0 \xrightarrow{\cong} S^0 \wedge S^0$$

of  $\Delta^\wedge$  at  $S^0$  is an isomorphism.

*Proof. Item 1, Monoidality:* We claim that  $\Delta^\wedge$  is indeed monoidal:

1. *Item 1a: Compatibility With Strong Monoidality Constraints:* We need to show that the diagram

$$\begin{array}{ccc} X \wedge Y & \xrightarrow{\Delta_X^\wedge \wedge \Delta_Y^\wedge} & (X \wedge X) \wedge (Y \wedge Y) \\ & \searrow \Delta_{X \wedge Y}^\wedge & \downarrow \wr \\ & & (X \wedge Y) \wedge (X \wedge Y) \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc} x \wedge y & \xrightarrow{\quad} & (x \wedge x) \wedge (y \wedge y) \\ & \searrow & \downarrow \\ & & (x \wedge y) \wedge (x \wedge y) \end{array}$$

and hence indeed commutes.

2. *Item 1b: Compatibility With Strong Unitality Constraints:* As shown in the proof of **Definition 5.5.1.1**, the inverse of the left unitor of  $\mathbf{Sets}_*$  with

respect to the smash product of pointed sets at  $(X, x_0) \in \text{Obj}(\text{Sets}_*)$  is given by

$$\lambda_X^{\text{Sets}_*, -1}(x) \stackrel{\text{def}}{=} 1 \wedge x$$

for each  $x \in X$ , so when  $X = S^0$ , we have

$$\lambda_{S^0}^{\text{Sets}_*, -1}(0) \stackrel{\text{def}}{=} 1 \wedge 0,$$

$$\lambda_{S^0}^{\text{Sets}_*, -1}(1) \stackrel{\text{def}}{=} 1 \wedge 1.$$

But since  $1 \wedge 0 = 0 \wedge 0$  and

$$\Delta_{S^0}^\wedge(0) \stackrel{\text{def}}{=} 0 \wedge 0,$$

$$\Delta_{S^0}^\wedge(1) \stackrel{\text{def}}{=} 1 \wedge 1,$$

it follows that we indeed have  $\Delta_{S^0}^\wedge = \lambda_{S^0}^{\text{Sets}_*, -1}$ .

This finishes the proof.

*Item 2, The Diagonal of the Unit:* This follows from *Item 1* and the invertibility of the left/right unitor of  $\text{Sets}_*$  with respect to  $\wedge$ , proved in the proof of *Definition 5.5.1.1* for the left unitor or the proof of *Definition 5.6.1.1* for the right unitor.  $\square$

## 00H0 5.9 The Monoidal Structure on Pointed Sets Associated to $\wedge$

**00H1 Proposition 5.9.1.1.** The category  $\text{Sets}_*$  admits a closed monoidal category with diagonals structure consisting of

- *The Underlying Category.* The category  $\text{Sets}_*$  of pointed sets;
- *The Monoidal Product.* The smash product functor

$$\wedge : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

of *Item 1* of *Proposition 5.1.1.9*;

- *The Internal Hom.* The internal Hom functor

$$\mathbf{Sets}_* : \text{Sets}_*^{\text{op}} \times \text{Sets}_* \rightarrow \text{Sets}_*$$

of *Item 1* of *Proposition 5.2.1.2*;

- *The Monoidal Unit.* The functor

$$\mathbb{1}^{\text{Sets}_*} : \text{pt} \rightarrow \text{Sets}_*$$

of *Definition 5.3.1.1*;

- *The Associators.* The natural isomorphism

$$\alpha^{\mathbf{Sets}_*} : \wedge \circ (\wedge \times \mathrm{id}_{\mathbf{Sets}_*}) \xrightarrow{\sim} \wedge \circ (\mathrm{id}_{\mathbf{Sets}_*} \times \wedge) \circ \alpha_{\mathbf{Sets}_*, \mathbf{Sets}_*, \mathbf{Sets}_*}^{\mathbf{Cats}}$$

of Definition 5.4.1.1;

- *The Left Unitors.* The natural isomorphism

$$\lambda^{\mathbf{Sets}_*} : \wedge \circ (\mathbb{1}^{\mathbf{Sets}_*} \times \mathrm{id}_{\mathbf{Sets}_*}) \xrightarrow{\sim} \lambda_{\mathbf{Sets}_*}^{\mathbf{Cats}_2}$$

of Definition 5.5.1.1;

- *The Right Unitors.* The natural isomorphism

$$\rho^{\mathbf{Sets}_*} : \wedge \circ (\mathrm{id} \times \mathbb{1}^{\mathbf{Sets}_*}) \xrightarrow{\sim} \rho_{\mathbf{Sets}_*}^{\mathbf{Cats}_2}$$

of Definition 5.6.1.1;

- *The Symmetry.* The natural isomorphism

$$\sigma^{\mathbf{Sets}_*} : \wedge \xrightarrow{\sim} \wedge \circ \sigma_{\mathbf{Sets}_*, \mathbf{Sets}_*}^{\mathbf{Cats}_2}$$

of Definition 5.7.1.1;

- *The Diagonals.* The monoidal natural transformation

$$\Delta^\wedge : \mathrm{id}_{\mathbf{Sets}_*} \Longrightarrow \wedge \circ \Delta_{\mathbf{Sets}_*}^{\mathbf{Cats}_2}$$

of Definition 5.8.1.1.

*Proof. The Pentagon Identity:* Let  $(W, w_0)$ ,  $(X, x_0)$ ,  $(Y, y_0)$  and  $(Z, z_0)$  be

pointed sets. We have to show that the diagram

$$\begin{array}{ccc}
 & (W \wedge (X \wedge Y)) \wedge Z & \\
 \alpha_{W,X,Y}^{\text{Sets}*} \wedge \text{id}_Z \nearrow & & \searrow \alpha_{W,X \wedge Y,Z}^{\text{Sets}*} \\
 ((W \wedge X) \wedge Y) \wedge Z & & W \wedge ((X \wedge Y) \wedge Z) \\
 \alpha_{W \wedge X,Y,Z}^{\text{Sets}*} \searrow & & \swarrow \text{id}_W \wedge \alpha_{X,Y,Z}^{\text{Sets}*} \\
 (W \wedge X) \wedge (Y \wedge Z) & \xrightarrow{\alpha_{W,X,Y \wedge Z}^{\text{Sets}*}} & W \wedge (X \wedge (Y \wedge Z))
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 & (w \wedge (x \wedge y)) \wedge z & \\
 \swarrow & & \searrow \\
 ((w \wedge x) \wedge y) \wedge z & & w \wedge ((x \wedge y) \wedge z) \\
 \swarrow & & \searrow \\
 (w \wedge x) \wedge (y \wedge z) & \longmapsto & w \wedge (x \wedge (y \wedge z))
 \end{array}$$

and thus we see that the pentagon identity is satisfied.

*The Triangle Identity:* Let  $(X, x_0)$  and  $(Y, y_0)$  be pointed sets. We have to

show that the diagram

$$\begin{array}{ccc}
 (X \wedge S^0) \wedge Y & \xrightarrow{\alpha_{X, S^0, Y}^{\text{Sets}_*}} & X \wedge (S^0 \wedge Y) \\
 \searrow \rho_X^{\text{Sets}_*} \wedge \text{id}_Y & & \swarrow \text{id}_X \wedge \lambda_Y^{\text{Sets}_*} \\
 & X \wedge Y &
 \end{array}$$

commutes. Indeed, this diagram acts on elements as

$$\begin{array}{ccc}
 (x \wedge 0) \wedge y & \xrightarrow{\quad} & x \wedge (0 \wedge y) \\
 \searrow & & \swarrow \\
 x_0 \wedge y & & x \wedge y_0
 \end{array}$$

and

$$\begin{array}{ccc}
 (x \wedge 1) \wedge y & \xrightarrow{\quad} & x \wedge (1 \wedge y) \\
 \searrow & & \swarrow \\
 & x \wedge y &
 \end{array}$$


and thus we see that the triangle identity is satisfied.

*The Left Hexagon Identity:* Let  $(X, x_0)$ ,  $(Y, y_0)$ , and  $(Z, z_0)$  be pointed sets.

We have to show that the diagram


$$\begin{array}{ccc}
 & (X \wedge Y) \wedge Z & \\
 \alpha_{X, Y, Z}^{\text{Sets}_*} \swarrow & & \searrow \beta_{X, Y}^{\text{Sets}_*} \wedge \text{id}_Z \\
 X \wedge (Y \wedge Z) & & (Y \wedge X) \wedge Z \\
 \downarrow \beta_{X, Y \wedge Z}^{\text{Sets}_*} & & \downarrow \alpha_{Y, X, Z}^{\text{Sets}_*} \\
 (Y \wedge Z) \wedge X & & Y \wedge (X \wedge Z) \\
 \alpha_{Y, Z, X}^{\text{Sets}_*} \swarrow & & \swarrow \text{id}_Y \wedge \beta_{X, Z}^{\text{Sets}_*} \\
 & Y \wedge (Z \wedge X) &
 \end{array}$$

commutes. Indeed, this diagram acts on elements as




and thus we see that the left hexagon identity is satisfied.

*The Right Hexagon Identity:* Let  $(X, x_0)$ ,  $(Y, y_0)$ , and  $(Z, z_0)$  be pointed sets. We have to show that the diagram





commutes. Indeed, this diagram acts on elements as



and thus we see that the right hexagon identity is satisfied.

*Monoidal Closedness:* This follows from **Item 2** of **Proposition 5.1.1.9**.

*Existence of Monoidal Diagonals:* This follows from **Items 1** and **2** of **Proposition 5.8.1.2**. □

## 00H2 5.10 Universal Properties of the Smash Product of Pointed Sets I

00H3 **Theorem 5.10.1.1.** The symmetric monoidal structure on the category  $\mathbf{Sets}_*$  is uniquely determined by the following requirements:

1. *Two-Sided Preservation of Colimits.* The smash product

$$\wedge : \mathbf{Sets}_* \times \mathbf{Sets}_* \rightarrow \mathbf{Sets}_*$$

of  $\mathbf{Sets}_*$  preserves colimits separately in each variable.

2. *The Unit Object Is  $S^0$ .* We have  $\mathbb{1}_{\mathbf{Sets}_*} = S^0$ .

*Proof.* Omitted. □

## 00H4 5.11 Universal Properties of the Smash Product of Pointed Sets II

00H5 **Theorem 5.11.1.1.** The symmetric monoidal structure on the category  $\mathbf{Sets}_*$  is the unique symmetric monoidal structure on  $\mathbf{Sets}_*$  such that the free pointed set functor

$$(-)^+ : \mathbf{Sets} \rightarrow \mathbf{Sets}_*$$

admits a symmetric monoidal structure.

*Proof.* See [GGN15, Theorem 5.1].  $\square$

## 00H6 5.12 Monoids With Respect to the Smash Product of Pointed Sets

00H7 **Proposition 5.12.1.1.** The category of monoids on  $(\mathbf{Sets}_*, \wedge, S^0)$  is isomorphic to the category of monoids with zero and morphisms between them.

*Proof.* See ??, in particular ??, ??, ??, and ??.  $\square$

## 00H8 5.13 Comonoids With Respect to the Smash Product of Pointed Sets

00H9 **Proposition 5.13.1.1.** The symmetric monoidal functor

$$\left((-)^+, (-)^{+, \times}, (-)_{\mathbb{1}}^{+, \times}\right): (\mathbf{Sets}, \times, \text{pt}) \rightarrow (\mathbf{Sets}_*, \wedge, S^0),$$

of **Pointed Sets**, **Item 4** of **Proposition 4.1.1.2** lifts to an equivalence of categories

$$\begin{aligned} \text{CoMon}(\mathbf{Sets}_*, \wedge, S^0) &\stackrel{\text{eq.}}{\cong} \text{CoMon}(\mathbf{Sets}, \times, \text{pt}) \\ &\cong \mathbf{Sets}. \end{aligned}$$

*Proof.* See [PS19, Lemma 2.4].  $\square$

## 00HA 6 Miscellany

### 00HB 6.1 The Smash Product of a Family of Pointed Sets

Let  $\{(X_i, x_0^i)\}_{i \in I}$  be a family of pointed sets.

00HC **Definition 6.1.1.1.** The **smash product of the family**  $\{(X_i, x_0^i)\}_{i \in I}$  is the pointed set  $\bigwedge_{i \in I} X_i$  consisting of:

- *The Underlying Set.* The set  $\bigwedge_{i \in I} X_i$  defined by

$$\bigwedge_{i \in I} X_i \stackrel{\text{def}}{=} \left( \prod_{i \in I} X_i \right) / \sim,$$

where  $\sim$  is the equivalence relation on  $\prod_{i \in I} X_i$  obtained by declaring

$$(x_i)_{i \in I} \sim (y_i)_{i \in I}$$

if there exist  $i_0 \in I$  such that  $x_{i_0} = x_0$  and  $y_{i_0} = y_0$ , for each  $(x_i)_{i \in I}, (y_i)_{i \in I} \in \prod_{i \in I} X_i$ .

- *The Basepoint.* The element  $[(x_0)_{i \in I}]$  of  $\bigwedge_{i \in I} X_i$ .

## Appendices

### A Other Chapters

#### Sets

1. [Sets](#)
2. [Constructions With Sets](#)
3. [Pointed Sets](#)
4. [Tensor Products of Pointed Sets](#)

#### 6. [Constructions With Relations](#)

#### 7. [Equivalence Relations and Apartness Relations](#)

#### Category Theory

#### 8. [Categories](#)

#### Bicategories

#### Relations

#### 5. [Relations](#)

#### 9. [Types of Morphisms in Bicat-](#) [egories](#)

### References

- [GGN15] David Gepner, Moritz Groth, and Thomas Nikolaus. “Universality of Multiplicative Infinite Loop Space Machines”. In: *Algebr. Geom. Topol.* 15.6 (2015), pp. 3107–3153. ISSN: 1472-2747. DOI: [10.2140/agt.2015.15.3107](https://doi.org/10.2140/agt.2015.15.3107). URL: <https://doi.org/10.2140/agt.2015.15.3107> (cit. on p. 98).
- [PS19] Maximilien Péroux and Brooke Shipley. “Coalgebras in Symmetric Monoidal Categories of Spectra”. In: *Homology Homotopy Appl.* 21.1 (2019), pp. 1–18. ISSN: 1532-0073. DOI: [10.4310/HHA.2019.v21.n1.a1](https://doi.org/10.4310/HHA.2019.v21.n1.a1). URL: <https://doi.org/10.4310/HHA.2019.v21.n1.a1> (cit. on p. 98).