

Internal Adjunctions

December 3, 2023

Create tags:

1. <https://www.google.com/search?q=mate+of+an+adjunction>
2. Moreover, by uniqueness of adjoints ([Internal Adjunctions](#), [Item 2 of Proposition 1.2.1.4](#)), this implies also that $S = f^{-1}$.
3. define bicategory $\text{Adj}(C)$
4. walking monad
5. proposition: 2-functors preserve unitors and associators
6. <https://ncatlab.org/nlab/show/2-category+of+adjunctions>. Is there a 3-category too?
7. <https://ncatlab.org/nlab/show/free+monad>
8. <https://ncatlab.org/nlab/show/CatAdj>
9. <https://ncatlab.org/nlab/show/Adj>
10. $\text{Adj}(\text{Adj}(C))$

Contents

1	Internal Adjunctions	2
1.1	The Walking Adjunction	2
1.2	Internal Adjunctions	4
1.3	Internal Adjoint Equivalences	10
1.4	Mates	12

2	Morphisms of Internal Adjunctions	15
2.1	Lax Morphisms of Internal Adjunctions	15
2.2	Oplax Morphisms of Internal Adjunctions	16
2.3	Strong Morphisms of Internal Adjunctions	17
2.4	Strict Morphisms of Internal Adjunctions	18
3	2-Morphisms Between Morphisms of Internal Adjunctions	18
3.1	2-Morphisms Between Lax Morphisms of Internal Adjunctions	18
3.2	2-Morphisms Between Oplax Morphisms of Internal Adjunctions	19
3.3	2-Morphisms Between Strong Morphisms of Internal Adjunctions	20
3.4	2-Morphisms Between Strict Morphisms of Internal Adjunctions	20
4	Bicategories of Internal Adjunctions in a Bicategory	21
A	Other Chapters	21

1 Internal Adjunctions

1.1 The Walking Adjunction

Definition 1.1.1.1. The **walking adjunction** is the bicategory Adj freely generated by¹

- *Objects.* A pair of objects A and B ;
- *Morphisms.* A pair of morphisms

$$L: A \rightarrow B,$$

$$R: B \rightarrow A;$$

- *2-Morphisms.* A pair of 2-morphisms

$$\eta: \text{id}_A \rightarrow L \circ R,$$

$$\epsilon: R \circ L \rightarrow \text{id}_B;$$

¹See [SS86] for an explicit description of the 2-category (as opposed to a bicategory) version of Adj in terms of finite ordinals, similar to the description of the 2-category version of the walking monad (??) as a subcategory of Δ .

subject to the equalities

$$\begin{array}{ccc}
 \begin{array}{c} B \xrightarrow{id_B} B \\ \swarrow L \quad \searrow R \\ A \xrightarrow{id_A} A \end{array} & \begin{array}{c} \uparrow \eta \\ \parallel \\ \downarrow \epsilon \end{array} & \begin{array}{c} B \xrightarrow{id_B} B \\ \swarrow L \quad \searrow L \\ A \xrightarrow{id_A} A \end{array} \\
 & = & \\
 \begin{array}{c} B \xrightarrow{id_B} B \\ \swarrow R \quad \searrow L \\ A \xrightarrow{id_A} A \end{array} & \begin{array}{c} \uparrow \epsilon \\ \parallel \\ \downarrow \eta \end{array} & \begin{array}{c} B \xrightarrow{id_B} B \\ \swarrow R \quad \searrow R \\ A \xrightarrow{id_A} A \end{array} \\
 & = &
 \end{array}$$

of pasting diagrams, which are equivalent to the following conditions:

1. *The Left Triangle Identity.* The diagram

$$\begin{array}{ccc}
 L \circ id_A & \xrightarrow{id_L \circ \eta} & L \circ (R \circ L) \xrightarrow{(\alpha_{L,R,L}^{Adj})^{-1}} (L \circ R) \circ L \\
 & \searrow \rho_L^{Adj} & \downarrow \epsilon \circ id_L \\
 & & id_B \circ L \\
 & & \downarrow \lambda_L^{Adj} \\
 & & L
 \end{array}$$

commutes.

2. *The Right Triangle Identity.* The diagram

$$\begin{array}{ccc}
 id_A \circ R & \xrightarrow{\eta \circ id_R} & (R \circ L) \circ R \xrightarrow{\alpha_{R,L,R}^{Adj}} R \circ (L \circ R) \\
 & \searrow \lambda_R^{Adj} & \downarrow id_R \circ \epsilon \\
 & & R \circ id_B \\
 & & \downarrow \rho_R^{Adj} \\
 & & R.
 \end{array}$$

1.2 Internal Adjunctions

Let C be a bicategory.

Definition 1.2.1.1. An **internal adjunction in $C^{2,3}$** is a 2-functor $\text{Adj} \rightarrow C$.

Remark 1.2.1.2. In detail, an **internal adjunction in C** consists of

- *Objects.* A pair of objects A and B of C ;
- *Morphisms.* A pair of morphisms

$$\begin{aligned} L &: A \rightarrow B, \\ R &: B \rightarrow A \end{aligned}$$

of C ;

- *2-Morphisms.* A pair of 2-morphisms

$$\begin{aligned} \eta &: \text{id}_A \rightarrow L \circ R, \\ \epsilon &: R \circ L \rightarrow \text{id}_B \end{aligned}$$

of C ;

subject to the equalities

²Further Terminology: Also called an **adjunction internal to C** .

³Further Terminology: In this situation, we also call (g, f) an **adjoint pair**, f the **left adjoint** of the pair, g the **right adjoint** of the pair, η the **unit** of the adjunction, and ϵ the **counit** of the adjunction.

of pasting diagrams in C , which are equivalent to the following conditions:⁴

1. *The Left Triangle Identity.* The diagram

$$\begin{array}{ccccc}
 L \circ \text{id}_A & \xrightarrow{\text{id}_L \circ \eta} & L \circ (R \circ L) & \xrightarrow{(\alpha_{L,R,L}^C)^{-1}} & (L \circ R) \circ L \\
 & \searrow \rho_L^C & & & \downarrow \epsilon \circ \text{id}_L \\
 & & & & \text{id}_B \circ L \\
 & & & & \downarrow \lambda_L^C \\
 & & & & L
 \end{array}$$

commutes.

2. *The Right Triangle Identity.* The diagram

$$\begin{array}{ccccc}
 \text{id}_A \circ R & \xrightarrow{\eta \circ \text{id}_R} & (R \circ L) \circ R & \xrightarrow{\alpha_{R,L,R}^C} & R \circ (L \circ R) \\
 & \searrow \lambda_R^C & & & \downarrow \text{id}_R \circ \epsilon \\
 & & & & R \circ \text{id}_B \\
 & & & & \downarrow \rho_R^C \\
 & & & & R.
 \end{array}$$

Example 1.2.1.3. Here are some examples of internal adjunctions.

1. *Internal Adjunctions in Cats_2 .* The internal adjunctions in the 2-category Cats_2 of categories, functors, and natural transformations are precisely the adjunctions of **Categories, ??**.

⁴When C is a 2-category, these diagrams take the following form:

$$\begin{array}{ccc}
 L & \xrightarrow{\text{id}_L \circ \eta} & L \circ R \circ L \\
 \searrow \text{id}_L & & \downarrow \epsilon \circ \text{id}_L \\
 & & L
 \end{array}
 \qquad
 \begin{array}{ccc}
 R & \xrightarrow{\text{id}_R \circ \eta} & R \circ L \circ R \\
 \searrow \text{id}_L & & \downarrow \epsilon \circ \text{id}_R \\
 & & R.
 \end{array}$$

2. *Internal Adjunctions in **Rel***. The internal adjunctions in **Rel** are precisely the relations of the form $\text{Gr}(f) \dashv f^{-1}$ with f a function; see [Relations, Item 4](#) of [Proposition 2.5.1.1](#).
3. *Internal Adjunctions in Span*. The internal adjunctions in Span are precisely the spans of the form

$$\begin{array}{ccc} & S & \\ \phi \swarrow & & \searrow g \\ A & & B \end{array}$$

with ϕ an isomorphism; see [Spans, Item 4](#) of [Proposition 2.5.1.1](#).

Proposition 1.2.1.4. Let C be a bicategory.

1. *Duality*. Let (f, g, η, ϵ) be an internal adjunction in C .
 - (a) The quadruple (g, f, η, ϵ) is an internal adjunction in C^{op} .
 - (b) The quadruple (g, f, ϵ, η) is an internal adjunction in C^{co} .
 - (c) The quadruple (f, g, η, ϵ) is an internal adjunction in C^{coop} .
2. *Uniqueness of Adjoints*. Let (f, g, η, ϵ) and $(f, g', \eta', \epsilon')$ be internal adjunctions in C . We have a canonical isomorphism⁵

$$g \xrightarrow{(\lambda_g^C)^{-1}} \text{id}_A \circ g \xrightarrow{\eta' \circ \text{id}_g} (g' \circ f) \circ g \xrightarrow{\alpha_{g', f, g}^C} g' \circ (f \circ g) \xrightarrow{\text{id}_{g'} \circ \epsilon} g' \circ \text{id}_B \xrightarrow{(\rho_{g'}^C)^{-1}} g'$$

with inverse


$$g' \xrightarrow{(\lambda_{g'}^C)^{-1}} \text{id}_B \circ g' \xrightarrow{\eta \circ \text{id}_{g'}} (g \circ f) \circ g' \xrightarrow{\alpha_{g, f, g'}^C} g \circ (f \circ g') \xrightarrow{\text{id}_g \circ \epsilon'} g \circ \text{id}_B \xrightarrow{(\lambda_g^C)^{-1}} g.$$

3. *Carrying Internal Adjunctions Through Pseudofunctors*. Let $F: C \rightarrow \mathcal{D}$ be a pseudofunctor and (f, g, η, ϵ) be an internal adjunction in C . There is an induced internal adjunction⁶

$$(F(f), F(g), \bar{\eta}, \bar{\epsilon})$$

in \mathcal{D} , where:

⁵*Slogan*: Left adjoints are unique up to canonical isomorphism. Dually, so are right adjoints.

⁶ *Warning*: Lax or oplax functors which are not pseudofunctors need not preserve internal adjunctions.

(a) The unit

$$\bar{\eta}: \text{id}_{F(A)} \Longrightarrow F(g) \circ F(f)$$

is the composition


$$\text{id}_{F(A)} \xrightarrow{F_A} F(\text{id}_A) \xrightarrow{F(\eta)} F(g \circ f) \xrightarrow{F_{g,f}^{-1}} F(g) \circ F(f).$$

(b) The counit

$$\bar{\epsilon}: F(f) \circ F(g) \Longrightarrow \text{id}_{F(B)}$$

is the composition

$$F(f) \circ F(g) \xrightarrow{F_{f,g}} F(f \circ g) \xrightarrow{F(\epsilon)} F(\text{id}_B) \xrightarrow{F_B} \text{id}_{F(B)}.$$

Proof. **Item 1, Duality:** Omitted.⁷**Item 2, Uniqueness of Adjoints:** ⁸ Consider the diagram (if you *really* want to consider it I fear you will need to zoom in)

In this diagram:

1. The morphisms in **green** are the composition $g \xRightarrow{\quad} g' \xRightarrow{\quad} g$;

⁷Reference: [Y21, Exercise 6.6.2].⁸Reference: [Y21, Lemma 6.1.6].

2. The morphisms in **red** are equal to λ_g^C by the right triangle identity for (f, g, η, ϵ) . Hence the composition of the morphism in **blue** with the morphisms in **red** is the identity;
3. Subdiagrams (1), (2), (10), (11), (29), (31), and (43) commute by the naturality of the left unitor of C and its inverse;
4. Subdiagrams (8), (19), and (21) commute by the naturality of the right unitor of C and its inverse;
5. Subdiagrams (6), (13), (17), (18), (20), (22), (32), (33), (36), (38), (40), (41), and (45) commute by the naturality of the associator of C and its inverse;
6. Subdiagrams (37), (39), and (42) commute by the pentagon identity for C ;
7. Subdiagrams (3), (4), (7), (12), (25), (30), and (48) commute by **Bicategories**, ?? of ??;
8. Subdiagrams (5), (14), (23), (24), (34), and (35) commute by middle-four exchange;
9. Subdiagrams (9), (15), (16), (27), (28), (44), (46), (49), and (50) commute trivially;
10. Subdiagram (26) commutes by **Bicategories**, ??? of ??;
11. Subdiagram (47) commutes by **Bicategories**, ?? of ?? and the naturality of the left unitor of right unitor of C .


Hence $g \cong g'$.

Item 3, Carrying Internal Adjunctions Through Pseudofunctors:⁹ We claim that the left and right triangle identities for $(F(f), F(g), \bar{\eta}, \bar{\epsilon})$ hold:

1. The left triangle identity for $(F(f), F(g), \bar{\eta}, \bar{\epsilon})$ is the condition that the boundary

⁹Reference: [Y21, Proposition 6.1.7].

diagram of the diagram (you may need to zoom in)




commutes. Since

- (a) Subdiagrams (1) and (7) commute by applying middle-four exchange twice,
- (b) Subdiagrams (2) and (8) commute by the left and right lax unity conditions for F ,
- (c) Subdiagrams (3) and (6) commute by the naturality of the lax functoriality constraints of F ,
- (d) Subdiagram (4) commutes by the lax associativity condition for F , and
- (e) Subdiagram (5) commutes by the left triangle identity for (f, g, η, ϵ) ,

so does the boundary diagram.

2. The right triangle identity for $(F(f), F(g), \bar{\eta}, \bar{\epsilon})$ is the condition that the boundary

diagram of the diagram (you may need to zoom in)



commutes. Since

- (a) Subdiagrams (1) and (7) commute by applying middle-four exchange twice,
- (b) Subdiagrams (2) and (8) commute by the left and right lax unity conditions for F ,
- (c) Subdiagrams (3) and (6) commute by the naturality of the lax functoriality constraints of F ,
- (d) Subdiagram (4) commutes by the lax associativity condition for F , and
- (e) Subdiagram (5) commutes by the right triangle identity for (f, g, η, ϵ) ,

so does the boundary diagram.

This finishes the proof. \square

1.3 Internal Adjoint Equivalences

Let \mathcal{C} be a bicategory.

Definition 1.3.1.1. An internal adjunction (f, g, η, ϵ) in \mathcal{C} is an **internal adjoint equivalence** if η and ϵ are isomorphisms in \mathcal{C} .

Example 1.3.1.2. Here are some examples of internal adjoint equivalences.

1. *Internal Adjoint Equivalences in \mathbf{Cats}_2 .* The internal adjoint equivalences in the 2-category \mathbf{Cats}_2 of categories, functors, and natural transformations are precisely the adjoint equivalences of **Categories**, ??.¹⁰
2. *Internal Adjoint Equivalences in \mathbf{Mod} .* The internal adjoint equivalences in \mathbf{Mod} are precisely the invertible R -modules; see ??.¹¹
3. *Internal Adjoint Equivalences in $\mathbf{PseudoFun}(C, \mathcal{D})$.* The internal adjoint equivalences in $\mathbf{PseudoFun}(C, \mathcal{D})$ are precisely the invertible strong transformations; see ??.¹²
4. *Internal Adjoint Equivalences in \mathbf{Rel} .* The internal adjoint equivalences in \mathbf{Rel} are precisely the relations of the form $\text{Gr}(f) \dashv f^{-1}$ with f an isomorphism; see ??.
5. *Internal Adjoint Equivalences in \mathbf{Span} .* The internal adjoint equivalences in \mathbf{Span} are precisely the spans of the form $A \xleftarrow{\phi} S \xrightarrow{\psi} B$ with ϕ and ψ isomorphisms; see ??.

Proposition 1.3.1.3. Let C be a bicategory.

1. *Carrying Internal Adjoint Equivalences Through Pseudofunctors.* Let $F: C \longrightarrow \mathcal{D}$ be a pseudofunctor and (f, g, η, ϵ) be an internal adjunction in C . If (f, g, η, ϵ) is an internal adjoint equivalence in C , then the induced internal adjunction

$$(F(f), F(g), \bar{\eta}, \bar{\epsilon})$$

in \mathcal{D} of **Item 3** of **Proposition 1.2.1.4** is an internal adjoint equivalence as well.

2. *Internal Adjunctions Always Refine to Internal Adjoint Equivalences.* Let (f, g, η, ϵ) be an internal adjunction in C . If f is an equivalence, then there exist 2-morphisms

$$\bar{\eta}: \text{id}_A \Longrightarrow g \circ f$$

$$\bar{\epsilon}: f \circ g \Longrightarrow \text{id}_B$$

of C such that $(f, g, \bar{\eta}, \bar{\epsilon})$ is an internal adjoint equivalence.

Proof. **Item 1**, *Carrying Internal Adjoint Equivalences Through Pseudofunctors*: See [Y21, Proposition 6.2.3].

Item 2, *Internal Adjunctions Always Refine to Internal Adjoint Equivalences*: See [Y21, Proposition 6.2.4]. □

¹⁰Reference: [Y21, Examples 6.2.5].

¹¹Reference: [Y21, Examples 6.2.6].

¹²Reference: [Y21, Examples 6.2.7].

1.4 Mates

Let \mathcal{C} be a bicategory, let (f, g, η, ϵ) and $(f', g', \eta', \epsilon')$ be adjunctions, and let h and k be morphisms of \mathcal{C} as in the diagram

$$\begin{array}{ccc}
 A & \begin{array}{c} \xrightarrow{f} \\ \perp \\ \xleftarrow{g} \end{array} & B \\
 h \downarrow & & \downarrow k \\
 C & \begin{array}{c} \xrightarrow{f'} \\ \perp \\ \xleftarrow{g'} \end{array} & D.
 \end{array}$$


Definition 1.4.1.1. The **mates** of a pair of 2-morphisms

$$\begin{array}{ccc}
 A \xrightarrow{f} B & & A \xleftarrow{g} B \\
 h \downarrow \quad \nearrow \omega & \omega : f' \circ h \Rightarrow k \circ f, & \Downarrow \nu \\
 C \xrightarrow{f'} D & \nu : h \circ g \Rightarrow g' \circ k & C \xleftarrow{g'} D
 \end{array}$$

are the 2-morphisms

$$\begin{array}{ccc}
 A \xleftarrow{g} B & & A \xrightarrow{f} B \\
 h \downarrow \quad \Downarrow \omega^\dagger & \omega^\dagger : h \circ g \Rightarrow g' \circ k, & \Downarrow \nu^\dagger \\
 C \xleftarrow{g'} D & \nu^\dagger : f' \circ h \Rightarrow k \circ f & C \xrightarrow{f'} D
 \end{array}$$

defined as the pastings of the diagrams¹³



Proposition 1.4.1.2. Let $\omega: f' \circ h \Rightarrow k \circ f$ and $\nu: h \circ g \Rightarrow g' \circ k$ be 2-morphisms.

1. *The Mate Correspondence.* The map


$$(-)^\dagger: \text{Hom}_{\text{Hom}_C(A,C)}(f' \circ h, k \circ f) \longrightarrow \text{Hom}_{\text{Hom}_C(B,D)}(h \circ g, g' \circ k)$$

$$\omega \longmapsto \omega^\dagger$$

is a bijection.

Proof. **Item 1, The Mate Correspondence:** Here we give a proof for 2-categories (which indirectly proves also the general case by **Bicategories**, ??). A proof for general bicategories can be found in [Y21, Lemma 6.1.13].

¹³If C is a 2-category, these pasting diagrams become the following:



Let

$$\nu: h \circ g \Rightarrow g' \circ k$$

be a 2-morphism of C . The mate ν^\dagger of ν is then given by

and the mate of ν^\dagger is the 2-morphism $(\nu^\dagger)^\dagger: f' \circ h \Rightarrow k \circ f$ given by

Similarly, $(\omega)^\dagger = \omega$.

□

2 Morphisms of Internal Adjunctions

2.1 Lax Morphisms of Internal Adjunctions

Let C be a bicategory and let $(A, B, F, G, \eta, \epsilon)$ and $(A', B', F', G', \eta', \epsilon')$ be internal adjunctions in C .

Definition 2.1.1.1. A **lax morphism of internal adjunctions** from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$ is a lax transformation between these viewed as 2-functors from the walking adjunction.

Remark 2.1.1.2. In detail, a **lax morphism of internal adjunctions** from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$ consists of

- *1-Morphisms.* A pair of 1-morphisms

$$\begin{aligned}\phi &: A \rightarrow A', \\ \psi &: B \rightarrow B'\end{aligned}$$

of C ;

- *2-Morphisms.* A pair of 2-morphisms

$$\begin{array}{ccc} \begin{array}{ccc} A & \xrightarrow{F} & B \\ \phi \downarrow & \alpha \swarrow \nearrow & \downarrow \psi \\ A' & \xrightarrow{F'} & B' \end{array} & \begin{array}{l} \alpha: F' \circ \phi \Rightarrow \psi \circ F, \\ \beta: G' \circ \phi \Rightarrow \psi \circ G \end{array} & \begin{array}{ccc} A & \xleftarrow{G} & B \\ \phi \downarrow & \beta \swarrow \nearrow & \downarrow \psi \\ A' & \xleftarrow{G'} & B' \end{array} \end{array}$$

of C ;

satisfying the following conditions:

1. *Compatibility With Units.* We have an equality

$$\begin{array}{ccc} \begin{array}{ccccc} & & F & & B \\ & & \nearrow & & \searrow G \\ A & \xrightarrow{\quad} & id_A & \xrightarrow{\quad} & A \\ \phi \downarrow & \lambda_\phi^C \nearrow & \phi & \searrow \rho_\phi^{C,-1} & \downarrow \phi \\ A' & \xrightarrow{\quad} & \phi & \xrightarrow{\quad} & A' \\ & & id_{A'} & & \end{array} & = & \begin{array}{ccccc} & & F & & B \\ & & \nearrow & & \searrow G \\ A & \xrightarrow{\quad} & \psi & \xrightarrow{\quad} & A \\ \phi \downarrow & \alpha \nearrow & \downarrow \psi & \searrow \beta & \downarrow \phi \\ A' & \xrightarrow{\quad} & B' & \xrightarrow{\quad} & A' \\ & & \eta' \uparrow & & \\ & & id_{A'} & & \end{array} \end{array}$$

of pasting diagrams in C ;

2. *Compatibility With Counits.* We have an equality

The diagram shows two equal pasting diagrams in a bicategory C .
 Left diagram: A square with vertices B (top-left), B (top-right), A (bottom-left), and A (bottom-right). The top edge is id_B . The left edge is ψ . The right edge is ψ . The bottom edge is ϕ . There are two internal arrows: G from B to A and F from A to B . There are also two 2-morphisms: β from G to $\psi \circ \phi$ and α from $\psi \circ \phi$ to F .
 Right diagram: A square with vertices B (top-left), B (top-right), B' (bottom-left), and B' (bottom-right). The top edge is id_B . The left edge is ψ . The right edge is ψ . The bottom edge is $\text{id}_{B'}$. There are two internal arrows: G' from B to B' and F' from B' to B . There are also two 2-morphisms: λ_ψ^C from G' to $\psi \circ \text{id}_{B'}$ and $\rho_\psi^{C,-1}$ from $\psi \circ \text{id}_{B'}$ to F' .
 The two diagrams are separated by an equals sign.

of pasting diagrams in C .

2.2 Oplax Morphisms of Internal Adjunctions

Let C be a bicategory and let $(A, B, F, G, \eta, \epsilon)$ and $(A', B', F', G', \eta', \epsilon')$ be internal adjunctions in C .

Definition 2.2.1.1. An **oplax morphism of internal adjunctions** from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$ is an oplax transformation between these viewed as 2-functors from the walking adjunction.

Remark 2.2.1.2. In detail, an **oplax morphism of internal adjunctions** from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$ consists of

- *1-Morphisms.* A pair of 1-morphisms

$$\begin{aligned}\phi &: A \rightarrow A', \\ \psi &: B \rightarrow B'\end{aligned}$$

of C ;

- *2-Morphisms.* A pair of 2-morphisms

The diagram shows two equations for 2-morphisms α and β .
 Left equation: A square with vertices A (top-left), B (top-right), A' (bottom-left), and B' (bottom-right). The top edge is F . The bottom edge is F' . The left edge is ϕ . The right edge is ψ . There are two 2-morphisms: α from $\psi \circ F$ to $F' \circ \phi$ and β from $\psi \circ G$ to $G' \circ \phi$.
 Right equation: A square with vertices A (top-left), B (top-right), A' (bottom-left), and B' (bottom-right). The top edge is G . The bottom edge is G' . The left edge is ϕ . The right edge is ψ . There are two 2-morphisms: α from $\psi \circ F$ to $F' \circ \phi$ and β from $\psi \circ G$ to $G' \circ \phi$.
 The two equations are separated by an equals sign.

of C ;

satisfying the following conditions:

1. *Compatibility With Units.* We have an equality

The diagram shows an equality between two pasting diagrams in category C . The left diagram has nodes B, A, B, B' with arrows $G: B \rightarrow A$, $F: A \rightarrow B$, $\text{id}_B: B \rightarrow B$, and $\text{id}_{B'}: B' \rightarrow B'$. It includes 2-morphisms $\lambda_\phi^{C,-1}: G \circ F \Rightarrow \text{id}_B$ and $\rho_\phi^C: \text{id}_{B'} \Rightarrow F' \circ G'$. The right diagram has nodes B, A, A', B' with arrows $G: B \rightarrow A$, $F: A \rightarrow B$, $\phi: A \rightarrow A'$, $\epsilon': A' \rightarrow B$, $G': B' \rightarrow A'$, and $F': A' \rightarrow B'$. It includes 2-morphisms $\beta: G \circ \phi \Rightarrow G'$ and $\alpha: \epsilon' \circ F \Rightarrow F'$. The two diagrams are equated.

of pasting diagrams in C ;

2. *Compatibility With Counits.* We have an equality

The diagram shows an equality between two pasting diagrams in category C . The left diagram has nodes A, B, A', B' with arrows $\text{id}_A: A \rightarrow A$, $\eta: A \rightarrow B$, $\phi: A \rightarrow A'$, $F: A \rightarrow B$, $G: B \rightarrow A'$, $F': A' \rightarrow B'$, and $G': B' \rightarrow A'$. It includes 2-morphisms $\alpha: \phi \circ F \Rightarrow G$ and $\beta: G \circ \phi \Rightarrow G'$. The right diagram has nodes A, A', B' with arrows $\text{id}_A: A \rightarrow A$, $\phi: A \rightarrow A'$, $\text{id}_{A'}: A' \rightarrow A'$, $\eta': A' \rightarrow B'$, $F': A' \rightarrow B'$, and $G': B' \rightarrow A'$. It includes 2-morphisms $\rho_\psi^C: \phi \circ \eta \Rightarrow \text{id}_{A'}$ and $\lambda_\psi^{C,-1}: \eta' \circ \phi \Rightarrow \text{id}_A$. The two diagrams are equated.

of pasting diagrams in C .

2.3 Strong Morphisms of Internal Adjunctions

Let C be a bicategory and let $(A, B, F, G, \eta, \epsilon)$ and $(A', B', F', G', \eta', \epsilon')$ be internal adjunctions in C .

Definition 2.3.1.1. A **strong morphism of internal adjunctions** from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$ is a strong transformation between these viewed as 2-functors from the walking adjunction.

Remark 2.3.1.2. In detail, a **strong morphism of internal adjunctions** from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$ is equivalently:

1. A lax morphism of internal adjunctions as in [Remark 2.1.1.2](#) whose 2-morphisms are invertible.
2. An oplax morphism of internal adjunctions as in [Remark 2.2.1.2](#) whose 2-morphisms are invertible.

2.4 Strict Morphisms of Internal Adjunctions

Let C be a bicategory and let $(A, B, F, G, \eta, \epsilon)$ and $(A', B', F', G', \eta', \epsilon')$ be internal adjunctions in C .

Definition 2.4.1.1. A **strict morphism of internal adjunctions** from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$ is a strict transformation between these viewed as 2-functors from the walking adjunction.

Remark 2.4.1.2. In detail, a **strict morphism of internal adjunctions** from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$ is equivalently:

1. A lax morphism of internal adjunctions as in [Remark 2.1.1.2](#) whose 2-morphisms are identities.
2. An oplax morphism of internal adjunctions as in [Remark 2.2.1.2](#) whose 2-morphisms are identities.

3 2-Morphisms Between Morphisms of Internal Adjunctions

3.1 2-Morphisms Between Lax Morphisms of Internal Adjunctions

Let C be a bicategory, let $(A, B, F, G, \eta, \epsilon)$ and $(A', B', F', G', \eta', \epsilon')$ be internal adjunctions in C , and let $(\phi_1, \psi_1, \alpha_1, \beta_1)$ and $(\phi_2, \psi_2, \alpha_2, \beta_2)$ be lax morphisms of internal adjunctions from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$.

Definition 3.1.1.1. A **2-morphism from** $(\phi_1, \psi_1, \alpha_1, \beta_1)$ **to** $(\phi_2, \psi_2, \alpha_2, \beta_2)$ is a modification between these viewed as lax transformations.

Remark 3.1.1.2. In detail, a **2-morphism from** $(\phi_1, \psi_1, \alpha_1, \beta_1)$ **to** $(\phi_2, \psi_2, \alpha_2, \beta_2)$ consist of 2-morphisms

$$\Gamma: \phi_1 \Rightarrow \phi_2$$

$$\Sigma: \psi_1 \Rightarrow \psi_2$$

of C such that we have equalities

$$\begin{array}{ccc}
 \begin{array}{c} A \xrightarrow{F} B \\ \phi_1 \left(\begin{array}{c} \Gamma \\ \Rightarrow \end{array} \right) \phi_2 \quad \begin{array}{c} \nearrow \alpha_2 \\ \parallel \end{array} \quad \psi_2 \\ A' \xrightarrow{F'} B' \end{array} & = & \begin{array}{c} A \xrightarrow{F} B \\ \phi_1 \left(\begin{array}{c} \nearrow \alpha_1 \\ \parallel \end{array} \right) \psi_1 \left(\begin{array}{c} \Sigma \\ \Rightarrow \end{array} \right) \psi_2 \\ A' \xrightarrow{F'} B' \end{array} \\
 \\
 \begin{array}{c} B \xrightarrow{G} A \\ \psi_1 \left(\begin{array}{c} \Sigma \\ \Rightarrow \end{array} \right) \psi_2 \quad \begin{array}{c} \nearrow \beta_2 \\ \parallel \end{array} \quad \phi_2 \\ B' \xrightarrow{G'} A' \end{array} & = & \begin{array}{c} B \xrightarrow{G} A \\ \psi_1 \left(\begin{array}{c} \nearrow \beta_1 \\ \parallel \end{array} \right) \phi_1 \left(\begin{array}{c} \Gamma \\ \Rightarrow \end{array} \right) \phi_2 \\ B' \xrightarrow{G'} A' \end{array}
 \end{array}$$

of pasting diagrams in C .

3.2 2-Morphisms Between Oplax Morphisms of Internal Adjunctions

Let C be a bicategory, let $(A, B, F, G, \eta, \epsilon)$ and $(A', B', F', G', \eta', \epsilon')$ be internal adjunctions in C , and let $(\phi_1, \psi_1, \alpha_1, \beta_1)$ and $(\phi_2, \psi_2, \alpha_2, \beta_2)$ be oplax morphisms of internal adjunctions from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$.

Definition 3.2.1.1. A **2-morphism from** $(\phi_1, \psi_1, \alpha_1, \beta_1)$ **to** $(\phi_2, \psi_2, \alpha_2, \beta_2)$ is a modification between these viewed as oplax transformations.

Remark 3.2.1.2. In detail, a **2-morphism from** $(\phi_1, \psi_1, \alpha_1, \beta_1)$ **to** $(\phi_2, \psi_2, \alpha_2, \beta_2)$ consist of 2-morphisms

$$\Gamma: \phi_1 \Rightarrow \phi_2$$

$$\Sigma: \psi_1 \Rightarrow \psi_2$$

of C such that we have equalities

$$\begin{array}{ccc}
 \begin{array}{c} A \xrightarrow{F} B \\ \left(\begin{array}{c} \phi_2 \left(\begin{array}{c} \lrcorner \\ \lrcorner \end{array} \right) \phi_1 \quad \alpha_1 \\ \downarrow \quad \downarrow \\ A' \xrightarrow{F'} B' \end{array} \right) \psi_1 \end{array} & = & \begin{array}{c} A \xrightarrow{F} B \\ \left(\begin{array}{c} \alpha_2 \quad \psi_2 \left(\begin{array}{c} \lrcorner \\ \lrcorner \end{array} \right) \end{array} \right) \psi_1 \\ \downarrow \quad \downarrow \\ A' \xrightarrow{F'} B' \end{array} \\
 \\
 \begin{array}{c} B \xrightarrow{G} A \\ \left(\begin{array}{c} \psi_2 \left(\begin{array}{c} \lrcorner \\ \lrcorner \end{array} \right) \psi_1 \quad \beta_1 \\ \downarrow \quad \downarrow \\ B' \xrightarrow{G'} A' \end{array} \right) \phi_1 \end{array} & = & \begin{array}{c} B \xrightarrow{G} A \\ \left(\begin{array}{c} \beta_2 \quad \phi_2 \left(\begin{array}{c} \lrcorner \\ \lrcorner \end{array} \right) \end{array} \right) \phi_1 \\ \downarrow \quad \downarrow \\ B' \xrightarrow{G'} A' \end{array}
 \end{array}$$

of pasting diagrams in C .

3.3 2-Morphisms Between Strong Morphisms of Internal Adjunctions

Let C be a bicategory, let $(A, B, F, G, \eta, \epsilon)$ and $(A', B', F', G', \eta', \epsilon')$ be internal adjunctions in C , and let $(\phi_1, \psi_1, \alpha_1, \beta_1)$ and $(\phi_2, \psi_2, \alpha_2, \beta_2)$ be strong morphisms of internal adjunctions from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$.

Definition 3.3.1.1. A **2-morphism from** $(\phi_1, \psi_1, \alpha_1, \beta_1)$ **to** $(\phi_2, \psi_2, \alpha_2, \beta_2)$ is a modification between these viewed as strong transformations.

Remark 3.3.1.2. In detail, a **2-morphism from** $(\phi_1, \psi_1, \alpha_1, \beta_1)$ **to** $(\phi_2, \psi_2, \alpha_2, \beta_2)$ is equivalently:

- A 2-morphism (Γ, Σ) from $(\phi_1, \psi_1, \alpha_1, \beta_1)$ to $(\phi_2, \psi_2, \alpha_2, \beta_2)$ viewed as lax transformations as in [Remark 3.1.1.2](#).
- A 2-morphism (Γ, Σ) from $(\phi_1, \psi_1, \alpha_1, \beta_1)$ to $(\phi_2, \psi_2, \alpha_2, \beta_2)$ viewed as oplax transformations as in [Remark 3.2.1.2](#).

3.4 2-Morphisms Between Strict Morphisms of Internal Adjunctions

Let C be a bicategory, let $(A, B, F, G, \eta, \epsilon)$ and $(A', B', F', G', \eta', \epsilon')$ be internal adjunctions in C , and let $(\phi_1, \psi_1, \alpha_1, \beta_1)$ and $(\phi_2, \psi_2, \alpha_2, \beta_2)$ be lax morphisms of internal adjunctions from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$.

Definition 3.4.1.1. A **2-morphism from** $(\phi_1, \psi_1, \alpha_1, \beta_1)$ **to** $(\phi_2, \psi_2, \alpha_2, \beta_2)$ is a modification between these viewed as strict transformations.

Remark 3.4.1.2. In detail, a **2-morphism from** $(\phi_1, \psi_1, \alpha_1, \beta_1)$ **to** $(\phi_2, \psi_2, \alpha_2, \beta_2)$ is equivalently:

- A 2-morphism (Γ, Σ) from $(\phi_1, \psi_1, \alpha_1, \beta_1)$ to $(\phi_2, \psi_2, \alpha_2, \beta_2)$ viewed as lax transformations as in [Remark 3.1.1.2](#).
- A 2-morphism (Γ, Σ) from $(\phi_1, \psi_1, \alpha_1, \beta_1)$ to $(\phi_2, \psi_2, \alpha_2, \beta_2)$ viewed as oplax transformations as in [Remark 3.2.1.2](#).
- A 2-morphism (Γ, Σ) from $(\phi_1, \psi_1, \alpha_1, \beta_1)$ to $(\phi_2, \psi_2, \alpha_2, \beta_2)$ viewed as strong transformations as in [Remark 3.3.1.2](#).

4 Bicategories of Internal Adjunctions in a Bicategory

Appendices

A Other Chapters

Set Theory

1. [Sets](#)
2. [Constructions With Sets](#)
3. [Pointed Sets](#)
4. [Tensor Products of Pointed Sets](#)
5. [Indexed and Fibred Sets](#)
6. [Relations](#)
7. [Spans](#)
8. [Posets](#)

Category Theory

9. [Categories](#)
10. [Constructions With Categories](#)

11. [Kan Extensions](#)

Bicategories

12. [Bicategories](#)
13. [Internal Adjunctions](#)

Internal Category Theory

14. [Internal Categories](#)

Cyclic Stuff

15. [The Cycle Category](#)

Cubical Stuff

16. [The Cube Category](#)

Globular Stuff

17. [The Globe Category](#)

Cellular Stuff

18. [The Cell Category](#)

Monoids

- 19. Monoids
- 20. Constructions With Monoids

Monoids With Zero

- 21. Monoids With Zero
- 22. Constructions With Monoids With Zero

Groups

- 23. Groups
- 24. Constructions With Groups

Hyper Algebra

- 25. Hypermonoids
- 26. Hypergroups
- 27. Hypersemirings and Hyperrings
- 28. Quantales

Near-Rings

- 29. Near-Semirings
- 30. Near-Rings

Real Analysis

- 31. Real Analysis in One Variable
- 32. Real Analysis in Several Variables

Measure Theory

- 33. Measurable Spaces
- 34. Measures and Integration

Probability Theory

- 34. Probability Theory

Stochastic Analysis

- 35. Stochastic Processes, Martingales, and Brownian Motion
- 36. Itô Calculus
- 37. Stochastic Differential Equations

Differential Geometry

- 38. Topological and Smooth Manifolds

Schemes

- 39. Schemes