

The Gigantic Mess Project

2017–2023

The Gigantic Mess Project Authors

Version 1d560bdd, compiled on Jan 05, 2024.

Contents

Part I Sets	1
1 Sets.....	2
1.1 The Enrichment of Sets in Classical Truth Values.....	2
1.A Other Chapters	8
1.2 Other Chapters	9
2 Constructions With Sets.....	11
2.1 Limits of Sets	12
2.2 Colimits of Sets	28
2.3 Operations With Sets	41
2.4 Powersets.....	56
2.A Other Chapters	80
2.2 Other Chapters	81
3 Pointed Sets.....	83
3.1 Pointed Sets	84
3.2 Limits of Pointed Sets.....	86
3.3 Colimits of Pointed Sets	87
3.4 Constructions With Pointed Sets	88
3.A Other Chapters	92
3.2 Other Chapters	93
4 Tensor Products of Pointed Sets	95
4.1 Bilinear Morphisms of Pointed Sets	96
4.2 Tensors and Cotensors of Pointed Sets by Sets.....	98
4.3 The Left Tensor Product of Pointed Sets.....	100
4.4 The Right Tensor Product of Pointed Sets.....	103
4.5 Smash Products of Pointed Sets.....	107
4.A Other Chapters	114
4.2 Other Chapters	115

5 Relations	117
5.1 Relations	120
5.2 Categories of Relations	126
5.3 Constructions With Relations.....	150
5.4 Equivalence Relations.....	169
5.5 Functoriality of Powersets	182
5.6 Relative Preorders	203
5.A Other Chapters	212
5.2 Other Chapters	214
6 Spans.....	216
6.1 Spans	222
6.2 Categories of Spans.....	225
6.3 Limits of Spans	236
6.4 Colimits of Spans.....	239
6.5 Constructions With Spans.....	239
6.6 Functoriality of Spans.....	241
6.7 Un/Straightening for Spans	241
6.8 Comparison of Spans to Functions and Relations.....	243
6.A Other Chapters	250
6.2 Other Chapters	251
7 Posets.....	253
7.1 Section.....	253
7.A Other Chapters	253
7.2 Other Chapters	255
Part II Indexed and Fibred Sets	257
8 Indexed Sets	258
8.1 Indexed Sets.....	259
8.2 Limits of Indexed Sets	261
8.3 Colimits of Indexed Sets.....	263
8.4 Constructions With Indexed Sets.....	264
8.A Other Chapters	267
8.2 Other Chapters	269
9 Fibred Sets	271
9.1 Fibred Sets	272
9.2 Constructions With Fibred Sets.....	275
9.A Other Chapters	294
9.2 Other Chapters	296

10	Un/Straightening for Indexed and Fibred Sets	298
10.1	Un/Straightening for Indexed and Fibred Sets	298
10.2	Miscellany.....	304
10.A	Other Chapters	305
10.2	Other Chapters	306
Part III Category Theory		308
11	Categories.....	309
11.1	Categories.....	311
11.2	The Quadruple Adjunction With Sets.....	318
11.3	Groupoids.....	324
11.4	Functors.....	330
11.5	Natural Transformations.....	339
11.6	Categories of Categories.....	349
11.7	Miscellany.....	354
11.A	Other Chapters	355
11.2	Other Chapters	357
12	Types of Morphisms in Categories.....	359
12.1	Monomorphisms	359
12.2	Epimorphisms	364
12.A	Other Chapters	367
12.2	Other Chapters	369
13	Adjunctions and the Yoneda Lemma	371
13.1	Other Chapters	371
14	Constructions With Categories.....	373
14.A	Other Chapters	373
14.2	Other Chapters	375
15	Profunctors	377
15.1	Profunctors	377
15.2	Operations With Profunctors	378
15.3	Categories of Profunctors.....	384
15.4	Other Chapters	386
16	Cartesian Closed Categories	388
16.1	Cartesian Closed Categories	389
16.A	Other Chapters	390
16.2	Other Chapters	391

17 Kan Extensions.....	393
17.A Other Chapters	393
17.2 Other Chapters	395
Part IV Bicategories	397
18 Bicategories.....	398
18.1 Monomorphisms in Bicategories	398
18.2 Epimorphisms in Bicategories.....	401
18.3 bicategories of spans	403
18.A Other Chapters	405
18.2 Other Chapters	406
19 Internal Adjunctions	408
19.1 Internal Adjunctions	409
19.2 Morphisms of Internal Adjunctions	420
19.3 2-Morphisms Between Morphisms of Internal Adjunctions	424
19.4 Bicategories of Internal Adjunctions in a Bicategory	426
19.A Other Chapters	426
19.2 Other Chapters	427
Part V Internal Category Theory	429
20 Internal Categories	430
20.A Other Chapters	430
20.2 Other Chapters	432
Part VI Cyclic Stuff	434
21 The Cycle Category.....	435
21.A Other Chapters	435
21.2 Other Chapters	437
Part VII Cubical Stuff	439
22 The Cube Category.....	440
22.A Other Chapters	440
22.2 Other Chapters	442
Part VIII Globular Stuff	444
23 The Globe Category.....	445
23.A Other Chapters	445
23.2 Other Chapters	447

Part IX Cellular Stuff	449
24 The Cell Category	450
24.A Other Chapters	450
24.2 Other Chapters	452
Part X Monoids	454
25 Monoids.....	455
25.A Other Chapters	455
25.2 Other Chapters	457
26 Constructions With Monoids.....	459
26.A Other Chapters	459
26.2 Other Chapters	461
Part XI Monoids With Zero	463
27 Monoids With Zero.....	464
27.A Other Chapters	464
27.2 Other Chapters	466
28 Constructions With Monoids With Zero	468
28.A Other Chapters	468
28.2 Other Chapters	470
Part XII Groups	472
29 Groups.....	473
29.A Other Chapters	473
29.2 Other Chapters	475
30 Constructions With Groups	477
30.A Other Chapters	477
30.2 Other Chapters	479
Part XIII Hyper Algebra	481
31 Hypermonoids	482
31.A Other Chapters	482
31.2 Other Chapters	484
32 Hypergroups	486
32.A Other Chapters	486
32.2 Other Chapters	488

33 Hypersemirings and Hyperrings	490
33.A Other Chapters	490
33.2 Other Chapters	492
34 Quantales	494
34.A Other Chapters	494
34.2 Other Chapters	496
Part XIV Near-Rings	498
35 Near-Semirings	499
35.A Other Chapters	499
35.2 Other Chapters	501
36 Near-Rings.....	503
36.A Other Chapters	503
36.2 Other Chapters	505
Part XV Real Analysis	507
37 Real Analysis in One Variable.....	508
37.A Other Chapters	508
37.2 Other Chapters	510
38 Real Analysis in Several Variables.....	512
38.A Other Chapters	512
38.2 Other Chapters	514
Part XVI Measure Theory	516
39 Measurable Spaces.....	517
39.A Other Chapters	517
39.2 Other Chapters	519
40 Measures and Integration	521
40.A Other Chapters	521
40.2 Other Chapters	523
Part XVII Probability Theory	525
41 Probability Theory.....	526
41.A Other Chapters	526
41.2 Other Chapters	528

Part XVIII Stochastic Analysis	530
42 Stochastic Processes, Martingales, and Brownian Motion	531
42.A Other Chapters	531
42.2 Other Chapters	533
43 Itô Calculus	535
43.A Other Chapters	535
43.2 Other Chapters	537
44 Stochastic Differential Equations	539
44.A Other Chapters	539
44.2 Other Chapters	541
Part XIX Differential Geometry	543
45 Topological and Smooth Manifolds	544
45.A Other Chapters	544
45.2 Other Chapters	546
Part XX Schemes	548
46 Schemes	549
46.1 Introduction	549
46.2 Other Chapters	549
Part XXI Secret Part	551
47 To Do List.....	552
47.1 Notes to Self	552
47.A Other Chapters	555
47.2 Other Chapters	556
Index of Notation.....	565
Index of Set Theory	568

Contents (detailed)

Part I Sets		1
1	Sets.....	2
1.1	The Enrichment of Sets in Classical Truth Values.....	2
1.1.1	(-2)-Categories	2
1.1.2	(-1)-Categories	2
1.1.3	0-Categories.....	6
1.1.4	Tables of Analogies Between Set Theory and Category Theory	6
1.A	Other Chapters	8
1.2	Other Chapters	9
2	Constructions With Sets.....	11
2.1	Limits of Sets	12
2.1.1	Products of Families of Sets	12
2.1.2	Binary Products of Sets	14
2.1.3	Pullbacks	19
2.1.4	Equalisers	24
2.2	Colimits of Sets	28
2.2.1	Coproducts of Families of Sets.....	28
2.2.2	Binary Coproducts	30
2.2.3	Pushouts	32
2.2.4	Coequalisers	38
2.3	Operations With Sets	41
2.3.1	The Empty Set.....	41
2.3.2	Singleton Sets.....	41
2.3.3	Pairings of Sets.....	41
2.3.4	Ordered Pairs	42
2.3.5	Unions of Families.....	42
2.3.6	Binary Unions	42
2.3.7	Intersections of Families.....	44
2.3.8	Binary Intersections	44
2.3.9	Differences	48

2.3.10	Complements.....	50
2.3.11	Symmetric Differences.....	52
2.4	Powersets.....	56
2.4.1	Characteristic Functions.....	56
2.4.2	The Yoneda Lemma for Sets.....	60
2.4.3	Powersets.....	60
2.4.4	Direct Images	65
2.4.5	Inverse Images	70
2.4.6	Direct Images With Compact Support	73
2.A	Other Chapters	80
2.2	Other Chapters	81
3	Pointed Sets.....	83
3.1	Pointed Sets.....	84
3.1.1	Foundations.....	84
3.1.2	Morphisms of Pointed Sets.....	84
3.1.3	The Category of Pointed Sets	85
3.1.4	Elementary Properties of Pointed Sets	86
3.2	Limits of Pointed Sets.....	86
3.2.1	Products.....	86
3.2.2	Equalisers	87
3.2.3	Pullbacks	87
3.3	Colimits of Pointed Sets	87
3.3.1	Coproducts	87
3.3.2	Pushouts	87
3.3.3	Coequalisers	87
3.4	Constructions With Pointed Sets	88
3.4.1	Internal Homs	88
3.4.2	Free Pointed Sets	88
3.4.3	Wedge Sums of Pointed Sets	89
3.A	Other Chapters	92
3.2	Other Chapters	93
4	Tensor Products of Pointed Sets	95
4.1	Bilinear Morphisms of Pointed Sets	96
4.1.1	Left Bilinear Morphisms of Pointed Sets	96
4.1.2	Right Bilinear Morphisms of Pointed Sets	96
4.1.3	Bilinear Morphisms of Pointed Sets	97
4.2	Tensors and Cotensors of Pointed Sets by Sets	98
4.2.1	Tensors of Pointed Sets by Sets	98
4.2.2	Cotensors of Pointed Sets by Sets.....	99
4.3	The Left Tensor Product of Pointed Sets.....	100
4.3.1	Foundations.....	100
4.3.2	The Skew Associator	101

4.3.3	The Skew Left Unitor	102
4.3.4	The Skew Right Unitor	102
4.3.5	The Left-Skew Monoidal Category Structure on Pointed Sets.	103
4.4	The Right Tensor Product of Pointed Sets.....	103
4.4.1	Foundations.....	103
4.4.2	The Skew Associator.....	105
4.4.3	The Skew Left Unitor	106
4.4.4	The Skew Right Unitor	106
4.4.5	The Right-Skew Monoidal Category Structure on Pointed Sets.....	107
4.5	Smash Products of Pointed Sets.....	107
4.5.1	Foundations.....	107
4.A	Other Chapters	114
4.2	Other Chapters	115
5	Relations	117
5.1	Relations	120
5.1.1	Foundations.....	120
5.1.2	Relations as Decategorifications of Profunctors	122
5.1.3	Examples of Relations.....	124
5.1.4	Functional Relations	125
5.1.5	Total Relations.....	126
5.2	Categories of Relations	126
5.2.1	The Category of Relations.....	126
5.2.2	The Closed Symmetric Monoidal Category of Relations.....	127
5.2.3	The 2-Category of Relations	131
5.2.4	The Double Category of Relations	132
5.2.5	Properties of the Category of Relations	138
5.3	Constructions With Relations.....	150
5.3.1	The Graph of a Function	150
5.3.2	The Inverse of a Function	154
5.3.3	Representable Relations	155
5.3.4	The Domain and Range of a Relation	156
5.3.5	Binary Unions of Relations.....	157
5.3.6	Unions of Families of Relations	158
5.3.7	Binary Intersections of Relations	158
5.3.8	Intersections of Families of Relations	159
5.3.9	Binary Products of Relations	160
5.3.10	Products of Families of Relations	162
5.3.11	The Inverse of a Relation	162
5.3.12	Composition of Relations	163
5.3.13	The Collage of a Relation.....	168
5.4	Equivalence Relations.....	169
5.4.1	Reflexive Relations	169

5.4.2	Symmetric Relations	172
5.4.3	Transitive Relations	174
5.4.4	Equivalence Relations.....	176
5.4.5	Quotients by Equivalence Relations.....	178
5.5	Functoriality of Powersets	182
5.5.1	Direct Images	182
5.5.2	Strong Inverse Images.....	187
5.5.3	Weak Inverse Images	192
5.5.4	Direct Images With Compact Support	196
5.5.5	Functoriality of Powersets	202
5.5.6	Functoriality of Powersets: Relations on Powersets	202
5.6	Relative Preorders	203
5.6.1	The Left Skew Monoidal Structure on $\mathbf{Rel}(A, B)$	203
5.6.2	Left Relative Preorders.....	206
5.6.3	The Right Skew Monoidal Structure on $\mathbf{Rel}(A, B)$	208
5.6.4	Right Relative Preorders.....	210
5.A	Other Chapters	212
5.2	Other Chapters	214
6	Spans.....	216
6.1	Spans	222
6.1.1	The Walking Span	222
6.1.2	Spans	222
6.1.3	Morphisms of Spans.....	223
6.1.4	Functional Spans	223
6.1.5	Total Spans.....	225
6.2	Categories of Spans.....	225
6.2.1	The Category of Spans Between Two Sets	225
6.2.2	The Bicategory of Spans	227
6.2.3	The Monoidal Bicategory of Spans	229
6.2.4	The Double Category of Spans.....	229
6.2.5	Properties of The Bicategory of Spans.....	233
6.3	Limits of Spans	236
6.3.1	tmp2	236
6.3.2	tmp	236
6.3.3	Left Kan Extensions	236
6.3.4	Right Kan Extensions	237
6.3.5	Right Kan Lifts.....	238
6.4	Colimits of Spans	239
6.5	Constructions With Spans	239
6.5.1	Representable Spans	239
6.5.2	Composition of Spans.....	239
6.5.3	Horizontal Composition of Morphisms of Spans	240
6.5.4	Properties of Composition of Spans	240

6.5.5	The Inverse of a Span	241
6.6	Functionality of Spans.....	241
6.6.1	Direct Images	241
6.6.2	Functionality of Spans on Powersets.....	241
6.7	Un/Straightening for Spans	241
6.7.1	Straightening for Spans	241
6.7.2	Unstraightening for Spans	242
6.7.3	The Un/Straightening Equivalence for Spans	243
6.8	Comparison of Spans to Functions and Relations.....	243
6.8.1	Comparison to Functions	243
6.8.2	Comparison to Relations: From Span to Rel	245
6.8.3	Comparison to Relations: From Rel to Span	247
6.8.4	Comparison to Relations: The Wehrheim–Woodward Con- struction.....	249
6.8.5	Comparison to Multirelations.....	249
6.8.6	Comparison to Relations via Double Categories	249
6.A	Other Chapters	250
6.2	Other Chapters	251
7	Posets.....	253
7.1	Section.....	253
7.1.1	Section.....	253
7.A	Other Chapters	253
7.2	Other Chapters	255

Part II Indexed and Fibred Sets **257**

8	Indexed Sets	258
8.1	Indexed Sets.....	259
8.1.1	Foundations.....	259
8.1.2	Morphisms of Indexed Sets.....	259
8.1.3	The Category of Sets Indexed by a Fixed Set	259
8.1.4	The Category of Indexed Sets.....	260
8.2	Limits of Indexed Sets	261
8.2.1	Products of K -Indexed Sets.....	261
8.2.2	Pullbacks of K -Indexed Sets.....	262
8.2.3	Equalisers of K -Indexed Sets	262
8.2.4	Products in ISets	262
8.2.5	Pullbacks in ISets	262
8.2.6	Equalisers in ISets	263
8.3	Colimits of Indexed Sets	263
8.3.1	Coproducts of K -Indexed Sets	263
8.3.2	Pushouts of K -Indexed Sets	263
8.3.3	Coequalisers of K -Indexed Sets	264

8.4	Constructions With Indexed Sets.....	264
8.4.1	Change of Indexing.....	264
8.4.2	Dependent Sums	265
8.4.3	Dependent Products	266
8.4.4	Internal Hom	267
8.4.5	Adjointness of Indexed Sets	267
8.A	Other Chapters	267
8.2	Other Chapters	269
9	Fibred Sets	271
9.1	Fibred Sets	272
9.1.1	Foundations.....	272
9.1.2	Morphisms of Fibred Sets	272
9.1.3	The Category of Fibred Sets Over a Fixed Base	273
9.1.4	The Category of Fibred Sets.....	274
9.2	Constructions With Fibred Sets.....	275
9.2.1	Change of Base	275
9.2.2	Dependent Sums	277
9.2.3	Dependent Products	278
9.2.4	Internal Hom	282
9.2.5	Adjointness for Fibred Sets.....	283
9.A	Other Chapters	294
9.2	Other Chapters	296
10	Un/Straightening for Indexed and Fibred Sets	298
10.1	Un/Straightening for Indexed and Fibred Sets	298
10.1.1	Straightening for Fibred Sets	298
10.1.2	Unstraightening for Indexed Sets	301
10.1.3	The Un/Straightening Equivalence.....	304
10.2	Miscellany.....	304
10.2.1	Other Kinds of Un/Straightening.....	304
10.A	Other Chapters	305
10.2	Other Chapters	306
Part III Category Theory		308
11	Categories.....	309
11.1	Categories.....	311
11.1.1	Foundations.....	311
11.1.2	Examples of Categories	313
11.1.3	Subcategories	315
11.1.4	Skeletons of Categories	316
11.1.5	Precomposition and Postcomposition.....	317

11.2	The Quadruple Adjunction With Sets.....	318
11.2.1	Statement	318
11.2.2	Connected Components of Categories.....	319
11.2.3	Sets of Connected Components of Categories	320
11.2.4	Connected Categories	321
11.2.5	Discrete Categories.....	321
11.2.6	Indiscrete Categories.....	323
11.3	Groupoids.....	324
11.3.1	Foundations.....	324
11.3.2	The Groupoid Completion of a Category.....	324
11.3.3	The Core of a Category	327
11.4	Functors.....	330
11.4.1	Foundations.....	330
11.4.2	Faithful Functors	332
11.4.3	Full Functors.....	333
11.4.4	Fully Faithful Functors.....	334
11.4.5	Essentially Surjective Functors.....	334
11.4.6	Conservative Functors	334
11.4.7	Equivalences of Categories	335
11.4.8	Isomorphisms of Categories.....	337
11.4.9	The Natural Transformation Associated to a Functor.....	338
11.5	Natural Transformations.....	339
11.5.1	Foundations.....	339
11.5.2	Vertical Composition of Natural Transformations	341
11.5.3	Horizontal Composition of Natural Transformations.....	344
11.5.4	Properties of Natural Transformations	347
11.5.5	Natural Isomorphisms	348
11.6	Categories of Categories	349
11.6.1	Functor Categories	349
11.6.2	The Category of Categories and Functors	352
11.6.3	The 2-Category of Categories, Functors, and Natural Transformations.....	353
11.6.4	The Category of Groupoids	354
11.6.5	The 2-Category of Groupoids	354
11.7	Miscellany.....	354
11.7.1	Concrete Categories.....	354
11.7.2	Balanced Categories	354
11.7.3	Monoid Actions on Objects of Categories	354
11.7.4	Group Actions on Objects of Categories	355
11.A	Other Chapters	355
11.2	Other Chapters	357

12 Types of Morphisms in Categories.....	359
12.1 Monomorphisms	359
12.1.1 Foundations.....	359
12.1.2 Monomorphism-Reflecting Functors	363
12.1.3 Split Monomorphisms	363
12.2 Epimorphisms	364
12.2.1 Foundations.....	364
12.2.2 Regular Epimorphisms.....	366
12.2.3 Effective Epimorphisms	367
12.2.4 Split Epimorphisms	367
12.A Other Chapters	367
12.B Other Chapters	369
13 Adjunctions and the Yoneda Lemma	371
13.1 Other Chapters	371
14 Constructions With Categories.....	373
14.1 Other Chapters	373
14.2 Other Chapters	375
15 Profunctors	377
15.1 Profunctors	377
15.1.1 Foundations.....	377
15.2 Operations With Profunctors	378
15.2.1 The Domain and Range of a Profunctor	378
15.2.2 Composition of Profunctors.....	379
15.2.3 Representable Profunctors	379
15.2.4 Collages	380
15.3 Categories of Profunctors.....	384
15.3.1 The Bicategory of Profunctors	384
15.3.2 Properties of Prof	384
15.4 Other Chapters	386
16 Cartesian Closed Categories.....	388
16.1 Cartesian Closed Categories	389
16.A Other Chapters	390
16.B Other Chapters	391
17 Kan Extensions.....	393
17.A Other Chapters	393
17.B Other Chapters	395

Part IV Bicategories	397
18 Bicategories.....	398
18.1 Monomorphisms in Bicategories	398
18.1.1 Faithful Monomorphisms	398
18.1.2 Full Monomorphisms.....	399
18.1.3 Fully Faithful Monomorphisms.....	400
18.1.4 Strict Monomorphisms	400
18.2 Epimorphisms in Bicategories.....	401
18.2.1 Faithful Epimorphisms	401
18.2.2 Full Epimorphisms.....	401
18.2.3 Fully Faithful Epimorphisms.....	402
18.2.4 Strict Epimorphisms	403
18.3 bicategories of spans	403
18.A Other Chapters	405
18.2 Other Chapters	406
19 Internal Adjunctions.....	408
19.1 Internal Adjunctions	409
19.1.1 The Walking Adjunction	409
19.1.2 Internal Adjunctions	410
19.1.3 Internal Adjoint Equivalences.....	416
19.1.4 Mates	418
19.2 Morphisms of Internal Adjunctions	420
19.2.1 Lax Morphisms of Internal Adjunctions	420
19.2.2 Oplax Morphisms of Internal Adjunctions.....	422
19.2.3 Strong Morphisms of Internal Adjunctions.....	423
19.2.4 Strict Morphisms of Internal Adjunctions	423
19.3 2-Morphisms Between Morphisms of Internal Adjunctions	424
19.3.1 2-Morphisms Between Lax Morphisms of Internal Adjunctions.....	424
19.3.2 2-Morphisms Between Oplax Morphisms of Internal Adjunctions.....	424
19.3.3 2-Morphisms Between Strong Morphisms of Internal Adjunctions.....	425
19.3.4 2-Morphisms Between Strict Morphisms of Internal Adjunctions.....	425
19.4 Bicategories of Internal Adjunctions in a Bicategory	426
19.A Other Chapters	426
19.2 Other Chapters	427

Part V Internal Category Theory	429
20 Internal Categories	430
20.A Other Chapters	430
20.2 Other Chapters	432
Part VI Cyclic Stuff	434
21 The Cycle Category.....	435
21.A Other Chapters	435
21.2 Other Chapters	437
Part VII Cubical Stuff	439
22 The Cube Category.....	440
22.A Other Chapters	440
22.2 Other Chapters	442
Part VIII Globular Stuff	444
23 The Globe Category.....	445
23.A Other Chapters	445
23.2 Other Chapters	447
Part IX Cellular Stuff	449
24 The Cell Category	450
24.A Other Chapters	450
24.2 Other Chapters	452
Part X Monoids	454
25 Monoids.....	455
25.A Other Chapters	455
25.2 Other Chapters	457
26 Constructions With Monoids.....	459
26.A Other Chapters	459
26.2 Other Chapters	461
Part XI Monoids With Zero	463
27 Monoids With Zero.....	464
27.A Other Chapters	464

27.2 Other Chapters	466
28 Constructions With Monoids With Zero	468
28.A Other Chapters	468
28.2 Other Chapters	470
Part XII Groups	472
29 Groups.....	473
29.A Other Chapters	473
29.2 Other Chapters	475
30 Constructions With Groups	477
30.A Other Chapters	477
30.2 Other Chapters	479
Part XIII Hyper Algebra	481
31 Hypermonoids	482
31.A Other Chapters	482
31.2 Other Chapters	484
32 Hypergroups	486
32.A Other Chapters	486
32.2 Other Chapters	488
33 Hypersemirings and Hyperrings	490
33.A Other Chapters	490
33.2 Other Chapters	492
34 Quantales	494
34.A Other Chapters	494
34.2 Other Chapters	496
Part XIV Near-Rings	498
35 Near-Semirings	499
35.A Other Chapters	499
35.2 Other Chapters	501
36 Near-Rings.....	503
36.A Other Chapters	503
36.2 Other Chapters	505

Part XV Real Analysis	507
37 Real Analysis in One Variable.....	508
37.A Other Chapters	508
37.2 Other Chapters	510
38 Real Analysis in Several Variables.....	512
38.A Other Chapters	512
38.2 Other Chapters	514
Part XVI Measure Theory	516
39 Measurable Spaces.....	517
39.A Other Chapters	517
39.2 Other Chapters	519
40 Measures and Integration.....	521
40.A Other Chapters	521
40.2 Other Chapters	523
Part XVII Probability Theory	525
41 Probability Theory.....	526
41.A Other Chapters	526
41.2 Other Chapters	528
Part XVIII Stochastic Analysis	530
42 Stochastic Processes, Martingales, and Brownian Motion	531
42.A Other Chapters	531
42.2 Other Chapters	533
43 Itô Calculus.....	535
43.A Other Chapters	535
43.2 Other Chapters	537
44 Stochastic Differential Equations	539
44.A Other Chapters	539
44.2 Other Chapters	541
Part XIX Differential Geometry	543
45 Topological and Smooth Manifolds	544
45.A Other Chapters	544
45.2 Other Chapters	546

Part XX Schemes	548
46 Schemes.....	549
46.1 Introduction	549
46.2 Other Chapters	549
Part XXI Secret Part	551
47 To Do List.....	552
47.1 Notes to Self	552
47.1.1 Things To Ask On MO/Zulip.....	552
47.1.2 Things To Explore/Add	553
47.1.3 Random Cool Papers	554
47.1.4 Omitted Proofs To Add.....	554
47.A Other Chapters	555
47.2 Other Chapters	556
Index of Notation.....	565
Index of Set Theory	568

Part I

Sets

Chapter 1

Sets

0000 This chapter (will eventually) contain material on axiomatic set theory, as well as a couple other things.

Contents

1.1	The Enrichment of Sets in Classical Truth Values.....	2
1.1.1	(-2)-Categories.....	2
1.1.2	(-1)-Categories.....	2
1.1.3	0-Categories	6
1.1.4	Tables of Analogies Between Set Theory and Category Theory	6
1.A	Other Chapters	8
1.2	Other Chapters	9

1.1 The Enrichment of Sets in Classical Truth Values

1.1.1 (-2)-Categories

0003 **Definition 1.1.1.1.** A (-2)-category is the “necessarily true” truth value.^{1,2,3}

1.1.2 (-1)-Categories

0005 **Definition 1.1.2.1.1.** A (-1)-category is a classical truth value.

0006 **Remark 1.1.2.1.2.** ⁴(-1)-categories should be thought of as being “categories enriched in (-2)-categories”, having a collection of objects and, for each pair of objects, a Hom-object $\text{Hom}(x, y)$ that is a (-2)-category (i.e. trivial).

¹Thus, there is only one (-2)-category.

²A ($-n$)-category for $n = 3, 4, \dots$ is also the “necessarily true” truth value, coinciding with a (-2)-category.

³For motivation, see [BS10, p. 13].

⁴For more motivation, see [BS10, p. 13].

Therefore, a (-1) -category C is either ([BS10, pp. 33–34]):

1. *Empty*, having no objects;
2. *Contractible*, having a collection of objects $\{a, b, c, \dots\}$, but with $\text{Hom}_C(a, b)$ being a (-2) -category (i.e. trivial) for all $a, b \in \text{Obj}(C)$, forcing all objects of C to be uniquely isomorphic to each other.

As such, there are only two (-1) -categories, up to equivalence:

- The (-1) -category **false** (the empty one);
- The (-1) -category **true** (the contractible one).

0007 Definition 1.1.2.1.3. The **poset of truth values**⁵ is the poset $(\{\text{true}, \text{false}\}, \leq)^6$ consisting of

- *The Underlying Set.* The set $\{\text{true}, \text{false}\}$ whose elements are the truth values **true** and **false**;
- *The Partial Order.* The partial order

$$\leq : \{\text{true}, \text{false}\} \times \{\text{true}, \text{false}\} \rightarrow \{\text{true}, \text{false}\}$$

on $\{\text{true}, \text{false}\}$ defined by⁷

$$\begin{aligned} \text{false} \leq \text{false} &\stackrel{\text{def}}{=} \text{true}, \\ \text{true} \leq \text{false} &\stackrel{\text{def}}{=} \text{false}, \\ \text{false} \leq \text{true} &\stackrel{\text{def}}{=} \text{true}, \\ \text{true} \leq \text{true} &\stackrel{\text{def}}{=} \text{true}. \end{aligned}$$

0008 Proposition 1.1.2.1.4. The poset of truth values $\{\text{t}, \text{f}\}$ is Cartesian closed with product given by⁸

$$\begin{aligned} \text{t} \times \text{t} &= \text{t}, \\ \text{t} \times \text{f} &= \text{f}, \\ \text{f} \times \text{t} &= \text{f}, \\ \text{f} \times \text{f} &= \text{f}, \end{aligned}$$

and internal Hom $\mathbf{Hom}_{\{\text{t}, \text{f}\}}$ given by the partial order of $\{\text{t}, \text{f}\}$, i.e. by

$$\begin{aligned} \mathbf{Hom}_{\{\text{t}, \text{f}\}}(\text{t}, \text{t}) &= \text{t}, \\ \mathbf{Hom}_{\{\text{t}, \text{f}\}}(\text{t}, \text{f}) &= \text{f}, \\ \mathbf{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{t}) &= \text{t}, \\ \mathbf{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{f}) &= \text{t}. \end{aligned}$$

⁵Further Terminology: Also called the **poset of (-1) -categories**.

⁶Further Notation: Also written $\{\text{t}, \text{f}\}$.

⁷This partial order coincides with logical implication.

⁸Note that \times coincides with the “and” operator, while $\mathbf{Hom}_{\{\text{t}, \text{f}\}}$ coincides with the logical implication

Proof. Existence of Products: We claim that the products $t \times t$, $t \times f$, $f \times t$, and $f \times f$ satisfy the universal property of the product in $\{t, f\}$. Indeed, consider the diagrams

$$\begin{array}{cccc} \begin{array}{c} P_1 \\ | \\ p_1^1 \swarrow \quad \searrow p_2^1 \\ t \xleftarrow{\text{pr}_1} \underbrace{t \times t}_{=t} \xrightarrow{\text{pr}_2} t \end{array} & \begin{array}{c} P_2 \\ | \\ p_1^2 \swarrow \quad \searrow p_2^2 \\ t \xleftarrow{\text{pr}_1} \underbrace{t \times f}_{=f} \xrightarrow{\text{pr}_2} f \end{array} & \begin{array}{c} P_3 \\ | \\ p_1^3 \swarrow \quad \searrow p_2^3 \\ f \xleftarrow{\text{pr}_1} \underbrace{f \times t}_{=f} \xrightarrow{\text{pr}_2} t \end{array} & \begin{array}{c} P_4 \\ | \\ p_1^4 \swarrow \quad \searrow p_2^4 \\ f \xleftarrow{\text{pr}_1} \underbrace{f \times f}_{=f} \xrightarrow{\text{pr}_2} f. \end{array} \end{array}$$

Here:

1. If $P_1 = t$, then $p_1^1 = p_2^1 = \text{id}_t$, and there's indeed a unique morphism from P_1 to t making the diagram commute, namely id_t ;
2. If $P_1 = f$, then $p_1^1 = p_2^1$ are given by the unique morphism from f to t , and there's indeed a unique morphism from P_1 to t making the diagram commute, namely the unique morphism from f to t ;
3. If $P_2 = t$, then there is no morphism p_2^2 .
4. If $P_2 = f$, then p_1^2 is the unique morphism from f to t while $p_2^2 = \text{id}_f$, and there's indeed a unique morphism from P_2 to f making the diagram commute, namely id_f ;
5. The proof for P_3 is similar to the one for P_2 ;
6. If $P_4 = t$, then there is no morphism p_1^4 or p_2^4 .
7. If $P_4 = f$, then $p_1^4 = p_2^4 = \text{id}_f$, and there's indeed a unique morphism from P_4 to f making the diagram commute, namely id_f .

Cartesian Closedness: We claim there's a bijection

$$\text{Hom}_{\{t,f\}}(A \times B, C) \cong \text{Hom}_{\{t,f\}}(A, \text{Hom}_{\{t,f\}}(B, C))$$

natural in $A, B, C \in \{t, f\}$. Indeed:

- For $(A, B, C) = (t, t, t)$, we have

$$\begin{aligned} \text{Hom}_{\{t,f\}}(t \times t, t) &\cong \text{Hom}_{\{t,f\}}(t, t) \\ &= \{\text{id}_{\text{true}}\} \\ &\cong \text{Hom}_{\{t,f\}}(t, t) \\ &\cong \text{Hom}_{\{t,f\}}(t, \text{Hom}_{\{t,f\}}(t, t)). \end{aligned}$$

operator.

- For $(A, B, C) = (\text{t}, \text{t}, \text{f})$, we have

$$\begin{aligned}\text{Hom}_{\{\text{t}, \text{f}\}}(\text{t} \times \text{t}, \text{f}) &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{t}, \text{f}) \\ &= \emptyset \\ &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{t}, \text{f}) \\ &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{t}, \mathbf{Hom}_{\{\text{t}, \text{f}\}}(\text{t}, \text{f})).\end{aligned}$$

- For $(A, B, C) = (\text{t}, \text{f}, \text{t})$, we have

$$\begin{aligned}\text{Hom}_{\{\text{t}, \text{f}\}}(\text{t} \times \text{f}, \text{t}) &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{t}) \\ &\cong \text{pt} \\ &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{t}) \\ &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \mathbf{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{t})).\end{aligned}$$

- For $(A, B, C) = (\text{t}, \text{f}, \text{f})$, we have

$$\begin{aligned}\text{Hom}_{\{\text{t}, \text{f}\}}(\text{t} \times \text{f}, \text{f}) &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{f}) \\ &\cong \{\text{id}_{\text{false}}\} \\ &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{f}) \\ &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{t}, \mathbf{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{f})).\end{aligned}$$

- For $(A, B, C) = (\text{f}, \text{t}, \text{t})$, we have

$$\begin{aligned}\text{Hom}_{\{\text{t}, \text{f}\}}(\text{f} \times \text{t}, \text{t}) &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{t}) \\ &\cong \text{pt} \\ &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{t}) \\ &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \mathbf{Hom}_{\{\text{t}, \text{f}\}}(\text{t}, \text{t})).\end{aligned}$$

- For $(A, B, C) = (\text{f}, \text{t}, \text{f})$, we have

$$\begin{aligned}\text{Hom}_{\{\text{t}, \text{f}\}}(\text{f} \times \text{t}, \text{f}) &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{f}) \\ &\cong \{\text{id}_{\text{false}}\} \\ &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{f}) \\ &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \mathbf{Hom}_{\{\text{t}, \text{f}\}}(\text{t}, \text{f})).\end{aligned}$$

- For $(A, B, C) = (\text{f}, \text{f}, \text{t})$, we have

$$\begin{aligned}\text{Hom}_{\{\text{t}, \text{f}\}}(\text{f} \times \text{f}, \text{t}) &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{t}) \\ &\cong \text{pt} \\ &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{t}) \\ &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \mathbf{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{t})).\end{aligned}$$

- For $(A, B, C) = (\text{f}, \text{f}, \text{f})$, we have

$$\begin{aligned}\text{Hom}_{\{\text{t}, \text{f}\}}(\text{f} \times \text{f}, \text{f}) &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{f}) \\ &= \{\text{id}_{\text{false}}\} \\ &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{f}) \\ &\cong \text{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \mathbf{Hom}_{\{\text{t}, \text{f}\}}(\text{f}, \text{f})).\end{aligned}$$

The proof of naturality is omitted. \square

1.1.3 0-Categories

000A Definition 1.1.3.1.1. A **0-category** is a poset.⁹

000B Definition 1.1.3.1.2. A **0-groupoid** is a 0-category in which every morphism is invertible.¹⁰

1.1.4 Tables of Analogies Between Set Theory and Category Theory

Here we record some analogies between notions in set theory and category theory. Note that the analogies relating to presheaves relate equally well to copresheaves, as the opposite X^{op} of a set X is just X again.

Basics:

SET THEORY	CATEGORY THEORY
Enrichment in {true, false}	Enrichment in Sets
Set X	Category C
Element $x \in X$	Object $X \in \text{Obj}(C)$
Function	Functor
Function $X \rightarrow \{\text{true}, \text{false}\}$	Functor $C \rightarrow \text{Sets}$
Function $X \rightarrow \{\text{true}, \text{false}\}$	Presheaf $C^{\text{op}} \rightarrow \text{Sets}$

Powersets and categories of presheaves:

⁹Motivation: A 0-category is precisely a category enriched in the poset of (-1) -categories.

¹⁰That is, a set.

SET THEORY	CATEGORY THEORY
Powerset $\mathcal{P}(X)$	Presheaf category $\text{PSh}(C)$
Characteristic function $\chi_{\{x\}}$	Representable presheaf h_X
Characteristic embedding $\chi_{(-)}: X \hookrightarrow \mathcal{P}(X)$	Yoneda embedding $\natural: C^{\text{op}} \hookrightarrow \text{PSh}(C)$
Characteristic relation $\chi_X(-_1, -_2)$	Hom profunctor $\text{Hom}_C(-_1, -_2)$
The Yoneda lemma for sets $\text{Hom}_{\mathcal{P}(X)}(\chi_x, \chi_U) = \chi_U(x)$	The Yoneda lemma for categories $\text{Nat}(h_X, \mathcal{F}) \cong \mathcal{F}(X)$
The characteristic embedding is fully faithful, $\text{Hom}_{\mathcal{P}(X)}(\chi_x, \chi_y) = \chi_X(x, y)$	The Yoneda embedding is fully faithful, $\text{Nat}(h_X, h_Y) \cong \text{Hom}_C(X, Y)$
Subsets are unions of their elements $U = \bigcup_{x \in U} \{x\}$ or $\chi_U = \underset{\chi_x \in \text{Sets}(U, \{\text{t}, \text{f}\})}{\text{colim}} (\chi_x)$	Presheaves are colimits of representables, $\mathcal{F} \cong \underset{h_X \in \int_C \mathcal{F}}{\text{colim}} (h_X)$

Categories of elements:

SET THEORY	CATEGORY THEORY
Assignment $U \mapsto \chi_U$	Assignment $\mathcal{F} \mapsto \int_C \mathcal{F}$ (the category of elements)
Assignment $U \mapsto \chi_U$ giving an isomorphism $\mathcal{P}(X) \cong \text{Sets}(X, \{\text{t}, \text{f}\})$	Assignment $\mathcal{F} \mapsto \int_C \mathcal{F}$ giving an equivalence $\text{PSh}(C) \xrightarrow{\text{eq.}} \text{DFib}(C)$

Functions between powersets and functors between presheaf categories:

SET THEORY	CATEGORY THEORY
Direct image function $f_*: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$	Inverse image functor $f^{-1}: \text{PSh}(C) \rightarrow \text{PSh}(\mathcal{D})$
Inverse image function $f^{-1}: \mathcal{P}(Y) \rightarrow \mathcal{P}(X)$	Direct image functor $f_*: \text{PSh}(\mathcal{D}) \rightarrow \text{PSh}(C)$
Direct image with compact support function $f_!: \mathcal{P}(X) \rightarrow \mathcal{P}(Y)$	Direct image with compact support functor $f_!: \text{PSh}(C) \rightarrow \text{PSh}(\mathcal{D})$

Relations and profunctors:

SET THEORY	CATEGORY THEORY
Relation $R: X \times Y \rightarrow \{t, f\}$	Profunctor $p: \mathcal{D}^{\text{op}} \times C \rightarrow \text{Sets}$
Relation $R: X \rightarrow \mathcal{P}(Y)$	Profunctor $p: C \rightarrow \text{PSh}(\mathcal{D})$
Relation as a cocontinuous morphism of posets $R: (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(Y), \subset)$	Profunctor as a colimit-preserving functor $p: \text{PSh}(C) \rightarrow \text{PSh}(\mathcal{D})$

Appendices

1.A Other Chapters

Sets

- 1. Sets
- 2. Constructions With Sets
- 3. Pointed Sets
- 4. Tensor Products of Pointed Sets
- 5. Relations
- 6. Spans
- 7. Posets

Indexed and Fibred Sets

- 7. Indexed Sets
- 8. Fibred Sets
- 9. Un/Straightening for Indexed and Fibred Sets

Category Theory

- 11. Categories
- 12. Types of Morphisms in Categories
- 13. Adjunctions and the Yoneda Lemma
- 14. Constructions With Categories
- 15. Profunctors

16. Cartesian Closed Categories

17. Kan Extensions

Bicategories

- 18. Bicategories
- 19. Internal Adjunctions

Internal Category Theory

- 20. Internal Categories

Cyclic Stuff

- 21. The Cycle Category

Cubical Stuff

- 22. The Cube Category

Globular Stuff

- 23. The Globe Category

Cellular Stuff

- 24. The Cell Category

Monoids

- 25. Monoids
- 26. Constructions With Monoids

Monoids With Zero

- 27. Monoids With Zero

-
28. Constructions With Monoids With Zero 38. Real Analysis in Several Variables

Groups

29. Groups
30. Constructions With Groups

Hyper Algebra

31. Hypermonoids
32. Hypergroups
33. Hypersemirings and Hyperrings
34. Quantales

Near-Rings

35. Near-Semirings
36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

Measure Theory

39. Measurable Spaces
40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales, and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

1.2 Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets
5. Relations
6. Spans
7. Posets

Indexed and Fibred Sets

7. Indexed Sets
8. Fibred Sets

9. Un/Straightening for Indexed and Fibred Sets

Category Theory

11. Categories
12. Types of Morphisms in Categories
13. Adjunctions and the Yoneda Lemma
14. Constructions With Categories
15. Profunctors
16. Cartesian Closed Categories
17. Kan Extensions

Bicategories

18. Bicategories

19. Internal Adjunctions	32. Hypergroups
Internal Category Theory	33. Hypersemirings and Hyperrings
20. Internal Categories	34. Quantales
Cyclic Stuff	Near-Rings
21. The Cycle Category	35. Near-Semirings
Cubical Stuff	36. Near-Rings
22. The Cube Category	Real Analysis
Globular Stuff	37. Real Analysis in One Variable
23. The Globe Category	38. Real Analysis in Several Variables
Cellular Stuff	Measure Theory
24. The Cell Category	39. Measurable Spaces
Monoids	40. Measures and Integration
25. Monoids	Probability Theory
26. Constructions With Monoids	40. Probability Theory
Monoids With Zero	Stochastic Analysis
27. Monoids With Zero	41. Stochastic Processes, Martingales, and Brownian Motion
28. Constructions With Monoids With Zero	42. Itô Calculus
Groups	43. Stochastic Differential Equations
29. Groups	Differential Geometry
30. Constructions With Groups	44. Topological and Smooth Manifolds
Hyper Algebra	Schemes
31. Hypermonoids	45. Schemes

Chapter 2

Constructions With Sets

000D This chapter contains some material relating to constructions with sets. Notably, it contains:

1. Explicit descriptions of the major types of co/limits in Sets, including in particular pushouts and coequalisers (see [Definitions 2.2.3.1.1](#) and [2.2.4.1.1](#) and [Remarks 2.2.3.1.2](#) and [2.2.4.1.2](#));
2. A discussion of powersets as decategorifications of categories of presheaves ([Remarks 2.4.1.1.2](#) and [2.4.3.1.2](#));
3. A lengthy discussion of the adjoint triple

$$f_* \dashv f^{-1} \dashv f_! : \mathcal{P}(A) \rightleftarrows \mathcal{P}(B)$$

of functors (morphisms of posets) between $\mathcal{P}(A)$ and $\mathcal{P}(B)$ induced by a map of sets $f : A \rightarrow B$, along with a discussion of the properties of f_* , f^{-1} , and $f_!$.

Contents

2.1	Limits of Sets.....	12
2.1.1	Products of Families of Sets.....	12
2.1.2	Binary Products of Sets.....	14
2.1.3	Pullbacks.....	19
2.1.4	Equalisers.....	24
2.2	Colimits of Sets.....	28
2.2.1	Coproducts of Families of Sets	28
2.2.2	Binary Coproducts.....	30
2.2.3	Pushouts	32
2.2.4	Coequalisers.....	38
2.3	Operations With Sets	41
2.3.1	The Empty Set	41

2.3.2	Singleton Sets	41
2.3.3	Pairings of Sets	41
2.3.4	Ordered Pairs	42
2.3.5	Unions of Families	42
2.3.6	Binary Unions	42
2.3.7	Intersections of Families	44
2.3.8	Binary Intersections	44
2.3.9	Differences	48
2.3.10	Complements	50
2.3.11	Symmetric Differences	52
2.4	Powersets	56
2.4.1	Characteristic Functions	56
2.4.2	The Yoneda Lemma for Sets	60
2.4.3	Powersets	60
2.4.4	Direct Images	65
2.4.5	Inverse Images	70
2.4.6	Direct Images With Compact Support	73
2.A	Other Chapters	80
2.2	Other Chapters	81

2.1 Limits of Sets

2.1.1 Products of Families of Sets

Let $\{A_i\}_{i \in I}$ be a family of sets.

000G Definition 2.1.1.1.1. The **product**¹ of $\{A_i\}_{i \in I}$ is the pair $(\prod_{i \in I} A_i, \{\text{pr}_i\}_{i \in I})$ consisting of

- *The Limit.* The set $\prod_{i \in I} A_i$ defined by²

$$\prod_{i \in I} A_i \stackrel{\text{def}}{=} \left\{ f \in \text{Sets}\left(I, \bigcup_{i \in I} A_i\right) \middle| \begin{array}{l} \text{for each } i \in I, \text{ we} \\ \text{have } f(i) \in A_i \end{array} \right\}.$$

- *The Cone.* The collection

$$\left\{ \text{pr}_i: \prod_{i \in I} A_i \rightarrow A_i \right\}_{i \in I}$$

¹Further Terminology: Also called the **Cartesian product** of $\{A_i\}_{i \in I}$.

²Less formally, $\prod_{i \in I} A_i$ is the set whose elements are I -indexed collections $(a_i)_{i \in I}$ with $a_i \in A_i$ for each $i \in I$.

of maps given by

$$\text{pr}_i(f) \stackrel{\text{def}}{=} f(i)$$

for each $f \in \prod_{i \in I} A_i$ and each $i \in I$.

Proof. We claim that $\prod_{i \in I} A_i$ is the categorical product of $\{A_i\}_{i \in I}$ in Sets. Indeed, suppose we have, for each $i \in I$, a diagram of the form

$$\begin{array}{ccc} P & & \\ & \searrow p_i & \\ & \prod_{i \in I} A_i & \xrightarrow{\text{pr}_i} A_i \end{array}$$

in Sets. Then there exists a unique map $\phi: P \rightarrow \prod_{i \in I} A_i$, uniquely determined by the condition $\text{pr}_i \circ \phi = p_i$ for each $i \in I$, being necessarily given by

$$\phi(x) = (p_i(x))_{i \in I}$$

for each $x \in P$. □

000H Proposition 2.1.1.1.2. Let $\{A_i\}_{i \in I}$ be a family of sets.

000J 1. *Functionality.* The assignment $\{A_i\}_{i \in I} \mapsto \prod_{i \in I} A_i$ defines a functor

$$\prod_{i \in I}: \text{Fun}(I_{\text{disc}}, \text{Sets}) \rightarrow \text{Sets}$$

where

- *Action on Objects.* For each $(A_i)_{i \in I} \in \text{Obj}(\text{Fun}(I_{\text{disc}}, \text{Sets}))$, we have

$$\left[\prod_{i \in I} \right] ((A_i)_{i \in I}) \stackrel{\text{def}}{=} \prod_{i \in I} A_i$$

- *Action on Morphisms.* For each $(A_i)_{i \in I}, (B_i)_{i \in I} \in \text{Obj}(\text{Fun}(I_{\text{disc}}, \text{Sets}))$, the action on Hom-sets

$$\left(\prod_{i \in I} \right)_{(A_i)_{i \in I}, (B_i)_{i \in I}} : \text{Nat}((A_i)_{i \in I}, (B_i)_{i \in I}) \rightarrow \text{Sets} \left(\prod_{i \in I} A_i, \prod_{i \in I} B_i \right)$$

of $\prod_{i \in I}$ at $((A_i)_{i \in I}, (B_i)_{i \in I})$ is defined by sending a map

$$\{f_i: A_i \rightarrow B_i\}_{i \in I}$$

in $\text{Nat}((A_i)_{i \in I}, (B_i)_{i \in I})$ to the map of sets

$$\prod_{i \in I} f_i: \prod_{i \in I} A_i \rightarrow \prod_{i \in I} B_i$$

defined by

$$\left[\prod_{i \in I} f_i \right] ((a_i)_{i \in I}) \stackrel{\text{def}}{=} (f_i(a_i))_{i \in I}$$

for each $(a_i)_{i \in I} \in \prod_{i \in I} A_i$.

Proof. **Item 1, Functoriality:** Clear. \square

2.1.2 Binary Products of Sets

Let A and B be sets.

000L Definition 2.1.2.1.1. The **product**³ of A and B is the pair $(A \times B, \{\text{pr}_1, \text{pr}_2\})$ consisting of

- *The Limit.* The set $A \times B$ defined by⁴

$$\begin{aligned} A \times B &\stackrel{\text{def}}{=} \prod_{z \in \{A, B\}} z \\ &\stackrel{\text{def}}{=} \{f \in \text{Sets}(\{0, 1\}, A \cup B) \mid \text{we have } f(0) \in A \text{ and } f(1) \in B\} \\ &\cong \{\{\{a\}, \{a, b\}\} \in \mathcal{P}(\mathcal{P}(A \cup B)) \mid \text{we have } a \in A \text{ and } b \in B\}. \end{aligned}$$

- *The Cone.* The maps

$$\begin{aligned} \text{pr}_1 &: A \times B \rightarrow A, \\ \text{pr}_2 &: A \times B \rightarrow B \end{aligned}$$

defined by

$$\begin{aligned} \text{pr}_1(a, b) &\stackrel{\text{def}}{=} a, \\ \text{pr}_2(a, b) &\stackrel{\text{def}}{=} b \end{aligned}$$

for each $(a, b) \in A \times B$.

Proof. We claim that $A \times B$ is the categorical product of A and B in Sets . Indeed, suppose we have a diagram of the form

$$\begin{array}{ccc} & P & \\ p_1 \swarrow & & \searrow p_2 \\ A & \xleftarrow{\text{pr}_1} & A \times B \xrightarrow{\text{pr}_2} B \end{array}$$

³Further Terminology: Also called the **Cartesian product of A and B** or the **binary Cartesian product of A and B** , for emphasis.

This can also be thought of as the $(\mathbb{E}_{-1}, \mathbb{E}_{-1})$ -**tensor product of A and B** .

⁴In other words, $A \times B$ is the set whose elements are ordered pairs (a, b) with $a \in A$ and $b \in B$ as in

in Sets . Then there exists a unique map $\phi: P \rightarrow A \times B$, uniquely determined by the conditions

$$\begin{aligned}\text{pr}_1 \circ \phi &= p_1, \\ \text{pr}_2 \circ \phi &= p_2,\end{aligned}$$

being necessarily given by

$$\phi(x) = (p_1(x), p_2(x))$$

for each $x \in P$. □

000M Proposition 2.1.2.1.2. Let A, B, C , and X be sets.

000N 1. *Functoriality.* The assignments $A, B, (A, B) \mapsto A \times B$ define functors

$$\begin{aligned}A \times -_2: \text{Sets} &\rightarrow \text{Sets}, \\ -_1 \times B: \text{Sets} &\rightarrow \text{Sets}, \\ -_1 \times -_2: \text{Sets} \times \text{Sets} &\rightarrow \text{Sets},\end{aligned}$$

where $-_1 \times -_2$ is the functor where

- *Action on Objects.* For each $(A, B) \in \text{Obj}(\text{Sets} \times \text{Sets})$, we have

$$[-_1 \times -_2](A, B) \stackrel{\text{def}}{=} A \times B;$$

- *Action on Morphisms.* For each $(A, B), (X, Y) \in \text{Obj}(\text{Sets})$, the action on Hom-sets

$$\times_{(A,B),(X,Y)}: \text{Sets}(A, X) \times \text{Sets}(B, Y) \rightarrow \text{Sets}(A \times B, X \times Y)$$

of \times at $((A, B), (X, Y))$ is defined by sending (f, g) to the function

$$f \times g: A \times B \rightarrow X \times Y$$

defined by

$$[f \times g](a, b) \stackrel{\text{def}}{=} (f(a), g(b))$$

for each $(a, b) \in A \times B$;

and where $A \times -$ and $- \times B$ are the partial functors of $-_1 \times -_2$ at $A, B \in \text{Obj}(\text{Sets})$.

000P 2. *Adjointness.* We have adjunctions

$$\begin{aligned}(A \times - \dashv \text{Sets}(A, -)): \text{Sets} &\xrightleftharpoons[\text{Sets}(A, -)]{\perp} \text{Sets}, \\ (- \times B \dashv \text{Sets}(B, -)): \text{Sets} &\xrightleftharpoons[\text{Sets}(B, -)]{\perp} \text{Sets},\end{aligned}$$

witnessed by bijections

$$\begin{aligned}\text{Sets}(A \times B, C) &\cong \text{Sets}(A, \text{Sets}(B, C)), \\ \text{Sets}(A \times B, C) &\cong \text{Sets}(B, \text{Sets}(A, C)),\end{aligned}$$

natural in $A, B, C \in \text{Obj}(\text{Sets})$.

- 000Q** 3. *Associativity.* We have an isomorphism of sets

$$(A \times B) \times C \cong A \times (B \times C),$$

natural in $A, B, C \in \text{Obj}(\text{Sets})$.

- 000R** 4. *Unitality.* We have isomorphisms of sets

$$\begin{aligned}\text{pt} \times A &\cong A, \\ A \times \text{pt} &\cong A,\end{aligned}$$

natural in $A \in \text{Obj}(\text{Sets})$.

- 000S** 5. *Commutativity.* We have an isomorphism of sets

$$A \times B \cong B \times A,$$

natural in $A, B \in \text{Obj}(\text{Sets})$.

- 000T** 6. *Annihilation With the Empty Set.* We have isomorphisms of sets

$$\begin{aligned}A \times \emptyset &\cong \emptyset, \\ \emptyset \times A &\cong \emptyset,\end{aligned}$$

natural in $A \in \text{Obj}(\text{Sets})$.

- 000U** 7. *Distributivity Over Unions.* We have isomorphisms of sets

$$\begin{aligned}A \times (B \cup C) &= (A \times B) \cup (A \times C), \\ (A \cup B) \times C &= (A \times C) \cup (B \times C).\end{aligned}$$

- 000V** 8. *Distributivity Over Intersections.* We have isomorphisms of sets

$$\begin{aligned}A \times (B \cap C) &= (A \times B) \cap (A \times C), \\ (A \cap B) \times C &= (A \times C) \cap (B \times C).\end{aligned}$$

- 000W** 9. *Middle-Four Exchange with Respect to Intersections.* We have an isomorphism of sets

$$(A \times B) \cap (C \times D) \cong (A \cap B) \times (C \cap D).$$

000X 10. *Distributivity Over Differences.* We have isomorphisms of sets

$$\begin{aligned} A \times (B \setminus C) &= (A \times B) \setminus (A \times C), \\ (A \setminus B) \times C &= (A \times C) \setminus (B \times C), \end{aligned}$$

natural in $A, B, C \in \text{Obj}(\text{Sets})$.

000Y 11. *Distributivity Over Symmetric Differences.* We have isomorphisms of sets

$$\begin{aligned} A \times (B \Delta C) &= (A \times B) \Delta (A \times C), \\ (A \Delta B) \times C &= (A \times C) \Delta (B \times C), \end{aligned}$$

natural in $A, B, C \in \text{Obj}(\text{Sets})$.

000Z 12. *Symmetric Monoidality.* The triple $(\text{Sets}, \times, \text{pt})$ is a symmetric monoidal category.

0010 13. *Symmetric Bimonoidality.* The quintuple $(\text{Sets}, \coprod, \emptyset, \times, \text{pt})$ is a symmetric bimonoidal category.

Proof. **Item 1, Functoriality:** This follows by applying associativity and unitality componentwise.

Item 2, Adjointness: We prove only that there's an adjunction $X \times - \dashv \text{Hom}_{\text{Sets}}(-, Z)$, witnessed by a bijection

$$\text{Hom}_{\text{Sets}}(X \times Y, Z) \cong \text{Hom}_{\text{Sets}}(X, \text{Hom}_{\text{Sets}}(Y, Z)),$$

natural in $Y, Z \in \text{Obj}(\text{Sets})$, as the proof of the existence of the adjunction $- \times Y \dashv \text{Hom}_{\text{Sets}}(-, Z)$ follows almost exactly in the same way.⁵

- *Map I.* We define a map

$$\Phi_{Y,Z}: \text{Hom}_{\text{Sets}}(X \times Y, Z) \rightarrow \text{Hom}_{\text{Sets}}(X, \text{Hom}_{\text{Sets}}(Y, Z)),$$

by sending a morphism $\xi: X \times Y \rightarrow Z$ to the morphism

$$\xi^\dagger: X \rightarrow \text{Hom}_{\text{Sets}}(Y, Z)$$

defined by

$$\xi^\dagger(x) \stackrel{\text{def}}{=} \xi_x$$

for each $x \in X$, where $\xi_x: Y \rightarrow Z$ is the map defined by

$$\xi_x(y) \stackrel{\text{def}}{=} \xi(x, y)$$

for each $y \in Y$.

Definition 2.3.4.1.1

⁵Here we sometimes denote a map $f: X \rightarrow Y$ by $[x \mapsto f(x)]$, similar to the lambda notation $\lambda x. f(x)$.

- *Map II.* We define a map

$$\Psi_{Y,Z}: \text{Hom}_{\text{Sets}}(X, \text{Hom}_{\text{Sets}}(Y, Z)), \rightarrow \text{Hom}_{\text{Sets}}(X \times Y, Z)$$

given by sending a map $\xi: X \rightarrow \text{Hom}_{\text{Sets}}(Y, Z)$ to the map

$$\xi^\dagger: X \times Y \rightarrow Z$$

defined by

$$\xi^\dagger(x, y) \stackrel{\text{def}}{=} [\xi(x)](y)$$

for each $(x, y) \in X \times Y$.

- *Naturality I.* We need to show that, given a function $g: Y \rightarrow Y'$, the diagram

$$\begin{array}{ccc} \text{Hom}_{\text{Sets}}(X \times Y', Z) & \xrightarrow{\Phi_{Y',Z}} & \text{Hom}_{\text{Sets}}(X, \text{Hom}_{\text{Sets}}(Y', Z)), \\ \text{id}_X \times g^* \downarrow & & \downarrow (g^*)_* \\ \text{Hom}_{\text{Sets}}(X \times Y, Z) & \xrightarrow{\Phi_{Y,Z}} & \text{Hom}_{\text{Sets}}(X, \text{Hom}_{\text{Sets}}(Y, Z)), \end{array}$$

commutes. Indeed, given a morphism $\xi: X' \times Y \rightarrow Z$, we have

$$\begin{aligned} [\Phi_{Y,Z} \circ (g^* \times \text{id}_Y)](\xi) &\stackrel{\text{def}}{=} (\xi(-_1, g(-_2)))^\dagger \\ &\stackrel{\text{def}}{=} \xi_{-1}(g(-_2)) \\ &\stackrel{\text{def}}{=} (g_*)^*(\xi_{-1}(-_2)) \\ &\stackrel{\text{def}}{=} (g_*)^*(\xi^\dagger) \\ &\stackrel{\text{def}}{=} [(g_*)^* \circ \Phi_{Y',Z}](\xi). \end{aligned}$$

- *Naturality II.* We need to show that, given a function $h: Z \rightarrow Z'$, the diagram

$$\begin{array}{ccc} \text{Hom}_{\text{Sets}}(X \times Y, Z) & \xrightarrow{\Phi_{Y,Z}} & \text{Hom}_{\text{Sets}}(X, \text{Hom}_{\text{Sets}}(Y, Z)), \\ h_* \downarrow & & \downarrow (h_*)_* \\ \text{Hom}_{\text{Sets}}(X \times Y, Z') & \xrightarrow{\Phi_{Y,Z'}} & \text{Hom}_{\text{Sets}}(X, \text{Hom}_{\text{Sets}}(Y, Z')), \end{array}$$

commutes. Indeed, given a morphism $\xi: X \times Y \rightarrow Z$, we have

$$\begin{aligned} [\Phi_{Y,Z} \circ h_*](\xi) &\stackrel{\text{def}}{=} (h(\xi(-_1, -_2)))^\dagger \\ &\stackrel{\text{def}}{=} [x \mapsto [y \mapsto h(\xi(x, y))]] \\ &\stackrel{\text{def}}{=} [x \mapsto h_*([y \mapsto \xi(x, y)])] \\ &\stackrel{\text{def}}{=} [x \mapsto h_*\left(\xi^\dagger(x)\right)] \\ &\stackrel{\text{def}}{=} h_*\left(\xi^\dagger\right) \\ &\stackrel{\text{def}}{=} [(h_*)_* \circ \Phi_{Y,Z}](\xi). \end{aligned}$$

- *Invertibility I.* We claim that

$$\Psi_{X,Y} \circ \Phi_{X,Y} = \text{id}_{\text{Hom}_{\text{Sets}}(X \times Y, Z)}.$$

Indeed, given a morphism $\xi: X \times Y \rightarrow Z$, we have

$$\begin{aligned} [\Psi_{X,Y} \circ \Phi_{X,Y}](\xi) &\stackrel{\text{def}}{=} \Psi_{X,Y}(\Phi_{X,Y}(\xi)) \\ &\stackrel{\text{def}}{=} \Psi_{X,Y}([x \mapsto [y \mapsto \xi(x, y)]]) \\ &\stackrel{\text{def}}{=} [(x, y) \mapsto \text{ev}_x([x \mapsto \text{ev}_y([y \mapsto \xi(x, y)])])] \\ &\stackrel{\text{def}}{=} [(x, y) \mapsto \text{ev}_x([x \mapsto \xi(x, y)])] \\ &\stackrel{\text{def}}{=} [(x, y) \mapsto \xi(x, y)] \\ &\stackrel{\text{def}}{=} \xi. \end{aligned}$$

- *Invertibility II.* We claim that

$$\Phi_{X,Y} \circ \Psi_{X,Y} = \text{id}_{\text{Hom}_{\text{Sets}}(X, \text{Hom}_{\text{Sets}}(Y, Z))}.$$

Indeed, given a morphism $\xi: X \rightarrow \text{Hom}_{\text{Sets}}(Y, Z)$, we have

$$\begin{aligned} [\Phi_{X,Y} \circ \Psi_{X,Y}](\xi) &\stackrel{\text{def}}{=} \Phi_{X,Y}(\Psi_{X,Y}(\xi)) \\ &\stackrel{\text{def}}{=} \Phi_{X,Y}(\Psi_{X,Y}([x \mapsto \xi(x)])) \\ &\stackrel{\text{def}}{=} \Phi_{X,Y}([(x, y) \mapsto \text{ev}_x([x \mapsto \text{ev}_y(\xi(x))])]) \\ &\stackrel{\text{def}}{=} \Phi_{X,Y}([(x, y) \mapsto \xi(x, y)]) \\ &\stackrel{\text{def}}{=} [x \mapsto [y \mapsto \xi(x, y)]] \\ &\stackrel{\text{def}}{=} \xi. \end{aligned}$$

Item 3, Associativity: See [Pro24a].

Item 4, Unitality: Clear.

Item 5, Commutativity: See [Pro24b].

Item 6, Annihilation With the Empty Set: See [Pro24f].

Item 7, Distributivity Over Unions: See [Pro24e].

Item 8, Distributivity Over Intersections: See [Pro24g, Corollary 1].

Item 9, Middle-Four Exchange With Respect to Intersections: See [Pro24g, Corollary 1].

Item 10, Distributivity Over Differences: See [Pro24c].

Item 11, Distributivity Over Symmetric Differences: See [Pro24d].

Item 12, Symmetric Monoidality: See [MO 382264].

Item 13, Symmetric Bimonoidality: Omitted. □

2.1.3 Pullbacks

Let A , B , and C be sets and let $f: A \rightarrow C$ and $g: B \rightarrow C$ be functions.

0012 Definition 2.1.3.1.1. The **pullback of A and B over C along f and g** ⁶ is the pair⁷ $(A \times_C B, \{\text{pr}_1, \text{pr}_2\})$ consisting of

- *The Limit.* The set $A \times_C B$ defined by

$$A \times_C B \stackrel{\text{def}}{=} \{(a, b) \in A \times B \mid f(a) = g(b)\}.$$

- *The Cone.* The maps

$$\text{pr}_1: A \times_C B \rightarrow A,$$

$$\text{pr}_2: A \times_C B \rightarrow B$$

defined by

$$\begin{aligned} \text{pr}_1(a, b) &\stackrel{\text{def}}{=} a, \\ \text{pr}_2(a, b) &\stackrel{\text{def}}{=} b \end{aligned}$$

for each $(a, b) \in A \times_C B$.

Proof. We claim that $A \times_C B$ is the categorical pullback of A and B over C with respect to (f, g) in Sets. First we need to check that the relevant pullback diagram commutes, i.e. that we have

$$\begin{array}{ccc} A \times_C B & \xrightarrow{\text{pr}_2} & B \\ \text{pr}_1 \downarrow & & \downarrow g \\ A & \xrightarrow{f} & C. \end{array}$$

Indeed, given $(a, b) \in A \times_C B$, we have

$$\begin{aligned} [f \circ \text{pr}_1](a, b) &= f(\text{pr}_1(a, b)) \\ &= f(a) \\ &= g(b) \\ &= g(\text{pr}_2(a, b)) \\ &= [g \circ \text{pr}_2](a, b), \end{aligned}$$

where $f(a) = g(b)$ since $(a, b) \in A \times_C B$. Next, we prove that $A \times_C B$ satisfies the universal property of the pullback. Suppose we have a diagram of the form

$$\begin{array}{ccccc} P & \xrightarrow{p_2} & A \times_C B & \xrightarrow{\text{pr}_2} & B \\ p_1 \swarrow & & \downarrow \lrcorner & & \downarrow g \\ A & \xrightarrow{f} & C & & \end{array}$$

⁶Further Terminology: Also called the **fibre product of A and B over C along f and g** .

⁷Further Notation: Also written $A \times_{f,C,g} B$.

in Sets. Then there exists a unique map $\phi: P \rightarrow A \times_C B$, uniquely determined by the conditions

$$\begin{aligned}\text{pr}_1 \circ \phi &= p_1, \\ \text{pr}_2 \circ \phi &= p_2,\end{aligned}$$

being necessarily given by

$$\phi(x) = (p_1(x), p_2(x))$$

for each $x \in P$, where we note that $(p_1(x), p_2(x)) \in A \times B$ indeed lies in $A \times_C B$ by the condition

$$f \circ p_1 = g \circ p_2,$$

which gives

$$f(p_1(x)) = g(p_2(x))$$

for each $x \in P$, so that $(p_1(x), p_2(x)) \in A \times_C B$. □

0013 Example 2.1.3.1.2. Here are some examples of pullbacks of sets.

0014 1. *Unions via Intersections.* Let $A, B \subset X$. We have a bijection of sets

$$\begin{array}{ccc} A \cap B & \longrightarrow & B \\ \downarrow \lrcorner & & \downarrow \iota_B \\ A & \xrightarrow{\iota_A} & A \cup B. \end{array}$$

Proof. **Item 1, Unions via Intersections:** Indeed, we have

$$\begin{aligned}A \times_{A \cup B} B &\cong \{(x, y) \in A \times B \mid x = y\} \\ &\cong A \cap B.\end{aligned}$$

This finishes the proof. □

0015 Proposition 2.1.3.1.3. Let A, B, C , and X be sets.

0016 1. *Functoriality.* The assignment $(A, B, C, f, g) \mapsto A \times_{f, C, g} B$ defines a functor

$$-_1 \times_{-3} -_1: \mathbf{Fun}(\mathcal{P}, \mathbf{Sets}) \rightarrow \mathbf{Sets},$$

where \mathcal{P} is the category that looks like this:

$$\begin{array}{ccc} & \bullet & \\ & \downarrow & \\ \bullet & \longrightarrow & \bullet. \end{array}$$

In particular, the action on morphisms of $-_1 \times_{-3} -_1$ is given by sending a morphism

$$\begin{array}{ccc}
 A \times_C B & \xrightarrow{\quad} & B \\
 \downarrow & \lrcorner & \downarrow g \quad \searrow \psi \\
 A' \times_{C'} B' & \xrightarrow{\quad} & B' \\
 \downarrow & \lrcorner & \downarrow \\
 A & \xrightarrow{f} & C \\
 \downarrow & \phi \searrow & \downarrow g' \quad \searrow \chi \\
 A' & \xrightarrow{f'} & C'
 \end{array}$$

in $\text{Fun}(\mathcal{P}, \text{Sets})$ to the map $\xi: A \times_C B \xrightarrow{\exists!} A' \times_{C'} B'$ given by

$$\xi(a, b) \stackrel{\text{def}}{=} (\phi(a), \psi(b))$$

for each $(a, b) \in A \times_C B$, which is the unique map making the diagram

$$\begin{array}{ccc}
 A \times_C B & \xrightarrow{\quad} & B \\
 \downarrow & \lrcorner \searrow & \downarrow g \quad \searrow \psi \\
 A' \times_{C'} B' & \xrightarrow{\quad} & B' \\
 \downarrow & \lrcorner & \downarrow \\
 A & \xrightarrow{f} & C \\
 \downarrow & \phi \searrow & \downarrow g' \quad \searrow \chi \\
 A' & \xrightarrow{f'} & C'
 \end{array}$$

commute.

0017 2. *Associativity.* Given a diagram

$$\begin{array}{ccccc}
 A & & B & & C \\
 & \searrow f & \swarrow g & \searrow h & \swarrow k \\
 & X & & Y &
 \end{array}$$

in Sets , we have isomorphisms

$$(A \times_X B) \times_Y C \cong (A \times_X B) \times_B (B \times_Y C) \cong A \times_X (B \times_Y C),$$

where these pullbacks are built as in the diagrams

$$\begin{array}{ccc}
 \begin{array}{c} (A \times_X B) \times_Y C \\ \downarrow \quad \searrow \\ A \times_X B \quad \searrow \\ \downarrow \quad \searrow \\ A \quad B \quad C \\ \searrow f \quad \searrow g \quad \searrow h \quad \searrow k \\ X \quad Y \quad C \end{array} & \begin{array}{c} (A \times_X B) \times_B (B \times_Y C) \\ \downarrow \quad \searrow \\ A \times_X B \quad \searrow \\ \downarrow \quad \searrow \\ A \quad B \quad B \times_Y C \quad C \\ \searrow f \quad \searrow g \quad \searrow h \quad \searrow k \\ X \quad Y \quad C \end{array} & \begin{array}{c} A \times_X (B \times_Y C) \\ \downarrow \quad \searrow \\ B \times_Y C \quad C \\ \downarrow \quad \searrow \\ B \quad C \\ \searrow h \quad \searrow k \\ Y \quad C \end{array} \\
 \end{array}$$

0018 3. *Unitality.* We have isomorphisms of sets

$$\begin{array}{ccc} A & \xlongequal{\quad} & A \\ f \downarrow & \lrcorner & \downarrow f \\ X & \xlongequal{\quad} & X \end{array} \quad \begin{array}{c} X \times_X A \cong A, \\ A \times_X X \cong A, \end{array} \quad \begin{array}{ccc} A & \xrightarrow{f} & X \\ \parallel & \lrcorner & \parallel \\ X & \xrightarrow{f} & X. \end{array}$$

0019 4. *Commutativity.* We have an isomorphism of sets

$$\begin{array}{ccc} A \times_X B & \longrightarrow & B \\ \downarrow & \lrcorner & \downarrow g \\ A & \xrightarrow{f} & X, \end{array} \quad A \times_X B \cong B \times_X A \quad \begin{array}{ccc} B \times_X A & \longrightarrow & A \\ \downarrow & \lrcorner & \downarrow f \\ B & \xrightarrow{g} & X. \end{array}$$

001A 5. *Annihilation With the Empty Set.* We have isomorphisms of sets

$$\begin{array}{ccc} \emptyset & \longrightarrow & \emptyset \\ \downarrow & \lrcorner & \downarrow \\ A & \xrightarrow{f} & X, \end{array} \quad \begin{array}{c} A \times_X \emptyset \cong \emptyset, \\ \emptyset \times_X A \cong \emptyset, \end{array} \quad \begin{array}{ccc} \emptyset & \longrightarrow & A \\ \downarrow & \lrcorner & \downarrow f \\ \emptyset & \longrightarrow & X. \end{array}$$

001B 6. *Interaction With Products.* We have

$$\begin{array}{ccc} A \times B & \longrightarrow & B \\ \downarrow & \lrcorner & \downarrow !_B \\ A & \xrightarrow{!_A} & \text{pt.} \end{array}$$

$$A \times_{\text{pt}} B \cong A \times B,$$

001C 7. *Symmetric Monoidality.* The triple $(\text{Sets}, \times_X, X)$ is a symmetric monoidal category.

Proof. Item 1, Functoriality: This is a special case of functoriality of co/limits, ?? of ??, with the explicit expression for ξ following from the commutativity of the cube pullback diagram.

Item 2, Associativity: Indeed, we have

$$\begin{aligned} (A \times_X B) \times_Y C &\cong \{((a, b), c) \in (A \times_X B) \times C \mid h(b) = k(c)\} \\ &\cong \{((a, b), c) \in (A \times B) \times C \mid f(a) = g(b) \text{ and } h(b) = k(c)\} \\ &\cong \{(a, (b, c)) \in A \times (B \times C) \mid f(a) = g(b) \text{ and } h(b) = k(c)\} \\ &\cong \{(a, (b, c)) \in A \times (B \times_Y C) \mid f(a) = g(b)\} \\ &\cong A \times_X (B \times_Y C) \end{aligned}$$

and

$$\begin{aligned}
 (A \times_X B) \times_B (B \times_Y C) &\cong \{(a, b), (b', c) \} \in (A \times_X B) \times (B \times_Y C) \mid b = b'\} \\
 &\cong \left\{ (a, b), (b', c) \in (A \times B) \times (B \times C) \mid \begin{array}{l} f(a) = g(b), b = b', \\ \text{and } h(b') = k(c) \end{array} \right\} \\
 &\cong \left\{ (a, (b, (b', c))) \in A \times (B \times (B \times C)) \mid \begin{array}{l} f(a) = g(b), b = b', \\ \text{and } h(b') = k(c) \end{array} \right\} \\
 &\cong \left\{ (a, ((b, b'), c)) \in A \times ((B \times B) \times C) \mid \begin{array}{l} f(a) = g(b), b = b', \\ \text{and } h(b') = k(c) \end{array} \right\} \\
 &\cong \left\{ (a, ((b, b'), c)) \in A \times ((B \times_B B) \times C) \mid \begin{array}{l} f(a) = g(b) \text{ and} \\ h(b') = k(c) \end{array} \right\} \\
 &\cong \{(a, (b, c)) \in A \times (B \times C) \mid f(a) = g(b) \text{ and } h(b) = k(c)\} \\
 &\cong A \times_X (B \times_Y C),
 \end{aligned}$$

where we have used [Item 3](#) for the isomorphism $B \times_B B \cong B$.

[Item 3, Unitality](#): Indeed, we have

$$\begin{aligned}
 X \times_X A &\cong \{(x, a) \in X \times A \mid f(a) = x\}, \\
 A \times_X X &\cong \{(a, x) \in X \times A \mid f(a) = x\},
 \end{aligned}$$

which are isomorphic to A via the maps $(x, a) \mapsto a$ and $(a, x) \mapsto a$.

[Item 4, Commutativity](#): Clear.

[Item 5, Annihilation With the Empty Set](#): Clear.

[Item 6, Interaction With Products](#): Clear.

[Item 7, Symmetric Monoidality](#): Omitted. □

2.1.4 Equalisers

Let A and B be sets and let $f, g: A \rightrightarrows B$ be functions.

001E Definition 2.1.4.1.1. The **equaliser of f and g** is the pair $(\text{Eq}(f, g), \text{eq}(f, g))$ consisting of

- *The Limit.* The set $\text{Eq}(f, g)$ defined by

$$\text{Eq}(f, g) \stackrel{\text{def}}{=} \{a \in A \mid f(a) = g(a)\}.$$

- *The Cone.* The inclusion map

$$\text{eq}(f, g): \text{Eq}(f, g) \hookrightarrow A.$$

Proof. We claim that $\text{Eq}(f, g)$ is the categorical equaliser of f and g in **Sets**. First we need to check that the relevant equaliser diagram commutes, i.e. that we have

$$f \circ \text{eq}(f, g) = g \circ \text{eq}(f, g),$$

which indeed holds by the definition of the set $\text{Eq}(f, g)$. Next, we prove that $\text{Eq}(f, g)$ satisfies the universal property of the equaliser. Suppose we have a diagram of the form

$$\begin{array}{ccccc} \text{Eq}(f, g) & \xrightarrow{\text{eq}(f, g)} & A & \xrightleftharpoons[f]{g} & B \\ & \nearrow e & & & \\ E & & & & \end{array}$$

in Sets. Then there exists a unique map $\phi: E \rightarrow \text{Eq}(f, g)$, uniquely determined by the condition

$$\text{eq}(f, g) \circ \phi = e$$

being necessarily given by

$$\phi(x) = e(x)$$

for each $x \in E$, where we note that $e(x) \in A$ indeed lies in $\text{Eq}(f, g)$ by the condition

$$f \circ e = g \circ e,$$

which gives

$$f(e(x)) = g(e(x))$$

for each $x \in E$, so that $e(x) \in \text{Eq}(f, g)$. □

001F Proposition 2.1.4.1.2. Let A, B , and C be sets.

001G 1. *Associativity.* We have an isomorphism of sets⁸

$$\underbrace{\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h))}_{=\text{Eq}(f \circ \text{eq}(g, h), h \circ \text{eq}(g, h))} \cong \text{Eq}(f, g, h) \cong \underbrace{\text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g))}_{=\text{Eq}(g \circ \text{eq}(f, g), h \circ \text{eq}(f, g))},$$

where $\text{Eq}(f, g, h)$ is the limit of the diagram

$$A \xrightarrow{\begin{matrix} f \\ -g \\ h \end{matrix}} B$$

in Sets , being explicitly given by

$$\text{Eq}(f, g, h) \cong \{a \in A \mid f(a) = g(a) = h(a)\}.$$

001H 4. *Unitality.* We have an isomorphism of sets

$$\text{Eq}(f, f) \cong A.$$

⁸That is, the following three ways of forming “the” equaliser of (f, g, h) agree:

1. Take the equaliser of (f, g, h) , i.e. the limit of the diagram

$$A \xrightarrow{\begin{matrix} f \\ -g \\ h \end{matrix}} B$$

in Sets .

2. First take the equaliser of f and g , forming a diagram

$$\text{Eq}(f, g) \xrightarrow{\text{eq}(f, g)} A \xrightarrow{\begin{matrix} f \\ g \end{matrix}} B$$

and then take the equaliser of the composition

$$\text{Eq}(f, g) \xrightarrow{\text{eq}(f, g)} A \xrightarrow{\begin{matrix} f \\ h \end{matrix}} B,$$

obtaining a subset

$$\text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g)) = \text{Eq}(g \circ \text{eq}(f, g), h \circ \text{eq}(f, g))$$

of $\text{Eq}(f, g)$.

3. First take the equaliser of g and h , forming a diagram

$$\text{Eq}(g, h) \xrightarrow{\text{eq}(g, h)} A \xrightarrow{\begin{matrix} g \\ h \end{matrix}} B$$

and then take the equaliser of the composition

$$\text{Eq}(g, h) \xrightarrow{\text{eq}(g, h)} A \xrightarrow{\begin{matrix} f \\ g \end{matrix}} B,$$

obtaining a subset

$$\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h)) = \text{Eq}(f \circ \text{eq}(g, h), h \circ \text{eq}(g, h))$$

of $\text{Eq}(g, h)$.

001J 5. *Commutativity.* We have an isomorphism of sets

$$\text{Eq}(f, g) \cong \text{Eq}(g, f).$$

001K 6. *Interaction With Composition.* Let

$$\begin{array}{c} A \xrightarrow{f} B \xrightarrow{h} C \\ \downarrow g \qquad \downarrow k \end{array}$$

be functions. We have an inclusion of sets

$$\text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g)) \subset \text{Eq}(h \circ f, k \circ g),$$

where $\text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g))$ is the equaliser of the composition

$$\begin{array}{c} \text{Eq}(f, g) \xrightarrow{\text{eq}(f, g)} A \xrightarrow{f} B \xrightarrow{h} C \\ \downarrow g \qquad \downarrow k \end{array}$$

Proof. **Item 1, Associativity:** We first prove that $\text{Eq}(f, g, h)$ is indeed given by

$$\text{Eq}(f, g, h) \cong \{a \in A \mid f(a) = g(a) = h(a)\}.$$

Indeed, suppose we have a diagram of the form

$$\begin{array}{ccc} \text{Eq}(f, g, h) & \xrightarrow{\text{eq}(f, g, h)} & A \xrightarrow{\begin{array}{c} f \\ g \\ h \end{array}} B \\ & \nearrow e & \\ E & & \end{array}$$

in Sets. Then there exists a unique map $\phi: E \rightarrow \text{Eq}(f, g, h)$, uniquely determined by the condition

$$\text{eq}(f, g) \circ \phi = e$$

being necessarily given by

$$\phi(x) = e(x)$$

for each $x \in E$, where we note that $e(x) \in A$ indeed lies in $\text{Eq}(f, g, h)$ by the condition

$$f \circ e = g \circ e = h \circ e,$$

which gives

$$f(e(x)) = g(e(x)) = h(e(x))$$

for each $x \in E$, so that $e(x) \in \text{Eq}(f, g, h)$.

We now check the equalities

$$\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h)) \cong \text{Eq}(f, g, h) \cong \text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g)).$$

Indeed, we have

$$\begin{aligned}\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h)) &\cong \{x \in \text{Eq}(g, h) \mid [f \circ \text{eq}(g, h)](a) = [g \circ \text{eq}(g, h)](a)\} \\ &\cong \{x \in \text{Eq}(g, h) \mid f(a) = g(a)\} \\ &\cong \{x \in A \mid f(a) = g(a) \text{ and } g(a) = h(a)\} \\ &\cong \{x \in A \mid f(a) = g(a) = h(a)\} \\ &\cong \text{Eq}(f, g, h).\end{aligned}$$

Similarly, we have

$$\begin{aligned}\text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g)) &\cong \{x \in \text{Eq}(f, g) \mid [f \circ \text{eq}(f, g)](a) = [h \circ \text{eq}(f, g)](a)\} \\ &\cong \{x \in \text{Eq}(f, g) \mid f(a) = h(a)\} \\ &\cong \{x \in A \mid f(a) = h(a) \text{ and } f(a) = g(a)\} \\ &\cong \{x \in A \mid f(a) = g(a) = h(a)\} \\ &\cong \text{Eq}(f, g, h).\end{aligned}$$

Item 4, Unitality: Clear.

Item 5, Commutativity: Clear.

Item 6, Interaction With Composition: Indeed, we have

$$\begin{aligned}\text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g)) &\cong \{a \in \text{Eq}(f, g) \mid h(f(a)) = k(g(a))\} \\ &\cong \{a \in A \mid f(a) = g(a) \text{ and } h(f(a)) = k(g(a))\}.\end{aligned}$$

and

$$\text{Eq}(h \circ f, k \circ g) \cong \{a \in A \mid h(f(a)) = k(g(a))\},$$

and thus there's an inclusion from $\text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g))$ to $\text{Eq}(h \circ f, k \circ g)$. \square

2.2 Colimits of Sets

2.2.1 Coproducts of Families of Sets

Let $\{A_i\}_{i \in I}$ be a family of sets.

001N Definition 2.2.1.1.1. The **disjoint union of the family** $\{A_i\}_{i \in I}$ is the pair $(\coprod_{i \in I} A_i, \{\text{inj}_i\}_{i \in I})$ consisting of

- *The Colimit.* The set $\coprod_{i \in I} A_i$ defined by

$$\coprod_{i \in I} A_i \stackrel{\text{def}}{=} \left\{ (i, x) \in I \times \left(\bigcup_{i \in I} A_i \right) \mid x \in A_i \right\}.$$

- *The Cocone.* The collection

$$\left\{ \text{inj}_i: A_i \rightarrow \coprod_{i \in I} A_i \right\}_{i \in I}$$

of maps given by

$$\text{inj}_i(x) \stackrel{\text{def}}{=} (i, x)$$

for each $x \in A_i$ and each $i \in I$.

Proof. We claim that $\coprod_{i \in I} A_i$ is the categorical coproduct of $\{A_i\}_{i \in I}$ in Sets. Indeed, suppose we have, for each $i \in I$, a diagram of the form

$$\begin{array}{ccc} & & C \\ & i_i \nearrow & \swarrow \\ A_i & \xrightarrow{\text{inj}_i} & \coprod_{i \in I} A_i \end{array}$$

in Sets. Then there exists a unique map $\phi: \coprod_{i \in I} A_i \rightarrow C$, uniquely determined by the condition $\phi \circ \text{inj}_i = i_i$ for each $i \in I$, being necessarily given by

$$\phi(i, x) = i_i(x)$$

for each $(i, x) \in \coprod_{i \in I} A_i$. □

001P Proposition 2.2.1.1.2. Let $\{A_i\}_{i \in I}$ be a family of sets.

001Q 1. Functoriality. The assignment $\{A_i\}_{i \in I} \mapsto \coprod_{i \in I} A_i$ defines a functor

$$\coprod_{i \in I}: \text{Fun}(I_{\text{disc}}, \text{Sets}) \rightarrow \text{Sets}$$

where

- *Action on Objects.* For each $(A_i)_{i \in I} \in \text{Obj}(\text{Fun}(I_{\text{disc}}, \text{Sets}))$, we have

$$\left[\coprod_{i \in I} \right] ((A_i)_{i \in I}) \stackrel{\text{def}}{=} \coprod_{i \in I} A_i$$

- *Action on Morphisms.* For each $(A_i)_{i \in I}, (B_i)_{i \in I} \in \text{Obj}(\text{Fun}(I_{\text{disc}}, \text{Sets}))$, the action on Hom-sets

$$\left(\coprod_{i \in I} \right)_{(A_i)_{i \in I}, (B_i)_{i \in I}} : \text{Nat}((A_i)_{i \in I}, (B_i)_{i \in I}) \rightarrow \text{Sets} \left(\coprod_{i \in I} A_i, \coprod_{i \in I} B_i \right)$$

of $\coprod_{i \in I}$ at $((A_i)_{i \in I}, (B_i)_{i \in I})$ is defined by sending a map

$$\{f_i: A_i \rightarrow B_i\}_{i \in I}$$

in $\text{Nat}((A_i)_{i \in I}, (B_i)_{i \in I})$ to the map of sets

$$\coprod_{i \in I} f_i: \coprod_{i \in I} A_i \rightarrow \coprod_{i \in I} B_i$$

defined by

$$\left[\coprod_{i \in I} f_i \right] (i, a) \stackrel{\text{def}}{=} f_i(a)$$

for each $(i, a) \in \coprod_{i \in I} A_i$.

Proof. **Item 1, Functoriality:** Clear. □

2.2.2 Binary Coproducts

Let A and B be sets.

001S Definition 2.2.2.1.1. The **coproduct⁹** of A and B is the pair $(A \coprod B, \{\text{inj}_1, \text{inj}_2\})$ consisting of

- *The Colimit.* The set $A \coprod B$ defined by

$$\begin{aligned} A \coprod B &\stackrel{\text{def}}{=} \coprod_{z \in \{A, B\}} z \\ &\cong \{(0, a) \mid a \in A\} \cup \{(1, b) \mid b \in B\}. \end{aligned}$$

- *The Cocone.* The maps

$$\begin{aligned} \text{inj}_1 : A &\rightarrow A \coprod B, \\ \text{inj}_2 : B &\rightarrow A \coprod B, \end{aligned}$$

given by

$$\begin{aligned} \text{inj}_1(a) &\stackrel{\text{def}}{=} (0, a), \\ \text{inj}_2(b) &\stackrel{\text{def}}{=} (1, b), \end{aligned}$$

for each $a \in A$ and each $b \in B$.

Proof. We claim that $A \coprod B$ is the categorical coproduct of A and B in Sets. Indeed, suppose we have a diagram of the form

$$\begin{array}{ccccc} & & C & & \\ & i_A \nearrow & & \swarrow i_B & \\ A & \xrightarrow{\text{inj}_A} & A \coprod B & \xleftarrow{\text{inj}_B} & B \end{array}$$

in Sets. Then there exists a unique map $\phi : A \coprod B \rightarrow C$, uniquely determined by the conditions

$$\begin{aligned} \phi \circ \text{inj}_A &= i_A, \\ \phi \circ \text{inj}_B &= i_B, \end{aligned}$$

being necessarily given by

$$\phi(x) = \begin{cases} i_A(x) & \text{if } x \in A, \\ i_B(x) & \text{if } x \in B \end{cases}$$

for each $x \in C$. □

⁹Further Terminology: Also called the **disjoint union** of A and B , or the **binary disjoint union** of A and B .

001T Proposition 2.2.2.1.2. Let A, B, C , and X be sets.

001U 1. *Functoriality.* The assignment $A, B, (A, B) \mapsto A \coprod B$ defines functors

$$\begin{aligned} A \coprod -_2 &: \text{Sets} \rightarrow \text{Sets}, \\ -_1 \coprod B &: \text{Sets} \rightarrow \text{Sets}, \\ -_1 \coprod -_2 &: \text{Sets} \times \text{Sets} \rightarrow \text{Sets}, \end{aligned}$$

where $-_1 \coprod -_2$ is the functor where

- *Action on Objects.* For each $(A, B) \in \text{Obj}(\text{Sets} \times \text{Sets})$, we have

$$[-_1 \coprod -_2](A, B) \stackrel{\text{def}}{=} A \coprod B;$$

- *Action on Morphisms.* For each $(A, B), (X, Y) \in \text{Obj}(\text{Sets})$, the action on Hom-sets

$$\coprod_{(A,B),(X,Y)} : \text{Sets}(A, X) \times \text{Sets}(B, Y) \rightarrow \text{Sets}(A \coprod B, X \coprod Y)$$

of \coprod at $((A, B), (X, Y))$ is defined by sending (f, g) to the function

$$f \coprod g : A \coprod B \rightarrow X \coprod Y$$

defined by

$$[f \coprod g](x) \stackrel{\text{def}}{=} \begin{cases} f(x) & \text{if } x \in A, \\ g(x) & \text{if } x \in B \end{cases}$$

for each $x \in A \coprod B$;

and where $A \coprod -$ and $- \coprod B$ are the partial functors of $-_1 \coprod -_2$ at $A, B \in \text{Obj}(\text{Sets})$.

001V 2. *Associativity.* We have an isomorphism of sets

$$(A \coprod B) \coprod C \cong A \coprod (B \coprod C),$$

natural in $A, B, C \in \text{Obj}(\text{Sets})$.

001W 3. *Unitality.* We have isomorphisms of sets

$$\begin{aligned} A \coprod \emptyset &\cong A, \\ \emptyset \coprod A &\cong A, \end{aligned}$$

natural in $A \in \text{Obj}(\text{Sets})$.

and B , for emphasis.

001X 4. *Commutativity.* We have an isomorphism of sets

$$A \coprod B \cong B \coprod A,$$

natural in $A, B \in \text{Obj}(\text{Sets})$.

001Y 5. *Symmetric Monoidality.* The triple $(\text{Sets}, \coprod, \emptyset)$ is a symmetric monoidal category.

Proof. **Item 1, Functoriality:** Clear.

Item 2, Associativity: Clear.

Item 3, Unitality: Clear.

Item 4, Commutativity: Clear.

Item 5, Symmetric Monoidality: Omitted. □

2.2.3 Pushouts

Let A, B , and C be sets and let $f: C \rightarrow A$ and $g: C \rightarrow B$ be functions.

0020 Definition 2.2.3.1.1. The **pushout of A and B over C along f and g** ¹⁰ is the pair $(A \coprod_C B, \{\text{inj}_1, \text{inj}_2\})$ consisting of

- *The Colimit.* The set $A \coprod_C B$ defined by

$$A \coprod_C B \stackrel{\text{def}}{=} A \coprod B / \sim_C,$$

where \sim_C is the equivalence relation on $A \coprod B$ generated by $(0, f(c)) \sim_C (1, g(c))$.

- *The Cocone.* The maps

$$\begin{aligned} \text{inj}_1: A &\rightarrow A \coprod_C B, \\ \text{inj}_2: B &\rightarrow A \coprod_C B \end{aligned}$$

given by

$$\begin{aligned} \text{inj}_1(a) &\stackrel{\text{def}}{=} [(0, a)] \\ \text{inj}_2(b) &\stackrel{\text{def}}{=} [(1, b)] \end{aligned}$$

for each $a \in A$ and each $b \in B$.

Proof. We claim that $A \coprod_C B$ is the categorical pushout of A and B over C with respect to (f, g) in Sets . First we need to check that the relevant pushout diagram commutes, i.e. that we have

$$\begin{array}{ccc} A \coprod_C B & \xleftarrow{\text{inj}_2} & B \\ \text{inj}_1 \uparrow & & \uparrow g \\ A & \xleftarrow{f} & C. \end{array}$$

¹⁰Further Terminology: Also called the **fibre coproduct of A and B over C along f and g** .

Indeed, given $c \in C$, we have

$$\begin{aligned} [\text{inj}_1 \circ f](c) &= \text{inj}_1(f(c)) \\ &= [(0, f(c))] \\ &= [(1, g(c))] \\ &= \text{inj}_2(g(c)) \\ &= [\text{inj}_2 \circ g](c), \end{aligned}$$

where $[(0, f(c))] = [(1, g(c))]$ by the definition of the relation \sim on B . Next, we prove that $A \coprod_C B$ satisfies the universal property of the pushout. Suppose we have a diagram of the form

$$\begin{array}{ccccc} & & P & & \\ & \swarrow i_1 & & \searrow i_2 & \\ A & \xrightarrow{\text{inj}_1} & A \coprod_C B & \xleftarrow{\text{inj}_2} & B \\ & \uparrow & \lrcorner & \uparrow g & \\ & & A & \xleftarrow{f} & C \end{array}$$

in Sets . Then there exists a unique map $\phi: A \coprod_C B \rightarrow P$, uniquely determined by the conditions

$$\begin{aligned} \phi \circ \text{inj}_1 &= i_1, \\ \phi \circ \text{inj}_2 &= i_2, \end{aligned}$$

being necessarily given by

$$\phi(x) = \begin{cases} i_1(a) & \text{if } x = [(0, a)], \\ i_2(b) & \text{if } x = [(1, b)] \end{cases}$$

for each $x \in A \coprod_C B$, where the well-definedness of ϕ is guaranteed by the equality $i_1 \circ f = i_2 \circ g$ and the definition of the relation \sim on $A \coprod B$ as follows.

1. *Case 1:* Suppose we have $x = [(0, a)] = [(0, a')]$ for some $a, a' \in A$. Then, by [Remark 2.2.3.1.2](#), we have a sequence

$$(0, a) \sim' x_1 \sim' \cdots \sim' x_n \sim' (0, a').$$

2. *Case 2:* Suppose we have $x = [(1, b)] = [(1, b')]$ for some $b, b' \in B$. Then, by [Remark 2.2.3.1.2](#), we have a sequence

$$(1, b) \sim' x_1 \sim' \cdots \sim' x_n \sim' (1, b').$$

3. *Case 3:* Suppose we have $x = [(0, a)] = [(1, b)]$ for some $a \in A$ and $b \in B$. Then, by Remark 2.2.3.1.2, we have a sequence

$$(0, a) \sim' x_1 \sim' \dots \sim' x_n \sim' (1, b).$$

In all these cases, we declare $x \sim' y$ iff there exists some $c \in C$ such that $x = (0, f(c))$ and $y = (1, g(c))$ or $x = (1, g(c))$ and $y = (0, f(c))$. Then, the equality $i_1 \circ f = i_2 \circ g$ gives

$$\begin{aligned} \phi([x]) &= \phi([(0, f(c))]) \\ &\stackrel{\text{def}}{=} i_1(f(c)) \\ &= i_2(g(c)) \\ &\stackrel{\text{def}}{=} \phi([(1, g(c))]) \\ &= \phi([y]), \end{aligned}$$

with the case where $x = (1, g(c))$ and $y = (0, f(c))$ similarly giving $\phi([x]) = \phi([y])$. Thus, if $x \sim' y$, then $\phi([x]) = \phi([y])$. Applying this equality pairwise to the sequences

$$\begin{aligned} (0, a) \sim' x_1 \sim' \dots \sim' x_n \sim' (0, a'), \\ (1, b) \sim' x_1 \sim' \dots \sim' x_n \sim' (1, b'), \\ (0, a) \sim' x_1 \sim' \dots \sim' x_n \sim' (1, b) \end{aligned}$$

gives

$$\begin{aligned} \phi([(0, a)]) &= \phi([(0, a')]), \\ \phi([(1, b)]) &= \phi([(1, b')]), \\ \phi([(0, a)]) &= \phi([(1, b)]), \end{aligned}$$

showing ϕ to be well-defined. \square

0021 Remark 2.2.3.1.2. In detail, by ??, the relation \sim of Definition 2.2.3.1.1 is given by declaring $a \sim b$ iff one of the following conditions is satisfied:

- We have $a, b \in A$ and $a = b$;
- We have $a, b \in B$ and $a = b$;
- There exist $x_1, \dots, x_n \in A \coprod B$ such that $a \sim' x_1 \sim' \dots \sim' x_n \sim' b$, where we declare $x \sim' y$ if one of the following conditions is satisfied:
 1. There exists $c \in C$ such that $x = (0, f(c))$ and $y = (1, g(c))$.
 2. There exists $c \in C$ such that $x = (1, g(c))$ and $y = (0, f(c))$.

That is: we require the following condition to be satisfied:

- (★) There exist $x_1, \dots, x_n \in A \coprod B$ satisfying the following conditions:

1. There exists $c_0 \in C$ satisfying one of the following conditions:
 - (a) We have $a = f(c_0)$ and $x_1 = g(c_0)$.
 - (b) We have $a = g(c_0)$ and $x_1 = f(c_0)$.
2. For each $1 \leq i \leq n - 1$, there exists $c_i \in C$ satisfying one of the following conditions:
 - (a) We have $x_i = f(c_i)$ and $x_{i+1} = g(c_i)$.
 - (b) We have $x_i = g(c_i)$ and $x_{i+1} = f(c_i)$.
3. There exists $c_n \in C$ satisfying one of the following conditions:
 - (a) We have $x_n = f(c_n)$ and $b = g(c_n)$.
 - (b) We have $x_n = g(c_n)$ and $b = f(c_n)$.

0022 Example 2.2.3.1.3. Here are some examples of pushouts of sets.

- 0023** 1. *Wedge Sums of Pointed Sets.* The wedge sum of two pointed sets of ?? is an example of a pushout of sets.
- 0024** 2. *Intersections via Unions.* Let $A, B \subset X$. We have a bijection of sets

$$\begin{array}{ccc} A \cup B & \xleftarrow{\quad} & B \\ \uparrow & \lrcorner & \uparrow \\ A \amalg_{A \cap B} B & & A \cap B. \end{array}$$

Proof. **Item 1, Wedge Sums of Pointed Sets:** Follows by definition.

Item 2, Intersections via Unions: Indeed, $A \amalg_{A \cap B} B$ is the quotient of $A \coprod B$ by the equivalence relation obtained by declaring $(0, a) \sim (1, b)$ iff $a = b \in A \cap B$, which is in bijection with $A \cup B$ via the map with $[(0, a)] \mapsto a$ and $[(1, b)] \mapsto b$. \square

0025 Proposition 2.2.3.1.4. Let A, B, C , and X be sets.

- 0026** 1. *Functionality.* The assignment $(A, B, C, f, g) \mapsto A \amalg_{f, g} B$ defines a functor

$$-_1 \amalg_{-_3} -_1 : \text{Fun}(\mathcal{P}, \text{Sets}) \rightarrow \text{Sets},$$

where \mathcal{P} is the category that looks like this:

$$\begin{array}{ccc} & \bullet & \\ & \uparrow & \\ \bullet & \longleftarrow & \bullet. \end{array}$$

In particular, the action on morphisms of $-_1 \amalg_{-_3} -_1$ is given by sending a mor-

phism

$$\begin{array}{ccccc}
 & A \coprod_C B & \xleftarrow{\quad \lrcorner \quad} & B & \\
 \uparrow & & & \uparrow \psi & \\
 & A' \coprod_{C'} B' & \xleftarrow{\quad \lrcorner \quad} & B' & \\
 \uparrow & & \downarrow g & & \uparrow g' \\
 & A & \xleftarrow{\quad f \quad} & C & \\
 \downarrow \phi & \downarrow & & \downarrow \chi & \downarrow \\
 A' & \xleftarrow{\quad f' \quad} & C' & &
 \end{array}$$

in $\text{Fun}(\mathcal{P}, \text{Sets})$ to the map $\xi: A \coprod_C B \xrightarrow{\exists!} A' \coprod_{C'} B'$ given by

$$\xi(x) \stackrel{\text{def}}{=} \begin{cases} \phi(a) & \text{if } x = [(0, a)], \\ \psi(b) & \text{if } x = [(1, b)] \end{cases}$$

for each $x \in A \coprod_C B$, which is the unique map making the diagram

$$\begin{array}{ccccc}
 & A \coprod_C B & \xleftarrow{\quad \lrcorner \quad} & B & \\
 \uparrow & & & \uparrow \psi & \\
 & A' \coprod_{C'} B' & \xleftarrow{\quad \lrcorner \quad} & B' & \\
 \uparrow & & \downarrow g & & \uparrow g' \\
 & A & \xleftarrow{\quad f \quad} & C & \\
 \downarrow \phi & \downarrow & & \downarrow \chi & \downarrow \\
 A' & \xleftarrow{\quad f' \quad} & C' & &
 \end{array}$$

commute.

0027 2. *Associativity.* Given a diagram

$$\begin{array}{ccccc}
 A & & B & & C \\
 \swarrow f & & \nearrow g & & \swarrow h \\
 X & & Y & & \nearrow k
 \end{array}$$

in Sets , we have isomorphisms

$$(A \coprod_X B) \coprod_Y C \cong (A \coprod_X B) \coprod_B (B \coprod_Y C) \cong A \coprod_X (B \coprod_Y C),$$

where these pullbacks are built as in the diagrams

$$\begin{array}{ccc}
 \begin{array}{c} (A \sqcup_X B) \sqcup_Y C \\ \uparrow \quad \swarrow \\ A \sqcup_X B \end{array} & \begin{array}{c} (A \sqcup_X B) \sqcup_B (B \sqcup_Y C) \\ \uparrow \quad \swarrow \\ A \sqcup_X B \quad B \sqcup_Y C \end{array} & \begin{array}{c} A \sqcup_X (B \sqcup_Y C) \\ \uparrow \quad \swarrow \\ B \sqcup_Y C \end{array} \\
 \begin{array}{ccccc} A & & & & C, \\ \nearrow f & \nearrow g & \nearrow h & \nearrow k & \nearrow \\ X & B & Y & C & \end{array} & \begin{array}{ccccc} A & & & & C, \\ \nearrow f & \nearrow g & \nearrow h & \nearrow k & \nearrow \\ X & B & Y & C & \end{array} & \begin{array}{ccccc} A & & & & C, \\ \nearrow f & \nearrow g & \nearrow h & \nearrow k & \nearrow \\ X & B & Y & C & \end{array}
 \end{array}$$

0028 3. *Unitality.* We have isomorphisms of sets

$$\begin{array}{ccc}
 \begin{array}{c} A \xlongequal{\quad} A \\ \uparrow f \qquad \downarrow f \\ X \xlongequal{\quad} X \end{array} & \begin{array}{c} X \sqcup_X A \cong A, \\ A \sqcup_X X \cong A, \end{array} & \begin{array}{c} A \xleftarrow{f} X \\ \parallel \qquad \parallel \\ X \xleftarrow{f} X. \end{array}
 \end{array}$$

0029 4. *Commutativity.* We have an isomorphism of sets

$$\begin{array}{ccc}
 \begin{array}{c} A \sqcup_X B \leftarrow B \\ \uparrow \lrcorner \qquad \uparrow g \\ A \xleftarrow{f} X, \end{array} & \begin{array}{c} A \sqcup_X B \cong B \sqcup_X A \end{array} & \begin{array}{c} B \sqcup_X A \leftarrow A \\ \uparrow \lrcorner \qquad \uparrow f \\ B \xleftarrow{g} X. \end{array}
 \end{array}$$

002A 5. *Interaction With Coproducts.* We have

$$\begin{array}{c}
 \begin{array}{c} A \sqcup B \leftarrow B \\ \uparrow \lrcorner \qquad \uparrow \iota_B \\ A \sqcup_{\emptyset} B \cong A \sqcup B, \end{array} \\
 \begin{array}{c} A \xleftarrow{\iota_A} \emptyset. \end{array}
 \end{array}$$

002B 6. *Symmetric Monoidality.* The triple $(\text{Sets}, \sqcup_X, \emptyset)$ is a symmetric monoidal category.

Proof. Item 1, Functoriality: This is a special case of functoriality of co/limits, ?? of ??, with the explicit expression for ξ following from the commutativity of the cube pushout diagram.

Item 2, Associativity: Omitted.

Item 3, Unitality: Omitted.

Item 4, Commutativity: Clear.

Item 5, Interaction With Coproducts: Clear.

Item 6, Symmetric Monoidality: Omitted. □

2.2.4 Coequalisers

Let A and B be sets and let $f, g: A \rightrightarrows B$ be functions.

002D Definition 2.2.4.1.1. The **coequaliser of f and g** is the pair $(\text{CoEq}(f, g), \text{coeq}(f, g))$ consisting of

- *The Colimit.* The set $\text{CoEq}(f, g)$ defined by

$$\text{CoEq}(f, g) \stackrel{\text{def}}{=} B/\sim,$$

where \sim is the equivalence relation on B generated by $f(a) \sim g(a)$.

- *The Cocone.* The map

$$\text{coeq}(f, g): B \rightarrow \text{CoEq}(f, g)$$

given by the quotient map $\pi: B \twoheadrightarrow B/\sim$ with respect to the equivalence relation generated by $f(a) \sim g(a)$.

Proof. We claim that $\text{CoEq}(f, g)$ is the categorical coequaliser of f and g in Sets. First we need to check that the relevant coequaliser diagram commutes, i.e. that we have

$$\text{coeq}(f, g) \circ f = \text{coeq}(f, g) \circ g.$$

Indeed, we have

$$\begin{aligned} [\text{coeq}(f, g) \circ f](a) &\stackrel{\text{def}}{=} [\text{coeq}(f, g)](f(a)) \\ &\stackrel{\text{def}}{=} [f(a)] \\ &= [g(a)] \\ &\stackrel{\text{def}}{=} [\text{coeq}(f, g)](g(a)) \\ &\stackrel{\text{def}}{=} [\text{coeq}(f, g) \circ g](a) \end{aligned}$$

for each $a \in A$. Next, we prove that $\text{CoEq}(f, g)$ satisfies the universal property of the coequaliser. Suppose we have a diagram of the form

$$\begin{array}{ccccc} & & f & & \\ & A & \rightrightarrows & B & \xleftarrow{\text{coeq}(f, g)} \\ & & g & & \\ & & & \searrow c & \\ & & & & C \end{array}$$

in Sets. Then, since $c(f(a)) = c(g(a))$ for each $a \in A$, it follows from ??? of ?? that there exists a unique map $\text{CoEq}(f, g) \xrightarrow{\exists!} C$ making the diagram

$$\begin{array}{ccccc} & & f & & \\ & A & \rightrightarrows & B & \xleftarrow{\text{coeq}(f, g)} \\ & & g & & \\ & & & \searrow c & \\ & & & & C \\ & & & \downarrow \exists! & \\ & & & & C \end{array}$$

commutes. □

002E Remark 2.2.4.1.2. In detail, by ??, the relation \sim of Definition 2.2.4.1.1 is given by declaring $a \sim b$ iff one of the following conditions is satisfied:

- We have $a = b$;
- There exist $x_1, \dots, x_n \in B$ such that $a \sim' x_1 \sim' \dots \sim' x_n \sim' b$, where we declare $x \sim' y$ if one of the following conditions is satisfied:
 1. There exists $z \in A$ such that $x = f(z)$ and $y = g(z)$.
 2. There exists $z \in A$ such that $x = g(z)$ and $y = f(z)$.

That is: we require the following condition to be satisfied:

(★) There exist $x_1, \dots, x_n \in B$ satisfying the following conditions:

1. There exists $z_0 \in A$ satisfying one of the following conditions:
 - (a) We have $a = f(z_0)$ and $x_1 = g(z_0)$.
 - (b) We have $a = g(z_0)$ and $x_1 = f(z_0)$.
2. For each $1 \leq i \leq n - 1$, there exists $z_i \in A$ satisfying one of the following conditions:
 - (a) We have $x_i = f(z_i)$ and $x_{i+1} = g(z_i)$.
 - (b) We have $x_i = g(z_i)$ and $x_{i+1} = f(z_i)$.
3. There exists $z_n \in A$ satisfying one of the following conditions:
 - (a) We have $x_n = f(z_n)$ and $b = g(z_n)$.
 - (b) We have $x_n = g(z_n)$ and $b = f(z_n)$.

002F Example 2.2.4.1.3. Here are some examples of coequalisers of sets.

002G 1. *Quotients by Equivalence Relations.* Let R be an equivalence relation on a set X . We have a bijection of sets

$$X/\sim_R \cong \text{CoEq}\left(R \hookrightarrow X \times X \xrightarrow[\text{pr}_2]{\text{pr}_1} X\right).$$

Proof. Item 1, Quotients by Equivalence Relations: See [Pro24z]. □

002H Proposition 2.2.4.1.4. Let A, B , and C be sets.

002J 1. *Associativity.* We have an isomorphism of sets¹¹

$$\underbrace{\text{CoEq}(\text{coeq}(f, g) \circ f, \text{coeq}(f, g) \circ h)}_{= \text{CoEq}(\text{coeq}(f, g) \circ g, \text{coeq}(f, g) \circ h)} \cong \text{CoEq}(f, g, h) \cong \underbrace{\text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ g)}_{= \text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ h)}$$

¹¹That is, the following three ways of forming “the” coequaliser of (f, g, h) agree:

where $\text{CoEq}(f, g, h)$ is the colimit of the diagram

$$A \xrightarrow{\begin{matrix} f \\ -g \\ h \end{matrix}} B$$

in Sets .

002K 4. *Unitality.* We have an isomorphism of sets

$$\text{CoEq}(f, f) \cong B.$$

002L 5. *Commutativity.* We have an isomorphism of sets

$$\text{CoEq}(f, g) \cong \text{CoEq}(g, f).$$

1. Take the coequaliser of (f, g, h) , i.e. the colimit of the diagram

$$A \xrightarrow{\begin{matrix} f \\ -g \\ h \end{matrix}} B$$

in Sets .

2. First take the coequaliser of f and g , forming a diagram

$$A \xrightarrow{\begin{matrix} f \\ g \end{matrix}} B \xrightarrow{\text{coeq}(f, g)} \text{CoEq}(f, g)$$

and then take the coequaliser of the composition

$$A \xrightarrow{\begin{matrix} f \\ h \end{matrix}} B \xrightarrow{\text{coeq}(f, g)} \text{CoEq}(f, g),$$

obtaining a quotient

$$\text{CoEq}(\text{coeq}(f, g) \circ f, \text{coeq}(f, g) \circ h) = \text{CoEq}(\text{coeq}(f, g) \circ g, \text{coeq}(f, g) \circ h)$$

of $\text{CoEq}(f, g)$

3. First take the coequaliser of g and h , forming a diagram

$$A \xrightarrow{\begin{matrix} g \\ h \end{matrix}} B \xrightarrow{\text{coeq}(g, h)} \text{CoEq}(g, h)$$

and then take the coequaliser of the composition

$$A \xrightarrow{\begin{matrix} f \\ g \end{matrix}} B \xrightarrow{\text{coeq}(g, h)} \text{CoEq}(g, h),$$

obtaining a quotient

$$\text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ g) = \text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ h)$$

of $\text{CoEq}(g, h)$.

002M 6. *Interaction With Composition.* Let

$$A \xrightarrow[g]{f} B \xrightarrow[k]{h} C$$

be functions. We have a surjection

$$\text{CoEq}(h \circ f, k \circ g) \twoheadrightarrow \text{CoEq}(\text{coeq}(h, k) \circ h \circ f, \text{coeq}(h, k) \circ k \circ g)$$

exhibiting $\text{CoEq}(\text{coeq}(h, k) \circ h \circ f, \text{coeq}(h, k) \circ k \circ g)$ as a quotient of $\text{CoEq}(h \circ f, k \circ g)$ by the relation generated by declaring $h(y) \sim k(y)$ for each $y \in B$.

Proof. **Item 1, Associativity:** Omitted.

Item 4, Unitality: Clear.

Item 5, Commutativity: Clear.

Item 6, Interaction With Composition: Omitted. □

2.3 Operations With Sets

2.3.1 The Empty Set

002Q Definition 2.3.1.1. The **empty set** is the set \emptyset defined by

$$\emptyset \stackrel{\text{def}}{=} \{x \in X \mid x \neq x\},$$

where A is the set in the set existence axiom, ?? of ??.

2.3.2 Singleton Sets

Let X be a set.

002S Definition 2.3.2.1. The **singleton set containing X** is the set $\{X\}$ defined by

$$\{X\} \stackrel{\text{def}}{=} \{X, X\},$$

where $\{X, X\}$ is the pairing of X with itself ([Definition 2.3.3.1.1](#)).

2.3.3 Pairings of Sets

Let X and Y be sets.

002U Definition 2.3.3.1.1. The **pairing of X and Y** is the set $\{X, Y\}$ defined by

$$\{X, Y\} \stackrel{\text{def}}{=} \{x \in A \mid x = X \text{ or } x = Y\},$$

where A is the set in the axiom of pairing, ?? of ??.

2.3.4 Ordered Pairs

Let A and B be sets.

002W Definition 2.3.4.1.1. The **ordered pair associated to A and B** is the set (A, B) defined by

$$(A, B) \stackrel{\text{def}}{=} \{\{A\}, \{A, B\}\}.$$

002X Proposition 2.3.4.1.2. Let A and B be sets.

002Y 1. *Uniqueness.* Let A, B, C , and D be sets. The following conditions are equivalent:

- (a) We have $(A, B) = (C, D)$.
- (b) We have $A = C$ and $B = D$.

Proof. **Item 1, Uniqueness:** See [Cie97, Theorem 1.2.3]. □

2.3.5 Unions of Families

Let $\{A_i\}_{i \in I}$ be a family of sets.

0030 Definition 2.3.5.1.1. The **union of the family** $\{A_i\}_{i \in I}$ is the set $\bigcup_{i \in I} A_i$ defined by

$$\bigcup_{i \in I} A_i \stackrel{\text{def}}{=} \{x \in F \mid \text{there exists some } i \in I \text{ such that } x \in A_i\},$$

where F is the set in the axiom of union, ?? of ??.

2.3.6 Binary Unions

Let A and B be sets.

0032 Definition 2.3.6.1.1. The **union¹² of A and B** is the set $A \cup B$ defined by

$$A \cup B \stackrel{\text{def}}{=} \bigcup_{z \in \{A, B\}} z.$$

0033 Proposition 2.3.6.1.2. Let X be a set.

0034 1. *Functionality.* The assignments $U, V, (U, V) \mapsto U \cup V$ define functors

$$\begin{aligned} U \cup -: (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ - \cup V: (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ -_1 \cup -_2: (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \subset) &\rightarrow (\mathcal{P}(X), \subset), \end{aligned}$$

where $-_1 \cup -_2$ is the functor where

¹²Further Terminology: Also called the **binary union of A and B** , for emphasis.

- *Action on Objects.* For each $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(X)$, we have

$$[-_1 \cup -_2](U, V) \stackrel{\text{def}}{=} U \cup V;$$

- *Action on Morphisms.* For each pair of morphisms

$$\begin{aligned} \iota_U &: U \hookrightarrow U', \\ \iota_V &: V \hookrightarrow V' \end{aligned}$$

of $\mathcal{P}(X) \times \mathcal{P}(X)$, the image

$$\iota_U \cup \iota_V : U \cup V \hookrightarrow U' \cup V'$$

of (ι_U, ι_V) by \cup is the inclusion

$$U \cup V \subset U' \cup V'$$

i.e. where we have

(★) If $U \subset U'$ and $V \subset V'$, then $U \cup V \subset U' \cup V'$;

and where $U \cup -$ and $- \cup V$ are the partial functors of $-_1 \cup -_2$ at $U, V \in \mathcal{P}(X)$.

0035 2. *Via Intersections and Symmetric Differences.* We have an equality of sets

$$U \cup V = (U \Delta V) \Delta (U \cap V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

0036 3. *Associativity.* We have an equality of sets

$$(U \cup V) \cup W = U \cup (V \cup W)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

0037 4. *Unitality.* We have equalities of sets

$$\begin{aligned} U \cup \emptyset &= U, \\ \emptyset \cup U &= U \end{aligned}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

0038 5. *Commutativity.* We have an equality of sets

$$U \cup V = V \cup U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

0039 6. *Idempotency.* We have an equality of sets

$$U \cup U = U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

003A 7. *Distributivity Over Intersections.* We have equalities of sets

$$\begin{aligned} U \cup (V \cap W) &= (U \cup V) \cap (U \cup W), \\ (U \cap V) \cup W &= (U \cup W) \cap (V \cup W) \end{aligned}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

003B 8. *Interaction With Characteristic Functions I.* We have

$$\chi_{U \cup V} = \max(\chi_U, \chi_V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

003C 9. *Interaction With Characteristic Functions II.* We have

$$\chi_{U \cup V} = \chi_U + \chi_V - \chi_{U \cap V}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

003D 10. *Interaction With Powersets and Semirings.* The quintuple $(\mathcal{P}(X), \cup, \cap, \emptyset, X)$ is an idempotent commutative semiring.

Proof. **Item 1**, Functoriality: See [Pro24an].

Item 2, Via Intersections and Symmetric Differences: See [Pro24ay].

Item 3, Associativity: See [Pro24ba].

Item 4, Unitality: This follows from [Pro24bd] and **Item 5**.

Item 5, Commutativity: See [Pro24bb].

Item 6, Idempotency: See [Pro24am].

Item 7, Distributivity Over Intersections: See [Pro24az].

Item 8, Interaction With Characteristic Functions I: See [Pro24k].

Item 9, Interaction With Characteristic Functions II: See [Pro24k].

Item 10, Interaction With Powersets and Semirings: This follows from **Items 3 to 6** and **Items 3 to 5, 7 and 8** of **Proposition 2.3.8.1.2.** \square

2.3.7 Intersections of Families

Let \mathcal{F} be a family of sets.

003F **Definition 2.3.7.1.1.** The **intersection of a family \mathcal{F} of sets** is the set $\bigcap_{X \in \mathcal{F}} X$ defined by

$$\bigcap_{X \in \mathcal{F}} X \stackrel{\text{def}}{=} \left\{ z \in \bigcup_{X \in \mathcal{F}} X \mid \text{for each } X \in \mathcal{F}, \text{ we have } z \in X \right\}.$$

2.3.8 Binary Intersections

Let X and Y be sets.

003H Definition 2.3.8.1.1. The **intersection**¹³ of X and Y is the set $X \cap Y$ defined by

$$X \cap Y \stackrel{\text{def}}{=} \bigcap_{z \in \{X, Y\}} z.$$

003J Proposition 2.3.8.1.2. Let X be a set.

003K 1. Functoriality. The assignments $U, V, (U, V) \mapsto U \cap V$ define functors

$$\begin{aligned} U \cap -: (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ - \cap V: (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ -_1 \cap -_2: (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \subset) &\rightarrow (\mathcal{P}(X), \subset), \end{aligned}$$

where $-_1 \cap -_2$ is the functor where

- *Action on Objects.* For each $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(X)$, we have

$$[-_1 \cap -_2](U, V) \stackrel{\text{def}}{=} U \cap V;$$

- *Action on Morphisms.* For each pair of morphisms

$$\begin{aligned} \iota_U: U &\hookrightarrow U', \\ \iota_V: V &\hookrightarrow V' \end{aligned}$$

of $\mathcal{P}(X) \times \mathcal{P}(X)$, the image

$$\iota_U \cap \iota_V: U \cap V \hookrightarrow U' \cap V'$$

of (ι_U, ι_V) by \cap is the inclusion

$$U \cap V \subset U' \cap V'$$

i.e. where we have

- (★) If $U \subset U'$ and $V \subset V'$, then $U \cap V \subset U' \cap V'$;

and where $U \cap -$ and $- \cap V$ are the partial functors of $-_1 \cap -_2$ at $U, V \in \mathcal{P}(X)$.

003L 2. Adjointness. We have adjunctions

$$\begin{aligned} (U \cap - \dashv \mathbf{Hom}_{\mathcal{P}(X)}(U, -)): \quad \mathcal{P}(X) &\xrightleftharpoons[\mathbf{Hom}_{\mathcal{P}(X)}(U, -)]{\perp} \mathcal{P}(X), \\ (- \cap V \dashv \mathbf{Hom}_{\mathcal{P}(X)}(V, -)): \quad \mathcal{P}(X) &\xrightleftharpoons[\mathbf{Hom}_{\mathcal{P}(X)}(V, -)]{\perp} \mathcal{P}(X), \end{aligned}$$

¹³Further Terminology: Also called the **binary intersection of X and Y** , for emphasis.

where

$$\mathbf{Hom}_{\mathcal{P}(X)}(-_1, -_2) : \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \rightarrow \mathcal{P}(X)$$

is the bifunctor defined by¹⁴

$$\mathbf{Hom}_{\mathcal{P}(X)}(U, V) \stackrel{\text{def}}{=} (X \setminus U) \cup V$$

witnessed by bijections

$$\begin{aligned} \text{Hom}_{\mathcal{P}(X)}(U \cap V, W) &\cong \text{Hom}_{\mathcal{P}(X)}(U, \mathbf{Hom}_{\mathcal{P}(X)}(V, W)), \\ \text{Hom}_{\mathcal{P}(X)}(U \cap V, W) &\cong \text{Hom}_{\mathcal{P}(X)}(V, \mathbf{Hom}_{\mathcal{P}(X)}(U, W)), \end{aligned}$$

natural in $U, V, W \in \mathcal{P}(X)$, i.e. where:

(a) The following conditions are equivalent:

- i. We have $U \cap V \subset W$.
- ii. We have $U \subset \mathbf{Hom}_{\mathcal{P}(X)}(V, W)$.
- iii. We have $U \subset (X \setminus V) \cup W$.

(b) The following conditions are equivalent:

- i. We have $V \cap U \subset W$.
- ii. We have $V \subset \mathbf{Hom}_{\mathcal{P}(X)}(U, W)$.
- iii. We have $V \subset (X \setminus U) \cup W$.

003M 3. *Associativity.* We have an equality of sets

$$(U \cap V) \cap W = U \cap (V \cap W)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

003N 4. *Unitality.* Let X be a set and let $U \in \mathcal{P}(X)$. We have equalities of sets

$$\begin{aligned} X \cap U &= U, \\ U \cap X &= U \end{aligned}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

003P 5. *Commutativity.* We have an equality of sets

$$U \cap V = V \cap U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

¹⁴*Intuition:* Since intersections are the products in $\mathcal{P}(X)$, the left adjoint $\mathbf{Hom}_{\mathcal{P}(X)}(U, V)$ works as a function type $U \rightarrow V$.

Now, under the Curry–Howard correspondence, the function type $U \rightarrow V$ corresponds to implication $U \implies V$, which is logically equivalent to the statement $\neg U \vee V$, which in turn corresponds to the set $U^c \vee V \stackrel{\text{def}}{=} (X \setminus U) \cup V$.

003Q 6. *Idempotency.* We have an equality of sets

$$U \cap U = U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

003R 7. *Distributivity Over Unions.* We have equalities of sets

$$\begin{aligned} U \cap (V \cup W) &= (U \cap V) \cup (U \cap W), \\ (U \cup V) \cap W &= (U \cap W) \cup (V \cap W) \end{aligned}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

003S 8. *Annihilation With the Empty Set.* We have an equality of sets

$$\begin{aligned} \emptyset \cap X &= \emptyset, \\ X \cap \emptyset &= \emptyset \end{aligned}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

003T 9. *Interaction With Characteristic Functions I.* We have

$$\chi_{U \cap V} = \chi_U \chi_V$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

003U 10. *Interaction With Characteristic Functions II.* We have

$$\chi_{U \cap V} = \min(\chi_U, \chi_V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

003V 11. *Interaction With Powersets and Monoids With Zero.* The quadruple $((\mathcal{P}(X), \emptyset), \cap, X)$ is a commutative monoid with zero.

003W 12. *Interaction With Powersets and Semirings.* The quintuple $(\mathcal{P}(X), \cup, \cap, \emptyset, X)$ is an idempotent commutative semiring.

Proof. **Item 1**, Functoriality: See [Pro24al].

Item 2, Adjointness: See [MSE 267469].

Item 3, Associativity: See [Pro24t].

Item 4, Unitality: This follows from [Pro24x] and **Item 5**.

Item 5, Commutativity: See [Pro24u].

Item 6, Idempotency: See [Pro24ak].

Item 7, Distributivity Over Unions: See [Pro24aj].

Item 8, Annihilation With the Empty Set: This follows from [Pro24v] and **Item 5**.

Item 9, Interaction With Characteristic Functions I: See [Pro24h].

Item 10, Interaction With Characteristic Functions II: See [Pro24h].

Item 11, Interaction With Powersets and Monoids With Zero: This follows from **Items 3** to **5** and **8**.

Item 12, Interaction With Powersets and Semirings: This follows from **Items 3** to **6** and **Items 3** to **5**, **7** and **8** of Proposition 2.3.8.1.2. \square

2.3.9 Differences

Let X and Y be sets.

003Y Definition 2.3.9.1.1. The **difference of X and Y** is the set $X \setminus Y$ defined by

$$X \setminus Y \stackrel{\text{def}}{=} \{a \in X \mid a \notin Y\}.$$

003Z Proposition 2.3.9.1.2. Let X be a set.

0040 1. *Functionality.* The assignments $U, V, (U, V) \mapsto U \cap V$ define functors

$$\begin{aligned} U \setminus - &: (\mathcal{P}(X), \supset) \rightarrow (\mathcal{P}(X), \subset), \\ - \setminus V &: (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(X), \subset), \\ -_1 \setminus -_2 &: (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \supset) \rightarrow (\mathcal{P}(X), \subset), \end{aligned}$$

where $-_1 \setminus -_2$ is the functor where

- *Action on Objects.* For each $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(X)$, we have

$$[-_1 \setminus -_2](U, V) \stackrel{\text{def}}{=} U \setminus V;$$

- *Action on Morphisms.* For each pair of morphisms

$$\begin{aligned} \iota_A &: A \hookrightarrow B, \\ \iota_U &: U \hookrightarrow V \end{aligned}$$

of $\mathcal{P}(X) \times \mathcal{P}(X)$, the image

$$\iota_U \setminus \iota_V: A \setminus V \hookrightarrow B \setminus U$$

of (ι_U, ι_V) by \setminus is the inclusion

$$A \setminus V \subset B \setminus U$$

i.e. where we have

- (★) If $A \subset B$ and $U \subset V$, then $A \setminus V \subset B \setminus U$;

and where $U \setminus -$ and $- \setminus V$ are the partial functors of $-_1 \setminus -_2$ at $U, V \in \mathcal{P}(X)$.

0041 2. *De Morgan's Laws.* We have equalities of sets

$$\begin{aligned} X \setminus (U \cup V) &= (X \setminus U) \cap (X \setminus V), \\ X \setminus (U \cap V) &= (X \setminus U) \cup (X \setminus V) \end{aligned}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

0042 3. *Interaction With Unions I.* We have equalities of sets

$$U \setminus (V \cup W) = (U \setminus V) \cap (U \setminus W)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

0043 4. *Interaction With Unions II.* We have equalities of sets

$$(U \setminus V) \cup W = (U \cup W) \setminus (V \setminus W)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

0044 5. *Interaction With Unions III.* We have equalities of sets

$$\begin{aligned} U \setminus (V \cup W) &= (U \cup W) \setminus (V \cup W) \\ &= (U \setminus V) \setminus W \\ &= (U \setminus W) \setminus V \end{aligned}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

0045 6. *Interaction With Unions IV.* We have equalities of sets

$$(U \cup V) \setminus W = (U \setminus W) \cup (V \setminus W)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

0046 7. *Interaction With Intersections.* We have equalities of sets

$$\begin{aligned} (U \setminus V) \cap W &= (U \cap W) \setminus V \\ &= U \cap (W \setminus V) \end{aligned}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

0047 8. *Interaction With Complements.* We have an equality of sets

$$U \setminus V = U \cap V^c$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

0048 9. *Interaction With Symmetric Differences.* We have an equality of sets

$$U \setminus V = U \Delta (U \cap V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

0049 10. *Triple Differences.* We have

$$U \setminus (V \setminus W) = (U \cap W) \cup (U \setminus V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

004A 11. *Left Annihilation.* We have

$$\emptyset \setminus U = \emptyset$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

004B 12. *Right Unitality.* We have

$$U \setminus \emptyset = U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

004C 13. *Invertibility.* We have

$$U \setminus U = \emptyset$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

004D 14. *Interaction With Containment.* The following conditions are equivalent:

- (a) We have $V \setminus U \subset W$.
- (b) We have $V \setminus W \subset U$.

004E 15. *Interaction With Characteristic Functions.* We have

$$\chi_{U \setminus V} = \chi_U - \chi_{U \cap V}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

Proof. **Item 1, Functoriality:** See [Pro24ad] and [Pro24ah].

Item 2, De Morgan's Laws: See [Pro24p].

Item 3, Interaction With Unions I: See [Pro24q].

Item 4, Interaction With Unions II: Omitted.

Item 5, Interaction With Unions III: See [Pro24ai].

Item 6, Interaction With Unions IV: See [Pro24ac].

Item 7, Interaction With Intersections: See [Pro24w].

Item 8, Interaction With Complements: See [Pro24aa].

Item 9, Interaction With Symmetric Differences: See [Pro24ab].

Item 10, Triple Differences: See [Pro24ag].

Item 11, Left Annihilation: Clear.

Item 12, Right Unitality: See [Pro24ae].

Item 13, Invertibility: See [Pro24af].

Item 14, Interaction With Containment: Omitted.

Item 15, Interaction With Characteristic Functions: See [Pro24i].

□

2.3.10 Complements

Let X be a set and let $U \in \mathcal{P}(X)$.

004G Definition 2.3.10.1.1. The **complement** of U is the set U^c defined by

$$\begin{aligned} U^c &\stackrel{\text{def}}{=} X \setminus U \\ &\stackrel{\text{def}}{=} \{a \in X \mid a \notin U\}. \end{aligned}$$

004H Proposition 2.3.10.1.2. Let X be a set.

004J 1. Functoriality. The assignment $U \mapsto U^c$ defines a functor

$$(-)^c : \mathcal{P}(X)^{\text{op}} \rightarrow \mathcal{P}(X),$$

where

- *Action on Objects.* For each $U \in \mathcal{P}(X)$, we have

$$[(-)^c](U) \stackrel{\text{def}}{=} U^c;$$

- *Action on Morphisms.* For each morphism $\iota_U : U \hookrightarrow V$ of $\mathcal{P}(X)$, the image

$$\iota_U^c : V^c \hookrightarrow U^c$$

of ι_U by $(-)^c$ is the inclusion

$$V^c \subset U^c$$

i.e. where we have

- (★) If $U \subset V$, then $V^c \subset U^c$.

004K 2. De Morgan's Laws. We have equalities of sets

$$\begin{aligned} (U \cup V)^c &= U^c \cap V^c, \\ (U \cap V)^c &= U^c \cup V^c \end{aligned}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

004L 3. Involutority. We have

$$(U^c)^c = U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

004M 4. Interaction With Characteristic Functions. We have

$$\chi_{U^c} = 1 - \chi_U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

Proof. **Item 1, Functoriality:** This follows from [Item 1 of Proposition 2.3.9.1.2](#).

Item 2, De Morgan's Laws: See [\[Pro24p\]](#).

Item 3, Involutority: See [\[Pro24l\]](#).

Item 4, Interaction With Characteristic Functions: Clear. □

2.3.11 Symmetric Differences

Let A and B be sets.

004P Definition 2.3.11.1.1. The **symmetric difference of A and B** is the set $A \Delta B$ defined by

$$A \Delta B \stackrel{\text{def}}{=} (A \setminus B) \cup (B \setminus A).$$

004Q Proposition 2.3.11.1.2. Let X be a set.

004R 1. Lack of Functoriality. The assignment $(U, V) \mapsto U \Delta V$ **need not** define functors

$$\begin{aligned} U \Delta -_2 &: (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(X), \subset), \\ -_1 \Delta V &: (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(X), \subset), \\ -_1 \Delta -_2 &: (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \subset) \rightarrow (\mathcal{P}(X), \subset). \end{aligned}$$

004S 2. Via Unions and Intersections. We have¹⁵

$$U \Delta V = (U \cup V) \setminus (U \cap V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

004T 3. Associativity. We have¹⁶

$$(U \Delta V) \Delta W = U \Delta (V \Delta W)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

004U 4. Commutativity. We have

$$U \Delta V = V \Delta U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

004V 5. Unitality. We have

$$U \Delta \emptyset = U,$$

$$\emptyset \Delta U = U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

¹⁵Illustration:



¹⁶Illustration:



004W 6. *Invertibility.* We have

$$U \Delta U = \emptyset$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

004X 7. *Interaction With Unions.* We have

$$(U \Delta V) \cup (V \Delta T) = (U \cup V \cup W) \setminus (U \cap V \cap W)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

004Y 8. *Interaction With Complements I.* We have

$$U \Delta U^c = X$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

004Z 9. *Interaction With Complements II.* We have

$$\begin{aligned} U \Delta X &= U^c, \\ X \Delta U &= U^c \end{aligned}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

0050 10. *Interaction With Complements III.* We have

$$U^c \Delta V^c = U \Delta V$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

0051 11. “*Transitivity*”. We have

$$(U \Delta V) \Delta (V \Delta W) = U \Delta W$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

0052 12. *The Triangle Inequality for Symmetric Differences.* We have

$$U \Delta W \subset U \Delta V \cup V \Delta W$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

0053 13. *Distributivity Over Intersections.* We have

$$\begin{aligned} U \cap (V \Delta W) &= (U \cap V) \Delta (U \cap W), \\ (U \Delta V) \cap W &= (U \cap W) \Delta (V \cap W) \end{aligned}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V, W \in \mathcal{P}(X)$.

0054 14. *Interaction With Characteristic Functions.* We have

$$\chi_{U \Delta V} = \chi_U + \chi_V - 2\chi_{U \cap V}$$

and thus, in particular, we have

$$\chi_{U \Delta V} \equiv \chi_U + \chi_V \pmod{2}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

0055 15. *Bijection.* Given $A, B \subset \mathcal{P}(X)$, the maps

$$\begin{aligned} A \Delta - &: \mathcal{P}(X) \rightarrow \mathcal{P}(X), \\ - \Delta B &: \mathcal{P}(X) \rightarrow \mathcal{P}(X) \end{aligned}$$

are bijections with inverses given by

$$\begin{aligned} (A \Delta -)^{-1} &= - \cup (A \cap -), \\ (- \Delta B)^{-1} &= - \cup (B \cap -). \end{aligned}$$

Moreover, the map

$$C \mapsto C \Delta (A \Delta B)$$

is a bijection of $\mathcal{P}(X)$ onto itself sending A to B and B to A .

0056 16. *Interaction With Powersets and Groups.* Let X be a set.

0057 (a) The quadruple $(\mathcal{P}(X), \Delta, \emptyset, \text{id}_{\mathcal{P}(X)})$ is an abelian group.¹⁷

0058 (b) Every element of $\mathcal{P}(X)$ has order 2 with respect to Δ , and thus $\mathcal{P}(X)$ is a Boolean group (i.e. an abelian 2-group).

0059 4. *Interaction With Powersets and Vector Spaces I.* The pair $(\mathcal{P}(X), \alpha_{\mathcal{P}(X)})$ consisting of

- The group $\mathcal{P}(X)$ of ??;

¹⁷Here are some examples:

1. When $X = \emptyset$, we have an isomorphism of groups between $\mathcal{P}(\emptyset)$ and the trivial group:

$$(\mathcal{P}(\emptyset), \Delta, \emptyset, \text{id}_{\mathcal{P}(\emptyset)}) \cong \text{pt}.$$

2. When $X = \text{pt}$, we have an isomorphism of groups between $\mathcal{P}(\text{pt})$ and $\mathbb{Z}/2$:

$$(\mathcal{P}(\text{pt}), \Delta, \emptyset, \text{id}_{\mathcal{P}(\text{pt})}) \cong \mathbb{Z}/2.$$

3. When $X = \{0, 1\}$, we have an isomorphism of groups between $\mathcal{P}(\{0, 1\})$ and $\mathbb{Z}/2 \times \mathbb{Z}/2$:

$$(\mathcal{P}(\{0, 1\}), \Delta, \emptyset, \text{id}_{\mathcal{P}(\{0, 1\})}) \cong \mathbb{Z}/2 \times \mathbb{Z}/2.$$

- The map $\alpha_{\mathcal{P}(X)} : \mathbb{F}_2 \times \mathcal{P}(X) \rightarrow \mathcal{P}(X)$ defined by

$$\begin{aligned} 0 \cdot U &\stackrel{\text{def}}{=} \emptyset, \\ 1 \cdot U &\stackrel{\text{def}}{=} U; \end{aligned}$$

is an \mathbb{F}_2 -vector space.

005A 5. *Interaction With Powersets and Vector Spaces II.* If X is finite, then:

- (a) The set of singletons sets on the elements of X forms a basis for the \mathbb{F}_2 -vector space $(\mathcal{P}(X), \alpha_{\mathcal{P}(X)})$ of Item 4.
- (b) We have

$$\dim(\mathcal{P}(X)) = \#\mathcal{P}(X).$$

005B 6. *Interaction With Powersets and Rings.* The quintuple $(\mathcal{P}(X), \Delta, \cap, \emptyset, X)$ is a commutative ring.¹⁸

Proof. Item 1, Lack of Functoriality: Omitted.

Item 2, Via Unions and Intersections: See [Pro24r].

Item 3, Associativity: See [Pro24ao].

Item 4, Commutativity: See [Pro24ap].

Item 5, Unitality: This follows from Item 4 and [Pro24at].

Item 6, Invertibility: See [Pro24av].

Item 7, Interaction With Unions: See [Pro24bc].

Item 8, Interaction With Complements I: See [Pro24as].

Item 9, Interaction With Complements II: This follows from Item 4 and [Pro24ax].

Item 10, Interaction With Complements III: See [Pro24aq].

Item 11, "Transitivity": We have

$$\begin{aligned} (U \Delta V) \Delta (V \Delta W) &= U \Delta (V \Delta (V \Delta W)) && (\text{by Item 3}) \\ &= U \Delta ((V \Delta V) \Delta W) && (\text{by Item 3}) \\ &= U \Delta (\emptyset \Delta W) && (\text{by Item 6}) \\ &= U \Delta W && (\text{by Item 5}) \end{aligned}$$

Item 12, The Triangle Inequality for Symmetric Differences: This follows from Items 2 and 11.

Item 13, Distributivity Over Intersections: See [Pro24s].

Item 14, Interaction With Characteristic Functions: See [Pro24j].

Item 15, Bijectivity: Clear.

Item 16, Interaction With Powersets and Groups: Item 16a follows from¹⁹ Items 3 to 6, while Item 3b follows from Item 6.

¹⁸  Warning: The analogous statement replacing intersections by unions (i.e. that the quintuple $(\mathcal{P}(X), \Delta, \cup, \emptyset, X)$ is a ring) is false, however. See [Pro24aw] for a proof.

END TEXTDBEND

¹⁹ Reference: [Pro24ar].

Item 4, Interaction With Powersets and Vector Spaces I: Clear.

Item 5, Interaction With Powersets and Vector Spaces II: Omitted.

Item 6, Interaction With Powersets and Rings: This follows from Items 8 and 11 of Proposition 2.3.8.1.2 and Items 13 and 16.²⁰ \square

2.4 Powersets

2.4.1 Characteristic Functions

Let X be a set.

005E **Definition 2.4.1.1.1.** Let $U \subset X$ and let $x \in X$.

005F 1. The **characteristic function of U** ²¹ is the function²²

$$\chi_U: X \rightarrow \{\text{t, f}\}$$

defined by

$$\chi_U(x) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x \in U, \\ \text{false} & \text{if } x \notin U \end{cases}$$

for each $x \in X$.

005G 2. The **characteristic function of x** is the function²³

$$\chi_x: X \rightarrow \{\text{t, f}\}$$

defined by

$$\chi_x \stackrel{\text{def}}{=} \chi_{\{x\}},$$

i.e. by

$$\chi_x(y) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x = y, \\ \text{false} & \text{if } x \neq y \end{cases}$$

for each $y \in X$.

005H 3. The **characteristic relation on X** ²⁴ is the relation²⁵

$$\chi_X(-_1, -_2): X \times X \rightarrow \{\text{t, f}\}$$

²⁰Reference: [Pro24au].

²¹Further Terminology: Also called the **indicator function of U** .

²²Further Notation: Also written $\chi_X(U, -)$ or $\chi_X(-, U)$.

²³Further Notation: Also written χ_x , $\chi_X(x, -)$, or $\chi_X(-, x)$.

²⁴Further Terminology: Also called the **identity relation on X** .

²⁵Further Notation: Also written χ_{-2}^{-1} , or \sim_{id} in the context of relations.

on X defined by²⁶

$$\chi_X(x, y) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x = y, \\ \text{false} & \text{if } x \neq y \end{cases}$$

for each $x, y \in X$.

005J 4. The **characteristic embedding** ²⁷ of X into $\mathcal{P}(X)$ is the function

$$\chi_{(-)} : X \hookrightarrow \mathcal{P}(X)$$

defined by

$$\chi_{(-)}(x) \stackrel{\text{def}}{=} \chi_x$$

for each $x \in X$.

005K Remark 2.4.1.1.2. The definitions in [Definition 2.4.1.1.1](#) are decategorifications of co/presheaves, representable co/presheaves, Hom profunctors, and the Yoneda embedding:²⁸

1. A function

$$f : X \rightarrow \{\text{t}, \text{f}\}$$

²⁶As a subset of $X \times X$, the relation χ_X corresponds to the diagonal $\Delta_X \subset X \times X$ of X .

²⁷The name “characteristic embedding” comes from the fact that there is an analogue of fully faithfulness for $\chi_{(-)}$: given a set X , we have

$$\text{Hom}_{\mathcal{P}(X)}(\chi_x, \chi_y) = \chi_X(x, y),$$

for each $x, y \in X$.

²⁸These statements can be made precise by using the embeddings

$$\begin{aligned} (-)_{\text{disc}} : \text{Sets} &\hookrightarrow \text{Cats}, \\ (-)_{\text{disc}} : \{\text{t}, \text{f}\}_{\text{disc}} &\hookrightarrow \text{Sets} \end{aligned}$$

of sets into categories and of classical truth values into sets.

For instance, in this approach the characteristic function

$$\chi_x : X \rightarrow \{\text{t}, \text{f}\}$$

of an element x of X , defined by

$$\chi_x(y) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x = y, \\ \text{false} & \text{if } x \neq y \end{cases}$$

for each $y \in X$, is recovered as the representable presheaf

$$\text{Hom}_{X_{\text{disc}}}(-, x) : X_{\text{disc}} \rightarrow \text{Sets}$$

of the corresponding object x of X_{disc} , defined on objects by

$$\text{Hom}_{X_{\text{disc}}}(y, x) \stackrel{\text{def}}{=} \begin{cases} \text{pt} & \text{if } x = y, \\ \emptyset & \text{if } x \neq y \end{cases}$$

for each $y \in \text{Obj}(X_{\text{disc}})$.

is a decategorification of a presheaf

$$\mathcal{F}: C^{\text{op}} \rightarrow \text{Sets},$$

with the characteristic functions χ_U of the subsets of X being the primordial examples (and, in fact, all examples) of these.

2. The characteristic function

$$\chi_x: X \rightarrow \{\text{t, f}\}$$

of an *element* x of X is a decategorification of the representable presheaf

$$h_X: C^{\text{op}} \rightarrow \text{Sets}$$

of an *object* x of a category C .

3. The characteristic relation

$$\chi_X(-_1, -_2): X \times X \rightarrow \{\text{t, f}\}$$

of X is a decategorification of the Hom profunctor

$$\text{Hom}_C(-_1, -_2): C^{\text{op}} \times C \rightarrow \text{Sets}$$

of a category C .

4. The characteristic embedding

$$\chi_{(-)}: X \hookrightarrow \mathcal{P}(X)$$

of X into $\mathcal{P}(X)$ is a decategorification of the Yoneda embedding

$$\mathfrak{J}: C^{\text{op}} \hookrightarrow \text{PSh}(C)$$

of a category C into $\text{PSh}(C)$.

5. There is also a direct parallel between unions and colimits:

- An element of $\mathcal{P}(X)$ is a union of elements of X , viewed as one-point subsets $\{x\} \in \mathcal{P}(A)$;
- An object of $\text{PSh}(C)$ is a colimit of objects of C , viewed as representable presheaves $h_X \in \text{Obj}(\text{PSh}(C))$.

005L Proposition 2.4.1.1.3. Let X be a set.

005M 1. *The Inclusion of Characteristic Relations Associated to a Function.* Let $f: A \rightarrow B$

be a function. We have an inclusion²⁹

$$\begin{array}{ccc} A \times A & \xrightarrow{f \times f} & B \times B \\ \chi_B \circ (f \times f) \subset \chi_A, & \swarrow \curvearrowright \searrow & \chi_B \\ & \{t, f\}. & \end{array}$$

005N 2. *Interaction With Unions I.* We have

$$\chi_{U \cup V} = \max(\chi_U, \chi_V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

005P 3. *Interaction With Unions II.* We have

$$\chi_{U \cup V} = \chi_U + \chi_V - \chi_{U \cap V}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

005Q 4. *Interaction With Intersections I.* We have

$$\chi_{U \cap V} = \chi_U \chi_V$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

005R 5. *Interaction With Intersections II.* We have

$$\chi_{U \cap V} = \min(\chi_U, \chi_V)$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

005S 6. *Interaction With Differences.* We have

$$\chi_{U \setminus V} = \chi_U - \chi_{U \cap V}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

005T 7. *Interaction With Complements.* We have

$$\chi_{U^c} = 1 - \chi_U$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U \in \mathcal{P}(X)$.

005U 8. *Interaction With Symmetric Differences.* We have

$$\chi_{U \Delta V} = \chi_U + \chi_V - 2\chi_{U \cap V}$$

and thus, in particular, we have

$$\chi_{U \Delta V} \equiv \chi_U + \chi_V \pmod{2}$$

for each $X \in \text{Obj}(\text{Sets})$ and each $U, V \in \mathcal{P}(X)$.

²⁹This is the 0-categorical version of ??.

Proof. **Item 1, The Inclusion of Characteristic Relations Associated to a Function:** The inclusion $\chi_B(f(a), f(b)) \subset \chi_A(a, b)$ is equivalent to the statement “if $a = b$, then $f(a) = f(b)$ ”, which is true.

Item 2, Interaction With Unions I: This is a repetition of **Item 8** of [Proposition 2.3.6.1.2](#) and is proved there.

Item 3, Interaction With Unions II: This is a repetition of **Item 9** of [Proposition 2.3.6.1.2](#) and is proved there.

Item 4, Interaction With Intersections I: This is a repetition of **Item 9** of [Proposition 2.3.8.1.2](#) and is proved there.

Item 5, Interaction With Intersections II: This is a repetition of **Item 10** of [Proposition 2.3.8.1.2](#) and is proved there.

Item 6, Interaction With Differences: This is a repetition of **Item 15** of [Proposition 2.3.9.1.2](#) and is proved there.

Item 7, Interaction With Complements: This is a repetition of **Item 4** of [Proposition 2.3.10.1.2](#) and is proved there.

Item 8, Interaction With Symmetric Differences: This is a repetition of **Item 14** of [Proposition 2.3.11.1.2](#) and is proved there. \square

2.4.2 The Yoneda Lemma for Sets

Let X be a set and let $U \subset X$ be a subset of X .

005W Proposition 2.4.2.1.1. We have

$$\chi_{\mathcal{P}(X)}(\chi_x, \chi_U) = \chi_U(x)$$

for each $x \in X$, giving an equality of functions

$$\chi_{\mathcal{P}(X)}(\chi_{(-)}, \chi_U) = \chi_U.$$

Proof. Clear. \square

005X Corollary 2.4.2.1.2. The characteristic embedding is fully faithful, i.e., we have

$$\chi_{\mathcal{P}(X)}(\chi_x, \chi_y) = \chi_X(x, y)$$

for each $x, y \in X$.

Proof. This follows from [Proposition 2.4.2.1.1](#). \square

2.4.3 Powersets

Let X be a set.

005Z Definition 2.4.3.1.1. The **powerset** of X is the set $\mathcal{P}(X)$ defined by

$$\mathcal{P}(X) \stackrel{\text{def}}{=} \{U \in P \mid U \subset X\},$$

where P is the set in the axiom of powerset, ?? of ??.

0060 Remark 2.4.3.1.2. The powerset of a set is a decategorification of the category of presheaves of a category: while³⁰

- The powerset of a set X is equivalently ([Item 6 of Proposition 2.4.3.1.3](#)) the set

$$\text{Sets}(X, \{\text{t}, \text{f}\})$$

of functions from X to the set $\{\text{t}, \text{f}\}$ of classical truth values;

- The category of presheaves on a category C is the category

$$\text{Fun}(C^{\text{op}}, \text{Sets})$$

of functors from C^{op} to the category Sets of sets.

0061 Proposition 2.4.3.1.3. Let X be a set.

0062 1. *Functionality.* The assignment $X \mapsto \mathcal{P}(X)$ defines functors

$$\begin{aligned} \mathcal{P}_*: \text{Sets} &\rightarrow \text{Sets}, \\ \mathcal{P}^{-1}: \text{Sets}^{\text{op}} &\rightarrow \text{Sets}, \\ \mathcal{P}_!: \text{Sets} &\rightarrow \text{Sets} \end{aligned}$$

where

- *Action on Objects.* For each $A \in \text{Obj}(\text{Sets})$, we have

$$\begin{aligned} \mathcal{P}_*(A) &\stackrel{\text{def}}{=} \mathcal{P}(A), \\ \mathcal{P}^{-1}(A) &\stackrel{\text{def}}{=} \mathcal{P}(A), \\ \mathcal{P}_!(A) &\stackrel{\text{def}}{=} \mathcal{P}(A); \end{aligned}$$

- *Action on Morphisms.* For each morphism $f: A \rightarrow B$ of Sets , the images

$$\begin{aligned} \mathcal{P}_*(f): \mathcal{P}(A) &\rightarrow \mathcal{P}(B), \\ \mathcal{P}^{-1}(f): \mathcal{P}(B) &\rightarrow \mathcal{P}(A), \\ \mathcal{P}_!(f): \mathcal{P}(A) &\rightarrow \mathcal{P}(B) \end{aligned}$$

³⁰This parallel is based on the following comparison:

- A category is enriched over the category

$$\text{Sets} \stackrel{\text{def}}{=} \text{Cats}_0$$

of sets (i.e. “0-categories”), with presheaves taking values on it;

- A set is enriched over the set

$$\{\text{t}, \text{f}\} \stackrel{\text{def}}{=} \text{Cats}_{-1}$$

of classical truth values (i.e. “(-1)-categories”), with characteristic functions taking values on it.

of f by \mathcal{P}_* , \mathcal{P}^{-1} , and $\mathcal{P}_!$ are defined by

$$\begin{aligned}\mathcal{P}_*(f) &\stackrel{\text{def}}{=} f_*, \\ \mathcal{P}^{-1}(f) &\stackrel{\text{def}}{=} f^{-1}, \\ \mathcal{P}_!(f) &\stackrel{\text{def}}{=} f_!,\end{aligned}$$

as in [Definitions 2.4.4.1.1, 2.4.5.1.1 and 2.4.6.1.1](#).

0063 2. *Adjointness I.* We have an adjunction

$$(\mathcal{P}^{-1} \dashv \mathcal{P}^{-1,\text{op}}): \quad \text{Sets}^{\text{op}} \begin{array}{c} \xrightarrow{\mathcal{P}^{-1}} \\ \perp \\ \xleftarrow{\mathcal{P}^{-1,\text{op}}} \end{array} \text{Sets},$$

witnessed by a bijection

$$\underbrace{\text{Sets}^{\text{op}}(\mathcal{P}(X), Y)}_{\stackrel{\text{def}}{=} \text{Sets}(Y, \mathcal{P}(X))} \cong \text{Sets}(X, \mathcal{P}(Y)),$$

natural in $X \in \text{Obj}(\text{Sets})$ and $Y \in \text{Obj}(\text{Sets}^{\text{op}})$.

0064 3. *Adjointness II.* We have an adjunction

$$(\text{Gr} \dashv \mathcal{P}_*): \quad \text{Sets} \begin{array}{c} \xrightarrow{\text{Gr}} \\ \perp \\ \xleftarrow{\mathcal{P}_*} \end{array} \text{Rel},$$

witnessed by a bijection of sets

$$\text{Rel}(\text{Gr}(A), B) \cong \text{Sets}(A, \mathcal{P}(B))$$

natural in $A \in \text{Obj}(\text{Sets})$ and $B \in \text{Obj}(\text{Rel})$, where Gr is the graph functor of ?? of ??.

0065 4. *Symmetric Strong Monoidality With Respect to Coproducts.* The powerset functor \mathcal{P}_* of [Item 1](#) has a symmetric strong monoidal structure

$$(\mathcal{P}_*, \mathcal{P}_*^{\coprod}, \mathcal{P}_{*\mid \mathbb{1}}^{\coprod}): (\text{Sets}, \coprod, \emptyset) \rightarrow (\text{Sets}, \times, \text{pt})$$

being equipped with isomorphisms

$$\begin{aligned}\mathcal{P}_{*\mid X,Y}^{\coprod}: \mathcal{P}(X) \times \mathcal{P}(Y) &\xrightarrow{\cong} \mathcal{P}(X \coprod Y), \\ \mathcal{P}_{*\mid \mathbb{1}}^{\coprod}: \text{pt} &\xrightarrow{=} \mathcal{P}(\emptyset),\end{aligned}$$

natural in $X, Y \in \text{Obj}(\text{Sets})$.

- 0066** 5. *Symmetric Lax Monoidality With Respect to Products.* The powerset functor \mathcal{P}_* of Item 1 has a symmetric lax monoidal structure

$$\left(\mathcal{P}_*, \mathcal{P}_*^\otimes, \mathcal{P}_{*|k}^\otimes \right) : (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Sets}, \times, \text{pt})$$

being equipped with isomorphisms

$$\begin{aligned} \mathcal{P}_{*|X,Y}^\otimes : \mathcal{P}(X) \times \mathcal{P}(Y) &\rightarrow \mathcal{P}(X \times Y), \\ \mathcal{P}_{*|k}^\otimes : \text{pt} &\xrightarrow{\cong} \mathcal{P}(\emptyset), \end{aligned}$$

natural in $X, Y \in \text{Obj}(\text{Sets})$, where $\mathcal{P}_{*|X,Y}^\otimes$ is given by

$$\mathcal{P}_{*|X,Y}^\otimes(U, V) \stackrel{\text{def}}{=} U \times V$$

for each $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(Y)$.

- 0067** 6. *Powersets as Sets of Functions.* The assignment $U \mapsto \chi_U$ defines a bijection³¹

$$\chi_{(-)} : \mathcal{P}(X) \xrightarrow{\cong} \text{Sets}(X, \{\text{t}, \text{f}\}),$$

natural in $X \in \text{Obj}(\text{Sets})$.

- 0068** 7. *Powersets as Sets of Relations.* We have bijections

$$\begin{aligned} \mathcal{P}(X) &\cong \text{Rel}(\text{pt}, X), \\ \mathcal{P}(X) &\cong \text{Rel}(X, \text{pt}), \end{aligned}$$

natural in $X \in \text{Obj}(\text{Sets})$.

- 0069** 8. *As a Free Cocompletion: Universal Property.* The pair $(\mathcal{P}(X), \chi_{(-)})$ consisting of

- The powerset $\mathcal{P}(X)$ of X ;
- The characteristic embedding $\chi_{(-)} : X \hookrightarrow \mathcal{P}(X)$ of X into $\mathcal{P}(X)$;

satisfies the following universal property:

- (★) Given another pair (Y, f) consisting of
- A cocomplete poset (Y, \leq) ;
 - A function $f : X \rightarrow Y$;

³¹This bijection is a decategorified form of the equivalence

$$\text{PSh}(C) \xrightarrow{\text{eq}} \text{DFib}(C)$$

of ?? of ??, with $\chi_{(-)}$ being a decategorified version of the category of elements construction of ??.
See also ?? of ??.

there exists a unique cocontinuous morphism of posets $(\mathcal{P}(X), \subset) \xrightarrow{\exists!} (Y, \leq)$ making the diagram

$$\begin{array}{ccc} & \mathcal{P}(X) & \\ \chi_X \nearrow & \downarrow \exists! & \\ X & \xrightarrow{f} & Y \end{array}$$

commute.

006A 9. As a Free Cocompletion: Adjointness. We have an adjunction³²

$$(\chi_{(-)} \dashv \text{忘}): \text{Sets} \begin{array}{c} \xrightarrow{\chi_{(-)}} \\ \perp \\ \xleftarrow{\text{忘}} \end{array} \text{Pos}^{\text{cocomp.}},$$

witnessed by a bijection

$$\text{Pos}^{\text{cocomp.}}((\mathcal{P}(X), \subset), (Y, \leq)) \cong \text{Sets}(X, Y),$$

natural in $X \in \text{Obj}(\text{Sets})$ and $(Y, \leq) \in \text{Obj}(\text{Pos})$, where

- We have a natural map

$$\chi_X^*: \text{Pos}^{\text{cocomp.}}((\mathcal{P}(X), \subset), (Y, \leq)) \rightarrow \text{Sets}(X, Y)$$

defined by

$$\chi_X^*(f) \stackrel{\text{def}}{=} f \circ \chi_X,$$

i.e. by sending a cocontinuous morphism of posets $f: \mathcal{P}(X) \rightarrow Y$ to the composition

$$X \xhookrightarrow{\chi_X} \mathcal{P}(X) \xrightarrow{f} Y;$$

- We have a natural map

$$\text{Lan}_{\chi_X}: \text{Sets}(X, Y) \rightarrow \text{Pos}^{\text{cocomp.}}((\mathcal{P}(X), \subset), (Y, \leq))$$

computed by

$$\begin{aligned} [\text{Lan}_{\chi_X}(f)](U) &\cong \int^{x \in X} \chi_{\mathcal{P}(X)}(\chi_x, U) \odot f(x) \\ &\cong \int^{x \in X} \chi_U(x) \odot f(x) \quad (\text{by Proposition 2.4.2.1.1}) \\ &\cong \bigvee_{x \in X} (\chi_U(x) \odot f(x)) \end{aligned}$$

for each $U \in \mathcal{P}(X)$, where:

³²In this sense, $\mathcal{P}(A)$ is the free cocompletion of A . (Note that, despite its name, however, this is not an

- \vee is the join in (Y, \leq) ;
- We have

$$\begin{aligned}\text{true} \odot f(x) &\stackrel{\text{def}}{=} f(x), \\ \text{false} \odot f(x) &\stackrel{\text{def}}{=} \emptyset_Y,\end{aligned}$$

where \emptyset_Y is the minimal element of (Y, \leq) .

Proof. **Item 1, Functoriality:** This follows from **Items 3 and 4 of Proposition 2.4.4.1.4**, **Items 3 and 4 of Proposition 2.4.5.1.4**, and **Items 3 and 4 of Proposition 2.4.6.1.6**.

Item 2, Adjointness I: Omitted.

Item 3, Adjointness II: We have

$$\begin{aligned}\text{Rel}(\text{Gr}(A), B) &= \mathcal{P}(A \times B) \\ &= \text{Sets}(A \times B, \{\text{t}, \text{f}\}) && (\text{by Item 6}) \\ &= \text{Sets}(A, \text{Sets}(B, \{\text{t}, \text{f}\})) && (\text{by Item 2 of Proposition 2.1.2.1.2}) \\ &= \text{Sets}(A, \mathcal{P}(B)) && (\text{by Item 6})\end{aligned}$$

with all bijections natural in A and B .

Item 4, Symmetric Strong Monoidality With Respect to Coproducts: Omitted.

Item 5, Symmetric Lax Monoidality With Respect to Products: Omitted.

Item 6, Powersets as Sets of Functions: Omitted.

Item 7, Powersets as Sets of Relations: Omitted.

Item 8, As a Free Cocompletion: Universal Property: This is a rephrasing of ??.

Item 9, As a Free Cocompletion: Adjointness: Omitted. □

2.4.4 Direct Images

Let A and B be sets and let $f: A \rightarrow B$ be a function.

006C Definition 2.4.4.1.1. The **direct image function associated to f** is the function³³

$$f_*: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

idempotent operation, as we have $\mathcal{P}(\mathcal{P}(A)) \neq \mathcal{P}(A)$.)

³³Further Notation: Also written $\exists_f: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$. This notation comes from the fact that the following statements are equivalent, where $b \in B$ and $U \in \mathcal{P}(A)$:

- We have $b \in \exists_f(U)$.
- There exists some $a \in U$ such that $f(a) = b$.

defined by^{34,35}

$$\begin{aligned} f_*(U) &\stackrel{\text{def}}{=} f(U) \\ &\stackrel{\text{def}}{=} \left\{ b \in B \mid \begin{array}{l} \text{there exists some } a \in U \\ \text{such that } b = f(a) \end{array} \right\} \\ &= \{f(a) \in B \mid a \in U\} \end{aligned}$$

for each $U \in \mathcal{P}(A)$.

006D Remark 2.4.4.1.2. Identifying subsets of A with functions from A to $\{\text{true}, \text{false}\}$ via Item 6 of Proposition 2.4.3.1.3, we see that the direct image function associated to f is equivalently the function

$$f_*: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by

$$\begin{aligned} f_*(\chi_U) &\stackrel{\text{def}}{=} \text{Lan}_f(\chi_U) \\ &= \text{colim} \left(\left(f \times \underline{(-_1)} \right) \xrightarrow{\text{pr}} A \xrightarrow{\chi_U} \{\text{t, f}\} \right) \\ &= \underset{\substack{a \in A \\ f(a) = -_1}}{\text{colim}} (\chi_U(a)) \\ &= \bigvee_{\substack{a \in A \\ f(a) = -_1}} (\chi_U(a)). \end{aligned}$$

So, in other words, we have

$$\begin{aligned} [f_*(\chi_U)](b) &= \bigvee_{\substack{a \in A \\ f(a) = b}} (\chi_U(a)) \\ &= \begin{cases} \text{true} & \text{if there exists some } a \in A \text{ such} \\ & \text{that } f(a) = b \text{ and } a \in U, \\ \text{false} & \text{otherwise} \end{cases} \\ &= \begin{cases} \text{true} & \text{if there exists some } a \in U \\ & \text{such that } f(a) = b, \\ \text{false} & \text{otherwise} \end{cases} \end{aligned}$$

for each $b \in B$.

006E Proposition 2.4.4.1.3. Let $f: A \rightarrow B$ be a function.

³⁴Further Terminology: The set $f(U)$ is called the **direct image of U by f** .

³⁵We also have

$$f_*(U) = B \setminus f_!(A \setminus U);$$

see Item 7 of Proposition 2.4.4.1.3.

006F 1. *Functionality.* The assignment $U \mapsto f_*(U)$ defines a functor

$$f_*: (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

where

- *Action on Objects.* For each $U \in \mathcal{P}(A)$, we have

$$[f_*](U) \stackrel{\text{def}}{=} f_*(U);$$

- *Action on Morphisms.* For each $U, V \in \mathcal{P}(A)$:

(★) If $U \subset V$, then $f_*(U) \subset f_*(V)$.

006G 2. *Triple Adjunction.* We have a triple adjunction

$$(f_* \dashv f^{-1} \dashv f_!): \quad \mathcal{P}(A) \begin{array}{c} \xleftarrow[f_*]{\perp} \\[-1ex] \xleftarrow[f^{-1}]{\perp} \\[-1ex] \xleftarrow[f_!]{\perp} \end{array} \mathcal{P}(B),$$

witnessed by bijections of sets

$$\begin{aligned} \text{Hom}_{\mathcal{P}(B)}(f_*(U), V) &\cong \text{Hom}_{\mathcal{P}(A)}(U, f^{-1}(V)), \\ \text{Hom}_{\mathcal{P}(A)}(f^{-1}(U), V) &\cong \text{Hom}_{\mathcal{P}(B)}(U, f_!(V)), \end{aligned}$$

natural in $U \in \mathcal{P}(A)$ and $V \in \mathcal{P}(B)$ and (respectively) $V \in \mathcal{P}(A)$ and $U \in \mathcal{P}(B)$,
i.e. where:

(a) The following conditions are equivalent:

- i. We have $f_*(U) \subset V$.
- ii. We have $U \subset f^{-1}(V)$.

(b) The following conditions are equivalent:

- i. We have $f^{-1}(U) \subset V$.
- ii. We have $U \subset f_!(V)$.

006H 3. *Preservation of Colimits.* We have an equality of sets

$$f_* \left(\bigcup_{i \in I} U_i \right) = \bigcup_{i \in I} f_*(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$. In particular, we have equalities

$$\begin{aligned} f_*(U) \cup f_*(V) &= f_*(U \cup V), \\ f_*(\emptyset) &= \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

006J 4. *Oplax Preservation of Limits.* We have an inclusion of sets

$$f_* \left(\bigcap_{i \in I} U_i \right) \subset \bigcap_{i \in I} f_*(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$. In particular, we have inclusions

$$\begin{aligned} f_*(U \cap V) &\subset f_*(U) \cap f_*(V), \\ f_*(A) &\subset B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

006K 5. *Symmetric Strict Monoidality With Respect to Unions.* The direct image function of **Item 1** has a symmetric strict monoidal structure

$$\left(f_*, f_*^\otimes, f_{*\mid \sharp}^\otimes \right): (\mathcal{P}(A), \cup, \emptyset) \rightarrow (\mathcal{P}(B), \cup, \emptyset),$$

being equipped with equalities

$$\begin{aligned} f_{*|U,V}^\otimes: f_*(U) \cup f_*(V) &\xrightarrow{=} f_*(U \cup V), \\ f_{*\mid \sharp}^\otimes: \emptyset &\xrightarrow{=} \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

006L 6. *Symmetric Oplax Monoidality With Respect to Intersections.* The direct image function of **Item 1** has a symmetric oplax monoidal structure

$$\left(f_*, f_*^\otimes, f_{*\mid \sharp}^\otimes \right): (\mathcal{P}(A), \cap, A) \rightarrow (\mathcal{P}(B), \cap, B),$$

being equipped with inclusions

$$\begin{aligned} f_{*|U,V}^\otimes: f_*(U \cap V) &\hookrightarrow f_*(U) \cap f_*(V), \\ f_{*\mid \sharp}^\otimes: f_*(A) &\hookrightarrow B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

006M 7. *Relation to Direct Images With Compact Support.* We have

$$f_*(U) = B \setminus f_!(A \setminus U)$$

for each $U \in \mathcal{P}(A)$.

Proof. **Item 1**, Functoriality: Clear.

Item 2, Triple Adjunctions: This follows from ?? of ??.

Item 3, Preservation of Colimits: This follows from **Item 2** and ?? of ??.

Item 4, Oplax Preservation of Limits: Omitted.

Item 5, Symmetric Strict Monoidality With Respect to Unions: This follows from **Item 3**.

Item 6, Symmetric Oplax Monoidality With Respect to Intersections: This follows from ??.

Item 7, Relation to Direct Images With Compact Support: Applying ?? of ?? to $A \setminus U$, we have

$$\begin{aligned} f_!(A \setminus U) &= B \setminus f_*(A \setminus (A \setminus U)) \\ &= B \setminus f_*(U). \end{aligned}$$

Taking complements, we then obtain

$$\begin{aligned} f_*(U) &= B \setminus (B \setminus f_*(U)), \\ &= B \setminus f_!(A \setminus U), \end{aligned}$$

which finishes the proof. \square

006N Proposition 2.4.4.1.4. Let $f: A \rightarrow B$ be a function.

006P 1. Functionality I. The assignment $f \mapsto f_*$ defines a function

$$(-)_{*|A,B}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B)).$$

006Q 2. Functionality II. The assignment $f \mapsto f_*$ defines a function

$$(-)_{*|A,B}: \text{Sets}(A, B) \rightarrow \text{Pos}((\mathcal{P}(A), \subset), (\mathcal{P}(B), \subset)).$$

006R 3. Interaction With Identities. For each $A \in \text{Obj}(\text{Sets})$, we have

$$(\text{id}_A)_* = \text{id}_{\mathcal{P}(A)};$$

006S 4. Interaction With Composition. For each pair of composable functions $f: A \rightarrow B$ and $g: B \rightarrow C$, we have

$$\begin{array}{ccc} \mathcal{P}(A) & \xrightarrow{f_*} & \mathcal{P}(B) \\ (g \circ f)_* = g_* \circ f_* & \searrow & \downarrow g_* \\ & & \mathcal{P}(C). \end{array}$$

Proof. **Item 1, Functionality I:** Clear.

Item 2, Functionality II: Clear.

Item 3, Interaction With Identities: This follows from ?? of ??.

Item 4, Interaction With Composition: This follows from ?? of ??.

\square

2.4.5 Inverse Images

Let A and B be sets and let $f: A \rightarrow B$ be a function.

006U Definition 2.4.5.1.1. The **inverse image function associated to f** is the function³⁶

$$f^{-1}: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

defined by³⁷

$$f^{-1}(V) \stackrel{\text{def}}{=} \{a \in A \mid \text{we have } f(a) \in V\}$$

for each $V \in \mathcal{P}(B)$.

006V Remark 2.4.5.1.2. Identifying subsets of B with functions from B to $\{\text{true, false}\}$ via Item 6 of Proposition 2.4.3.1.3, we see that the inverse image function associated to f is equivalently the function

$$f^*: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

defined by

$$f^*(\chi_V) \stackrel{\text{def}}{=} \chi_V \circ f$$

for each $\chi_V \in \mathcal{P}(B)$, where $\chi_V \circ f$ is the composition

$$A \xrightarrow{f} B \xrightarrow{\chi_V} \{\text{true, false}\}$$

in Sets.

006W Proposition 2.4.5.1.3. Let $f: A \rightarrow B$ be a function.

006X 1. Functoriality. The assignment $V \mapsto f^{-1}(V)$ defines a functor

$$f^{-1}: (\mathcal{P}(B), \subset) \rightarrow (\mathcal{P}(A), \subset)$$

where

- *Action on Objects.* For each $V \in \mathcal{P}(B)$, we have

$$[f^{-1}](V) \stackrel{\text{def}}{=} f^{-1}(V);$$

- *Action on Morphisms.* For each $U, V \in \mathcal{P}(B)$:

(★) If $U \subset V$, then $f^{-1}(U) \subset f^{-1}(V)$.

³⁶Further Notation: Also written $f^*: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$.

³⁷Further Terminology: The set $f^{-1}(V)$ is called the **inverse image of V by f** .

006Y 2. *Triple Adjunctions.* We have a triple adjunction

$$(f_* \dashv f^{-1} \dashv f_!): \quad \mathcal{P}(A) \begin{array}{c} \xleftarrow{\perp} \\[-1ex] \xleftarrow{f^{-1}} \\[-1ex] \xleftarrow{\perp} \end{array} \mathcal{P}(B),$$

witnessed by bijections of sets

$$\begin{aligned} \text{Hom}_{\mathcal{P}(B)}(f_*(U), V) &\cong \text{Hom}_{\mathcal{P}(A)}(U, f^{-1}(V)), \\ \text{Hom}_{\mathcal{P}(A)}(f^{-1}(U), V) &\cong \text{Hom}_{\mathcal{P}(A)}(U, f_!(V)), \end{aligned}$$

natural in $U \in \mathcal{P}(A)$ and $V \in \mathcal{P}(B)$ and (respectively) $V \in \mathcal{P}(A)$ and $U \in \mathcal{P}(B)$, i.e. where:

(a) The following conditions are equivalent:

- i. We have $f_*(U) \subset V$;
- ii. We have $U \subset f^{-1}(V)$;

(b) The following conditions are equivalent:

- i. We have $f^{-1}(U) \subset V$.
- ii. We have $U \subset f_!(V)$.

006Z 3. *Preservation of Colimits.* We have an equality of sets

$$f^{-1}\left(\bigcup_{i \in I} U_i\right) = \bigcup_{i \in I} f^{-1}(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(B)^{\times I}$. In particular, we have equalities

$$\begin{aligned} f^{-1}(U) \cup f^{-1}(V) &= f^{-1}(U \cup V), \\ f^{-1}(\emptyset) &= \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

0070 4. *Preservation of Limits.* We have an equality of sets

$$f^{-1}\left(\bigcap_{i \in I} U_i\right) = \bigcap_{i \in I} f^{-1}(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(B)^{\times I}$. In particular, we have equalities

$$\begin{aligned} f^{-1}(U) \cap f^{-1}(V) &= f^{-1}(U \cap V), \\ f^{-1}(B) &= A, \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

- 0071** 5. *Symmetric Strict Monoidality With Respect to Unions.* The inverse image function of **Item 1** has a symmetric strict monoidal structure

$$\left(f^{-1}, f^{-1,\otimes}, f_{\sharp}^{-1,\otimes} \right) : (\mathcal{P}(B), \cup, \emptyset) \rightarrow (\mathcal{P}(A), \cup, \emptyset),$$

being equipped with equalities

$$\begin{aligned} f_{U,V}^{-1,\otimes} &: f^{-1}(U) \cup f^{-1}(V) \xrightarrow{\cong} f^{-1}(U \cup V), \\ f_{\sharp}^{-1,\otimes} &: \emptyset \xrightarrow{\cong} f^{-1}(\emptyset), \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

- 0072** 6. *Symmetric Strict Monoidality With Respect to Intersections.* The inverse image function of **Item 1** has a symmetric strict monoidal structure

$$\left(f^{-1}, f^{-1,\otimes}, f_{\sharp}^{-1,\otimes} \right) : (\mathcal{P}(B), \cap, B) \rightarrow (\mathcal{P}(A), \cap, A),$$

being equipped with equalities

$$\begin{aligned} f_{U,V}^{-1,\otimes} &: f^{-1}(U) \cap f^{-1}(V) \xrightarrow{\cong} f^{-1}(U \cap V), \\ f_{\sharp}^{-1,\otimes} &: A \xrightarrow{\cong} f^{-1}(B), \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

Proof. **Item 1, Functoriality:** Clear.

Item 2, Triple Adjointness: This follows from ?? of ??.

Item 3, Preservation of Colimits: This follows from **Item 2** and ?? of ??.

Item 4, Preservation of Limits: This follows from **Item 2** and ?? of ??.

Item 5, Symmetric Strict Monoidality With Respect to Unions: This follows from **Item 3**.

Item 6, Symmetric Strict Monoidality With Respect to Intersections: This follows from **Item 4.**

□

- 0073** **Proposition 2.4.5.1.4.** Let $f: A \rightarrow B$ be a function.

- 0074** 1. *Functionality I.* The assignment $f \mapsto f^{-1}$ defines a function

$$(-)^{-1}_{A,B}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(B), \mathcal{P}(A)).$$

- 0075** 2. *Functionality II.* The assignment $f \mapsto f^{-1}$ defines a function

$$(-)^{-1}_{A,B}: \text{Sets}(A, B) \rightarrow \text{Pos}((\mathcal{P}(B), \subset), (\mathcal{P}(A), \subset)).$$

- 0076** 3. *Interaction With Identities.* For each $A \in \text{Obj}(\text{Sets})$, we have

$$\text{id}_A^{-1} = \text{id}_{\mathcal{P}(A)};$$

- 0077** 4. *Interaction With Composition.* For each pair of composable functions $f: A \rightarrow B$ and $g: B \rightarrow C$, we have

$$\begin{array}{ccc} \mathcal{P}(C) & \xrightarrow{g^{-1}} & \mathcal{P}(B) \\ (g \circ f)^{-1} = f^{-1} \circ g^{-1}, & \searrow & \downarrow f^{-1} \\ & (g \circ f)^{-1} & \mathcal{P}(A). \end{array}$$

Proof. **Item 1, Functionality I:** Clear.

Item 2, Functionality II: Clear.

Item 3, Interaction With Identities: This follows from ?? of ??.

Item 4, Interaction With Composition: This follows from ?? of ??.

□

2.4.6 Direct Images With Compact Support

Let A and B be sets and let $f: A \rightarrow B$ be a function.

- 0079 Definition 2.4.6.1.1.** The **direct image with compact support function associated to f** is the function³⁸

$$f_!: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by^{39,40}

$$\begin{aligned} f_!(U) &\stackrel{\text{def}}{=} \left\{ b \in B \mid \begin{array}{l} \text{for each } a \in A, \text{ if we have} \\ f(a) = b, \text{ then } a \in U \end{array} \right\} \\ &= \left\{ b \in B \mid \text{we have } f^{-1}(b) \subset U \right\} \end{aligned}$$

for each $U \in \mathcal{P}(A)$.

- 007A Remark 2.4.6.1.2.** Identifying subsets of A with functions from A to $\{\text{true}, \text{false}\}$ via **Item 6 of Proposition 2.4.3.1.3**, we see that the direct image with compact support function associated to f is equivalently the function

$$f_!: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

³⁸Further Notation: Also written $\forall_f: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$. This notation comes from the fact that the following statements are equivalent, where $b \in B$ and $U \in \mathcal{P}(A)$:

- We have $b \in \forall_f(U)$.
- For each $a \in A$, if $b = f(a)$, then $a \in U$.

³⁹Further Terminology: The set $f_!(U)$ is called the **direct image with compact support of U by f** .

⁴⁰We also have

$$f_!(U) = B \setminus f_*(A \setminus U);$$

see **Item 7 of Proposition 2.4.6.1.5**.

defined by

$$\begin{aligned}
 f_!(\chi_U) &\stackrel{\text{def}}{=} \text{Ran}_f(\chi_U) \\
 &= \lim \left(\left(\underline{(-_1)} \xrightarrow{\vec{x}} f \right) \xrightarrow{\text{pr}} A \xrightarrow{\chi_U} \{\text{true}, \text{false}\} \right) \\
 &= \lim_{\substack{a \in A \\ f(a) = -_1}} (\chi_U(a)) \\
 &= \bigwedge_{\substack{a \in A \\ f(a) = -_1}} (\chi_U(a)).
 \end{aligned}$$

So, in other words, we have

$$\begin{aligned}
 [f_!(\chi_U)](b) &= \bigwedge_{\substack{a \in A \\ f(a) = b}} (\chi_U(a)) \\
 &= \begin{cases} \text{true} & \text{if, for each } a \in A \text{ such that} \\ & f(a) = b, \text{ we have } a \in U, \\ \text{false} & \text{otherwise} \end{cases} \\
 &= \begin{cases} \text{true} & \text{if } f^{-1}(b) \subset U \\ \text{false} & \text{otherwise} \end{cases}
 \end{aligned}$$

for each $b \in B$.

007B Definition 2.4.6.1.3. Let U be a subset of A .^{41,42}

1. The **image part of the direct image with compact support** $f_!(U)$ of U is the set $f_{!,\text{im}}(U)$ defined by

$$\begin{aligned}
 f_{!,\text{im}}(U) &\stackrel{\text{def}}{=} f_!(U) \cap \text{Im}(f) \\
 &= \left\{ b \in B \mid \begin{array}{l} \text{we have } f^{-1}(b) \subset U \\ \text{and } f^{-1}(b) \neq \emptyset \end{array} \right\}.
 \end{aligned}$$

⁴¹Note that we have

$$f_!(U) = f_{!,\text{im}}(U) \cup f_{!,\text{cp}}(U),$$

as

$$\begin{aligned}
 f_!(U) &= f_!(U) \cap B \\
 &= f_!(U) \cap (\text{Im}(f) \cup (B \setminus \text{Im}(f))) \\
 &= (f_!(U) \cap \text{Im}(f)) \cup (f_!(U) \cap (B \setminus \text{Im}(f))) \\
 &\stackrel{\text{def}}{=} f_{!,\text{im}}(U) \cup f_{!,\text{cp}}(U).
 \end{aligned}$$

⁴²In terms of the meet computation of $f_!(U)$ of Remark 2.4.6.1.2, namely

$$f_!(\chi_U) = \bigwedge_{\substack{a \in A \\ f(a) = -_1}} (\chi_U(a)),$$

2. The **complement part of the direct image with compact support** $f_!(U)$ of U is the set $f_{!,cp}(U)$ defined by

$$\begin{aligned} f_{!,cp}(U) &\stackrel{\text{def}}{=} f_!(U) \cap (B \setminus \text{Im}(f)) \\ &= B \setminus \text{Im}(f) \\ &= \left\{ b \in B \left| \begin{array}{l} \text{we have } f^{-1}(b) \subset U \\ \text{and } f^{-1}(b) = \emptyset \end{array} \right. \right\} \\ &= \left\{ b \in B \mid f^{-1}(b) = \emptyset \right\}. \end{aligned}$$

007C Example 2.4.6.1.4. Here are some examples of direct images with compact support.

1. *The Multiplication by Two Map on the Natural Numbers.* Consider the function $f: \mathbb{N} \rightarrow \mathbb{N}$ given by

$$f(n) \stackrel{\text{def}}{=} 2n$$

for each $n \in \mathbb{N}$. Since f is injective, we have

$$\begin{aligned} f_{!,im}(U) &= f_*(U) \\ f_{!,cp}(U) &= \{\text{odd natural numbers}\} \end{aligned}$$

for any $U \subset \mathbb{N}$.

2. *Parabolas.* Consider the function $f: \mathbb{R} \rightarrow \mathbb{R}$ given by

$$f(x) \stackrel{\text{def}}{=} x^2$$

for each $x \in \mathbb{R}$. We have

$$f_{!,cp}(U) = \mathbb{R}_{<0}$$

for any $U \subset \mathbb{R}$. Moreover, since $f^{-1}(x) = \{-\sqrt{x}, \sqrt{x}\}$, we have e.g.:

$$\begin{aligned} f_{!,im}([0, 1]) &= \{0\}, \\ f_{!,im}([-1, 1]) &= [0, 1], \\ f_{!,im}([1, 2]) &= \emptyset, \\ f_{!,im}([-2, -1] \cup [1, 2]) &= [1, 4]. \end{aligned}$$

3. *Circles.* Consider the function $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ given by

$$f(x, y) \stackrel{\text{def}}{=} x^2 + y^2$$

for each $(x, y) \in \mathbb{R}^2$. We have

$$f_{!,cp}(U) = \mathbb{R}_{<0}$$

for any $U \subset \mathbb{R}^2$, and since

$$f^{-1}(r) = \begin{cases} \text{a circle of radius } r \text{ about the origin} & \text{if } r > 0, \\ \{(0, 0)\} & \text{if } r = 0, \\ \emptyset & \text{if } r < 0, \end{cases}$$

we have e.g.:

$$\begin{aligned} f_{!,\text{im}}([-1, 1] \times [-1, 1]) &= [0, 1], \\ f_{!,\text{im}}(([[-1, 1] \times [-1, 1]] \setminus [-1, 1] \times \{0\})) &= \emptyset. \end{aligned}$$

007D Proposition 2.4.6.1.5. Let $f: A \rightarrow B$ be a function.

007E 1. *Functoriality.* The assignment $U \mapsto f_!(U)$ defines a functor

$$f_!: (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

where

- *Action on Objects.* For each $U \in \mathcal{P}(A)$, we have

$$[f_!](U) \stackrel{\text{def}}{=} f_!(U);$$

- *Action on Morphisms.* For each $U, V \in \mathcal{P}(A)$:

(★) If $U \subset V$, then $f_!(U) \subset f_!(V)$.

007F 2. *Triple Adjunction.* We have a triple adjunction

$$(f_* \dashv f^{-1} \dashv f_!): \quad \mathcal{P}(A) \begin{array}{c} \xleftarrow{f_*} \\ \perp \\ \xrightarrow{f^{-1}} \end{array} \mathcal{P}(B),$$

witnessed by bijections of sets

$$\begin{aligned} \text{Hom}_{\mathcal{P}(B)}(f_*(U), V) &\cong \text{Hom}_{\mathcal{P}(A)}(U, f^{-1}(V)), \\ \text{Hom}_{\mathcal{P}(A)}(f^{-1}(U), V) &\cong \text{Hom}_{\mathcal{P}(B)}(U, f_!(V)), \end{aligned}$$

natural in $U \in \mathcal{P}(A)$ and $V \in \mathcal{P}(B)$ and (respectively) $V \in \mathcal{P}(A)$ and $U \in \mathcal{P}(B)$,
i.e. where:

- (a) The following conditions are equivalent:

i. We have $f_*(U) \subset V$;

we see that $f_{!,\text{im}}$ corresponds to meets indexed over nonempty sets, while $f_{!,\text{cp}}$ corresponds to meets indexed over the empty set.

- ii. We have $U \subset f^{-1}(V)$;
- (b) The following conditions are equivalent:
- We have $f^{-1}(U) \subset V$.
 - We have $U \subset f_!(V)$.

007G 3. *Lax Preservation of Colimits.* We have an inclusion of sets

$$\bigcup_{i \in I} f_!(U_i) \subset f_!\left(\bigcup_{i \in I} U_i\right),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$. In particular, we have inclusions

$$\begin{aligned} f_!(U) \cup f_!(V) &\hookrightarrow f_!(U \cup V), \\ \emptyset &\hookrightarrow f_!(\emptyset), \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

007H 4. *Preservation of Limits.* We have an equality of sets

$$f_!\left(\bigcap_{i \in I} U_i\right) = \bigcap_{i \in I} f_!(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$. In particular, we have equalities

$$\begin{aligned} f^{-1}(U \cap V) &= f_!(U) \cap f^{-1}(V), \\ f_!(A) &= B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

007J 5. *Symmetric Lax Monoidality With Respect to Unions.* The direct image with compact support function of **Item 1** has a symmetric lax monoidal structure

$$\left(f_!, f_!^\otimes, f_{!|*}^\otimes\right): (\mathcal{P}(A), \cup, \emptyset) \rightarrow (\mathcal{P}(B), \cup, \emptyset),$$

being equipped with inclusions

$$\begin{aligned} f_{!|U,V}^\otimes: f_!(U) \cup f_!(V) &\hookrightarrow f_!(U \cup V), \\ f_{!|*}^\otimes: \emptyset &\hookrightarrow f_!(\emptyset), \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

007K 6. *Symmetric Strict Monoidality With Respect to Intersections.* The direct image function of **Item 1** has a symmetric strict monoidal structure

$$\left(f_!, f_!^\otimes, f_{!|*}^\otimes\right): (\mathcal{P}(A), \cap, A) \rightarrow (\mathcal{P}(B), \cap, B),$$

being equipped with equalities

$$\begin{aligned} f_{!|U,V}^{\otimes} : f_!(U \cap V) &\xrightarrow{=} f_!(U) \cap f_!(V), \\ f_{!|*}^{\otimes} : f_!(A) &\xrightarrow{=} B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

007L 7. *Relation to Direct Images.* We have

$$f_!(U) = B \setminus f_*(A \setminus U)$$

for each $U \in \mathcal{P}(A)$.

007M 8. *Interaction With Injections.* If f is injective, then we have

$$\begin{aligned} f_{!,\text{im}}(U) &= f_*(U), \\ f_{!,\text{cp}}(U) &= B \setminus \text{Im}(f), \\ f_!(U) &= f_{!,\text{im}}(U) \cup f_{!,\text{cp}}(U) \\ &= f_*(U) \cup (B \setminus \text{Im}(f)) \end{aligned}$$

for each $U \in \mathcal{P}(A)$.

007N 9. *Interaction With Surjections.* If f is surjective, then we have

$$\begin{aligned} f_{!,\text{im}}(U) &\subset f_*(U), \\ f_{!,\text{cp}}(U) &= \emptyset, \\ f_!(U) &\subset f_*(U) \end{aligned}$$

for each $U \in \mathcal{P}(A)$.

Proof. **Item 1, Functoriality:** Clear.

Item 2, Triple Adjointness: This follows from ?? of ??.

Item 3, Lax Preservation of Colimits: Omitted.

Item 4, Preservation of Limits: Omitted. This follows from **Item 2** and ?? of ??.

Item 5, Symmetric Lax Monoidality With Respect to Unions: This follows from ??.

Item 6, Symmetric Strict Monoidality With Respect to Intersections: This follows from **Item 4**.

Item 7, Relation to Direct Images: We claim that $f_!(U) = B \setminus f_*(A \setminus U)$.

- *The First Implication.* We claim that

$$f_!(U) \subset B \setminus f_*(A \setminus U).$$

Let $b \in f_!(U)$. We need to show that $b \notin f_*(A \setminus U)$, i.e. that there is no $a \in A \setminus U$ such that $f(a) = b$.

This is indeed the case, as otherwise we would have $a \in f^{-1}(b)$ and $a \notin U$, contradicting $f^{-1}(b) \subset U$ (which holds since $b \in f_!(U)$).

Thus $b \in B \setminus f_*(A \setminus U)$.

- *The Second Implication.* We claim that

$$B \setminus f_*(A \setminus U) \subset f_!(U).$$

Let $b \in B \setminus f_*(A \setminus U)$. We need to show that $b \in f_!(U)$, i.e. that $f^{-1}(b) \subset U$.

Since $b \notin f_*(A \setminus U)$, there exists no $a \in A \setminus U$ such that $b = f(a)$, and hence $f^{-1}(b) \subset U$.

Thus $b \in f_!(U)$.

This finishes the proof of **Item 7**.

Item 8, Interaction With Injections: Clear.

Item 9, Interaction With Surjections: Clear. \square

007P Proposition 2.4.6.1.6. Let $f: A \rightarrow B$ be a function.

007Q 1. *Functionality I.* The assignment $f \mapsto f_!$ defines a function

$$(-)_{!|A,B}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B)).$$

007R 2. *Functionality II.* The assignment $f \mapsto f_!$ defines a function

$$(-)_{!|A,B}: \text{Sets}(A, B) \rightarrow \text{Pos}((\mathcal{P}(A), \subset), (\mathcal{P}(B), \subset)).$$

007S 3. *Interaction With Identities.* For each $A \in \text{Obj}(\text{Sets})$, we have

$$(\text{id}_A)_! = \text{id}_{\mathcal{P}(A)};$$

007T 4. *Interaction With Composition.* For each pair of composable functions $f: A \rightarrow B$ and $g: B \rightarrow C$, we have

$$\begin{array}{ccc} \mathcal{P}(A) & \xrightarrow{f_!} & \mathcal{P}(B) \\ (g \circ f)_! & \searrow & \downarrow g_! \\ & & \mathcal{P}(C). \end{array}$$

Proof. **Item 1, Functionality I:** Clear.

Item 2, Functionality II: Clear.

Item 3, Interaction With Identities: This follows from ?? of ??.

Item 4, Interaction With Composition: This follows from ?? of ??.

\square

Appendices

2.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets
5. Relations
6. Spans
7. Posets

Indexed and Fibred Sets

7. Indexed Sets
8. Fibred Sets
9. Un/Straightening for Indexed and Fibred Sets

Category Theory

11. Categories
12. Types of Morphisms in Categories
13. Adjunctions and the Yoneda Lemma
14. Constructions With Categories
15. Profunctors
16. Cartesian Closed Categories
17. Kan Extensions

Bicategories

18. Bicategories
19. Internal Adjunctions

Internal Category Theory

20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

41. Stochastic Processes, Martingales, and Brownian Motion

Measure Theory

39. Measurable Spaces

42. Itô Calculus

40. Measures and Integration

43. Stochastic Differential Equations

Probability Theory

40. Probability Theory

Differential Geometry

44. Topological and Smooth Manifolds

Stochastic Analysis

Schemes

45. Schemes

2.2 Other Chapters

Sets

15. Profunctors

1. Sets

16. Cartesian Closed Categories

2. Constructions With Sets

17. Kan Extensions

3. Pointed Sets

Bicategories

4. Tensor Products of Pointed Sets

18. Bicategories

5. Relations

19. Internal Adjunctions

6. Spans

Internal Category Theory

7. Posets

20. Internal Categories

Indexed and Fibred Sets

Cyclic Stuff

7. Indexed Sets

21. The Cycle Category

8. Fibred Sets

Cubical Stuff

9. Un/Straightening for Indexed and Fibred Sets

22. The Cube Category

Category Theory

Globular Stuff

11. Categories

23. The Globe Category

12. Types of Morphisms in Categories

Cellular Stuff

13. Adjunctions and the Yoneda Lemma

24. The Cell Category

14. Constructions With Categories

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero	37. Real Analysis in One Variable
27. Monoids With Zero	38. Real Analysis in Several Variables
28. Constructions With Monoids With Zero	Measure Theory
Groups	39. Measurable Spaces
29. Groups	40. Measures and Integration
30. Constructions With Groups	Probability Theory
Hyper Algebra	40. Probability Theory
31. Hypermonoids	Stochastic Analysis
32. Hypergroups	41. Stochastic Processes, Martingales, and Brownian Motion
33. Hypersemirings and Hyperrings	42. Itô Calculus
34. Quantales	43. Stochastic Differential Equations
Near-Rings	Differential Geometry
35. Near-Semirings	44. Topological and Smooth Manifolds
36. Near-Rings	Schemes
Real Analysis	45. Schemes

Chapter 3

Pointed Sets

007U This chapter contains some foundational material on pointed sets.

Contents

3.1	Pointed Sets	84
3.1.1	Foundations	84
3.1.2	Morphisms of Pointed Sets	84
3.1.3	The Category of Pointed Sets.....	85
3.1.4	Elementary Properties of Pointed Sets.....	86
3.2	Limits of Pointed Sets.....	86
3.2.1	Products	86
3.2.2	Equalisers.....	87
3.2.3	Pullbacks.....	87
3.3	Colimits of Pointed Sets.....	87
3.3.1	Coproducts	87
3.3.2	Pushouts	87
3.3.3	Coequalisers.....	87
3.4	Constructions With Pointed Sets	88
3.4.1	Internal Homs	88
3.4.2	Free Pointed Sets.....	88
3.4.3	Wedge Sums of Pointed Sets	89
3.A	Other Chapters	92
3.2	Other Chapters	93

3.1 Pointed Sets

3.1.1 Foundations

007X Definition 3.1.1.1. A **pointed set**¹ is equivalently

- An \mathbb{E}_0 -monoid in $(N_\bullet(\text{Sets}), \text{pt})$;
- A pointed object in (Sets, pt) .

007Y Remark 3.1.1.2. In detail, a **pointed set** is a pair (X, x_0) consisting of

- *The Underlying Set.* A set X , called the **underlying set of** (X, x_0) ;
- *The Basepoint.* A morphism

$$[x_0] : \text{pt} \rightarrow X$$

in Sets , determining an element $x_0 \in X$, called the **basepoint of** X .

007Z Example 3.1.1.3. The **0-sphere**² is the pointed set $(S^0, 0)$ ³ consisting of

- *The Underlying Set.* The set S^0 defined by

$$S^0 \stackrel{\text{def}}{=} \{0, 1\};$$

- *The Basepoint.* The element 0 of S^0 .

0080 Example 3.1.1.4. The **trivial pointed set** is the pointed set (pt, \star) consisting of

- *The Underlying Set.* The punctual set $\text{pt} \stackrel{\text{def}}{=} \{\star\}$;
- *The Basepoint.* The element \star of pt .

0081 Example 3.1.1.5. The **underlying pointed set** of a semimodule (M, α_M) is the pointed set $(M, 0_M)$.

0082 Example 3.1.1.6. The **underlying pointed set** of a module (M, α_M) is the pointed set $(M, 0_M)$.

3.1.2 Morphisms of Pointed Sets

0084 Definition 3.1.2.1.1. A **morphism of pointed sets**⁴ is equivalently

- A morphism of \mathbb{E}_0 -monoids in $(N_\bullet(\text{Sets}), \text{pt})$.

¹Further Terminology: Also called an \mathbb{F}_1 -**module**.

²Further Terminology: Also called the **underlying pointed set of the field with one element**.

³Further Notation: Also denoted $(\mathbb{F}_1, 0)$.

⁴Further Terminology: Also called a **pointed function** or a **morphism of \mathbb{F}_1 -modules**.

- A morphism of pointed objects in (Sets, pt) .

0085 Remark 3.1.2.1.2. In detail, a **morphism of pointed sets** $f: (X, x_0) \rightarrow (Y, y_0)$ is a morphism of sets $f: X \rightarrow Y$ such that the diagram

$$\begin{array}{ccc} & \text{pt} & \\ [x_0] & \swarrow & \searrow [y_0] \\ X & \xrightarrow{f} & Y \end{array}$$

commutes, i.e. such that

$$f(x_0) = y_0.$$

3.1.3 The Category of Pointed Sets

0087 Definition 3.1.3.1.1. The **category of pointed sets** is the category Sets_* defined equivalently as

- The homotopy category of the ∞ -category $\text{Mon}_{\mathbb{E}_0}(\text{N}_*(\text{Sets}), \text{pt})$ of ??;
- The category Sets_* of ??.

0088 Remark 3.1.3.1.2. In detail, the **category of pointed sets** is the category Sets_* where

- *Objects.* The objects of Sets_* are pointed sets;
- *Morphisms.* The morphisms of Sets_* are morphisms of pointed sets;
- *Identities.* For each $(X, x_0) \in \text{Obj}(\text{Sets}_*)$, the unit map

$$\text{id}_{(X, x_0)}^{\text{Sets}_*}: \text{pt} \rightarrow \text{Sets}_*((X, x_0), (X, x_0))$$

of Sets_* at (X, x_0) is defined by⁵

$$\text{id}_{(X, x_0)}^{\text{Sets}_*} \stackrel{\text{def}}{=} \text{id}_X;$$

- *Composition.* For each $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$, the composition map

$$\circ_{(X, x_0), (Y, y_0), (Z, z_0)}^{\text{Sets}_*}: \text{Sets}_*((Y, y_0), (Z, z_0)) \times \text{Sets}_*((X, x_0), (Y, y_0)) \rightarrow \text{Sets}_*((X, x_0), (Z, z_0))$$

of Sets_* at $((X, x_0), (Y, y_0), (Z, z_0))$ is defined by⁶

$$g \circ_{(X, x_0), (Y, y_0), (Z, z_0)}^{\text{Sets}_*} f \stackrel{\text{def}}{=} g \circ f.$$

⁵Note that id_X is indeed a morphism of pointed sets, as we have $\text{id}_X(x_0) = x_0$.

⁶Note that the composition of two morphisms of pointed sets is indeed a morphism of pointed sets, as

3.1.4 Elementary Properties of Pointed Sets

008A Proposition 3.1.4.1.1. Let (X, x_0) be a pointed set.

- 008B** 1. *Completeness.* The category Sets_* of pointed sets and morphisms between them is complete, having in particular products (Definition 3.2.1.1), pullbacks (Definition 3.2.3.1.1), and equalisers (Definition 3.2.2.1.1).
- 008C** 2. *Cocompleteness.* The category Sets_* of pointed sets and morphisms between them is cocomplete, having in particular coproducts (Definition 3.3.1.1), pushouts (Definition 3.3.2.1.1), and coequalisers (Definition 3.3.3.1.1).
- 008D** 3. *Failure To Be Cartesian Closed.* The category Sets_* is not Cartesian closed.
- 008E** 4. *Relation to Partial Functions.* We have an equivalence of categories⁷

$$\text{Sets}_* \xrightarrow{\text{eq.}} \text{Sets}^{\text{part.}}$$

between the category of pointed sets and pointed functions between them and the category of sets and partial functions between them.

Proof. Item 1, Completeness: Omitted.

Item 2, Cocompleteness: Omitted.

Item 3, Failure To Be Cartesian Closed: See [MSE2855868].

Item 4, Relation to Partial Functions: Omitted. □

3.2 Limits of Pointed Sets

3.2.1 Products

Let (X, x_0) and (Y, y_0) be pointed sets.

008H Definition 3.2.1.1.1. The **product** of (X, x_0) and (Y, y_0) is the pointed set $(X \times Y, (x_0, y_0))$.

we have

$$\begin{aligned} g(f(x_0)) &= g(y_0) \\ &= z_0, \end{aligned}$$

or

$$\begin{array}{ccccc} & & \text{pt} & & \\ & \swarrow & \downarrow & \searrow & \\ [x_0] & & [y_0] & & [z_0] \\ \downarrow & & \downarrow & & \downarrow \\ X & \xrightarrow{f} & Y & \xrightarrow{g} & Z \end{array}$$

in terms of diagrams.

⁷ 

Warning: This is not an isomorphism of categories, only an equivalence.
END TEXTDBEND

3.2.2 Equalisers

Let $f, g: (X, x_0) \rightrightarrows (Y, y_0)$ be morphisms of pointed sets.

008K Definition 3.2.2.1.1. The **equaliser of** (f, g) is the pointed set $(\text{Eq}_*(f, g), x_0)$ consisting of

- *The Underlying Set.* The set $\text{Eq}_*(f, g)$ defined by

$$\text{Eq}_*(f, g) \stackrel{\text{def}}{=} \{x \in X \mid f(x) = y_0 = g(x)\};$$

- *The Basepoint.* The element x_0 of $\text{Eq}_*(f, g)$.

3.2.3 Pullbacks

Let (X, x_0) , (Y, y_0) , and (Z, z_0) be pointed sets and let $f: (X, x_0) \rightarrow (Z, z_0)$ and $g: (Y, y_0) \rightarrow (Z, z_0)$ be morphisms of pointed sets.

008M Definition 3.2.3.1.1. The **pullback of** (X, x_0) and (Y, y_0) over (Z, z_0) along (f, g) is the pointed set $((X, x_0) \times_{(Z, z_0)} (Y, y_0), p_0)$ consisting of

- *The Underlying Set.* The set $(X, x_0) \times_{(Z, z_0)} (Y, y_0)$ defined by

$$(X, x_0) \times_{(Z, z_0)} (Y, y_0) \stackrel{\text{def}}{=} \{(x, y) \in X \times Y \mid f(x) = z_0 = g(y)\};$$

- *The Basepoint.* The element (x_0, y_0) of $(X, x_0) \times_{(Z, z_0)} (Y, y_0)$.

3.3 Colimits of Pointed Sets

3.3.1 Coproducts

Let (X, x_0) and (Y, y_0) be pointed sets.

008Q Definition 3.3.1.1.1. The **coproduct of** (X, x_0) and (Y, y_0) is their wedge sum $(X \vee Y, p_0)$ of [Definition 3.4.3.1.1](#).

3.3.2 Pushouts

Let (X, x_0) , (Y, y_0) , and (Z, z_0) be pointed sets and let $f: (Z, z_0) \rightarrow (X, x_0)$ and $g: (Z, z_0) \rightarrow (Y, y_0)$ be morphisms of pointed sets.

008S Definition 3.3.2.1.1. The **pushout of** (X, x_0) and (Y, y_0) over (Z, z_0) along (f, g) is the pointed set $(X \coprod_{f, Z, g} Y, p_0)$, where $p_0 = [x_0] = [y_0]$.

3.3.3 Coequalisers

Let $f, g: (X, x_0) \rightrightarrows (Y, y_0)$ be morphisms of pointed sets.

008U Definition 3.3.3.1.1. The **coequaliser of** (f, g) is the pointed set $(\text{CoEq}(f, g), x_0)$.

3.4 Constructions With Pointed Sets

3.4.1 Internal Homs

Let (X, x_0) and (Y, y_0) be pointed sets.

008X Definition 3.4.1.1.1. The **pointed set of morphisms of pointed sets from** (X, x_0) **to** (Y, y_0) is the pointed set $\mathbf{Sets}_*((X, x_0), (Y, y_0))$ consisting of

- *The Underlying Set.* The set $\mathbf{Sets}_*((X, x_0), (Y, y_0))$ of morphisms of pointed sets from (X, x_0) to (Y, y_0) ;
- *The Basepoint.* The element

$$\Delta_{y_0}: (X, x_0) \rightarrow (Y, y_0)$$

of $\mathbf{Sets}_*((X, x_0), (Y, y_0))$.

3.4.2 Free Pointed Sets

Let X be a set.

008Z Definition 3.4.2.1.1. The **free pointed set on** X is the pointed set X^+ consisting of

- *The Underlying Set.* The set X^+ defined by

$$X^+ \stackrel{\text{def}}{=} X \coprod \text{pt};$$

- *The Basepoint.* The element \star of X^+ .

0090 Proposition 3.4.2.1.2. Let X be a set.

0091 1. Functoriality. The assignment $X \mapsto X^+$ defines a functor

$$(-)^+: \mathbf{Sets} \rightarrow \mathbf{Sets}_*,$$

where

- *Action on Objects.* For each $X \in \text{Obj}(\mathbf{Sets})$, we have

$$[(-)^+](X) \stackrel{\text{def}}{=} X_+,$$

where X_+ is the pointed set of **Definition 3.4.2.1.1**;

- *Action on Morphisms.* For each morphism $f: X \rightarrow Y$ of \mathbf{Sets} , the image

$$f_+: X_+ \rightarrow Y_+$$

of f by $(-)^+$ is the map of pointed sets defined by

$$f^+(x) \stackrel{\text{def}}{=} \begin{cases} f(x) & \text{if } x \in X, \\ \star & \text{if } x = \star. \end{cases}$$

0092 2. *Adjointness.* We have an adjunction

$$((-)^+ \dashv \overline{\text{忘}}): \text{Sets} \begin{array}{c} \xrightarrow{(-)^+} \\[-1ex] \perp \\[-1ex] \xleftarrow{\text{忘}} \end{array} \text{Sets}_*,$$

witnessed by a bijection of sets

$$\text{Sets}_*((X_+, \star), (Y, y_0)) \cong \text{Sets}(X, Y),$$

natural in $X \in \text{Obj}(\text{Sets})$ and $(Y, y_0) \in \text{Obj}(\text{Sets}_*)$.

0093 3. *Symmetric Strong Monoidality With Respect to Wedge Sums.* The free pointed set functor of [Item 1](#) has a symmetric strong monoidal structure

$$((-)^+, (-)^+, \coprod, (-)_\sharp^\pm, \coprod): (\text{Sets}, \coprod, \emptyset) \rightarrow (\text{Sets}_*, \vee, \text{pt}),$$

being equipped with isomorphisms of pointed sets

$$\begin{aligned} (-)_{X,Y}^{+, \coprod}: X^+ \vee Y^+ &\xrightarrow{\cong} (X \coprod Y)^+, \\ (-)_\sharp^\pm, \coprod: \text{pt} &\xrightarrow{\cong} \emptyset^+, \end{aligned}$$

natural in $X, Y \in \text{Obj}(\text{Sets})$.

0094 4. *Symmetric Strong Monoidality With Respect to Smash Products.* The free pointed set functor of [Item 1](#) has a symmetric strong monoidal structure

$$((-)^+, (-)^{+, \times}, (-)_\sharp^{+, \times}): (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Sets}_*, \wedge, S^0),$$

being equipped with isomorphisms of pointed sets

$$\begin{aligned} (-)_{X,Y}^{+, \times}: X^+ \wedge Y^+ &\xrightarrow{\cong} (X \times Y)^+, \\ (-)_\sharp^{+, \times}: S^0 &\xrightarrow{\cong} \text{pt}^+, \end{aligned}$$

natural in $X, Y \in \text{Obj}(\text{Sets})$.

Proof. [Item 1](#), *Functionality:* Clear.

[Item 2](#), *Adjointness:* Clear.

[Item 3](#), *Symmetric Strong Monoidality With Respect to Wedge Sums:* Omitted.

[Item 4](#), *Symmetric Strong Monoidality With Respect to Smash Products:* Omitted. \square

3.4.3 Wedge Sums of Pointed Sets

Let (X, x_0) and (Y, y_0) be pointed sets.

0096 **Definition 3.4.3.1.1.** The **wedge sum of X and Y** is the pointed set $(X \vee Y, p_0)$ consisting of

- *The Underlying Set.* The set $X \vee Y$ defined by⁸

$$(X \vee Y, p_0) \stackrel{\text{def}}{=} (X, x_0) \coprod (Y, y_0) \quad \begin{array}{ccc} X \vee Y & \xleftarrow{\lrcorner} & Y \\ \uparrow & & \uparrow [y_0] \\ X & \xleftarrow{[x_0]} & \text{pt}, \end{array}$$

$$\cong (X \coprod_{\text{pt}} Y, p_0)$$

$$\cong (X \coprod Y/\sim, p_0),$$

where \sim is the equivalence relation on $X \coprod Y$ given by $x_0 \sim y_0$;

- *The Basepoint.* The element p_0 of $X \vee Y$ defined by

$$p_0 \stackrel{\text{def}}{=} [x_0]$$

$$= [y_0].$$

0097 Proposition 3.4.3.1.2. Let (X, x_0) and (Y, y_0) be pointed sets.

0098 1. *Functionality.* The assignments $(X, x_0), (Y, y_0), ((X, x_0), (Y, y_0)) \mapsto (X \vee Y, p_0)$ define functors

$$\begin{aligned} X \vee - &: \text{Sets}_* \rightarrow \text{Sets}_*, \\ - \vee Y &: \text{Sets}_* \rightarrow \text{Sets}_*, \\ -_1 \vee -_2 &: \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*. \end{aligned}$$

0099 2. *Associativity.* We have an isomorphism of pointed sets

$$(X \vee Y) \vee Z \cong X \vee (Y \vee Z),$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Sets}_*$.

009A 3. *Unitality.* We have isomorphisms of pointed sets

$$\begin{aligned} \text{pt} \vee X &\cong X, \\ X \vee \text{pt} &\cong X, \end{aligned}$$

natural in $(X, x_0) \in \text{Sets}_*$.

009B 4. *Commutativity.* We have an isomorphism of pointed sets

$$X \vee Y \cong Y \vee X,$$

natural in $(X, x_0), (Y, y_0) \in \text{Sets}_*$.

009C 5. *Symmetric Monoidality.* The triple $(\text{Sets}_*, \vee, \text{pt})$ is a symmetric monoidal category.

⁸Here $(X, x_0) \coprod (Y, y_0)$ is the coproduct of (X, x_0) and (Y, y_0) in Sets_* .

- 009D** 6. *Symmetric Strong Monoidality With Respect to Free Pointed Sets.* The free pointed set functor of [Item 1](#) of [Proposition 3.4.2.1.2](#) has a symmetric strong monoidal structure

$$\left((-)^+, (-)^+, \coprod, (-)_{\sharp}^+, \coprod\right): (\text{Sets}, \coprod, \emptyset) \rightarrow (\text{Sets}_*, \vee, \text{pt}),$$

being equipped with isomorphisms of pointed sets

$$\begin{aligned} (-)_{X,Y}^{+, \coprod}: X^+ \vee Y^+ &\xrightarrow{\cong} (X \coprod Y)^+, \\ (-)_{\sharp}^{+, \coprod}: \text{pt} &\xrightarrow{\cong} \emptyset^+, \end{aligned}$$

natural in $X, Y \in \text{Obj}(\text{Sets})$.

- 009E** 7. *The Fold Map.* We have a natural transformation

$$\nabla: \vee \circ \Delta_{\text{Sets}_*}^{\text{Cats}} \Rightarrow \text{id}_{\text{Sets}_*},$$

```

    \begin{CD}
    Sets_* \times Sets_*
    @V \Delta_{Sets_*}^{\text{Cats}} VV \\
    Sets_* @= Sets_*
    @VV \nabla V \\
    Sets_* @>>> Sets_*
    @V id_{Sets_*} VV
    \end{CD}
  
```

called the **fold map**, whose component

$$\nabla_X: X \vee X \rightarrow X$$

at X is given by the composition

$$\begin{aligned} X &\xrightarrow{\Delta_X} X \times X \\ &\longrightarrow X \times X / \sim \\ &\xrightarrow{\text{def}} X \vee X. \end{aligned}$$

Proof. [Item 1](#), *Functionality:* Omitted.

[Item 2](#), *Associativity:* Omitted.

[Item 3](#), *Unitality:* Omitted.

[Item 4](#), *Commutativity:* Omitted.

[Item 5](#), *Symmetric Monoidality:* Omitted.

[Item 6](#), *Symmetric Strong Monoidality With Respect to Free Pointed Sets:* Omitted.

[Item 7](#), *The Fold Map:* Omitted. □

Appendices

3.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets
5. Relations
6. Spans
7. Posets

Indexed and Fibred Sets

7. Indexed Sets
8. Fibred Sets
9. Un/Straightening for Indexed and Fibred Sets

Category Theory

11. Categories
12. Types of Morphisms in Categories
13. Adjunctions and the Yoneda Lemma
14. Constructions With Categories
15. Profunctors
16. Cartesian Closed Categories
17. Kan Extensions

Bicategories

18. Bicategories
19. Internal Adjunctions

Internal Category Theory

20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

41. Stochastic Processes, Martingales, and Brownian Motion

Measure Theory

39. Measurable Spaces

42. Itô Calculus

40. Measures and Integration

43. Stochastic Differential Equations

Probability Theory

40. Probability Theory

Differential Geometry

44. Topological and Smooth Manifolds

Stochastic Analysis

Schemes

45. Schemes

3.2 Other Chapters

Sets

15. Profunctors

1. Sets

16. Cartesian Closed Categories

2. Constructions With Sets

17. Kan Extensions

3. Pointed Sets

Bicategories

4. Tensor Products of Pointed Sets

18. Bicategories

5. Relations

19. Internal Adjunctions

6. Spans

Internal Category Theory

7. Posets

20. Internal Categories

Indexed and Fibred Sets

Cyclic Stuff

7. Indexed Sets

21. The Cycle Category

8. Fibred Sets

Cubical Stuff

9. Un/Straightening for Indexed and Fibred Sets

22. The Cube Category

Category Theory

Globular Stuff

11. Categories

23. The Globe Category

12. Types of Morphisms in Categories

Cellular Stuff

13. Adjunctions and the Yoneda Lemma

24. The Cell Category

14. Constructions With Categories

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero	37. Real Analysis in One Variable
27. Monoids With Zero	38. Real Analysis in Several Variables
28. Constructions With Monoids With Zero	Measure Theory
Groups	39. Measurable Spaces
29. Groups	40. Measures and Integration
30. Constructions With Groups	Probability Theory
Hyper Algebra	40. Probability Theory
31. Hypermonoids	Stochastic Analysis
32. Hypergroups	41. Stochastic Processes, Martingales, and Brownian Motion
33. Hypersemirings and Hyperrings	42. Itô Calculus
34. Quantales	43. Stochastic Differential Equations
Near-Rings	Differential Geometry
35. Near-Semirings	44. Topological and Smooth Manifolds
36. Near-Rings	Schemes
Real Analysis	45. Schemes

Chapter 4

Tensor Products of Pointed Sets

009F This chapter contains some material on tensor products of pointed sets.

Contents

4.1	Bilinear Morphisms of Pointed Sets.....	96
4.1.1	Left Bilinear Morphisms of Pointed Sets	96
4.1.2	Right Bilinear Morphisms of Pointed Sets.....	96
4.1.3	Bilinear Morphisms of Pointed Sets.....	97
4.2	Tensors and Cotensors of Pointed Sets by Sets	98
4.2.1	Tensors of Pointed Sets by Sets.....	98
4.2.2	Cotensors of Pointed Sets by Sets	99
4.3	The Left Tensor Product of Pointed Sets	100
4.3.1	Foundations	100
4.3.2	The Skew Associator	101
4.3.3	The Skew Left Unitor.....	102
4.3.4	The Skew Right Unitor.....	102
4.3.5	The Left-Skew Monoidal Category Structure on Pointed Sets	103
4.4	The Right Tensor Product of Pointed Sets	103
4.4.1	Foundations	103
4.4.2	The Skew Associator	105
4.4.3	The Skew Left Unitor.....	106
4.4.4	The Skew Right Unitor.....	106
4.4.5	The Right-Skew Monoidal Category Structure on Pointed Sets	107
4.5	Smash Products of Pointed Sets.....	107
4.5.1	Foundations	107
4.A	Other Chapters	114
4.2	Other Chapters	115

4.1 Bilinear Morphisms of Pointed Sets

4.1.1 Left Bilinear Morphisms of Pointed Sets

Let (X, x_0) , (Y, y_0) , and (Z, z_0) be pointed sets.

009J Definition 4.1.1.1. A left bilinear morphism of pointed sets from $(X \times Y, (x_0, y_0))$ to (Z, z_0) is a map of sets

$$f: (X \times Y, (x_0, y_0)) \rightarrow (Z, z_0)$$

satisfying the following condition:^{1,2}

(★) *Left Unital Bilinearity.* The diagram

$$\begin{array}{ccc} & \text{pt} \times \text{pt} & \\ \text{id}_{\text{pt}} \times \epsilon_Y \nearrow & \swarrow \curvearrowright & \\ \text{pt} \times Y & & \text{pt} \\ \downarrow [x_0] \times \text{id}_Y & & \downarrow [z_0] \\ X \times Y & \xrightarrow{f} & Z \end{array}$$

commutes, i.e. for each $y \in Y$, we have

$$f(x_0, y) = z_0.$$

009K Definition 4.1.1.2. The set of left bilinear morphisms of pointed sets from $(X \times Y, (x_0, y_0))$ to (Z, z_0) is the set $\text{Hom}_{\text{Sets}_*}^{\otimes, L}(X \times Y, Z)$ defined by

$$\text{Hom}_{\text{Sets}_*}^{\otimes, L}(X \times Y, Z) \stackrel{\text{def}}{=} \{f \in \text{Sets}_*(A \times B, C) \mid f \text{ is left bilinear}\}.$$

4.1.2 Right Bilinear Morphisms of Pointed Sets

Let (X, x_0) , (Y, y_0) , and (Z, z_0) be pointed sets.

009M Definition 4.1.2.1.1. A right bilinear morphism of pointed sets from $(X \times Y, (x_0, y_0))$ to (Z, z_0) is a map of sets

$$f: (X \times Y, (x_0, y_0)) \rightarrow (Z, z_0)$$

satisfying the following condition:^{3,4}

¹Slogan: f is left bilinear if it preserves basepoints in its first argument.

²Succinctly, f is bilinear if we have

$$f(x_0, y) = z_0$$

for each $y \in Y$.

³Slogan: f is right bilinear if it preserves basepoints in its second argument.

⁴Succinctly, f is bilinear if we have

$$f(x, y_0) = z_0$$

(★) *Right Unital Bilinearity.* The diagram

$$\begin{array}{ccc}
 & \text{pt} \times \text{pt} & \\
 \swarrow_{\epsilon_X \times \text{id}_{\text{pt}}} & & \searrow \curvearrowright \\
 X \times \text{pt} & & \text{pt} \\
 \downarrow \text{id}_X \times [y_0] & & \downarrow [z_0] \\
 X \times Y & \xrightarrow{f} & Z
 \end{array}$$

commutes, i.e. for each $x \in X$, we have

$$f(x, y_0) = z_0.$$

009N Definition 4.1.2.1.2. The **set of right bilinear morphisms of pointed sets from** $(X \times Y, (x_0, y_0))$ **to** (Z, z_0) is the set $\text{Hom}_{\text{Sets}_*}^{\otimes, R}(X \times Y, Z)$ defined by

$$\text{Hom}_{\text{Sets}_*}^{\otimes, R}(X \times Y, Z) \stackrel{\text{def}}{=} \{f \in \text{Sets}_*(A \times B, C) \mid f \text{ is right bilinear}\}.$$

4.1.3 Bilinear Morphisms of Pointed Sets

Let (X, x_0) , (Y, y_0) , and (Z, z_0) be pointed sets.

009Q Definition 4.1.3.1.1. A **bilinear morphism of pointed sets from** $(X \times Y, (x_0, y_0))$ **to** (Z, z_0) is a map of sets

$$f: (X \times Y, (x_0, y_0)) \rightarrow (Z, z_0)$$

that is both left bilinear and right bilinear.

009R Remark 4.1.3.1.2. In detail, a **bilinear morphism of pointed sets from** $(X \times Y, (x_0, y_0))$ **to** (Z, z_0) is a map of sets

$$f: (X \times Y, (x_0, y_0)) \rightarrow (Z, z_0)$$

satisfying the following conditions:^{5,6}

for each $x \in X$.

⁵Slogan: f is bilinear if it preserves basepoints in each argument.

⁶Succinctly, f is bilinear if we have

$$\begin{aligned}
 f(x_0, y) &= z_0, \\
 f(x, y_0) &= z_0
 \end{aligned}$$

for each $x \in X$ and each $y \in Y$.

1. *Left Unital Bilinearity.* The diagram

$$\begin{array}{ccc}
 & \text{pt} \times \text{pt} & \\
 \text{id}_{\text{pt}} \times \epsilon_Y \nearrow & \swarrow \text{dashed} & \\
 \text{pt} \times Y & & \text{pt} \\
 \downarrow [x_0] \times \text{id}_Y & & \downarrow [z_0] \\
 X \times Y & \xrightarrow{f} & Z
 \end{array}$$

commutes, i.e. for each $y \in Y$, we have

$$f(x_0, y) = z_0.$$

2. *Right Unital Bilinearity.* The diagram

$$\begin{array}{ccc}
 & \text{pt} \times \text{pt} & \\
 \epsilon_X \times \text{id}_{\text{pt}} \nearrow & \swarrow \text{dashed} & \\
 X \times \text{pt} & & \text{pt} \\
 \downarrow \text{id}_X \times [y_0] & & \downarrow [z_0] \\
 X \times Y & \xrightarrow{f} & Z
 \end{array}$$

commutes, i.e. for each $x \in X$, we have

$$f(x, y_0) = z_0.$$

009S Definition 4.1.3.1.3. The **set of bilinear morphisms of pointed sets from** $(X \times Y, (x_0, y_0))$ **to** (Z, z_0) is the set $\text{Hom}_{\text{Sets}_*}^\otimes(X \times Y, Z)$ defined by

$$\text{Hom}_{\text{Sets}_*}^\otimes(X \times Y, Z) \stackrel{\text{def}}{=} \{f \in \text{Sets}_*(A \times B, C) \mid f \text{ is bilinear}\}.$$

4.2 Tensors and Cotensors of Pointed Sets by Sets

4.2.1 Tensors of Pointed Sets by Sets

Let (X, x_0) be a pointed set and let A be a set.

009V Definition 4.2.1.1.1. The **tensor of** (X, x_0) **by** A is the pointed set $A \odot (X, x_0)$ satisfying the following universal property:

(UP) We have a bijection

$$\text{Sets}_*(A \odot X, K) \cong \text{Sets}(A, \text{Sets}_*(X, K)),$$

natural in $(K, k_0) \in \text{Obj}(\text{Sets}_*)$.

009W Remark 4.2.1.1.2. The tensor of (X, x_0) by A satisfies the following universal property:

$$\text{Sets}_*(A \odot X, K) \cong \text{Sets}_{\mathbb{B}_0}^\otimes(A \times X, K),$$

where $\text{Sets}_{\mathbb{B}_0}^\otimes(A \times X, K)$ is the set defined by

$$\text{Sets}_{\mathbb{B}_0}^\otimes(A \times X, K) \stackrel{\text{def}}{=} \left\{ f \in \text{Sets}(A \times X, K) \mid \begin{array}{l} \text{for each } a \in A, \text{ we have} \\ f(a, x_0) = k_0 \end{array} \right\}.$$

009X Construction 4.2.1.1.3. Concretely, the **tensor of** (X, x_0) **by** A is the pointed set $A \odot (X, x_0)$ consisting of

- *The Underlying Set.* The set $A \odot X$ given by

$$A \odot X \cong \bigvee_{a \in A} (X, x_0);$$

- *The Basepoint.* The point $[x_0]$ of $\bigvee_{a \in A} (X, x_0)$.

4.2.2 Cotensors of Pointed Sets by Sets

Let (X, x_0) be a pointed set and let A be a set.

009Z Definition 4.2.2.1.1. The **cotensor of** (X, x_0) **by** A is the pointed set $A \pitchfork (X, x_0)$ satisfying the following universal property:

(UP) We have a bijection

$$\text{Sets}_*(K, A \pitchfork X) \cong \text{Sets}(A, \text{Sets}_*(K, X)),$$

natural in $(K, k_0) \in \text{Obj}(\text{Sets}_*)$.

00A0 Remark 4.2.2.1.2. The cotensor of (X, x_0) by A satisfies the following universal property:

$$\text{Sets}_*(K, A \pitchfork X) \cong \text{Sets}_{\mathbb{B}_0}^\otimes(A \times K, X),$$

where $\text{Sets}_{\mathbb{B}_0}^\otimes(A \times K, X)$ is the set defined by

$$\text{Sets}_{\mathbb{B}_0}^\otimes(A \times K, X) \stackrel{\text{def}}{=} \left\{ f \in \text{Sets}(A \times K, X) \mid \begin{array}{l} \text{for each } a \in A, \text{ we have} \\ f(a, k_0) = x_0 \end{array} \right\}.$$

00A1 Construction 4.2.2.1.3. Concretely, the **cotensor of** (X, x_0) **by** A is the pointed set $A \pitchfork (X, x_0)$ consisting of

- *The Underlying Set.* The set $A \pitchfork X$ given by

$$A \pitchfork X \cong \bigwedge_{a \in A} (X, x_0);$$

- *The Basepoint.* The point $[(x_0, x_0, x_0, \dots)]$ of $\bigwedge_{a \in A} (X, x_0)$.

4.3 The Left Tensor Product of Pointed Sets

4.3.1 Foundations

Let (X, x_0) and (Y, y_0) be pointed sets.

00A4 Definition 4.3.1.1.1. The **left tensor product of pointed sets** is the functor

$$\triangleleft_{\text{Sets}_*} : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

defined as the composition

$$\text{Sets}_* \times \text{Sets}_* \xrightarrow{\text{id} \times \overline{\triangleleft}} \text{Sets}_* \times \text{Sets} \xrightarrow{\beta_{\text{Sets}_*, \text{Sets}}^{\text{Cats}_2}} \text{Sets} \times \text{Sets}_* \xrightarrow{\odot} \text{Sets}_*.$$

00A5 Remark 4.3.1.1.2. The left tensor product of pointed sets satisfies the following universal property:⁷

$$\text{Sets}_*(X \triangleleft_{\text{Sets}_*} Y, Z) \cong \text{Hom}_{\text{Sets}_*}^{\otimes, L}(X \times Y, Z).$$

00A6 Remark 4.3.1.1.3. In detail, the **left tensor product of** (X, x_0) and (Y, y_0) is the pointed set $(X \triangleleft_{\text{Sets}_*} Y, [x_0])$ consisting of⁸

- *The Underlying Set.* The set $X \triangleleft_{\text{Sets}_*} Y$ defined by

$$\begin{aligned} X \triangleleft_{\text{Sets}_*} Y &\stackrel{\text{def}}{=} |Y| \odot X \\ &\cong \bigvee_{y \in Y} (X, x_0); \end{aligned}$$

- *The Underlying Basepoint.* The point $[x_0]$ of $\bigvee_{y \in Y} (X, x_0)$.

00A7 Proposition 4.3.1.1.4. Let (X, x_0) and (Y, y_0) be pointed sets.

00A8 1. Functoriality. The assignments $X, Y, (X, Y) \mapsto X \triangleleft_{\text{Sets}_*} Y$ define functors

$$\begin{aligned} X \triangleleft_{\text{Sets}_*} - &: \text{Sets}_* \rightarrow \text{Sets}_*, \\ - \triangleleft_{\text{Sets}_*} Y &: \text{Sets}_* \rightarrow \text{Sets}_*, \\ -_1 \triangleleft_{\text{Sets}_*} -_2 &: \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*. \end{aligned}$$

Proof. Item 1, Functoriality: Omitted. □

⁷Namely, a pointed map $f : X \triangleleft_{\text{Sets}_*} Y \rightarrow Z$ is the same as a map $f^\dagger : X \times Y \rightarrow Z$ such that

$$f^\dagger(x_0, y) = z_0$$

for each $y \in Y$.

⁸Further Notation: We write $x \triangleleft_{\text{Sets}_*} y$ for the image of (x, y) under the map

$$X \times Y \rightarrow \underbrace{X \triangleleft_{\text{Sets}_*} Y}_{\cong \bigvee_{y \in Y} (X, x_0)}.$$

4.3.2 The Skew Associator

00AA Definition 4.3.2.1.1. The **skew associator of the left tensor product of pointed sets** is the natural isomorphism

$$\alpha^{\text{Sets}_*, \triangleleft} : \triangleleft_{\text{Sets}_*} \circ (\triangleleft_{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}) \xrightarrow{\cong} \triangleleft_{\text{Sets}_*} \circ (\text{id}_{\text{Sets}_*} \times \triangleleft_{\text{Sets}_*}),$$

whose component

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft} : (X \triangleleft_{\text{Sets}_*} Y) \triangleleft_{\text{Sets}_*} Z \xrightarrow{\cong} X \triangleleft_{\text{Sets}_*} (Y \triangleleft_{\text{Sets}_*} Z)$$

at (X, Y, Z) is given by the composition⁹

$$\begin{aligned} (X \triangleleft_{\text{Sets}_*} Y) \triangleleft_{\text{Sets}_*} Z &\stackrel{\text{def}}{=} |Z| \odot (X \triangleleft_{\text{Sets}_*} Y) \\ &\stackrel{\text{def}}{=} |Z| \odot (|Y| \odot X) \\ &\cong \bigvee_{z \in Z} (|Y| \odot X, [x_0]) \\ &\stackrel{\text{def}}{=} \bigvee_{z \in Z} \left(\bigvee_{y \in Y} (X, x_0) \right) \\ &\cong \bigvee_{(z,y) \in \bigvee_{z \in Z} (Y, x_0)} (X, x_0) \\ &\stackrel{\text{def}}{=} \bigvee_{(z,y) \in |Z| \odot Y} (X, x_0) \\ &\cong |Z| \odot Y \odot X \\ &\stackrel{\text{def}}{=} |Y \triangleleft_{\text{Sets}_*} Z| \odot X \\ &\stackrel{\text{def}}{=} X \triangleleft_{\text{Sets}_*} (Y \triangleleft_{\text{Sets}_*} Z), \end{aligned}$$

where the isomorphism

$$\bigvee_{z \in Z} \left(\bigvee_{y \in Y} (X, x_0) \right) \cong \bigvee_{(y,z) \in \bigvee_{z \in Z} (Y, x_0)} (X, x_0)$$

is given by $[(z, (y, x))] \mapsto [(z, y), x]$.

sending (x, y) to the element $x \in X$ in the y th copy of X in $\bigvee_{y \in Y} (X, x_0)$. Note that we have

$$x_0 \triangleleft_{\text{Sets}_*} y = x_0 \triangleleft_{\text{Sets}_*} y',$$

for each $y, y' \in Y$.

⁹In other words, $\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft}$ acts on elements as

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleleft} ((x \triangleleft_{\text{Sets}_*} y) \triangleleft_{\text{Sets}_*} z) \stackrel{\text{def}}{=} x \triangleleft_{\text{Sets}_*} (y \triangleleft_{\text{Sets}_*} z)$$

for each $(x \triangleleft_{\text{Sets}_*} y) \triangleleft_{\text{Sets}_*} z \in (X \triangleleft_{\text{Sets}_*} Y) \triangleleft_{\text{Sets}_*} Z$.

4.3.3 The Skew Left Unitor

00AC Definition 4.3.3.1.1. The **skew left unitor of the left tensor product of pointed sets** is the natural transformation

$$\lambda^{\text{Sets}_*, \triangleleft} : \triangleleft_{\text{Sets}_*} \circ (\text{id}_{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}) \Rightarrow \text{id}_{\text{Sets}_*},$$

whose component

$$\lambda_X^{\text{Sets}_*, \triangleleft} : S^0 \triangleleft_{\text{Sets}_*} X \rightarrow X$$

at X is given by the composition¹⁰

$$\begin{aligned} S^0 \triangleleft_{\text{Sets}_*} X &\cong |X| \odot S^0 \\ &\cong \bigvee_{x \in X} S^0 \\ &\rightarrow X \end{aligned}$$

where $\bigvee_{x \in X} S^0 \rightarrow X$ is the map given by

$$\begin{aligned} (x, 0) &\mapsto x, \\ (x, 1) &\mapsto x. \end{aligned}$$

4.3.4 The Skew Right Unitor

00AE Definition 4.3.4.1.1. The **skew right unitor of the left tensor product of pointed sets** is the natural transformation

$$\rho^{\text{Sets}_*, \triangleleft} : \text{id}_{\text{Sets}_*} \Rightarrow \triangleleft_{\text{Sets}_*} \circ (\text{id}_{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}),$$

whose component

$$\rho_X^{\text{Sets}_*, \triangleleft} : X \rightarrow X \triangleleft_{\text{Sets}_*} S^0$$

at X is given by the composition¹¹

$$\begin{aligned} X &\rightarrow X \vee X \\ &\cong |S^0| \odot X \\ &\cong S^0 \triangleleft_{\text{Sets}_*} X, \end{aligned}$$

where $X \rightarrow X \vee X$ is the map sending X to the first factor of X in $X \vee X$.

¹⁰In other words, $\lambda_X^{\text{Sets}_*, \triangleleft}$ acts on elements as

$$\begin{aligned} \lambda_X^{\text{Sets}_*, \triangleleft}(x \triangleleft_{\text{Sets}_*} 0) &\stackrel{\text{def}}{=} x, \\ \lambda_X^{\text{Sets}_*, \triangleleft}(x \triangleleft_{\text{Sets}_*} 1) &\stackrel{\text{def}}{=} x, \end{aligned}$$

for each $x \in X$.

¹¹In other words, $\rho_X^{\text{Sets}_*, \triangleleft}$ acts on elements as

$$\rho_X^{\text{Sets}_*, \triangleleft}(x) \stackrel{\text{def}}{=} x \triangleleft_{\text{Sets}_*} 0$$

4.3.5 The Left-Skew Monoidal Category Structure on Pointed Sets

00AG Proposition 4.3.5.1.1. The category Sets_* admits a left-skew monoidal category structure consisting of¹²

- *The Skew Monoidal Product.* The left tensor product functor

$$\triangleleft_{\text{Sets}_*} : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*,$$

of [Proposition 4.3.1.1.4](#);

- *The Skew Monoidal Unit.* The functor

$$\mathbb{1}_{\text{Sets}_*} : \text{pt} \rightarrow \text{Sets}_*$$

defined by

$$\mathbb{1}_{\text{Sets}_*} \stackrel{\text{def}}{=} S^0;$$

- *The Skew Associators.* The natural isomorphism

$$\alpha^{\text{Sets}_*, \triangleleft} : \triangleleft_{\text{Sets}_*} \circ (\triangleleft_{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}) \xrightarrow{\cong} \triangleleft_{\text{Sets}_*} \circ (\text{id}_{\text{Sets}_*} \times \triangleleft_{\text{Sets}_*}),$$

of [Definition 4.3.2.1.1](#);

- *The Skew Left Unitors.* The natural transformation

$$\lambda^{\text{Sets}_*, \triangleleft} : \triangleleft_{\text{Sets}_*} \circ (\mathbb{1}_{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}) \Rightarrow \text{id}_{\text{Sets}_*},$$

of [Definition 4.3.3.1.1](#);

- *The Skew Right Unitors.* The natural transformation

$$\rho^{\text{Sets}_*, \triangleleft} : \text{id}_{\text{Sets}_*} \Rightarrow \triangleleft_{\text{Sets}_*} \circ (\text{id}_{\text{Sets}_*} \times \mathbb{1}_{\text{Sets}_*}),$$

of [Definition 4.3.4.1.1](#).

Proof. Omitted. □

4.4 The Right Tensor Product of Pointed Sets

4.4.1 Foundations

Let (X, x_0) and (Y, y_0) be pointed sets.

for each $x \in X$.

¹²Note in particular that, differently from general left-skew monoidal categories, the skew associator of $(\text{Sets}_*, \triangleleft_{\text{Sets}_*}, S^0)$ is a natural isomorphism.

00AK Definition 4.4.1.1.1. The **right tensor product of pointed sets** is the functor

$$\triangleright_{\text{Sets}_*} : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

defined as the composition

$$\text{Sets}_* \times \text{Sets}_* \xrightarrow{\cong \otimes \times \text{id}} \text{Sets} \times \text{Sets}_* \xrightarrow{\odot} \text{Sets}_*.$$

00AL Remark 4.4.1.1.2. The right tensor product of pointed sets satisfies the following universal property:¹³

$$\text{Sets}_*(X \triangleright_{\text{Sets}_*} Y, Z) \cong \text{Hom}_{\text{Sets}_*}^{\otimes, R}(X \times Y, Z).$$

00AM Remark 4.4.1.1.3. In detail, the **right tensor product of** (X, x_0) **and** (Y, y_0) is the pointed set $(X \triangleright_{\text{Sets}_*} Y, [y_0])$ consisting of¹⁴

- *The Underlying Set.* The set $X \triangleright_{\text{Sets}_*} Y$ defined by

$$\begin{aligned} X \triangleright_{\text{Sets}_*} Y &\stackrel{\text{def}}{=} |X| \odot Y \\ &\cong \bigvee_{x \in X} (Y, y_0); \end{aligned}$$

- *The Underlying Basepoint.* The point $[y_0]$ of $\bigvee_{x \in X} (Y, y_0)$.

00AN Proposition 4.4.1.1.4. Let (X, x_0) and (Y, y_0) be pointed sets.

00AP 1. Functoriality. The assignments $X, Y, (X, Y) \mapsto X \triangleright_{\text{Sets}_*} Y$ define functors

$$\begin{aligned} X \triangleright_{\text{Sets}_*} - &: \text{Sets}_* \rightarrow \text{Sets}_*, \\ - \triangleright_{\text{Sets}_*} Y &: \text{Sets}_* \rightarrow \text{Sets}_*, \\ -_1 \triangleright_{\text{Sets}_*} -_2 &: \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*. \end{aligned}$$

Proof. Item 1, Functoriality: Omitted. □

¹³Namely, a pointed map $f: X \triangleleft_{\text{Sets}_*} Y \rightarrow Z$ is the same as a map $f^\dagger: X \times Y \rightarrow Z$ such that

$$f^\dagger(x, y_0) = z_0$$

for each $y \in Y$.

¹⁴Further Notation: We write $x \triangleright_{\text{Sets}_*} y$ for the image of (x, y) under the map

$$\begin{aligned} X \times Y \rightarrow & \underbrace{X \triangleright_{\text{Sets}_*} Y}_{\cong \bigvee_{x \in X} (Y, y_0)} . \end{aligned}$$

sending (x, y) to the element $y \in Y$ in the x th copy of Y in $\bigvee_{x \in X} (Y, y_0)$. Note that we have

$$x \triangleright_{\text{Sets}_*} y_0 = x' \triangleright_{\text{Sets}_*} y_0,$$

for each $x, x' \in X$.

4.4.2 The Skew Associator

00AR Definition 4.4.2.1.1. The **skew associator of the right tensor product of pointed sets** is the natural isomorphism

$$\alpha^{\text{Sets}_*, \triangleright} : \triangleright_{\text{Sets}_*} \circ (\text{id}_{\text{Sets}_*} \times \triangleright_{\text{Sets}_*}) \xrightarrow{\cong} \triangleright_{\text{Sets}_*} \circ (\triangleright_{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}),$$

whose component

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleright} : X \triangleright_{\text{Sets}_*} (Y \triangleright_{\text{Sets}_*} Z) \xrightarrow{\cong} (X \triangleright_{\text{Sets}_*} Y) \triangleright_{\text{Sets}_*} Z$$

at (X, Y, Z) is given by the composition¹⁵

$$\begin{aligned} X \triangleright_{\text{Sets}_*} (Y \triangleright_{\text{Sets}_*} Z) &\stackrel{\text{def}}{=} |X| \odot (Y \triangleright_{\text{Sets}_*} Z) \\ &\stackrel{\text{def}}{=} |X| \odot (|Y| \odot Z) \\ &\cong |X| \odot \left(\bigvee_{y \in Y} (Z, z_0) \right) \\ &\cong \bigvee_{x \in X} \left(\bigvee_{y \in Y} (Z, z_0) \right) \\ &\cong \bigvee_{(x,y) \in \bigvee_{x \in X} (Y, y_0)} (Z, z_0) \\ &\cong \left| \bigvee_{x \in X} (Y, y_0) \right| \odot Z \\ &\stackrel{\text{def}}{=} |X \odot Y| \odot Z \\ &\stackrel{\text{def}}{=} |X \triangleright_{\text{Sets}_*} Y| \odot Z \\ &\stackrel{\text{def}}{=} (X \triangleright_{\text{Sets}_*} Y) \triangleright_{\text{Sets}_*} Z \end{aligned}$$

where the isomorphism

$$\bigvee_{x \in X} \left(\bigvee_{y \in Y} (Z, z_0) \right) \cong \bigvee_{(x,y) \in \bigvee_{x \in X} (Y, y_0)} (Z, z_0)$$

is given by $[(x, (y, z))] \mapsto [(x, y), z]$.

¹⁵In other words, $\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleright}$ acts on elements as

$$\alpha_{X,Y,Z}^{\text{Sets}_*, \triangleright} (x \triangleright_{\text{Sets}_*} (y \triangleright_{\text{Sets}_*} z)) \stackrel{\text{def}}{=} (x \triangleright_{\text{Sets}_*} y) \triangleright_{\text{Sets}_*} z$$

for each $x \triangleright_{\text{Sets}_*} (y \triangleright_{\text{Sets}_*} z) \in X \triangleright_{\text{Sets}_*} (Y \triangleright_{\text{Sets}_*} Z)$.

4.4.3 The Skew Left Unitor

00AT Definition 4.4.3.1.1. The **skew left unitor of the right tensor product of pointed sets** is the natural transformation

$$\lambda^{\text{Sets}_*, \triangleright} : \text{id}_{\text{Sets}_*} \Rightarrow \triangleright_{\text{Sets}_*} \circ (\text{id}_{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}),$$

whose component

$$\lambda_X^{\text{Sets}_*, \triangleright} : X \rightarrow S^0 \triangleright_{\text{Sets}_*} X$$

at X is given by the composition¹⁶

$$\begin{aligned} X &\rightarrow X \vee X \\ &\cong |S^0| \odot X \\ &\cong S^0 \triangleright_{\text{Sets}_*} X, \end{aligned}$$

where $X \rightarrow X \vee X$ is the map sending X to the first factor of X in $X \vee X$.

4.4.4 The Skew Right Unitor

00AV Definition 4.4.4.1.1. The **skew right unitor of the right tensor product of pointed sets** is the natural transformation

$$\rho^{\text{Sets}_*, \triangleright} : \triangleright_{\text{Sets}_*} \circ (\text{id}_{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}) \Rightarrow \text{id}_{\text{Sets}_*},$$

whose component¹⁷

$$\rho_X^{\text{Sets}_*, \triangleright} : X \triangleright_{\text{Sets}_*} S^0 \rightarrow X$$

at X is given by the composition

$$\begin{aligned} X \triangleright_{\text{Sets}_*} S^0 &\cong |X| \odot S^0 \\ &\cong \bigvee_{x \in X} S^0 \\ &\rightarrow X \end{aligned}$$

where $\bigvee_{x \in X} S^0 \rightarrow X$ is the map given by

$$\begin{aligned} (x, 0) &\mapsto x, \\ (x, 1) &\mapsto x. \end{aligned}$$

¹⁶In other words, $\lambda_X^{\text{Sets}_*, \triangleright}$ acts on elements as

$$\lambda_X^{\text{Sets}_*, \triangleright}(x) \stackrel{\text{def}}{=} 0 \triangleright_{\text{Sets}_*} x$$

for each $x \in X$.

¹⁷In other words, $\rho_X^{\text{Sets}_*, \triangleright}$ acts on elements as

$$\begin{aligned} \rho_X^{\text{Sets}_*, \triangleright}(x \triangleright_{\text{Sets}_*} 0) &\stackrel{\text{def}}{=} x, \\ \rho_X^{\text{Sets}_*, \triangleright}(x \triangleright_{\text{Sets}_*} 1) &\stackrel{\text{def}}{=} x \end{aligned}$$

4.4.5 The Right-Skew Monoidal Category Structure on Pointed Sets

00AX Proposition 4.4.5.1.1. The category Sets_* admits a right-skew monoidal category structure consisting of¹⁸

- *The Skew Monoidal Product.* The right tensor product functor

$$\triangleright_{\text{Sets}_*} : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*,$$

of Item 1;

- *The Skew Monoidal Unit.* The functor

$$\mathbb{1}_{\text{Sets}_*}^{\text{Sets}_*} : \text{pt} \rightarrow \text{Sets}_*$$

defined by

$$\mathbb{1}_{\text{Sets}_*}^{\text{Sets}_*} \stackrel{\text{def}}{=} S^0;$$

- *The Skew Associators.* The natural isomorphism

$$\alpha^{\text{Sets}_*, \triangleright} : \triangleright_{\text{Sets}_*} \circ (\triangleright_{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}) \xrightarrow{\cong} \triangleright_{\text{Sets}_*} \circ (\text{id}_{\text{Sets}_*} \times \triangleright_{\text{Sets}_*}),$$

of Definition 4.4.2.1.1;

- *The Skew Left Unitors.* The natural transformation

$$\lambda^{\text{Sets}_*, \triangleright} : \text{id}_{\text{Sets}_*} \Longrightarrow \triangleright_{\text{Sets}_*} \circ (\mathbb{1}_{\text{Sets}_*}^{\text{Sets}_*} \times \text{id}_{\text{Sets}_*}),$$

of Definition 4.3.3.1.1;

- *The Skew Right Unitors.* The natural transformation

$$\rho^{\text{Sets}_*, \triangleright} : \triangleright_{\text{Sets}_*} \circ (\text{id}_{\text{Sets}_*} \times \mathbb{1}_{\text{Sets}_*}^{\text{Sets}_*}) \Longrightarrow \text{id}_{\text{Sets}_*},$$

of Definition 4.3.4.1.1.

Proof. Omitted. □

4.5 Smash Products of Pointed Sets

4.5.1 Foundations

Let (X, x_0) and (Y, y_0) be pointed sets.

for each $x \in X$.

¹⁸Note in particular that, differently from general right-skew monoidal categories, the skew associator of $(\text{Sets}_*, \triangleright_{\text{Sets}_*}, S^0)$ is a natural isomorphism.

00B0 Definition 4.5.1.1.1. The **smash product of** (X, x_0) **and** (Y, y_0) ¹⁹ is the pointed set $X \wedge Y$ ²⁰ such that we have a bijection

$$\text{Sets}_*(X \wedge Y, Z) \cong \text{Hom}_{\text{Sets}_*}^\otimes(X \times Y, Z),$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$.

00B1 Remark 4.5.1.1.2. In detail, the **smash product of** (X, x_0) **and** (Y, y_0) is the pair $((X \wedge Y, [(x_0, y_0)]), \iota)$ consisting of

- A pointed set $(X \wedge Y, [(x_0, y_0)])$;
- A bilinear morphism of pointed sets $\iota: (X \times Y, (x_0, y_0)) \rightarrow X \wedge Y$;

satisfying the following universal property:

(UP) Given another such pair $((Z, z_0), f)$ consisting of

- A pointed set (Z, z_0) ;
- A bilinear morphism of pointed sets $f: (X \times Y, (x_0, y_0)) \rightarrow Z$;

there exists a unique morphism of pointed sets $X \wedge Y \xrightarrow{\exists!} Z$ making the diagram

$$\begin{array}{ccc} & X \wedge Y & \\ \iota \nearrow & \downarrow \exists! & \\ X \times Y & \xrightarrow{f} & Z \end{array}$$

commute.

00B2 Construction 4.5.1.1.3. Concretely, the **smash product of** (X, x_0) **and** (Y, y_0) is the pointed set $(X \wedge Y, [(x_0, y_0)])$ consisting of²¹

¹⁹Further Terminology: Also called the **tensor product of \mathbb{F}_1 -modules of** (X, x_0) **and** (Y, y_0) or the **tensor product of** (X, x_0) **and** (Y, y_0) **over** \mathbb{F}_1 .

²⁰Further Notation: Also written $X \otimes_{\mathbb{F}_1} Y$.

²¹Further Notation: We write $x \wedge y$ for the image of (x, y) under the quotient map

$$X \times Y \twoheadrightarrow \frac{X \times Y}{\underbrace{X \vee Y}_{\stackrel{\text{def}}{=} X \wedge Y}}$$

Note that we have

$$\begin{aligned} x \wedge y_0 &= x' \wedge y_0, \\ x_0 \wedge y &= x_0 \wedge y' \end{aligned}$$

for each $x, x' \in X$ and each $y, y' \in Y$.

- *The Underlying Set.* The set $X \wedge Y$ defined by

$$\begin{aligned} X \wedge Y &\cong \text{pt} \coprod_{X \vee Y} (X \times Y) \\ &\stackrel{\text{def}}{=} \frac{X \times Y}{X \vee Y} \\ &\cong X \times Y / \sim, \end{aligned} \quad \begin{array}{ccc} X \wedge Y & \xleftarrow{\lrcorner} & X \times Y \\ \uparrow & & \uparrow \\ \text{pt} & \xleftarrow{!} & X \vee Y, \end{array}$$

where \sim is the equivalence relation of $X \times Y$ obtained by declaring $(x, y) \sim (x', y')$ iff $(x, y), (x', y') \in X \vee Y$, i.e. by declaring

$$\begin{aligned} (x_0, y) &\sim (x_0, y'), \\ (x, y_0) &\sim (x', y_0) \end{aligned}$$

for all $x \in X$ and all $y \in Y$;

- *The Basepoint.* The element $[(x_0, y_0)]$ of $X \wedge Y$ given by the equivalence class of (x_0, y_0) under the equivalence relation \sim on $X \times Y$.

Proof. Clear. □

00B3 Example 4.5.1.1.4. Here are some examples of smash products of pointed sets.

1. *Smashing With S^0 .* For any pointed set X , we have isomorphisms of pointed sets

$$\begin{aligned} S^0 \wedge X &\cong X, \\ X \wedge S^0 &\cong X. \end{aligned}$$

00B4 Proposition 4.5.1.1.5. Let (X, x_0) and (Y, y_0) be pointed sets.

00B5 1. *Functoriality.* The assignments $(X, x_0), (Y, y_0), ((X, x_0), (Y, y_0)) \mapsto X \wedge Y$ define functors

$$\begin{aligned} X \wedge -: \text{Sets}_* &\rightarrow \text{Sets}_*, \\ - \wedge Y: \text{Sets}_* &\rightarrow \text{Sets}_*, \\ -_1 \wedge -_2: \text{Sets}_* \times \text{Sets}_* &\rightarrow \text{Sets}_*. \end{aligned}$$

00B6 2. *Adjointness.* We have adjunctions

$$\begin{aligned} (X \wedge - \dashv \text{Sets}_*(X, -)): \quad \text{Sets}_* &\begin{array}{c} \xrightarrow{X \wedge -} \\ \perp \\ \xleftarrow{\text{Sets}_*(X, -)} \end{array} \text{Sets}_*, \\ (- \wedge Y \dashv \text{Sets}_*(Y, -)): \quad \text{Sets}_* &\begin{array}{c} \xrightarrow{- \wedge Y} \\ \perp \\ \xleftarrow{\text{Sets}_*(Y, -)} \end{array} \text{Sets}_*, \end{aligned}$$

witnessed by bijections

$$\begin{aligned}\text{Sets}_*(X \wedge Y, Z) &\cong \text{Sets}_*(X, \mathbf{Sets}_*(Y, Z)), \\ \text{Sets}_*(X \wedge Y, Z) &\cong \text{Sets}_*(X, \mathbf{Sets}_*(A, Z)),\end{aligned}$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$, which internalise to isomorphisms of pointed sets

$$\begin{aligned}\mathbf{Sets}_*(X \wedge Y, Z) &\cong \mathbf{Sets}_*(X, \mathbf{Sets}_*(Y, Z)), \\ \mathbf{Sets}_*(X \wedge Y, Z) &\cong \mathbf{Sets}_*(X, \mathbf{Sets}_*(A, Z)),\end{aligned}$$

again natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$.

00B7 3. *Closed Symmetric Monoidality.* The quadruple $(\text{Sets}_*, \wedge, S^0, \mathbf{Sets}_*)$ is a closed symmetric monoidal category.

00B8 4. *Morphisms From the Monoidal Unit.* We have a bijection of sets²²

$$\text{Sets}_*(S^0, X) \cong X,$$

natural in $(X, x_0) \in \text{Obj}(\text{Sets}_*)$, internalising also to an isomorphism of pointed sets

$$\mathbf{Sets}_*(S^0, X) \cong (X, x_0),$$

again natural in $(X, x_0) \in \text{Obj}(\text{Sets}_*)$.

00B9 5. *Symmetric Strong Monoidality With Respect to Free Pointed Sets.* The free pointed set functor of ?? of ?? has a symmetric strong monoidal structure

$$((-)^+, (-)^{+, \times}, (-)_\sharp^{+, \times}): (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Sets}_*, \wedge, S^0),$$

being equipped with isomorphisms of pointed sets

$$\begin{aligned}(-)^{+, \times}_{X, Y}: X^+ \wedge Y^+ &\xrightarrow{\cong} (X \times Y)^+, \\ (-)_\sharp^{+, \times}: S^0 &\xrightarrow{\cong} \text{pt}^+,\end{aligned}$$

natural in $X, Y \in \text{Obj}(\text{Sets})$.

00BA 6. *Distributivity Over Wedge Sums.* We have isomorphisms of pointed sets

$$\begin{aligned}X \wedge (Y \vee Z) &\cong (X \wedge Y) \vee (X \wedge Z), \\ (X \vee Y) \wedge Z &\cong (X \wedge Z) \vee (Y \wedge Z),\end{aligned}$$

natural in $(X, x_0), (Y, y_0), (Z, z_0) \in \text{Obj}(\text{Sets}_*)$.

²²In other words, the forgetful functor

$$\mathfrak{U}: \text{Sets}_* \rightarrow \text{Sets}$$

00BB 7. *Universal Property I.* The symmetric monoidal structure on the category Sets_* is uniquely determined by the following requirements:

- (a) *Two-Sided Preservation of Colimits.* The smash product

$$\wedge : \text{Sets}_* \times \text{Sets}_* \rightarrow \text{Sets}_*$$

of Sets_* preserves colimits separately in each variable.

- (b) *The Unit Object Is S^0 .* We have $\mathbb{1}_{\text{Sets}_*} = S^0$.

00BC 8. *Universal Property II.* The symmetric monoidal structure on the category Sets_* is the unique symmetric monoidal structure on Sets_* such that the free pointed set functor

$$(-)^+ : \text{Sets} \rightarrow \text{Sets}_*$$

admits a symmetric monoidal structure.

00BD 9. *Existence of Monoidal Diagonals.* The triple $(\text{Sets}_*, \wedge, S^0)$ is a monoidal category with diagonals:

- (a) *Monoidal Diagonals.* The natural transformation

$$\Delta : \text{id}_{\text{Sets}_*} \Longrightarrow \wedge \circ \Delta_{\text{Sets}_*}^{\text{Cats}_2},$$

whose component

$$\Delta_X : (X, x_0) \rightarrow (X \wedge X, [(x_0, x_0)])$$

at $(X, x_0) \in \text{Obj}(\text{Sets}_*)$ is given by the composition

$$\begin{aligned} (X, x_0) &\xrightarrow{\Delta_X} (X \times X, (x_0, x_0)) \\ &\longrightarrow (\frac{X \times X}{X \vee X}, [(x_0, x_0)]) \\ &\xrightarrow{\text{def}} (X \wedge X, [(x_0, x_0)]) \end{aligned}$$

in Sets_* , is a monoidal natural transformation:

- i. *Naturality.* For each morphism $f : X \rightarrow Y$ of pointed sets, the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \Delta_X \downarrow & & \downarrow \Delta_Y \\ X \wedge X & \xrightarrow{f \wedge f} & Y \wedge Y \end{array}$$

commutes.

- ii. *Compatibility With Strong Monoidality Constraints.* For each $(X, x_0), (Y, y_0) \in \text{Obj}(\text{Sets}_*)$, the diagram

$$\begin{array}{ccc} X \wedge Y & \xrightarrow{\Delta_X \wedge \Delta_Y} & (X \wedge X) \wedge (Y \wedge Y) \\ \parallel & & \downarrow \\ X \wedge Y & \xrightarrow{\Delta_{X \wedge Y}} & (X \wedge Y) \wedge (X \wedge Y) \end{array}$$

commutes.

- iii. *Compatibility With Strong Unitality Constraints.* The diagram

$$\begin{array}{ccc} S^0 & & \\ \parallel & \searrow & (\lambda_{S^0}^{\text{Sets}_*})^{-1} = (\rho_{S^0}^{\text{Sets}_*})^{-1} \\ S^0 & \xrightarrow{\Delta_{S^0}} & S^0 \wedge S^0 \end{array}$$

commutes.

- (b) *The Diagonal of the Unit.* The component

$$\Delta_{S^0}^{\text{Sets}_*} : S^0 \xrightarrow{\cong} S^0 \wedge S^0$$

of the diagonal natural transformation of Sets_* at S^0 is an isomorphism.

- 00BE** 10. *Comonoids in Sets_* .* The symmetric monoidal functor

$$((-)^+, (-)^{+,\times}, (-)_\times^{+,\times}) : (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Sets}_*, \wedge, S^0),$$

of ?? of ?? lifts to an equivalence of categories

$$\begin{aligned} \text{CoMon}(\text{Sets}_*, \wedge, S^0) &\stackrel{\text{eq.}}{\cong} \text{CoMon}(\text{Sets}, \times, \text{pt}) \\ &\cong \text{Sets}. \end{aligned}$$

Proof. **Item 1, Functoriality:** Omitted.

Item 2, Adjointness: Omitted.

Item 3, Closed Symmetric Monoidality: Omitted.

Item 4, Morphisms From the Monoidal Unit: Omitted.

Item 5, Symmetric Strong Monoidality With Respect to Free Pointed Sets: Omitted.

Item 6, Distributivity Over Wedge Sums: This follows from **Item 3**, ?? of ??, and the fact that \vee is the coproduct in Sets_* .

Item 7, Universal Property I: Omitted.

Item 8, Universal Property II: See [GGN15, Theorem 5.1].

Item 9, Existence of Monoidal Diagonals: Omitted.

Item 10, Comonoids in Sets_* : See [PS19, Lemma 2.4].

□

Appendices

defined on objects by sending a pointed set to its underlying set is corepresentable by S^0 .

4.A Other Chapters

Sets

- 1. Sets
- 2. Constructions With Sets
- 3. Pointed Sets
- 4. Tensor Products of Pointed Sets
- 5. Relations
- 6. Spans
- 7. Posets

Indexed and Fibred Sets

- 7. Indexed Sets
- 8. Fibred Sets
- 9. Un/Straightening for Indexed and Fibred Sets

Category Theory

- 11. Categories
- 12. Types of Morphisms in Categories
- 13. Adjunctions and the Yoneda Lemma
- 14. Constructions With Categories
- 15. Profunctors
- 16. Cartesian Closed Categories
- 17. Kan Extensions

Bicategories

- 18. Bicategories
- 19. Internal Adjunctions

Internal Category Theory

- 20. Internal Categories

Cyclic Stuff

- 21. The Cycle Category

Cubical Stuff

- 22. The Cube Category

Globular Stuff

- 23. The Globe Category

Cellular Stuff

- 24. The Cell Category

Monoids

- 25. Monoids
- 26. Constructions With Monoids

Monoids With Zero

- 27. Monoids With Zero
- 28. Constructions With Monoids With Zero

Groups

- 29. Groups
- 30. Constructions With Groups

Hyper Algebra

- 31. Hypermonoids
- 32. Hypergroups
- 33. Hypersemirings and Hyperrings
- 34. Quantales

Near-Rings

- 35. Near-Semirings
- 36. Near-Rings

Real Analysis

- 37. Real Analysis in One Variable

38. Real Analysis in Several Variables

41. Stochastic Processes, Martingales, and Brownian Motion

Measure Theory

39. Measurable Spaces

42. Itô Calculus

40. Measures and Integration

43. Stochastic Differential Equations

Probability Theory

40. Probability Theory

Differential Geometry

44. Topological and Smooth Manifolds

Stochastic Analysis

Schemes

45. Schemes

4.2 Other Chapters

Sets

15. Profunctors

1. Sets

16. Cartesian Closed Categories

2. Constructions With Sets

17. Kan Extensions

3. Pointed Sets

Bicategories

4. Tensor Products of Pointed Sets

18. Bicategories

5. Relations

19. Internal Adjunctions

6. Spans

Internal Category Theory

7. Posets

20. Internal Categories

Indexed and Fibred Sets

Cyclic Stuff

7. Indexed Sets

21. The Cycle Category

8. Fibred Sets

Cubical Stuff

9. Un/Straightening for Indexed and Fibred Sets

22. The Cube Category

Category Theory

Globular Stuff

11. Categories

23. The Globe Category

12. Types of Morphisms in Categories

Cellular Stuff

13. Adjunctions and the Yoneda Lemma

24. The Cell Category

14. Constructions With Categories

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero	37. Real Analysis in One Variable
27. Monoids With Zero	38. Real Analysis in Several Variables
28. Constructions With Monoids With Zero	Measure Theory
Groups	39. Measurable Spaces
29. Groups	40. Measures and Integration
30. Constructions With Groups	Probability Theory
Hyper Algebra	40. Probability Theory
31. Hypermonoids	Stochastic Analysis
32. Hypergroups	41. Stochastic Processes, Martingales, and Brownian Motion
33. Hypersemirings and Hyperrings	42. Itô Calculus
34. Quantales	43. Stochastic Differential Equations
Near-Rings	Differential Geometry
35. Near-Semirings	44. Topological and Smooth Manifolds
36. Near-Rings	Schemes
Real Analysis	45. Schemes

Chapter 5

Relations

00BF This chapter contains some material about relations and constructions with them. Notably, we discuss and explore:

1. The definition of relations ([Section 5.1.1](#)).
2. How relations may be viewed as decategorification of profunctors ([Section 5.1.2](#)).
3. The various kind of categories that relations form, namely:
 - (a) A category ([Section 5.2.1](#)),
 - (b) A monoidal category ([Section 5.2.2](#)),
 - (c) A 2-category ([Section 5.2.3](#)), and
 - (d) A double category ([Section 5.2.4](#)).
4. The various categorical properties of the 2-category of relations, including ([Section 5.2.5](#)):
 - (a) The self-duality of **Rel** and **Rel** ([Items 1 and 2 of Proposition 5.2.5.1.1](#));
 - (b) Identifications of equivalences and isomorphisms in **Rel** with bijections ([Item 3 of Proposition 5.2.5.1.1](#));
 - (c) Identifications of adjunctions in **Rel** with functions ([Item 4 of Proposition 5.2.5.1.1](#));
 - (d) Identifications of monads in **Rel** with preorders ([Item 5 of Proposition 5.2.5.1.1](#));
 - (e) Identifications of comonads in **Rel** with subsets ([Item 6 of Proposition 5.2.5.1.1](#));
 - (f) Characterisations of monomorphisms in **Rel** ([Item 7 of Proposition 5.2.5.1.1](#));
 - (g) Characterisations of epimorphisms in **Rel** ([Item 8 of Proposition 5.2.5.1.1](#));
 - (h) The partial co/completeness of **Rel** ([Item 10 of Proposition 5.2.5.1.1](#));
 - (i) The existence of right Kan extensions and right Kan lifts in **Rel** ([Items 11 and 12 of Proposition 5.2.5.1.1](#));
 - (j) The closedness of **Rel** ([Item 13 of Proposition 5.2.5.1.1](#)).

-
5. The various kinds of constructions involving relations, such as graphs, domains, ranges, unions, intersections, products, inverse relations, composition of relations, and collages ([Section 5.3](#)).
 6. Equivalence relations ([Section 5.4](#)) and quotient sets ([Section 5.4.5](#)).
 7. The adjoint pairs

$$\begin{aligned} R_* \dashv R_{-1} : \mathcal{P}(A) \rightleftarrows \mathcal{P}(B), \\ R^{-1} \dashv R_! : \mathcal{P}(B) \rightleftarrows \mathcal{P}(A) \end{aligned}$$

of functors (morphisms of posets) between $\mathcal{P}(A)$ and $\mathcal{P}(B)$ induced by a relation $R: A \rightarrow B$, as well as the properties of R_* , R_{-1} , R^{-1} , and $R_!$ ([Section 5.5](#)).

Here we also note that:

- (a) These two pairs of adjoint functors are the counterpart for relations of the adjoint triple $f_* \dashv f^{-1} \dashv f_!$ induced by a function $f: A \rightarrow B$ studied in [??](#);
- (b) We have $R_{-1} = R^{-1}$ iff R is total and functional ([Item 8 of Proposition 5.5.2.1.3](#)).
- (c) As a consequence of the previous item, when R comes from a function f the pair of adjunctions

$$R_* \dashv R_{-1} = R^{-1} \dashv R_!$$

reduces to the triple adjunction

$$f_* \dashv f^{-1} \dashv f_!$$

from before.

- (d) The pairs $R_* \dashv R_{-1}$ and $R^{-1} \dashv R_!$ later make an appearance in the context of continuous, open, and closed relations between topological spaces ([??](#)).

8. A notion of *relative preorder* that is to that of a preorder as relative monads are to monads, extending the identifications of monads in Rel with preorders of [Item 5 of Proposition 5.2.5.1.1](#) to “relative monads in Rel”.

Contents

5.1 Relations	120
5.1.1 Foundations	120
5.1.2 Relations as Decategorifications of Profunctors	122
5.1.3 Examples of Relations	124
5.1.4 Functional Relations	125
5.1.5 Total Relations	126
5.2 Categories of Relations	126
5.2.1 The Category of Relations	126

5.2.2	The Closed Symmetric Monoidal Category of Relations .	127
5.2.3	The 2-Category of Relations.....	131
5.2.4	The Double Category of Relations.....	132
5.2.5	Properties of the Category of Relations.....	138
5.3	Constructions With Relations.....	150
5.3.1	The Graph of a Function.....	150
5.3.2	The Inverse of a Function.....	154
5.3.3	Representable Relations.....	155
5.3.4	The Domain and Range of a Relation.....	156
5.3.5	Binary Unions of Relations	157
5.3.6	Unions of Families of Relations.....	158
5.3.7	Binary Intersections of Relations.....	158
5.3.8	Intersections of Families of Relations	159
5.3.9	Binary Products of Relations.....	160
5.3.10	Products of Families of Relations.....	162
5.3.11	The Inverse of a Relation.....	162
5.3.12	Composition of Relations.....	163
5.3.13	The Collage of a Relation	168
5.4	Equivalence Relations	169
5.4.1	Reflexive Relations	169
5.4.2	Symmetric Relations	172
5.4.3	Transitive Relations.....	174
5.4.4	Equivalence Relations	176
5.4.5	Quotients by Equivalence Relations	178
5.5	Functoriality of Powersets	182
5.5.1	Direct Images.....	182
5.5.2	Strong Inverse Images	187
5.5.3	Weak Inverse Images.....	192
5.5.4	Direct Images With Compact Support.....	196
5.5.5	Functoriality of Powersets	202
5.5.6	Functoriality of Powersets: Relations on Powersets	202
5.6	Relative Preorders.....	203
5.6.1	The Left Skew Monoidal Structure on $\mathbf{Rel}(A, B)$	203
5.6.2	Left Relative Preorders	206
5.6.3	The Right Skew Monoidal Structure on $\mathbf{Rel}(A, B)$	208
5.6.4	Right Relative Preorders	210
5.A	Other Chapters	212
5.2	Other Chapters	214

5.1 Relations

5.1.1 Foundations

Let A and B be sets.

00BJ Definition 5.1.1.1. A **relation** $R: A \rightarrow B$ from A to B ^{1,2} is a subset R of $A \times B$.³

00BK Definition 5.1.1.2. Let A and B be sets.

00BL 1. The **set of relations from A to B** is the set $\text{Rel}(A, B)$ defined by

$$\text{Rel}(A, B) \stackrel{\text{def}}{=} \{\text{Relations from } A \text{ to } B\}.$$

00BM 2. The **poset of relations from A to B** is the poset

$$\mathbf{Rel}(A, B) \stackrel{\text{def}}{=} (\text{Rel}(A, B), \subset)$$

consisting of

- *The Underlying Set.* The set $\text{Rel}(A, B)$ of **Item 1**;
- *The Partial Order.* The partial order

$$\subset: \text{Rel}(A, B) \times \text{Rel}(A, B) \rightarrow \{\text{true, false}\}$$

on $\text{Rel}(A, B)$ given by inclusion of relations.

00BN Remark 5.1.1.3. A relation from A to B is equivalently:⁴

00BP 1. A subset of $A \times B$;

00BQ 2. A function from $A \times B$ to $\{\text{true, false}\}$;

00BR 3. A function from A to $\mathcal{P}(B)$;

00BS 4. A function from B to $\mathcal{P}(A)$;

00BT 5. A cocontinuous morphism of posets from $(\mathcal{P}(A), \subset)$ to $(\mathcal{P}(B), \subset)$.

That is: we have bijections of sets

$$\begin{aligned} \text{Rel}(A, B) &\stackrel{\text{def}}{=} \mathcal{P}(A \times B), \\ &\cong \text{Sets}(A \times B, \{\text{true, false}\}), \\ &\cong \text{Sets}(A, \mathcal{P}(B)), \\ &\cong \text{Sets}(B, \mathcal{P}(A)), \\ &\cong \text{Hom}_{\mathbf{Pos}}^{\text{cocont}}(\mathcal{P}(A), \mathcal{P}(B)), \end{aligned}$$

natural in $A, B \in \text{Obj}(\text{Sets})$.

¹Further Terminology: Also called a **multivalued function** from A to B , a **relation over A and B** , **relation on A and B** , a **binary relation over A and B** , or a **binary relation on A and B** .

²Further Terminology: When $A = B$, we also call $R \subset A \times A$ a **relation on A** .

³Further Notation: Given elements $a \in A$ and $b \in B$, we write $a \sim_R b$ to mean $(a, b) \in R$.

⁴Intuition: In particular, we may think of a relation $R: A \rightarrow \mathcal{P}(B)$ from A to B as a multivalued function

Proof. We claim that **Items 1** to **5** are indeed equivalent:

- **Item 1** \iff **Item 2**: This is a special case of ?? of ??.
- **Item 2** \iff **Item 3**: This is an instance of currying, following from the bijections

$$\begin{aligned} \text{Sets}(A \times B, \{\text{true}, \text{false}\}) &\cong \text{Sets}(A, \text{Sets}(B, \{\text{true}, \text{false}\})) \\ &\cong \text{Sets}(A, \mathcal{P}(B)), \end{aligned}$$

where the last bijection is from ?? of ??.

- **Item 2** \iff **Item 4**: This is also an instance of currying, following from the bijections

$$\begin{aligned} \text{Sets}(A \times B, \{\text{true}, \text{false}\}) &\cong \text{Sets}(B, \text{Sets}(B, \{\text{true}, \text{false}\})) \\ &\cong \text{Sets}(B, \mathcal{P}(A)), \end{aligned}$$

where again the last bijection is from ?? of ??.

- **Item 2** \iff **Item 5**: This follows from the universal property of the powerset $\mathcal{P}(X)$ of a set X as the free cocompletion of X via the characteristic embedding

$$\chi_X: X \hookrightarrow \mathcal{P}(X)$$

of X into $\mathcal{P}(X)$ (?? of ??).

In particular, the bijection

$$\text{Rel}(A, B) \cong \text{Hom}_{\text{Pos}}^{\text{cocont}}(\mathcal{P}(A), \mathcal{P}(B))$$

is given by taking a relation $R: A \rightarrow B$, passing to its associated function $f: A \rightarrow \mathcal{P}(B)$ from A to B and then extending f from A to all of $\mathcal{P}(A)$ by taking its left Kan extension along χ_X .

This coincides with the direct image function $f_*: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$ of ??.

This finishes the proof. □

00BU Proposition 5.1.1.4. Let A and B be sets.

00BV 1. *End Formula for The Poset of Relations.* Let $R, S: A \rightarrow B$ be relations. We have

$$\text{Hom}_{\text{Rel}(A, B)}(R, S) \cong \int_{a \in A} \int_{b \in B} \text{Hom}_{\{\text{t}, \text{f}\}}(R_b^a, S_b^a).$$

Proof. Item 1, End Formula for The Poset of Relations: Unwinding the expression inside

from A to B (including the possibility of a given $a \in A$ having no value at all).

the end on the right hand side, we have

$$\int_{a \in A} \int_{b \in B} \text{Hom}_{\{\text{t,f}\}}(R_b^a, S_b^a) \cong \begin{cases} \text{pt} & \text{if for each } (a, b) \in A \times B, \\ & \text{if } a \sim_R b, \text{ then } a \sim_S b, \\ \emptyset & \text{otherwise.} \end{cases}$$

On the left hand-side, we have

$$\text{Hom}_{\mathbf{Rel}(A,B)}(R, S) \cong \begin{cases} \text{pt} & \text{if } R \subset S, \\ \emptyset & \text{otherwise.} \end{cases}$$

It is then clear that the conditions for each set to evaluate to pt are equivalent, implying that those two sets are isomorphic, finishing the proof. \square

5.1.2 Relations as Decategorifications of Profunctors

00BX Remark 5.1.2.1.1. The notion of a relation is a decategorification of that of a profunctor:

1. A profunctor from a category C to a category \mathcal{D} is a functor

$$p: \mathcal{D}^{\text{op}} \times C \rightarrow \text{Sets}.$$

2. A relation on sets A and B is a function

$$R: A \times B \rightarrow \{\text{true, false}\}.$$

Here we notice that:

- The opposite X^{op} of a set X is itself, as $(-)^\text{op}: \text{Cats} \rightarrow \text{Cats}$ restricts to the identity endofunctor on Sets ;
- The values that profunctors and relations take are directly related in relation to decategorification:
 - A category is enriched over the category

$$\text{Sets} \stackrel{\text{def}}{=} \text{Cats}_0$$

of sets, with profunctors taking values on it;

– A set is enriched over the set

$$\{\text{true, false}\} \stackrel{\text{def}}{=} \text{Cats}_{-1}$$

of classical truth values, with relations taking values on it;

00BY Remark 5.1.2.1.2. Extending Remark 5.1.2.1.1, the equivalent definitions of relations in Remark 5.1.1.1.3 are also related to the corresponding ones for profunctors (??), which state that a profunctor $p: C \nrightarrow D$ is equivalently:

- 00BZ 1. A functor $p: D^{\text{op}} \times C \rightarrow \text{Sets}$;
- 00C0 2. A functor $p: C \rightarrow \text{PSh}(D)$;
- 00C1 3. A functor $p: D^{\text{op}} \rightarrow \text{Fun}(C, \text{Sets})$;
- 00C2 4. A colimit-preserving functor $p: \text{PSh}(C) \rightarrow \text{PSh}(D)$.

Indeed:

- The equivalence between Items 1 and 2 (and also that between Items 1 and 3, which is proved analogously) is an instance of currying, both for profunctors as well as for relations, using the isomorphisms

$$\begin{aligned} \text{Sets}(A \times B, \{\text{true}, \text{false}\}) &\cong \text{Sets}(A, \text{Sets}(B, \{\text{true}, \text{false}\})) \\ &\cong \text{Sets}(A, \mathcal{P}(B)), \\ \text{Fun}(D^{\text{op}} \times D, \text{Sets}) &\cong \text{Fun}(C, \text{Fun}(D^{\text{op}}, \text{Sets})) \\ &\cong \text{Fun}(C, \text{PSh}(D)). \end{aligned}$$

- The equivalence between Items 1 and 3 follows from the universal properties of:
 - The powerset $\mathcal{P}(X)$ of a set X as the free cocompletion of X via the characteristic embedding

$$\chi_{(-)}: X \hookrightarrow \mathcal{P}(X)$$

of X into $\mathcal{P}(X)$ (?? of ??);

- The category $\text{PSh}(C)$ of presheaves on a category C as the free cocompletion of C via the Yoneda embedding

$$\mathfrak{y}: C \hookrightarrow \text{PSh}(C)$$

of C into $\text{PSh}(C)$ (?? of ??).

5.1.3 Examples of Relations

00C4 Example 5.1.3.1.1. The **trivial relation on A and B** is the relation \sim_{triv} defined by^{5,6,7}

$$\sim_{\text{triv}} \stackrel{\text{def}}{=} A \times A.$$

00C5 Example 5.1.3.1.2. The **cotrivial relation on A and B** is the relation \sim_{cotriv} defined by^{8,9,10}

$$\sim_{\text{cotriv}} \stackrel{\text{def}}{=} \emptyset.$$

00C6 Example 5.1.3.1.3. The characteristic relation on A of ?? of ?? is another example of a relation. It is in fact the unique relation on A making the following conditions equivalent, for each $a, b \in A$:

1. We have $a \sim_{\text{id}} b$.
2. We have $a = b$.

00C7 Example 5.1.3.1.4. Square roots are examples of relations:

1. *Square Roots in \mathbb{R} .* The assignment $x \mapsto \sqrt{x}$ defines a relation

$$\sqrt{-} : \mathbb{R} \rightarrow \mathcal{P}(\mathbb{R})$$

from \mathbb{R} to itself, being explicitly given by

$$\sqrt{x} \stackrel{\text{def}}{=} \begin{cases} 0 & \text{if } x = 0, \\ \{-\sqrt{|x|}, \sqrt{|x|}\} & \text{if } x \neq 0. \end{cases}$$

⁵This is the unique relation R on A and B such that we have $a \sim_R b$ for all $a \in A$ and all $b \in B$.

⁶As a function from $A \times A$ to $\{\text{true, false}\}$, the relation \sim_{triv} is the constant function

$$\Delta_{\text{true}} : A \times B \rightarrow \{\text{true, false}\}$$

from $A \times B$ to $\{\text{true, false}\}$ taking value true.

⁷As a function from A to $\mathcal{P}(B)$, the relation \sim_{triv} is the function

$$\Delta_{\text{true}} : A \rightarrow \mathcal{P}(B)$$

defined by

$$\Delta_{\text{true}}(a) \stackrel{\text{def}}{=} B$$

for each $a \in A$.

⁸This is the unique relation R on A and B such that we have $a \sim_R b$ for no $a \in A$ and no $b \in B$.

⁹As a function from $A \times B$ to $\{\text{true, false}\}$, the relation \sim_{cotriv} is the constant function

$$\Delta_{\text{false}} : A \times B \rightarrow \{\text{true, false}\}$$

from $A \times B$ to $\{\text{true, false}\}$ taking value false.

¹⁰As a function from A to $\mathcal{P}(A)$, the relation \sim_{cotriv} is the function

$$\Delta_{\text{false}} : A \rightarrow \mathcal{P}(A)$$

2. *Square Roots in \mathbb{Q} .* Square roots in \mathbb{Q} are similar to square roots in \mathbb{R} , though now additionally it may also occur that $\sqrt{-}: \mathbb{Q} \rightarrow \mathcal{P}(\mathbb{Q})$ sends a rational number x (e.g. 2) to the empty set (since $\sqrt{2} \notin \mathbb{Q}$).

00C8 Example 5.1.3.1.5. The complex logarithm defines a relation

$$\log: \mathbb{C} \rightarrow \mathcal{P}(\mathbb{C})$$

from \mathbb{C} to itself, where we have

$$\log(a + bi) \stackrel{\text{def}}{=} \left\{ \log\left(\sqrt{a^2 + b^2}\right) + i \arg(a + bi) + (2\pi i)k \mid k \in \mathbb{Z} \right\}$$

for each $a + bi \in \mathbb{C}$.

00C9 Example 5.1.3.1.6. See [[wikipedia:multivalued-functions](#)] for more examples of relations, such as antiderivation, inverse trigonometric functions, and inverse hyperbolic functions.

5.1.4 Functional Relations

Let A and B be sets.

00CB Definition 5.1.4.1.1. A relation $R: A \rightarrow B$ is **functional** if, for each $a \in A$, the set $R(a)$ is either empty or a singleton.

00CC Proposition 5.1.4.1.2. Let $R: A \rightarrow B$ be a relation.

00CD 1. *Characterisations.* The following conditions are equivalent:

00CE (a) The relation R is functional.

00CF (b) We have $R \diamond R^\dagger \subset \chi_B$.

Proof. **Item 1, Characterisations:** We claim that **Items 1a** and **1b** are indeed equivalent:

- **Item 1a** \implies **Item 1b**: Let $(b, b') \in B \times B$. We need to show that

$$[R \diamond R^\dagger](b, b') \leq_{\{\text{t,f}\}} \chi_B(b, b'),$$

i.e. that if there exists some $a \in A$ such that $b \sim_{R^\dagger} a$ and $a \sim_R b'$, then $b = b'$. But since $b \sim_{R^\dagger} a$ is the same as $a \sim_R b$, we have both $a \sim_R b$ and $a \sim_R b'$ at the same time, which implies $b = b'$ since R is functional.

- **Item 1b** \implies **Item 1a**: Suppose that we have $a \sim_R b$ and $a \sim_R b'$ for $b, b' \in B$. We claim that $b = b'$:

defined by

$$\Delta_{\text{true}}(a) \stackrel{\text{def}}{=} \emptyset$$

for each $a \in A$.

1. Since $a \sim_R b$, we have $b \sim_{R^\dagger} a$.
2. Since $R \diamond R^\dagger \subset \chi_B$, we have

$$[R \diamond R^\dagger](b, b') \leq_{\{\text{t,f}\}} \chi_B(b, b'),$$

and since $b \sim_{R^\dagger} a$ and $a \sim_R b'$, it follows that $[R \diamond R^\dagger](b, b') = \text{true}$, and thus $\chi_B(b, b') = \text{true}$ as well, i.e. $b = b'$.

This finishes the proof. \square

5.1.5 Total Relations

Let A and B be sets.

00CH Definition 5.1.5.1.1. A relation $R: A \rightarrow B$ is **total** if, for each $a \in A$, we have $R(a) \neq \emptyset$.

00CJ Proposition 5.1.5.1.2. Let $R: A \rightarrow B$ be a relation.

- 00CK** 1. *Characterisations.* The following conditions are equivalent:
- 00CL** (a) The relation R is total.
 - 00CM** (b) We have $\chi_A \subset R^\dagger \diamond R$.

Proof. **Item 1, Characterisations:** We claim that **Items 1a** and **1b** are indeed equivalent:

- **Item 1a** \implies **Item 1b**: We have to show that, for each $(a, a') \in A$, we have

$$\chi_A(a, a') \leq_{\{\text{t,f}\}} [R^\dagger \diamond R](a, a'),$$

i.e. that if $a = a'$, then there exists some $b \in B$ such that $a \sim_R b$ and $b \sim_{R^\dagger} a'$ (i.e. $a \sim_R b$ again), which follows from the totality of R .

- **Item 1b** \implies **Item 1a**: Given $a \in A$, since $\chi_A \subset R^\dagger \diamond R$, we must have

$$\{a\} \subset [R^\dagger \diamond R](a),$$

implying that there must exist some $b \in B$ such that $a \sim_R b$ and $b \sim_{R^\dagger} a$ (i.e. $a \sim_R b$) and thus $R(a) \neq \emptyset$, as $b \in R(a)$.

This finishes the proof. \square

5.2 Categories of Relations

5.2.1 The Category of Relations

00CQ Definition 5.2.1.1.1. The **category of relations** is the category Rel where

- *Objects.* The objects of Rel are sets;

- *Morphisms.* For each $A, B \in \text{Obj}(\text{Sets})$, we have

$$\text{Rel}(A, B) \stackrel{\text{def}}{=} \text{Rel}(A, B);$$

- *Identities.* For each $A \in \text{Obj}(\text{Rel})$, the unit map

$$\mathbb{1}_A^{\text{Rel}} : \text{pt} \rightarrow \text{Rel}(A, A)$$

of Rel at A is defined by

$$\text{id}_A^{\text{Rel}} \stackrel{\text{def}}{=} \chi_A(-_1, -_2),$$

where $\chi_A(-_1, -_2)$ is the characteristic relation of A of ?? of ??;

- *Composition.* For each $A, B, C \in \text{Obj}(\text{Rel})$, the composition map

$$\circ_{A,B,C}^{\text{Rel}} : \text{Rel}(B, C) \times \text{Rel}(A, B) \rightarrow \text{Rel}(A, C)$$

of Rel at (A, B, C) is defined by

$$S \circ_{A,B,C}^{\text{Rel}} R \stackrel{\text{def}}{=} S \diamond R$$

for each $(S, R) \in \text{Rel}(B, C) \times \text{Rel}(A, B)$, where $S \diamond R$ is the composition of S and R of [Definition 5.3.12.1.1](#).

5.2.2 The Closed Symmetric Monoidal Category of Relations

5.2.2.1 The Monoidal Product

00CT Definition 5.2.2.1.1. The **monoidal product** of Rel is the functor

$$\times : \text{Rel} \times \text{Rel} \rightarrow \text{Rel}$$

where

- *Action on Objects.* We have

$$\times(A, B) \stackrel{\text{def}}{=} A \times B,$$

where $A \times B$ is the Cartesian product of sets of ??;

- *Action on Morphisms.* For each $(A, C), (B, D) \in \text{Obj}(\text{Rel} \times \text{Rel})$, the action on morphisms

$$\times_{(A,C),(B,D)} : \text{Rel}(A, B) \times \text{Rel}(C, D) \rightarrow \text{Rel}(A \times C, B \times D)$$

of \times is given by sending a pair of morphisms (R, S) of the form

$$R : A \rightarrow B,$$

$$S : C \rightarrow D$$

to the relation

$$R \times S : A \times C \rightarrow B \times D$$

of [Definition 5.3.9.1.1](#).

5.2.2.2 The Monoidal Unit

00CV **Definition 5.2.2.2.1.** The **monoidal unit of Rel** is the functor

$$\mathbb{M}^{\text{Rel}} : \text{pt} \rightarrow \text{Rel}$$

picking the set

$$\mathbb{M}_{\text{Rel}} \stackrel{\text{def}}{=} \text{pt}$$

of Rel.

5.2.2.3 The Associator

00CX **Definition 5.2.2.3.1.** The **associator of Rel** is the natural isomorphism

$$\begin{array}{ccc} \alpha^{\text{Rel}} : \times \circ ((\times) \times \text{id}) & \xrightarrow{\cong} & \times \circ (\text{id} \times (\times)), \\ \text{id} \times (\times) \times \text{id} & \swarrow & \downarrow \text{id} \times (\times) \\ \text{Rel} \times \text{Rel} \times \text{Rel} & \xrightarrow{\quad \text{id} \times (\times) \quad} & \text{Rel} \times \text{Rel} \\ \downarrow (\times) \times \text{id} & & \downarrow \times \\ \text{Rel} \times \text{Rel} & \xrightarrow{\quad \alpha^{\text{Rel}} \quad} & \text{Rel}, \\ \downarrow \times & & \downarrow \times \end{array}$$

whose component

$$\alpha_{A,B,C}^{\text{Rel}} : (A \times B) \times C \xrightarrow{\sim} A \times (B \times C)$$

at (A, B, C) is defined by declaring

$$((a, b), c) \sim_{\alpha_{A,B,C}^{\text{Rel}}} (a', (b', c'))$$

iff $a = a'$, $b = b'$, and $c = c'$.

5.2.2.4 The Left Unitor

00CZ **Definition 5.2.2.4.1.** The **left unitor of Rel** is the natural isomorphism

$$\begin{array}{ccc} \text{id} \times \text{id} : \text{id} \times \text{id} & \xrightarrow{\cong} & \text{id} \times \text{id} \\ \text{id} \times \text{id} & \swarrow & \downarrow \text{id} \times \text{id} \\ \text{pt} \times \text{Rel} & \xrightarrow{\mathbb{M}^{\text{Rel}} \times \text{id}} & \text{Rel} \times \text{Rel} \\ \downarrow \lambda^{\text{Rel}} & & \downarrow \times \\ \text{pt} \times \text{Rel} & \xrightarrow{\quad \lambda^{\text{Rel}} \quad} & \text{Rel} \times \text{Rel} \\ \downarrow \lambda_{\text{Rel}}^{\text{Cats}_2} & & \downarrow \times \\ \text{pt} \times \text{Rel} & \xrightarrow{\quad \lambda_{\text{Rel}}^{\text{Cats}_2} \quad} & \text{Rel}, \end{array}$$

whose component

$$\lambda_A^{\text{Rel}} : \mathbb{M}_{\text{Rel}} \times A \xrightarrow{\sim} A$$

at A is defined by declaring

$$(\star, a) \sim_{\lambda_A^{\text{Rel}}} b$$

iff $a = b$.

5.2.2.5 The Right Unitor

00D1 **Definition 5.2.2.5.1.** The **right unitor of Rel** is the natural isomorphism

$$\begin{array}{ccc}
 \text{Rel} \times \text{pt} & \xrightarrow{\text{id} \times \mathbb{1}^{\text{Rel}}} & \text{Rel} \times \text{Rel} \\
 \downarrow & \nearrow \rho_{\text{Rel}}^{\text{Cats}_2} & \downarrow \times \\
 \rho_{\text{Rel}}^{\text{Rel}} : \times \circ (\text{id} \times \mathbb{1}^{\text{Rel}}) & \xrightarrow{\cong} & \rho_{\text{Rel}}^{\text{Cats}_2}, \\
 & \searrow \rho_{\text{Rel}}^{\text{Rel}} & \downarrow \\
 & & \text{Rel},
 \end{array}$$

whose component

$$\rho_A^{\text{Rel}} : A \times \mathbb{1}^{\text{Rel}} \rightarrow A$$

at A is defined by declaring

$$(a, \star) \sim_{\rho_A^{\text{Rel}}} b$$

iff $a = b$.

5.2.2.6 The Symmetry

00D3 **Definition 5.2.2.6.1.** The **symmetry of Rel** is the natural isomorphism

$$\begin{array}{ccc}
 \text{Rel} \times \text{Rel} & \xrightarrow{\times} & \text{Rel}, \\
 \sigma_{\text{Rel}, \text{Rel}}^{\text{Cats}_2} & \swarrow \sigma_{\text{Rel}, \text{Rel}}^{\text{Rel}} & \downarrow \parallel \\
 & & \text{Rel} \times \text{Rel}
 \end{array}$$

whose component

$$\sigma_{A, B}^{\text{Rel}} : A \times B \rightarrow B \times A$$

at (A, B) is defined by declaring

$$(a, b) \sim_{\sigma_{A, B}^{\text{Rel}}} (b', a')$$

iff $a = a'$ and $b = b'$.

5.2.2.7 The Internal Hom

00D5 **Definition 5.2.2.7.1.** The **internal Hom of Rel** is the functor

$$\mathbf{Hom}_{\text{Rel}} : \text{Rel}^{\text{op}} \times \text{Rel} \rightarrow \text{Rel}$$

defined by

$$\mathbf{Hom}_{\text{Rel}}(A, B) \stackrel{\text{def}}{=} A \times B$$

for each $A, B \in \text{Obj}(\text{Rel})$.

00D6 Proposition 5.2.2.7.2. Let $A, B, C \in \text{Obj}(\text{Rel})$.

00D7 1. *Via Self-Duality.* The internal Hom $\mathbf{Hom}_{\text{Rel}}$ of Rel is given by the composition

$$\text{Rel}^{\text{op}} \times \text{Rel} \xrightarrow{\cong} \text{Rel} \times \text{Rel} \xrightarrow{\times} \text{Rel},$$

where the self-duality equivalence $\text{Rel}^{\text{op}} \cong \text{Rel}$ comes from Item 1 of Proposition 5.2.5.1.1.

00D8 2. *Adjointness.* We have adjunctions

$$(A \times - \dashv \mathbf{Hom}_{\text{Rel}}(A, -)): \begin{array}{c} \text{Rel} \xrightarrow{\quad \perp \quad} \text{Rel}, \\ \mathbf{Hom}_{\text{Rel}}(A, -) \end{array}$$

$$(- \times B \dashv \mathbf{Hom}_{\text{Rel}}(B, -)): \begin{array}{c} \text{Rel} \xrightarrow{\quad \perp \quad} \text{Rel}, \\ \mathbf{Hom}_{\text{Rel}}(B, -) \end{array}$$

witnessed by bijections

$$\begin{aligned} \text{Rel}(A \times B, C) &\cong \text{Rel}(A, \mathbf{Hom}_{\text{Rel}}(B, C)) \\ &\stackrel{\text{def}}{=} \text{Rel}(A, B \times C), \\ \text{Rel}(A \times B, C) &\cong \text{Rel}(B, \mathbf{Hom}_{\text{Rel}}(A, C)) \\ &\stackrel{\text{def}}{=} \text{Rel}(B, A \times C), \end{aligned}$$

natural in $A, B, C \in \text{Obj}(\text{Rel})$.

Proof. Item 1, Via Self-Duality: Omitted.

Item 2, Adjointness: Indeed, we have

$$\begin{aligned} \text{Rel}(A \times B, C) &\stackrel{\text{def}}{=} \text{Sets}(A \times B \times C, \{\text{true}, \text{false}\}) \\ &\stackrel{\text{def}}{=} \text{Rel}(A, B \times C) \\ &\stackrel{\text{def}}{=} \text{Rel}(A, \mathbf{Hom}_{\text{Rel}}(B, C)), \end{aligned}$$

and similarly for the bijection $\text{Rel}(A \times B, C) \cong \text{Rel}(B, \mathbf{Hom}_{\text{Rel}}(A, C))$. \square

5.2.2.8 The Closed Symmetric Monoidal Category of Relations

00DA Definition 5.2.2.8.1. The **closed symmetric monoidal category of relations** is the closed symmetric monoidal category

$$(\text{Rel}, \times, \mathbb{1}_{\text{Rel}}, \alpha^{\text{Rel}}, \lambda^{\text{Rel}}, \rho^{\text{Rel}}, \sigma^{\text{Rel}}, \mathbf{Hom}_{\text{Rel}})$$

consisting of

- *The Underlying Category.* The category Rel of sets and relations of Definition 5.2.1.1.1;

- *The Monoidal Product.* The functor

$$\times: \text{Rel} \times \text{Rel} \rightarrow \text{Rel}$$

of [Definition 5.2.2.1.1](#);

- *The Monoidal Unit.* The functor $\mathbb{1}^{\text{Rel}}$ of [Definition 5.2.2.1](#);
- *The Associator.* The natural isomorphism α^{Rel} of [Definition 5.2.2.3.1](#);
- *The Left Unitor.* The natural isomorphism λ^{Rel} of [Definition 5.2.2.4.1](#);
- *The Right Unitor.* The natural isomorphism ρ^{Rel} of [Definition 5.2.2.5.1](#);
- *The Symmetry.* The natural isomorphism σ^{Rel} of [Definition 5.2.2.6.1](#);
- *The Internal Hom.* The functor

$$\mathbf{Hom}_{\text{Rel}}: \text{Rel}^{\text{op}} \times \text{Rel} \rightarrow \text{Rel}$$

of [Definition 5.2.2.7.1](#).

5.2.3 The 2-Category of Relations

00DC Definition 5.2.3.1.1. The **2-category of relations** is the locally posetal 2-category **Rel** where

- *Objects.* The objects of **Rel** are sets;
- *Hom-Objects.* For each $A, B \in \text{Obj}(\text{Sets})$, we have

$$\begin{aligned} \mathbf{Hom}_{\text{Rel}}(A, B) &\stackrel{\text{def}}{=} \mathbf{Rel}(A, B) \\ &\stackrel{\text{def}}{=} (\text{Rel}(A, B), \subset); \end{aligned}$$

- *Identities.* For each $A \in \text{Obj}(\mathbf{Rel})$, the unit map

$$\mathbb{1}_A^{\text{Rel}}: \text{pt} \rightarrow \mathbf{Rel}(A, A)$$

of **Rel** at A is defined by

$$\text{id}_A^{\text{Rel}} \stackrel{\text{def}}{=} \chi_A(-_1, -_2),$$

where $\chi_A(-_1, -_2)$ is the characteristic relation of A of ?? of ??;

- *Composition.* For each $A, B, C \in \text{Obj}(\mathbf{Rel})$, the composition map¹¹

$$\circ_{A,B,C}^{\text{Rel}}: \mathbf{Rel}(B, C) \times \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(A, C)$$

¹¹Note that this is indeed a morphism of posets: given relations $R_1, R_2 \in \mathbf{Rel}(A, B)$ and $S_1, S_2 \in$

of \mathbf{Rel} at (A, B, C) is defined by

$$S \circ_{A,B,C}^{\mathbf{Rel}} R \stackrel{\text{def}}{=} S \diamond R$$

for each $(S, R) \in \mathbf{Rel}(B, C) \times \mathbf{Rel}(A, B)$, where $S \diamond R$ is the composition of S and R of [Definition 5.3.12.1.1](#).

5.2.4 The Double Category of Relations

5.2.4.1 The Double Category of Relations

00DF Definition 5.2.4.1.1. The **double category of relations** is the locally posetal double category $\mathbf{Rel}^{\text{dbl}}$ where

- *Objects.* The objects of $\mathbf{Rel}^{\text{dbl}}$ are sets;
- *Vertical Morphisms.* The vertical morphisms of $\mathbf{Rel}^{\text{dbl}}$ are maps of sets $f: A \rightarrow B$;
- *Horizontal Morphisms.* The horizontal morphisms of $\mathbf{Rel}^{\text{dbl}}$ are relations $R: A \rightarrow B$;
- *2-Morphisms.* A 2-cell

$$\begin{array}{ccc} A & \xrightarrow{R} & B \\ f \downarrow & \Downarrow \alpha & \downarrow g \\ X & \xrightarrow{S} & Y \end{array}$$

of $\mathbf{Rel}^{\text{dbl}}$ is either non-existent or an inclusion of relations of the form

$$\begin{array}{ccc} A \times B & \xrightarrow{R} & \{\text{true, false}\} \\ R \subset S \circ (f \times g), \quad f \times g \downarrow & \curvearrowleft & \downarrow \text{id}_{\{\text{true, false}\}} \\ X \times Y & \xrightarrow{S} & \{\text{true, false}\}; \end{array}$$

- *Horizontal Identities.* The horizontal unit functor of $\mathbf{Rel}^{\text{dbl}}$ is the functor of [Definition 5.2.4.2.1](#);
- *Vertical Identities.* For each $A \in \text{Obj}(\mathbf{Rel}^{\text{dbl}})$, we have

$$\text{id}_A^{\mathbf{Rel}^{\text{dbl}}} \stackrel{\text{def}}{=} \text{id}_A;$$

$\mathbf{Rel}(B, C)$ such that

$$\begin{aligned} R_1 &\subset R_2, \\ S_1 &\subset S_2, \end{aligned}$$

- *Identity 2-Morphisms.* For each horizontal morphism $R: A \rightarrow B$ of Rel^{dbl} , the identity 2-morphism

$$\begin{array}{ccc} A & \xrightarrow{R} & B \\ \downarrow \text{id}_A & \parallel & \downarrow \text{id}_B \\ A & \xrightarrow{R} & B \end{array}$$

of R is the identity inclusion

$$\begin{array}{ccc} B \times A & \xrightarrow{R} & \{\text{true, false}\} \\ R \subset R, \quad \text{id}_B \times \text{id}_A \downarrow & \curvearrowleft & \downarrow \text{id}_{\{\text{true, false}\}} \\ B \times A & \xrightarrow{R} & \{\text{true, false}\}; \end{array}$$

- *Horizontal Composition.* The horizontal composition functor of Rel^{dbl} is the functor of [Definition 5.2.4.3.1](#);
- *Vertical Composition of 1-Morphisms.* For each composable pair $A \xrightarrow{F} B \xrightarrow{G} C$ of vertical morphisms of Rel^{dbl} , i.e. maps of sets, we have

$$g \circ^{\text{Rel}^{\text{dbl}}} f \stackrel{\text{def}}{=} g \circ f;$$

- *Vertical Composition of 2-Morphisms.* The vertical composition of 2-morphisms in Rel^{dbl} is defined as in [Definition 5.2.4.4.1](#);
- *Associators.* The associators of Rel^{dbl} is defined as in [Definition 5.2.4.5.1](#);
- *Left Unitors.* The left unitors of Rel^{dbl} is defined as in [Definition 5.2.4.6.1](#);
- *Right Unitors.* The right unitors of Rel^{dbl} is defined as in [Definition 5.2.4.7.1](#).

5.2.4.2 Horizontal Identities

00DH Definition 5.2.4.2.1. The **horizontal unit functor** of Rel^{dbl} is the functor

$$\mu^{\text{Rel}^{\text{dbl}}} : \text{Rel}_0^{\text{dbl}} \rightarrow \text{Rel}_1^{\text{dbl}}$$

of Rel^{dbl} is the functor where

- *Action on Objects.* For each $A \in \text{Obj}(\text{Rel}_0^{\text{dbl}})$, we have

$$\mu_A \stackrel{\text{def}}{=} \chi_A(-_1, -_2);$$

we have also $S_1 \diamond R_1 \subset S_2 \diamond R_2$.

- *Action on Morphisms.* For each vertical morphism $f: A \rightarrow B$ of Rel^{dbl} , i.e. each map of sets f from A to B , the identity 2-morphism

$$\begin{array}{ccc} A & \xrightarrow{\quad \#_A \quad} & A \\ f \downarrow & \parallel & \downarrow f \\ B & \xrightarrow{\quad \#_B \quad} & B \end{array}$$

of f is the inclusion

$$\begin{array}{ccc} A \times A & \xrightarrow{\chi_A(-_1, -_2)} & \{\text{true, false}\} \\ f \times f \downarrow & \subset & \downarrow \text{id}_{\{\text{true, false}\}} \\ B \times B & \xrightarrow{\chi_B(-_1, -_2)} & \{\text{true, false}\} \end{array}$$

of ?? of ??.

5.2.4.3 Horizontal Composition

00DK Definition 5.2.4.3.1. The **horizontal composition functor** of Rel^{dbl} is the functor

$$\odot^{\text{Rel}^{\text{dbl}}} : \text{Rel}_1^{\text{dbl}} \times_{\text{Rel}_0^{\text{dbl}}} \text{Rel}_1^{\text{dbl}} \rightarrow \text{Rel}_1^{\text{dbl}}$$

of Rel^{dbl} is the functor where

- *Action on Objects.* For each composable pair $A \xrightarrow{R} B \xrightarrow{S} C$ of horizontal morphisms of Rel^{dbl} , we have

$$S \odot R \stackrel{\text{def}}{=} S \diamond R,$$

where $S \diamond R$ is the composition of R and S of [Definition 5.3.12.1.1](#);

- *Action on Morphisms.* For each horizontally composable pair

$$\begin{array}{ccc} A & \xrightarrow{\quad R \quad} & B & \quad & B & \xrightarrow{\quad S \quad} & C \\ f \downarrow & \parallel & \downarrow g & \quad & g \downarrow & \parallel & \downarrow h \\ X & \xrightarrow{\quad T \quad} & Y & \quad & Y & \xrightarrow{\quad U \quad} & Z \end{array}$$

of 2-morphisms of Rel^{dbl} , i.e. for each pair

$$\begin{array}{ccc} A \times B & \xrightarrow{\quad R \quad} & \{\text{true, false}\} & \quad & B \times C & \xrightarrow{\quad S \quad} & \{\text{true, false}\} \\ f \times g \downarrow & \subset & \downarrow \text{id}_{\{\text{true, false}\}} & \quad & g \times h \downarrow & \subset & \downarrow \text{id}_{\{\text{true, false}\}} \\ X \times Y & \xrightarrow{\quad T \quad} & \{\text{true, false}\} & \quad & Y \times Z & \xrightarrow{\quad U \quad} & \{\text{true, false}\} \end{array}$$

of inclusions of relations, the horizontal composition

$$\begin{array}{ccc} A & \xrightarrow{S \circ R} & C \\ f \downarrow & \beta \circ \alpha \Downarrow & \downarrow h \\ X & \xrightarrow{U \circ T} & Z \end{array}$$

of α and β is the inclusion of relations¹²

$$\begin{array}{ccc} A \times C & \xrightarrow{S \circ R} & \{\text{true, false}\} \\ (U \diamond T) \circ (f \times h) \subset (S \diamond R) & f \times h \downarrow & \curvearrowleft \qquad \downarrow \text{id}_{\{\text{true, false}\}} \\ X \times Z & \xrightarrow{U \diamond T} & \{\text{true, false}\}. \end{array}$$

5.2.4.4 Vertical Composition of 2-Morphisms

00DM Definition 5.2.4.4.1. The **vertical composition** in Rel^{dbl} is defined as follows: for each vertically composable pair

$$\begin{array}{ccc} A & \xrightarrow{R} & X \\ f \downarrow & \alpha \Downarrow & \downarrow g \\ B & \xrightarrow[S]{\quad} & Y \end{array} \qquad \begin{array}{ccc} B & \xrightarrow{S} & Y \\ h \downarrow & \beta \Downarrow & \downarrow k \\ C & \xrightarrow[T]{\quad} & Z \end{array}$$

of 2-morphisms of Rel^{dbl} , i.e. for each each pair

$$\begin{array}{ccc} A \times X & \xrightarrow{R} & \{\text{true, false}\} \\ f \times g \downarrow & \curvearrowleft & \downarrow \text{id}_{\{\text{true, false}\}} \\ B \times Y & \xrightarrow[S]{\quad} & \{\text{true, false}\} \end{array} \qquad \begin{array}{ccc} B \times Y & \xrightarrow{S} & \{\text{true, false}\} \\ h \times k \downarrow & \curvearrowleft & \downarrow \text{id}_{\{\text{true, false}\}} \\ C \times Z & \xrightarrow[T]{\quad} & \{\text{true, false}\} \end{array}$$

¹²This is justified by noting that, given $(a, c) \in A \times C$, the statement

- We have $a \sim_{(U \diamond T) \circ (f \times h)} c$, i.e. $f(a) \sim_{U \diamond T} h(c)$, i.e. there exists some $y \in Y$ such that:
 1. We have $f(a) \sim_T y$;
 2. We have $y \sim_U h(c)$;

is implied by the statement

- We have $a \sim_{S \circ R} c$, i.e. there exists some $b \in B$ such that:
 1. We have $a \sim_R b$;
 2. We have $b \sim_S c$;

of inclusions of relations, we define the vertical composition

$$\begin{array}{ccc} A & \xrightarrow{R} & X \\ h \circ f \downarrow & \parallel & \downarrow k \circ g \\ C & \xrightarrow{T} & Z \end{array}$$

of α and β as the inclusion of relations

$$\begin{array}{ccc} A \times X & \xrightarrow{R} & \{\text{true, false}\} \\ T \circ [(h \circ f) \times (k \circ g)] \subset R, \quad (h \circ f) \times (k \circ g) \downarrow & \curvearrowleft & \downarrow \text{id}_{\{\text{true, false}\}} \\ C \times Z & \xrightarrow{T} & \{\text{true, false}\} \end{array}$$

given by the pasting of inclusions¹³

$$\begin{array}{ccc} A \times X & \xrightarrow{R} & \{\text{true, false}\} \\ f \times g \downarrow & \curvearrowleft & \downarrow \text{id}_{\{\text{true, false}\}} \\ B \times Y & \xrightarrow{s} & \{\text{true, false}\} \\ h \times k \downarrow & \curvearrowleft & \downarrow \text{id}_{\{\text{true, false}\}} \\ C \times Z & \xrightarrow{T} & \{\text{true, false}\}. \end{array}$$

5.2.4.5 The Associators

00DP Definition 5.2.4.5.1. For each composable triple $A \xrightarrow{R} B \xrightarrow{S} C \xrightarrow{T} D$ of horizontal morphisms of Rel^{dbl} , the component

$$\alpha_{T,S,R}^{\text{Rel}^{\text{dbl}}} : (T \odot S) \odot R \xrightarrow{\cong} T \odot (S \odot R), \quad \begin{array}{c} A \xrightarrow{R} B \xrightarrow{S} C \xrightarrow{T} D \\ \text{id}_A \downarrow \quad \alpha_{T,S,R}^{\text{Rel}^{\text{dbl}}} \Downarrow \quad \downarrow \text{id}_D \\ A \xrightarrow{R} B \xrightarrow{S} C \xrightarrow{T} D \end{array}$$

since:

- If $a \sim_R b$, then $f(a) \sim_T g(b)$, as $T \circ (f \times g) \subset R$;
- If $b \sim_S c$, then $g(b) \sim_U h(c)$, as $U \circ (g \times h) \subset S$;

¹³This is justified by noting that, given $(a, x) \in A \times X$, the statement

- We have $h(f(a)) \sim_T k(g(x))$;

of the associator of Rel^{dbl} at (R, S, T) is the identity inclusion¹⁴

$$\begin{array}{ccc} A \times B & \xrightarrow{(T \diamond S) \circ R} & \{\text{true, false}\} \\ \parallel & \approx & \downarrow \text{id}_{\{\text{true, false}\}} \\ (T \diamond S) \diamond R = T \diamond (S \diamond R) & & \\ A \times B & \xrightarrow{T \diamond (S \diamond R)} & \{\text{true, false}\}. \end{array}$$

5.2.4.6 The Left Unitors

00DR Definition 5.2.4.6.1. For each horizontal morphism $R: A \dashrightarrow B$ of Rel^{dbl} , the component

$$\lambda_R^{\text{Rel}^{\text{dbl}}}: \mathbb{M}_B \odot R \xrightarrow{\cong} R, \quad \begin{array}{ccccc} A & \xrightarrow{R} & B & \xrightarrow{\mathbb{M}_B} & B \\ \text{id}_A \downarrow & & \lambda_R^{\text{Rel}^{\text{dbl}}} \downarrow & & \downarrow \text{id}_B \\ A & \xrightarrow{R} & B & & \end{array}$$

of the left unit of Rel^{dbl} at R is the identity inclusion¹⁵

$$\begin{array}{ccc} A \times B & \xrightarrow{\chi_B \diamond R} & \{\text{true, false}\} \\ \parallel & \approx & \downarrow \text{id}_{\{\text{true, false}\}} \\ R = \chi_B \diamond R, & & \\ A \times B & \xrightarrow{R} & \{\text{true, false}\}. \end{array}$$

5.2.4.7 The Right Unitors

00DT Definition 5.2.4.7.1. For each horizontal morphism $R: A \dashrightarrow B$ of Rel^{dbl} , the component

$$\rho_R^{\text{Rel}^{\text{dbl}}}: R \odot \mathbb{M}_A \xrightarrow{\cong} R, \quad \begin{array}{ccccc} A & \xrightarrow{\mathbb{M}_A} & A & \xrightarrow{R} & B \\ \text{id}_A \downarrow & & \rho_R^{\text{Rel}^{\text{dbl}}} \downarrow & & \downarrow \text{id}_B \\ A & \xrightarrow{R} & B & & \end{array}$$

is implied by the statement

- We have $a \sim_R x$;

since

- If $a \sim_R x$, then $f(a) \sim_S g(x)$, as $S \circ (f \times g) \subset R$;
- If $b \sim_S y$, then $h(b) \sim_T k(y)$, as $T \circ (h \times k) \subset S$, and thus, in particular:
 - If $f(a) \sim_S g(x)$, then $h(f(a)) \sim_T k(g(x))$;

¹⁴This is justified by Item 2 of Proposition 5.3.12.1.3.

¹⁵This is justified by Item 3 of Proposition 5.3.12.1.3.

of the right unit of $\mathbf{Rel}^{\text{dbl}}$ at R is the identity inclusion¹⁶

$$\begin{array}{ccc} A \times B & \xrightarrow{R \diamond \chi_A} & \{\text{true, false}\} \\ R = R \diamond \chi_A, & \parallel & \cong \\ & & \downarrow \text{id}_{\{\text{true, false}\}} \\ A \times B & \xrightarrow{R} & \{\text{true, false}\}. \end{array}$$

5.2.5 Properties of the Category of Relations

00DV **Proposition 5.2.5.1.1.** Let A and B be sets.

00DW 1. *Self-Duality I.* The category \mathbf{Rel} is self-dual, i.e. we have an equivalence

$$\mathbf{Rel}^{\text{op}} \stackrel{\text{eq.}}{\cong} \mathbf{Rel}$$

of categories.

00DX 2. *Self-Duality II.* The bicategory \mathbf{Rel} is self-dual, i.e. we have a biequivalence

$$\mathbf{Rel}^{\text{op}} \stackrel{\text{eq.}}{\cong} \mathbf{Rel}$$

of bicategories.

00DY 3. *Equivalences and Isomorphisms in Rel.* Let $R: A \nrightarrow B$ be a relation from A to B . The following conditions are equivalent:

00DZ (a) The relation $R: A \nrightarrow B$ is an equivalence in \mathbf{Rel} , i.e. there exists a relation $R^{-1}: B \nrightarrow A$ from B to A together with isomorphisms

$$\begin{aligned} R^{-1} \diamond R &\cong \chi_A, \\ R \diamond R^{-1} &\cong \chi_B. \end{aligned}$$

00E0 (b) The relation $R: A \nrightarrow B$ is an isomorphism in \mathbf{Rel} , i.e. there exists a relation $R^{-1}: B \nrightarrow A$ from B to A such that we have

$$\begin{aligned} R^{-1} \diamond R &= \chi_A, \\ R \diamond R^{-1} &= \chi_B. \end{aligned}$$

00E1 (c) There exists a bijection $f: A \xrightarrow{\cong} B$ with $R = \text{Gr}(f)$.

00E2 4. *Adjunctions in Rel.* We have a natural bijection

$$\left\{ \begin{array}{l} \text{Adjunctions in } \mathbf{Rel} \\ \text{from } A \text{ to } B \end{array} \right\} \cong \left\{ \begin{array}{l} \text{Functions} \\ \text{from } A \text{ to } B \end{array} \right\},$$

with every adjunction in \mathbf{Rel} being of the form $\text{Gr}(f) \dashv f^{-1}$ for some function f .

¹⁶This is justified by Item 3 of Proposition 5.3.12.1.3.

00E3 5. *Monads in Rel.* We have a natural bijection¹⁷

$$\left\{ \begin{array}{l} \text{Monads in} \\ \text{Rel on } A \end{array} \right\} \cong \{\text{Preorders on } A\}.$$

00E4 6. *Comonads in Rel.* We have a natural bijection

$$\left\{ \begin{array}{l} \text{Comonads in} \\ \text{Rel on } A \end{array} \right\} \cong \{\text{Subsets of } A\}.$$

00E5 7. *Characterisations of Monomorphisms in Rel.* Let $R: A \rightarrow B$ be a relation. The following conditions are equivalent:

00E6 (a) The relation R is a monomorphism in Rel.

00E7 (b) The direct image function

$$R_*: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

associated to R is injective.

00E8 (c) The direct image with compact support function

$$R_!: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

associated to R is injective.

Moreover, if R is a monomorphism, then it satisfies the following condition, and the converse holds if R is total:

(★) For each $a, a' \in A$, if there exists some $b \in B$ such that $a \sim_R b$ and $a' \sim_R b$, then $a = a'$.

00E9 8. *Epimorphisms in Rel.* Let $R: A \rightarrow B$ be a relation. The following conditions are equivalent:

00EA (a) The relation R is an epimorphism in Rel.

00EB (b) The weak inverse image function

$$R^{-1}: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

associated to R is injective.

00EC (c) The strong inverse image function

$$R_{-1}: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

associated to R is injective.

¹⁷See also [Section 5.6](#) for an extension of this correspondence to “relative monads on Rel”.

- 00ED** (d) The function $R: A \rightarrow \mathcal{P}(B)$ is “surjective on singletons”:
 (★) For each $b \in B$, there exists some $a \in A$ such that $R(a) = \{b\}$.

- 00EE** 9. *As a Kleisli Category.* We have an isomorphism of categories

$$\mathbf{Rel} \cong \mathbf{FreeAlg}_{\mathcal{P}},$$

where \mathcal{P} is the powerset monad of \mathbf{Set} .

- 00EF** 10. *Co/Completeness (Or Lack Thereof).* The category \mathbf{Rel} is not co/complete, but admits some co/limits:

- (a) *Zero Objects.* The category \mathbf{Rel} has a zero object, the empty set \emptyset .
- (b) *Co/Products.* The category \mathbf{Rel} has co/products, both given by disjoint union of sets.
- (c) *Lack of Co/Equalisers.* The category \mathbf{Rel} does not have co/equalisers.
- (d) *Limits of Graphs of Functions.* The category \mathbf{Rel} has limits whose arrows are all graphs of functions.
- (e) *Colimits of Graphs of Functions.* The category \mathbf{Rel} has colimits whose arrows are all graphs of functions, and these agree with the corresponding limits in \mathbf{Sets} .

- 00EG** 11. *Existence of Right Kan Extensions.* The right Kan extension

$$\text{Ran}_R: \mathbf{Rel}(A, X) \rightarrow \mathbf{Rel}(B, X)$$

along a relation $R: A \rightarrow B$ in \mathbf{Rel} exists and is given by

$$\text{Ran}_R(S) \stackrel{\text{def}}{=} \int_{a \in A} \mathbf{Hom}_{\{\text{t}, \text{f}\}}(R_{-1}^a, S_{-2}^a)$$

for each $S \in \mathbf{Rel}(A, X)$, so that the following conditions are equivalent:

- (a) We have $b \sim_{\text{Ran}_R(S)} x$.
- (b) For each $a \in A$, if $a \sim_R b$, then $a \sim_S x$.

- 00EH** 12. *Existence of Right Kan Lifts.* The right Kan lift

$$\text{Rift}_R: \mathbf{Rel}(X, B) \rightarrow \mathbf{Rel}(X, A)$$

along a relation $R: A \rightarrow B$ in \mathbf{Rel} exists and is given by

$$\text{Rift}_R(S) \stackrel{\text{def}}{=} \int_{b \in B} \mathbf{Hom}_{\{\text{t}, \text{f}\}}(R_b^{-2}, S_b^{-1})$$

for each $S \in \mathbf{Rel}(X, B)$, so that the following conditions are equivalent:

- (a) We have $x \sim_{\text{Rift}_R(S)} a$.

(b) For each $b \in B$, if $a \sim_R b$, then $x \sim_S b$.

- 00EJ** 13. *Closedness.* The bicategory **Rel** is a closed bicategory, there being, for each $R: A \dashrightarrow B$ and set X , a pair of adjunctions

$$(R^* \dashv \text{Ran}_R): \quad \text{Rel}(B, X) \begin{array}{c} \xrightarrow{R^*} \\ \perp \\ \xleftarrow{\text{Ran}_R} \end{array} \text{Rel}(A, X),$$

$$(R_* \dashv \text{Rift}_R): \quad \text{Rel}(X, A) \begin{array}{c} \xrightarrow{R_*} \\ \perp \\ \xleftarrow{\text{Rift}_R} \end{array} \text{Rel}(X, B),$$

witnessed by bijections

$$\mathbf{Rel}(S \diamond R, T) \cong \mathbf{Rel}(S, \text{Ran}_R(T)),$$

$$\mathbf{Rel}(R \diamond U, V) \cong \mathbf{Rel}(U, \text{Rift}_R(V)),$$

natural in $S \in \text{Rel}(B, X)$, $T \in \text{Rel}(A, X)$, $U \in \text{Rel}(X, A)$, and $V \in \text{Rel}(X, B)$.

Proof. **Item 1, Self-Duality I:** Omitted.

Item 2, Self-Duality II: Omitted.

Item 3, Equivalences and Isomorphisms in Rel: We claim that **Items 3a** to **3c** are indeed equivalent:

- **Item 3a** \iff **Item 3b**: This follows from the fact that **Rel** is locally posetal, so that natural isomorphisms and equalities of 1-cells in **Rel** coincide.
- **Item 3b** \implies **Item 3c**: The equalities in **Item 3b** imply $R \dashv R^{-1}$, and thus by **Item 4**, there exists a function $f_R: A \rightarrow B$ associated to R , where, for each $a \in A$, the image $f_R(a)$ of a by f_R is the unique element of $R(a)$, which implies $R = \text{Gr}(f_R)$ in particular. Furthermore, we have $R^{-1} = f_R^{-1}$ (as in **Definition 5.3.2.1.1**). The conditions from **Item 3b** then become the following:

$$\begin{aligned} f_R^{-1} \diamond f_R &= \chi_A, \\ f_R \diamond f_R^{-1} &= \chi_B. \end{aligned}$$

All that is left is to show then is that f_R is a bijection:

- *The Function f_R Is Injective.* Let $a, b \in A$ and suppose that $f_R(a) = f_R(b)$. Since $a \sim_R f_R(a)$ and $f_R(a) = f_R(b) \sim_{R^{-1}} b$, the condition $f_R^{-1} \diamond f_R = \chi_A$ implies that $a = b$, showing f_R to be injective.
- *The Function f_R Is Surjective.* Let $b \in B$. Applying the condition $f_R \diamond f_R^{-1} = \chi_B$ to (b, b) , it follows that there exists some $a \in A$ such that $f_R^{-1}(b) = a$ and $f_R(a) = b$. This shows f_R to be surjective.
- **Item 3c** \implies **Item 3b**: By **Item 2**, we have an adjunction $\text{Gr}(f) \dashv f^{-1}$, giving

inclusions

$$\begin{aligned}\chi_A &\subset f^{-1} \diamond \text{Gr}(f), \\ \text{Gr}(f) \diamond f^{-1} &\subset \chi_B.\end{aligned}$$

We claim the reverse inclusions are also true:

- $f^{-1} \diamond \text{Gr}(f) \subset \chi_A$: This is equivalent to the statement that if $f(a) = b$ and $f^{-1}(b) = a'$, then $a = a'$, which follows from the injectivity of f .
- $\chi_B \subset \text{Gr}(f) \diamond f^{-1}$: This is equivalent to the statement that given $b \in B$ there exists some $a \in A$ such that $f^{-1}(b) = a$ and $f(a) = b$, which follows from the surjectivity of f .

*Item 4, Adjunctions in **Rel**:* We proceed step by step:

1. *From Adjunctions in **Rel** to Functions.* An adjunction in **Rel** from A to B consists of a pair of relations

$$\begin{aligned}R: A &\rightarrow B, \\ S: B &\rightarrow A,\end{aligned}$$

together with inclusions

$$\begin{aligned}\chi_A &\subset S \diamond R, \\ R \diamond S &\subset \chi_B.\end{aligned}$$

We claim that these conditions imply that R is total and functional, i.e. that $R(a)$ is a singleton for each $a \in A$:

- (a) *$R(a)$ Has an Element.* Given $a \in A$, since $\chi_A \subset S \diamond R$, we must have $\{a\} \subset S(R(a))$, implying that there exists some $b \in B$ such that $a \sim_R b$ and $b \sim_S a$, and thus $R(a) \neq \emptyset$, as $b \in R(a)$.
- (b) *$R(a)$ Has No More Than One Element.* Suppose that we have $a \sim_R b$ and $a \sim_R b'$ for $b, b' \in B$. We claim that $b = b'$.
 - i. Since $\chi_A \subset S \diamond R$, there exists some $k \in B$ such that $a \sim_R k$ and $k \sim_S a$.
 - ii. Since $R \diamond S \subset \chi_B$, if $b'' \sim_S a'$ and $a' \sim_R b'''$, then $b'' = b'''$.
 - iii. Applying the above to $b'' = k$, $b''' = b$, and $a' = a$, since $k \sim_S a$ and $a \sim_R b'$, we have $k = b$.
 - iv. Similarly $k = b'$.
 - v. Thus $b = b'$.

Together, the above two items show $R(a)$ to be a singleton, being thus given by $\text{Gr}(f)$ for some function $f: A \rightarrow B$, which gives a map

$$\left\{ \begin{array}{c} \text{Adjunctions in } \mathbf{Rel} \\ \text{from } A \text{ to } B \end{array} \right\} \rightarrow \left\{ \begin{array}{c} \text{Functions} \\ \text{from } A \text{ to } B \end{array} \right\}.$$

Moreover, by uniqueness of adjoints (?? of ??), this implies also that $S = f^{-1}$.

2. *From Functions to Adjunctions in Rel.* By Item 2 of Proposition 5.3.1.1.2, every function $f: A \rightarrow B$ gives rise to an adjunction $\text{Gr}(f) \dashv f^{-1}$ in **Rel**, giving a map

$$\left\{ \begin{array}{l} \text{Functions} \\ \text{from } A \text{ to } B \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \text{Adjunctions in } \mathbf{Rel} \\ \text{from } A \text{ to } B \end{array} \right\}.$$

3. *Invertibility: From Functions to Adjunctions Back to Functions.* We need to show that starting with a function $f: A \rightarrow B$, passing to $\text{Gr}(f) \dashv f^{-1}$, and then passing again to a function gives f again. This is clear however, since we have $a \sim_{\text{Gr}(f)} b$ iff $f(a) = b$.
4. *Invertibility: From Adjunctions to Functions Back to Adjunctions.* We need to show that, given an adjunction $R \dashv S$ in **Rel** giving rise to a function $f_{R,S}: A \rightarrow B$, we have

$$\begin{aligned} \text{Gr}(f_{R,S}) &= R, \\ f_{R,S}^{-1} &= S. \end{aligned}$$

We check these explicitly:

- $\text{Gr}(f_{R,S}) = R$. We have

$$\begin{aligned} \text{Gr}(f_{R,S}) &\stackrel{\text{def}}{=} \{(a, f_{R,S}(a)) \in A \times B \mid a \in A\} \\ &\stackrel{\text{def}}{=} \{(a, R(a)) \in A \times B \mid a \in A\} \\ &= R. \end{aligned}$$

- $f_{R,S}^{-1} = S$. We first claim that, given $a \in A$ and $b \in B$, the following conditions are equivalent:

- We have $a \sim_R b$.
- We have $b \sim_S a$.

Indeed:

- If $a \sim_R b$, then $b \sim_S a$: Since $\chi_A \subset S \diamond R$, there exists $k \in B$ such that $a \sim_R k$ and $k \sim_S a$, but since $a \sim_R b$ and R is functional, we have $k = b$ and thus $b \sim_S a$.
- If $b \sim_S a$, then $a \sim_R b$: First note that since R is total we have $a \sim_R b'$ for some $b' \in B$. Now, since $R \diamond S \subset \chi_B$, $b \sim_S a$, and $a \sim_R b'$, we have $b = b'$, and thus $a \sim_R b$.

Having shown this, we now have

$$\begin{aligned} f_{R,S}^{-1}(b) &\stackrel{\text{def}}{=} \{a \in A \mid f_{R,S}(a) = b\} \\ &\stackrel{\text{def}}{=} \{a \in A \mid a \sim_R b\} \\ &= \{a \in A \mid b \sim_S a\} \\ &\stackrel{\text{def}}{=} S(b). \end{aligned}$$

for each $b \in B$, showing $f_{R,S}^{-1} = S$.

This finishes the proof.

Item 5, Monads in \mathbf{Rel} : A monad in \mathbf{Rel} on A consists of a relation $R: A \rightarrowtail A$ together with maps

$$\begin{aligned}\mu_R: R \diamond R &\subset R, \\ \eta_R: \chi_A &\subset R\end{aligned}$$

making the diagrams

$$\begin{array}{ccccc} \chi_A \diamond R & \xrightarrow{\eta_R \diamond \text{id}_R} & R \diamond R & \xrightarrow{\alpha_{R,R,R}^{\mathbf{Rel}(A,B)}} & R \diamond (R \diamond R) \\ \lambda_R^{\mathbf{Rel}(A,B)} \swarrow \searrow & & \downarrow \mu_R & & \downarrow \text{id}_R \diamond \mu_R \\ & & (R \diamond R) \diamond R & & R \diamond R \\ & & \downarrow \mu_R \diamond \text{id}_R & & \downarrow \mu_R \\ & & R \diamond R & \xrightarrow{\mu_R} & R \end{array}$$

commute. However, since all morphisms involved are inclusions, the commutativity of the above diagrams is automatic, and hence all that is left is the data of the two maps μ_R and η_R , which correspond respectively to the following conditions:

1. For each $a, b, c \in A$, if $a \sim_R b$ and $b \sim_R c$, then $a \sim_R c$.
2. For each $a \in A$, we have $a \sim_R a$.

These are exactly the requirements for R to be a preorder (??). Conversely any preorder \leq gives rise to a pair of maps μ_\leq and η_\leq , forming a monad on A .

Item 6, Comonads in \mathbf{Rel} : A comonad in \mathbf{Rel} on A consists of a relation $R: A \rightarrowtail A$ together with maps

$$\begin{aligned}\Delta_R: R &\subset R \diamond R, \\ \epsilon_R: R &\subset \chi_A\end{aligned}$$

making the diagrams

$$\begin{array}{ccccc} R & \xrightarrow{\Delta_R} & R \diamond R & \xrightarrow{\text{id}_R \diamond \Delta_R} & R \diamond R \\ \lambda_R^{\mathbf{Rel}(A,B), -1} \swarrow \searrow & & \downarrow \epsilon_R \diamond \text{id}_R & & \downarrow \text{id}_R \diamond \epsilon_R \\ & & \chi_A \diamond R & & R \diamond \chi_A \\ & & \downarrow \Delta_R & & \downarrow \rho_R^{\mathbf{Rel}(A,B), -1} \\ & & R \diamond R & \xrightarrow{\Delta_R \diamond \text{id}_R} & (R \diamond R) \diamond R \\ & & & & \downarrow \alpha_{R,R,R}^{\mathbf{Rel}(A,B), -1} \end{array}$$

commute. However, since all morphisms involved are inclusions, the commutativity of the above diagrams is automatic, and hence all that is left is the data of the two maps Δ_R and ϵ_R , which correspond respectively to the following conditions:

1. For each $a, b \in A$, if $a \sim_R b$, then there exists some $k \in A$ such that $a \sim_R k$ and $k \sim_R b$.
2. For each $a, b \in A$, if $a \sim_R b$, then $a = b$.

Taking $k = b$ in the first condition above shows it to be trivially satisfied, while the second condition implies $R \subset \Delta_A$, i.e. R must be a subset of A . Conversely, any subset U of A satisfies $U \subset \Delta_A$, defining a comonad as above.

Item 7, Monomorphisms in Rel: Firstly note that **Items 7b** and **7c** are equivalent by **Item 7 of Proposition 5.5.1.1.3**. We then claim that **Items 7a** and **7b** are also equivalent:

- **Item 7a \implies Item 7b:** Let $U, V \in \mathcal{P}(A)$ and consider the diagram

$$\text{pt} \xrightarrow[U]{V} A \xrightarrow{R} B.$$

By **Remark 5.5.1.1.2**, we have

$$\begin{aligned} R_*(U) &= R \diamond U, \\ R_*(V) &= R \diamond V. \end{aligned}$$

Now, if $R \diamond U = R \diamond V$, i.e. $R_*(U) = R_*(V)$, then $U = V$ since R is assumed to be a monomorphism, showing R_* to be injective.

- **Item 7b \implies Item 7a:** Conversely, suppose that R_* is injective, consider the diagram

$$K \xrightarrow[S]{T} A \xrightarrow{R} B,$$

and suppose that $R \diamond S = R \diamond T$. Note that, since R_* is injective, given a diagram of the form

$$\text{pt} \xrightarrow[U]{V} A \xrightarrow{R} B,$$

if $R_*(U) = R \diamond U = R \diamond V = R_*(V)$, then $U = V$. In particular, for each $k \in K$, we may consider the diagram

$$\text{pt} \xrightarrow{[k]} K \xrightarrow[S]{T} A \xrightarrow{R} B,$$

for which we have $R \diamond S \diamond [k] = R \diamond T \diamond [k]$, implying that we have

$$S(k) = S \diamond [k] = T \diamond [k] = T(k)$$

for each $k \in K$, implying $S = T$, and thus R is a monomorphism.

We can also prove this in a more abstract way, following [[MSE 350788](#)]:

- *Item 7a* \implies *Item 7b*: Assume that R is a monomorphism.

- We first notice that the functor $\text{Rel}(\text{pt}, -) : \text{Rel} \rightarrow \text{Sets}$ maps R to R_* by [Remark 5.5.1.1.2](#).
- Since $\text{Rel}(\text{pt}, -)$ preserves all limits by ?? of ??, it follows by ?? of ?? that $\text{Rel}(\text{pt}, -)$ also preserves monomorphisms.
- Since R is a monomorphism and $\text{Rel}(\text{pt}, -)$ maps R to R_* , it follows that R_* is also a monomorphism.
- Since the monomorphisms in Sets are precisely the injections (?? of ??), it follows that R_* is injective.

- *Item 7b* \implies *Item 7a*: Assume that R_* is injective.

- We first notice that the functor $\text{Rel}(\text{pt}, -) : \text{Rel} \rightarrow \text{Sets}$ maps R to R_* by [Remark 5.5.1.1.2](#).
- Since the monomorphisms in Sets are precisely the injections (?? of ??), it follows that R_* is a monomorphism.
- Since $\text{Rel}(\text{pt}, -)$ is faithful, it follows by ?? of ?? that $\text{Rel}(\text{pt}, -)$ reflects monomorphisms.
- Since R_* is a monomorphism and $\text{Rel}(\text{pt}, -)$ maps R to R_* , it follows that R is also a monomorphism.

Finally, we prove the second part of the statement. Assume that R is a monomorphism, let $a, a' \in A$ such that $a \sim_R b$ and $a' \sim_R b$ for some $b \in B$, and consider the diagram

$$\begin{array}{ccc} \text{pt} & \xrightarrow{\quad [a] \quad} & A \xrightarrow{\quad R \quad} B \\ & \parallel & \\ & \xrightarrow{\quad [a'] \quad} & \end{array}$$

Since $\star \sim_{[a]} a$ and $a \sim_R b$, we have $\star \sim_{R \diamond [a]} b$. Similarly, $\star \sim_{R \diamond [a']} b$. Thus $R \diamond [a] = R \diamond [a']$, and since R is a monomorphism, we have $[a] = [a']$, i.e. $a = a'$.

Conversely, assume the condition

- (★) For each $a, a' \in A$, if there exists some $b \in B$ such that $a \sim_R b$ and $a' \sim_R b$, then $a = a'$,

consider the diagram

$$\begin{array}{ccc} K & \xrightarrow{\quad S \quad} & A \xrightarrow{\quad R \quad} B \\ & \parallel & \\ & \xrightarrow{\quad T \quad} & \end{array}$$

and let $(k, a) \in S$. Since R is total and $a \in A$, there exists some $b \in B$ such that $a \sim_R b$. In this case, we have $k \sim_{R \diamond S} b$, and since $R \diamond S = R \diamond T$, we have also $k \sim_{R \diamond T} b$. Thus there must exist some $a' \in A$ such that $k \sim_T a'$ and $a' \sim_R b$. However, since $a, a' \sim_R b$, we must have $a = a'$, and thus $(k, a) \in T$ as well.

A similar argument shows that if $(k, a) \in T$, then $(k, a) \in S$, and thus $S = T$ and R is a monomorphism.

Item 8, Epimorphisms in Rel: Firstly note that [Items 8b](#) and [8c](#) are equivalent by [Item 7](#) of [Proposition 5.5.2.1.3](#). We then claim that [Items 8a](#) and [8b](#) are also equivalent:

- *Item 8a* \implies *Item 8b*: Let $U, V \in \mathcal{P}(A)$ and consider the diagram

$$\begin{array}{ccccc} A & \xrightarrow{R} & B & \xrightarrow[U]{V} & \text{pt.} \end{array}$$

By Remark 5.5.1.1.2, we have

$$\begin{aligned} R^{-1}(U) &= U \diamond R, \\ R^{-1}(V) &= V \diamond R. \end{aligned}$$

Now, if $U \diamond R = V \diamond R$, i.e. $R^{-1}(U) = R^{-1}(V)$, then $U = V$ since R is assumed to be an epimorphism, showing R^{-1} to be injective.

- *Item 8b* \implies *Item 8a*: Conversely, suppose that R^{-1} is injective, consider the diagram

$$\begin{array}{ccccc} A & \xrightarrow{R} & B & \xrightarrow[S]{T} & K, \end{array}$$

and suppose that $S \diamond R = T \diamond R$. Note that, since R^{-1} is injective, given a diagram of the form

$$\begin{array}{ccccc} A & \xrightarrow{R} & B & \xrightarrow[U]{V} & \text{pt}, \end{array}$$

if $R^{-1}(U) = U \diamond R = V \diamond R = R^{-1}(V)$, then $U = V$. In particular, for each $k \in K$, we may consider the diagram

$$\begin{array}{ccccc} A & \xrightarrow{R} & B & \xrightarrow[S]{T} & K \xrightarrow{[k]} \text{pt}, \end{array}$$

for which we have $[k] \diamond S \diamond R = [k] \diamond T \diamond R$, implying that we have

$$S^{-1}(k) = [k] \diamond S = [k] \diamond T = T^{-1}(k)$$

for each $k \in K$, implying $S = T$, and thus R is an epimorphism.

We can also prove this in a more abstract way, following [MSE 350788]:

- *Item 8a* \implies *Item 8b*: Assume that R is an epimorphism.

- We first notice that the functor $\text{Rel}(-, \text{pt}) : \text{Rel}^{\text{op}} \rightarrow \text{Sets}$ maps R to R^{-1} by Remark 5.5.3.1.2.
- Since $\text{Rel}(-, \text{pt})$ preserves limits by ?? of ??, it follows by ?? of ?? that $\text{Rel}(-, \text{pt})$ also preserves monomorphisms.
- That is: $\text{Rel}(-, \text{pt})$ sends monomorphisms in Rel^{op} to monomorphisms in Sets .
- The monomorphisms Rel^{op} are precisely the epimorphisms in Rel by ?? of ??.

- Since R is an epimorphism and $\text{Rel}(-, \text{pt})$ maps R to R^{-1} , it follows that R^{-1} is a monomorphism.
- Since the monomorphisms in Sets are precisely the injections (?? of ??), it follows that R^{-1} is injective.
- **Item 8b \implies Item 8a:** Assume that R^{-1} is injective.
 - We first notice that the functor $\text{Rel}(-, \text{pt}) : \text{Rel}^{\text{op}} \rightarrow \text{Sets}$ maps R to R^{-1} by Remark 5.5.3.1.2.
 - Since the monomorphisms in Sets are precisely the injections (?? of ??), it follows that R^{-1} is a monomorphism.
 - Since $\text{Rel}(-, \text{pt})$ is faithful, it follows by ?? of ?? that $\text{Rel}(-, \text{pt})$ reflects monomorphisms.
 - That is: $\text{Rel}(-, \text{pt})$ reflects monomorphisms in Sets to monomorphisms in Rel^{op} .
 - The monomorphisms Rel^{op} are precisely the epimorphisms in Rel by ?? of ??.
 - Since R^{-1} is a monomorphism and $\text{Rel}(-, \text{pt})$ maps R to R^{-1} , it follows that R is an epimorphism.

Finally, we claim that Items 8b and 8d are also equivalent, following [MO 350788]:

- **Item 8b \implies Item 8d:** Since $B \setminus \{b\} \subset B$ and R^{-1} is injective, we have $R^{-1}(B \setminus \{b\}) \subseteq R^{-1}(B)$. So taking some $a \in R^{-1}(B) \setminus R^{-1}(B \setminus \{b\})$ we get an element of A such that $R(a) = \{b\}$.
- **Item 8d \implies Item 8b:** Let $U, V \subset B$ with $U \neq V$. Without loss of generality, we can assume $U \setminus V \neq \emptyset$; otherwise just swap U and V . Let then $b \in U \setminus V$. By assumption, there exists an $a \in A$ with $R(a) = \{b\}$. Then $a \in R^{-1}(U)$ but $a \notin R^{-1}(V)$, and thus $R^{-1}(U) \neq R^{-1}(V)$, showing R^{-1} to be injective.

Item 9, As a Kleisli Category: Omitted.

Item 10, Co/Completeness (Or Lack Thereof): Omitted.

Item 11, Existence of Right Kan Extensions: We have

$$\begin{aligned}
 \text{Hom}_{\mathbf{Rel}(A,X)}(S \diamond R, T) &\cong \int_{a \in A} \int_{x \in X} \mathbf{Hom}_{\{\text{t,f}\}}((S \diamond R)_x^a, T_x^a) \\
 &\cong \int_{a \in A} \int_{x \in X} \mathbf{Hom}_{\{\text{t,f}\}}\left(\left(\int^{b \in B} S_x^b \times R_b^a\right), T_x^a\right) \\
 &\cong \int_{a \in A} \int_{x \in X} \int_{b \in B} \mathbf{Hom}_{\{\text{t,f}\}}(S_x^b \times R_b^a, T_x^a) \\
 &\cong \int_{a \in A} \int_{x \in X} \int_{b \in B} \mathbf{Hom}_{\{\text{t,f}\}}(S_x^b, \mathbf{Hom}_{\{\text{t,f}\}}(R_b^a, T_x^a)) \\
 &\cong \int_{b \in B} \int_{x \in X} \int_{a \in A} \mathbf{Hom}_{\{\text{t,f}\}}(S_x^b, \mathbf{Hom}_{\{\text{t,f}\}}(R_b^a, T_x^a)) \\
 &\cong \int_{b \in B} \int_{x \in X} \mathbf{Hom}_{\{\text{t,f}\}}\left(S_x^b, \int_{a \in A} \mathbf{Hom}_{\{\text{t,f}\}}(R_b^a, T_x^a)\right) \\
 &\cong \text{Hom}_{\mathbf{Rel}(B,X)}\left(S, \int_{a \in A} \mathbf{Hom}_{\{\text{t,f}\}}(R_{-1}^a, T_{-2}^a)\right)
 \end{aligned}$$

naturally in each $S \in \mathbf{Rel}(B, X)$ and each $T \in \mathbf{Rel}(A, X)$, showing that

$$\int_{a \in A} \mathbf{Hom}_{\{\text{t,f}\}}(R_{-1}^a, T_{-2}^a)$$

is right adjoint to the precomposition functor $- \diamond R$, being thus the right Kan extension along R . Here we have used the following results, respectively (i.e. for each \cong sign):

1. Item 1 of Proposition 5.1.1.4;
2. Definition 5.3.12.1.1;
3. ?? of ??;
4. ??;
5. ?? of ??;
6. ?? of ??;
7. Item 1 of Proposition 5.1.1.4.

Item 12, Existence of Right Kan Lifts: We have

$$\begin{aligned}
 \text{Hom}_{\mathbf{Rel}(X,B)}(R \diamond S, T) &\cong \int_{x \in X} \int_{b \in B} \mathbf{Hom}_{\{\text{t,f}\}}((R \diamond S)_b^x, T_b^x) \\
 &\cong \int_{x \in X} \int_{b \in B} \mathbf{Hom}_{\{\text{t,f}\}}\left(\left(\int^{a \in A} R_b^a \times S_a^x\right), T_b^x\right) \\
 &\cong \int_{x \in X} \int_{b \in B} \int_{a \in A} \mathbf{Hom}_{\{\text{t,f}\}}(R_b^a \times S_a^x, T_b^x) \\
 &\cong \int_{x \in X} \int_{b \in B} \int_{a \in A} \mathbf{Hom}_{\{\text{t,f}\}}(S_a^x, \mathbf{Hom}_{\{\text{t,f}\}}(R_b^a, T_b^x)) \\
 &\cong \int_{x \in X} \int_{a \in A} \int_{b \in B} \mathbf{Hom}_{\{\text{t,f}\}}(S_a^x, \mathbf{Hom}_{\{\text{t,f}\}}(R_b^a, T_b^x)) \\
 &\cong \int_{x \in X} \int_{a \in A} \mathbf{Hom}_{\{\text{t,f}\}}\left(S_a^x, \int_{b \in B} \mathbf{Hom}_{\{\text{t,f}\}}(R_b^a, T_b^x)\right) \\
 &\cong \text{Hom}_{\mathbf{Rel}(X,A)}\left(S, \int_{b \in B} \mathbf{Hom}_{\{\text{t,f}\}}(R_b^{-2}, T_b^{-1})\right)
 \end{aligned}$$

naturally in each $S \in \mathbf{Rel}(X, A)$ and each $T \in \mathbf{Rel}(X, B)$, showing that

$$\int_{b \in B} \mathbf{Hom}_{\{\text{t,f}\}}(R_b^{-2}, T_b^{-1})$$

is right adjoint to the postcomposition functor $R \diamond -$, being thus the right Kan lift along R . Here we have used the following results, respectively (i.e. for each \cong sign):

1. Item 1 of Proposition 5.1.1.4;
2. Definition 5.3.12.1.1;
3. ?? of ??;
4. ??;
5. ?? of ??;
6. ?? of ??;
7. Item 1 of Proposition 5.1.1.4.

Item 13, Closedness: This has been proved as part of the proof of Items 11 and 12. \square

5.3 Constructions With Relations

5.3.1 The Graph of a Function

Let $f: A \rightarrow B$ be a function.

00EM Definition 5.3.1.1.1. The **graph** of f is the relation $\text{Gr}(f) : A \rightarrow B$ defined as follows:¹⁸

- Viewing relations from A to B as subsets of $A \times B$, we define

$$\text{Gr}(f) \stackrel{\text{def}}{=} \{(a, f(a)) \in A \times B \mid a \in A\};$$

- Viewing relations from A to B as functions $A \times B \rightarrow \{\text{true}, \text{false}\}$, we define

$$[\text{Gr}(f)](a, b) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } b = f(a), \\ \text{false} & \text{otherwise} \end{cases}$$

for each $(a, b) \in A \times B$;

- Viewing relations from A to B as functions $A \rightarrow \mathcal{P}(B)$, we define

$$[\text{Gr}(f)](a) \stackrel{\text{def}}{=} \{f(a)\}$$

for each $a \in A$, i.e. we define $\text{Gr}(f)$ as the composition

$$A \xrightarrow{f} B \xrightarrow{\chi_B} \mathcal{P}(B).$$

00EN Proposition 5.3.1.1.2. Let $f : A \rightarrow B$ be a function.

00EP 1. *Functoriality.* The assignment $A \mapsto \text{Gr}(A)$ defines a functor

$$\text{Gr} : \text{Sets} \rightarrow \text{Rel}$$

where

- *Action on Objects.* For each $A \in \text{Obj}(\text{Sets})$, we have

$$\text{Gr}(A) \stackrel{\text{def}}{=} A;$$

- *Action on Morphisms.* For each $A, B \in \text{Obj}(\text{Sets})$, the action on Hom-sets

$$\text{Gr}_{A,B} : \text{Sets}(A, B) \rightarrow \underbrace{\text{Rel}(\text{Gr}(A), \text{Gr}(B))}_{\stackrel{\text{def}}{=} \text{Rel}(A, B)}$$

of Gr at (A, B) is defined by

$$\text{Gr}_{A,B}(f) \stackrel{\text{def}}{=} \text{Gr}(f),$$

where $\text{Gr}(f)$ is the graph of f as in **Definition 5.3.1.1.1.**

In particular:

¹⁸Further Notation: We write $\text{Gr}(A)$ for $\text{Gr}(\text{id}_A)$, and call it the **graph** of A .

- *Preservation of Identities.* We have

$$\text{Gr}(\text{id}_A) = \chi_A$$

for each $A \in \text{Obj}(\text{Sets})$.

- *Preservation of Composition.* We have

$$\text{Gr}(g \circ f) = \text{Gr}(g) \diamond \text{Gr}(f)$$

for each pair of functions $f: A \rightarrow B$ and $g: B \rightarrow C$.

00EQ 2. *Adjointness Inside Rel.* We have an adjunction

$$(\text{Gr}(f) \dashv f^{-1}): \quad \begin{array}{c} \text{Gr}(f) \\ A \rightleftarrows B \\ f^{-1} \end{array}$$

in **Rel**, where f^{-1} is the inverse of f of [Definition 5.3.2.1.1](#).

00ER 3. *Adjointness.* We have an adjunction

$$(\text{Gr} \dashv \mathcal{P}_*): \quad \begin{array}{c} \text{Gr} \\ \text{Sets} \rightleftarrows \text{Rel} \\ \mathcal{P}_* \end{array}$$

witnessed by a bijection of sets

$$\text{Rel}(\text{Gr}(A), B) \cong \text{Sets}(A, \mathcal{P}(B))$$

natural in $A \in \text{Obj}(\text{Sets})$ and $B \in \text{Obj}(\text{Rel})$.

00ES 4. *Interaction With Inverses.* We have

$$\begin{aligned} \text{Gr}(f)^\dagger &= f^{-1}, \\ (f^{-1})^\dagger &= \text{Gr}(f). \end{aligned}$$

00ET 5. *Cocontinuity.* The functor $\text{Gr}: \text{Sets} \rightarrow \text{Rel}$ of [Item 1](#) preserves colimits.

00EU 6. *Characterisations.* Let $R: A \rightarrow B$ be a relation. The following conditions are equivalent:

- 00EV** (a) There exists a function $f: A \rightarrow B$ such that $R = \text{Gr}(f)$.
- 00EW** (b) The relation R is total and functional.
- 00EX** (c) The weak and strong inverse images of R agree, i.e. we have $R^{-1} = R_{-1}$.
- 00EY** (d) The relation R has a right adjoint R^\dagger in **Rel**.

Proof. **Item 1, Functoriality:** Clear.

Item 2, Adjointness Inside Rel: We need to check that there are inclusions

$$\begin{aligned}\chi_A &\subset f^{-1} \diamond \text{Gr}(f), \\ \text{Gr}(f) \diamond f^{-1} &\subset \chi_B.\end{aligned}$$

These correspond respectively to the following conditions:

1. For each $a \in A$, there exists some $b \in B$ such that $a \sim_{\text{Gr}(f)} b$ and $b \sim_{f^{-1}} a$.
2. For each $a, b \in A$, if $a \sim_{\text{Gr}(f)} b$ and $b \sim_{f^{-1}} a$, then $a = b$.

In other words, the first condition states that the image of any $a \in A$ by f is nonempty, whereas the second condition states that f is not multivalued. As f is a function, both of these statements are true, and we are done.

Item 3, Adjointness: The stated bijection follows from Remark 5.1.1.3, with naturality being clear.

Item 4, Interaction With Inverses: Clear.

Item 5, Cocontinuity: Omitted.

Item 6, Characterisations: We claim that Items 6a to 6d are indeed equivalent:

- **Item 6a \iff Item 6b.** This is shown in the proof of Item 4 of Proposition 5.2.5.1.1.
- **Item 6b \implies Item 6c.** If R is total and functional, then, for each $a \in A$, the set $R(a)$ is a singleton, implying that

$$\begin{aligned}R^{-1}(V) &\stackrel{\text{def}}{=} \{a \in A \mid R(a) \cap V \neq \emptyset\}, \\ R_{-1}(V) &\stackrel{\text{def}}{=} \{a \in A \mid R(a) \subset V\}\end{aligned}$$

are equal for all $V \in \mathcal{P}(B)$, as the conditions $R(a) \cap V \neq \emptyset$ and $R(a) \subset V$ are equivalent when $R(a)$ is a singleton.

- **Item 6c \implies Item 6b.** We claim that R is indeed total and functional:

- **Totality.** If we had $R(a) = \emptyset$ for some $a \in A$, then we would have $a \in R_{-1}(\emptyset)$, so that $R_{-1}(\emptyset) \neq \emptyset$. But since $R^{-1}(\emptyset) = \emptyset$, this would imply $R_{-1}(\emptyset) \neq R^{-1}(\emptyset)$, a contradiction. Thus $R(a) \neq \emptyset$ for all $a \in A$ and R is total.
- **Functionality.** If $R^{-1} = R_{-1}$, then we have

$$\begin{aligned}\{a\} &= R^{-1}(\{b\}) \\ &= R_{-1}(\{b\})\end{aligned}$$

for each $b \in R(a)$ and each $a \in A$, and thus $R(a) \subset \{b\}$. But since R is total, we must have $R(a) = \{b\}$, and thus we see that R is functional.

- **Item 6a \iff Item 6d.** This follows from Item 4 of Proposition 5.2.5.1.1.

This finishes the proof. □

5.3.2 The Inverse of a Function

Let $f: A \rightarrow B$ be a function.

00F0 Definition 5.3.2.1.1. The **inverse of f** is the relation $f^{-1}: B \rightarrow A$ defined as follows:

- Viewing relations from B to A as subsets of $B \times A$, we define

$$f^{-1} \stackrel{\text{def}}{=} \{(b, f^{-1}(b)) \in B \times A \mid a \in A\},$$

where

$$f^{-1}(b) \stackrel{\text{def}}{=} \{a \in A \mid f(a) = b\}$$

for each $b \in B$.

- Viewing relations from B to A as functions $B \times A \rightarrow \{\text{true, false}\}$, we define

$$f^{-1}(b, a) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if there exists } a \in A \text{ with } f(a) = b, \\ \text{false} & \text{otherwise} \end{cases}$$

for each $(b, a) \in B \times A$;

- Viewing relations from B to A as functions $B \rightarrow \mathcal{P}(A)$, we define

$$f^{-1}(b) \stackrel{\text{def}}{=} \{a \in A \mid f(a) = b\}$$

for each $b \in B$.

00F1 Proposition 5.3.2.1.2. Let $f: A \rightarrow B$ be a function.

00F2 1. Functoriality. The assignment $A \mapsto A$, $f \mapsto f^{-1}$ defines a functor

$$(-)^{-1}: \mathbf{Sets} \rightarrow \mathbf{Rel}$$

where

- *Action on Objects.* For each $A \in \mathbf{Obj}(\mathbf{Sets})$, we have

$$[(-)^{-1}](A) \stackrel{\text{def}}{=} A;$$

- *Action on Morphisms.* For each $A, B \in \mathbf{Obj}(\mathbf{Sets})$, the action on Hom-sets

$$(-)^{-1}_{A,B}: \mathbf{Sets}(A, B) \rightarrow \mathbf{Rel}(A, B)$$

of $(-)^{-1}$ at (A, B) is defined by

$$(-)^{-1}_{A,B}(f) \stackrel{\text{def}}{=} [(-)^{-1}](f),$$

where f^{-1} is the inverse of f as in [Definition 5.3.2.1.1](#).

In particular:

- *Preservation of Identities.* We have

$$\text{id}_A^{-1} = \chi_A$$

for each $A \in \text{Obj}(\text{Sets})$.

- *Preservation of Composition.* We have

$$(g \circ f)^{-1} = g^{-1} \diamond f^{-1}$$

for pair of functions $f: A \rightarrow B$ and $g: B \rightarrow C$.

00F3 2. *Adjointness Inside Rel.* We have an adjunction

$$(\text{Gr}(f) \dashv f^{-1}): \quad \begin{array}{ccc} & \text{Gr}(f) & \\ A & \begin{array}{c} \swarrow \downarrow \searrow \\ \perp \end{array} & B \\ & f^{-1} & \end{array}$$

in **Rel**.

00F4 3. *Interaction With Inverses of Relations.* We have

$$\begin{aligned} (f^{-1})^\dagger &= \text{Gr}(f), \\ \text{Gr}(f)^\dagger &= f^{-1}. \end{aligned}$$

Proof. **Item 1, Functoriality:** Clear.

Item 2, Adjointness Inside Rel: This is proved in **Item 2 of Proposition 5.3.1.1.2.**

Item 3, Interaction With Inverses of Relations: Clear. \square

5.3.3 Representable Relations

Let A and B be sets.

00F6 Definition 5.3.3.1.1. Let $f: A \rightarrow B$ and $g: B \rightarrow A$ be functions.¹⁹

1. The **representable relation associated to f** is the relation $\chi_f: A \rightarrow B$ defined as the composition

$$A \times B \xrightarrow{f \times \text{id}_B} B \times B \xrightarrow{\chi_B} \{\text{true, false}\},$$

i.e. given by declaring $a \sim_{\chi_f} b$ iff $f(a) = b$.

¹⁹More generally, given functions

$$\begin{aligned} f: A &\rightarrow C, \\ g: B &\rightarrow D \end{aligned}$$

2. The **corepresentable relation associated to g** is the relation $\chi^g: B \rightarrow A$ defined as the composition

$$B \times A \xrightarrow{g \times \text{id}_A} A \times A \xrightarrow{\chi_A} \{\text{true}, \text{false}\},$$

i.e. given by declaring $b \sim_{\chi^g} a$ iff $g(b) = a$.

5.3.4 The Domain and Range of a Relation

Let A and B be sets.

00F8 Definition 5.3.4.1.1. Let $R \subset A \times B$ be a relation.^{20,21}

1. The **domain of R** is the subset $\text{dom}(R)$ of A defined by

$$\text{dom}(R) \stackrel{\text{def}}{=} \left\{ a \in A \middle| \begin{array}{l} \text{there exists some } b \in B \\ \text{such that } a \sim_R b \end{array} \right\}.$$

2. The **range of R** is the subset $\text{range}(R)$ of B defined by

$$\text{range}(R) \stackrel{\text{def}}{=} \left\{ b \in B \middle| \begin{array}{l} \text{there exists some } a \in A \\ \text{such that } a \sim_R b \end{array} \right\}.$$

and a relation $B \rightarrow D$, we may consider the composite relation

$$A \times B \xrightarrow{f \times g} C \times D \xrightarrow{R} \{\text{true}, \text{false}\},$$

for which we have $a \sim_{R \circ (f \times g)} b$ iff $f(a) \sim_R g(b)$.

²⁰Following ??, we may compute the (characteristic functions associated to the) domain and range of a relation using the following colimit formulas:

$$\begin{aligned} \chi_{\text{dom}(R)}(a) &\cong \underset{b \in B}{\text{colim}}(R_b^a) & (a \in A) \\ &\cong \bigvee_{b \in B} R_b^a, \\ \chi_{\text{range}(R)}(b) &\cong \underset{a \in A}{\text{colim}}(R_b^a) & (b \in B) \\ &\cong \bigvee_{a \in A} R_b^a, \end{aligned}$$

where the join \vee is taken in the poset $(\{\text{true}, \text{false}\}, \leq)$ of ??.

²¹Viewing R as a function $R: A \rightarrow \mathcal{P}(B)$, we have

$$\begin{aligned} \text{dom}(R) &\cong \underset{y \in Y}{\text{colim}}(R(y)) \\ &\cong \bigcup_{y \in Y} R(y), \\ \text{range}(R) &\cong \underset{x \in X}{\text{colim}}(R(x)) \\ &\cong \bigcup_{x \in X} R(x), \end{aligned}$$

5.3.5 Binary Unions of Relations

Let A and B be sets and let R and S be relations from A to B .

00FA Definition 5.3.5.1.1. The **union of R and S** ²² is the relation $R \cup S$ from A to B defined as follows:

- Viewing relations from A to B as subsets of $A \times B$, we define²³

$$R \cup S \stackrel{\text{def}}{=} \{(a, b) \in B \times A \mid \text{we have } a \sim_R b \text{ or } a \sim_S b\}.$$

- Viewing relations from A to B as functions $A \rightarrow \mathcal{P}(B)$, we define

$$[R \cup S](a) \stackrel{\text{def}}{=} R(a) \cup S(a)$$

for each $a \in A$.

00FB Proposition 5.3.5.1.2. Let R, S, R_1 , and R_2 be relations from A to B , and let S_1 and S_2 be relations from B to C .

00FW 1. *Interaction With Inverses.* We have

$$(R \cup S)^\dagger = R^\dagger \cup S^\dagger.$$

00FX 2. *Interaction With Composition.* We have

$$(S_1 \diamond R_1) \cup (S_2 \diamond R_2) \stackrel{\text{poss.}}{\neq} (S_1 \cup S_2) \diamond (R_1 \cup R_2).$$

Proof. **Item 1, Interaction With Inverses:** Clear.

Item 2, Interaction With Composition: Unwinding the definitions, we see that:

1. The condition for $(S_1 \diamond R_1) \cup (S_2 \diamond R_2)$ is:

- (a) There exists some $b \in B$ such that:

i. $a \sim_{R_1} b$ and $b \sim_{S_1} c$;

or

i. $a \sim_{R_2} b$ and $b \sim_{S_2} c$;

3. The condition for $(S_1 \cup S_2) \diamond (R_1 \cup R_2)$ is:

- (a) There exists some $b \in B$ such that:

i. $a \sim_{R_1} b$ or $a \sim_{R_2} b$;

and

i. $b \sim_{S_1} c$ or $b \sim_{S_2} c$.

These two conditions may fail to agree (counterexample omitted), and thus the two resulting relations on $A \times C$ may differ. \square

²²Further Terminology: Also called the **binary union of R and S** , for emphasis.

²³This is the same as the union of R and S as subsets of $A \times B$.

5.3.6 Unions of Families of Relations

Let A and B be sets and let $\{R_i\}_{i \in I}$ be a family of relations from A to B .

00FF Definition 5.3.6.1.1. The **union of the family** $\{R_i\}_{i \in I}$ is the relation $\bigcup_{i \in I} R_i$ from A to B defined as follows:

- Viewing relations from A to B as subsets of $A \times B$, we define²⁴

$$\bigcup_{i \in I} R_i \stackrel{\text{def}}{=} \left\{ (a, b) \in (A \times B)^{\times I} \mid \begin{array}{l} \text{there exists some } i \in I \\ \text{such that } a \sim_{R_i} b \end{array} \right\}.$$

- Viewing relations from A to B as functions $A \rightarrow \mathcal{P}(B)$, we define

$$\left[\bigcup_{i \in I} R_i \right](a) \stackrel{\text{def}}{=} \bigcup_{i \in I} R_i(a)$$

for each $a \in A$.

00FG Proposition 5.3.6.1.2. Let A and B be sets and let $\{R_i\}_{i \in I}$ be a family of relations from A to B .

00FH 1. *Interaction With Inverses.* We have

$$\left(\bigcup_{i \in I} R_i \right)^{\dagger} = \bigcup_{i \in I} R_i^{\dagger}.$$

Proof. Item 1, Interaction With Inverses: Clear. □

5.3.7 Binary Intersections of Relations

Let A and B be sets and let R and S be relations from A to B .

00FK Definition 5.3.7.1.1. The **intersection of R and S** ²⁵ is the relation $R \cap S$ from A to B defined as follows:

- Viewing relations from A to B as subsets of $A \times B$, we define²⁶

$$R \cap S \stackrel{\text{def}}{=} \{(a, b) \in B \times A \mid \text{we have } a \sim_R b \text{ and } a \sim_S b\}.$$

- Viewing relations from A to B as functions $A \rightarrow \mathcal{P}(B)$, we define

$$[R \cap S](a) \stackrel{\text{def}}{=} R(a) \cap S(a)$$

for each $a \in A$.

²⁴This is the same as the union of $\{R_i\}_{i \in I}$ as a collection of subsets of $A \times B$.

²⁵Further Terminology: Also called the **binary intersection of R and S** , for emphasis.

²⁶This is the same as the intersection of R and S as subsets of $A \times B$.

00FL Proposition 5.3.7.1.2. Let R, S, R_1 , and R_2 be relations from A to B , and let S_1 and S_2 be relations from B to C .

00FM 1. *Interaction With Inverses.* We have

$$(R \cap S)^\dagger = R^\dagger \cap S^\dagger.$$

00FN 2. *Interaction With Composition.* We have

$$(S_1 \diamond R_1) \cap (S_2 \diamond R_2) = (S_1 \cap S_2) \diamond (R_1 \cap R_2).$$

Proof. **Item 1, Interaction With Inverses:** Clear.

Item 2, Interaction With Composition: Unwinding the definitions, we see that:

1. The condition for $(S_1 \diamond R_1) \cap (S_2 \diamond R_2)$ is:

(a) There exists some $b \in B$ such that:

i. $a \sim_{R_1} b$ and $b \sim_{S_1} c$;

and

i. $a \sim_{R_2} b$ and $b \sim_{S_2} c$;

3. The condition for $(S_1 \cap S_2) \diamond (R_1 \cap R_2)$ is:

(a) There exists some $b \in B$ such that:

i. $a \sim_{R_1} b$ and $a \sim_{R_2} b$;

and

i. $b \sim_{S_1} c$ and $b \sim_{S_2} c$.

These two conditions agree, and thus so do the two resulting relations on $A \times C$. \square

5.3.8 Intersections of Families of Relations

Let A and B be sets and let $\{R_i\}_{i \in I}$ be a family of relations from A to B .

Definition 5.3.8.1.1. The **intersection of the family** $\{R_i\}_{i \in I}$ is the relation $\bigcup_{i \in I} R_i$ defined as follows:

- Viewing relations from A to B as subsets of $A \times B$, we define²⁷

$$\bigcup_{i \in I} R_i \stackrel{\text{def}}{=} \left\{ (a, b) \in (A \times B)^{\times I} \mid \begin{array}{l} \text{for each } i \in I, \\ \text{we have } a \sim_{R_i} b \end{array} \right\}.$$

²⁷This is the same as the intersection of $\{R_i\}_{i \in I}$ as a collection of subsets of $A \times B$.

- Viewing relations from A to B as functions $A \rightarrow \mathcal{P}(B)$, we define

$$\left[\bigcap_{i \in I} R_i \right](a) \stackrel{\text{def}}{=} \bigcap_{i \in I} R_i(a)$$

for each $a \in A$.

00FR Proposition 5.3.8.1.2. Let A and B be sets and let $\{R_i\}_{i \in I}$ be a family of relations from A to B .

00FS 1. *Interaction With Inverses.* We have

$$\left(\bigcap_{i \in I} R_i \right)^{\dagger} = \bigcap_{i \in I} R_i^{\dagger}.$$

Proof. **Item 1, Interaction With Inverses:** Clear. \square

5.3.9 Binary Products of Relations

Let A, B, X , and Y be sets, let $R: A \rightarrow B$ be a relation from A to B , and let $S: X \rightarrow Y$ be a relation from X to Y .

00FU Definition 5.3.9.1.1. The **product of R and S** ²⁸ is the relation $R \times S$ from $A \times X$ to $B \times Y$ defined as follows:

- Viewing relations from $A \times X$ to $B \times Y$ as subsets of $(A \times X) \times (B \times Y)$, we define $R \times S$ as the Cartesian product of R and S as subsets of $A \times X$ and $B \times Y$;²⁹
- Viewing relations from $A \times X$ to $B \times Y$ as functions $A \times X \rightarrow \mathcal{P}(B \times Y)$, we define $R \times S$ as the composition

$$A \times X \xrightarrow{R \times S} \mathcal{P}(B) \times \mathcal{P}(Y) \xrightarrow{\mathcal{P}_{B,Y}^{\otimes}} \mathcal{P}(B \times Y)$$

in Sets, i.e. by

$$[R \times S](a, x) \stackrel{\text{def}}{=} R(a) \times S(x)$$

for each $(a, x) \in A \times X$.

00FV Proposition 5.3.9.1.2. Let A, B, X , and Y be sets.

00FW 1. *Interaction With Inverses.* Let

$$\begin{aligned} R: A &\rightarrow A, \\ S: X &\rightarrow X \end{aligned}$$

We have

$$(R \times S)^{\dagger} = R^{\dagger} \times S^{\dagger}.$$

²⁸Further Terminology: Also called the **binary product of R and S** , for emphasis.

²⁹That is, $R \times S$ is the relation given by declaring $(a, x) \sim_{R \times S} (b, y)$ iff $a \sim_R b$ and $x \sim_S y$.

00FX 2. *Interaction With Composition.* Let

$$\begin{aligned} R_1 &: A \rightarrow B, \\ S_1 &: B \rightarrow C, \\ R_2 &: X \rightarrow Y, \\ S_2 &: Y \rightarrow Z \end{aligned}$$

be relations. We have

$$(S_1 \diamond R_1) \times (S_2 \diamond R_2) = (S_1 \times S_2) \diamond (R_1 \times R_2).$$

Proof. **Item 1, Interaction With Inverses:** Unwinding the definitions, we see that:

1. We have $(a, x) \sim_{(R \times S)^{\dagger}} (b, y)$ iff :

- We have $(b, y) \sim_{R \times S} (a, x)$, i.e. iff :
 - We have $b \sim_R a$;
 - We have $y \sim_S x$;

2. We have $(a, x) \sim_{R^{\dagger} \times S^{\dagger}} (b, y)$ iff :

- We have $a \sim_{R^{\dagger}} b$ and $x \sim_{S^{\dagger}} y$, i.e. iff :
 - We have $b \sim_R a$;
 - We have $y \sim_S x$.

These two conditions agree, and thus the two resulting relations on $A \times X$ are equal.

Item 2, Interaction With Composition: Unwinding the definitions, we see that:

1. We have $(a, x) \sim_{(S_1 \diamond R_1) \times (S_2 \diamond R_2)} (c, z)$ iff :

- (a) We have $a \sim_{S_1 \diamond R_1} c$ and $x \sim_{S_2 \diamond R_2} z$, i.e. iff :
 - i. There exists some $b \in B$ such that $a \sim_{R_1} b$ and $b \sim_{S_1} c$;
 - ii. There exists some $y \in Y$ such that $x \sim_{R_2} y$ and $y \sim_{S_2} z$;

2. We have $(a, x) \sim_{(S_1 \times S_2) \diamond (R_1 \times R_2)} (c, z)$ iff :

- (a) There exists some $(b, y) \in B \times Y$ such that $(a, x) \sim_{R_1 \times R_2} (b, y)$ and $(b, y) \sim_{S_1 \times S_2} (c, z)$, i.e. such that:
 - i. We have $a \sim_{R_1} b$ and $x \sim_{R_2} y$;
 - ii. We have $b \sim_{S_1} c$ and $y \sim_{S_2} z$.

These two conditions agree, and thus the two resulting relations from $A \times X$ to $C \times Z$ are equal. \square

5.3.10 Products of Families of Relations

Let $\{A_i\}_{i \in I}$ and $\{B_i\}_{i \in I}$ be families of sets, and let $\{R_i : A_i \rightarrow B_i\}_{i \in I}$ be a family of relations.

00FZ Definition 5.3.10.1.1. The **product of the family** $\{R_i\}_{i \in I}$ is the relation $\prod_{i \in I} R_i$ from $\prod_{i \in I} A_i$ to $\prod_{i \in I} B_i$ defined as follows:

- Viewing relations as subsets, we define $\prod_{i \in I} R_i$ as its product as a family of sets, i.e. we have

$$\prod_{i \in I} R_i \stackrel{\text{def}}{=} \left\{ (a_i, b_i)_{i \in I} \in \prod_{i \in I} (A_i \times B_i) \mid \begin{array}{l} \text{for each } i \in I, \\ \text{we have } a_i \sim_{R_i} b_i \end{array} \right\}.$$

- Viewing relations as functions to powersets, we define

$$\left[\prod_{i \in I} R_i \right] ((a_i)_{i \in I}) \stackrel{\text{def}}{=} \prod_{i \in I} R_i(a_i)$$

for each $(a_i)_{i \in I} \in \prod_{i \in I} R_i$.

5.3.11 The Inverse of a Relation

Let A , B , and C be sets and let $R \subset A \times B$ be a relation.

00G1 Definition 5.3.11.1.1. The **inverse of R** ³⁰ is the relation R^\dagger defined as follows:

- Viewing relations as subsets, we define

$$R^\dagger \stackrel{\text{def}}{=} \{(b, a) \in B \times A \mid \text{we have } b \sim_R a\}.$$

- Viewing relations as functions $A \times B \rightarrow \{\text{true}, \text{false}\}$, we define

$$[R^\dagger]_a^b \stackrel{\text{def}}{=} R_b^a$$

for each $(b, a) \in B \times A$.

- Viewing relations as functions $A \rightarrow \mathcal{P}(B)$, we define

$$\begin{aligned} [R^\dagger](b) &\stackrel{\text{def}}{=} R^\dagger(\{b\}) \\ &\stackrel{\text{def}}{=} \{a \in A \mid b \in R(a)\} \end{aligned}$$

for each $b \in B$, where $R^\dagger(\{b\})$ is the fibre of R over $\{b\}$.

00G2 Example 5.3.11.1.2. Here are some examples of inverses of relations.

00G3 1. *Less Than Equal Signs.* We have $(\leq)^\dagger = \geq$.

³⁰Further Terminology: Also called the **opposite of R** , the **transpose of R** , or the **converse of R** .

00G4 2. *Greater Than Equal Signs.* Dually to ??, we have $(\geq)^\dagger = \leq$.

00G5 3. *Functions.* Let $f: A \rightarrow B$ be a function. We have

$$\begin{aligned}\text{Gr}(f)^\dagger &= f^{-1}, \\ (f^{-1})^\dagger &= \text{Gr}(f).\end{aligned}$$

00G6 **Proposition 5.3.11.1.3.** Let $R: A \rightarrow B$ and $S: B \rightarrow C$ be relations.

00G7 1. *Interaction With Ranges and Domains.* We have

$$\begin{aligned}\text{dom}(R^\dagger) &= \text{range}(R), \\ \text{range}(R^\dagger) &= \text{dom}(R).\end{aligned}$$

00G8 2. *Interaction With Composition I.* We have

$$(S \diamond R)^\dagger = R^\dagger \diamond S^\dagger.$$

00G9 3. *Interaction With Composition II.* We have

$$\begin{aligned}\chi_B(-_1, -_2) &\subset R \diamond R^\dagger, \\ \chi_A(-_1, -_2) &\subset R^\dagger \diamond R.\end{aligned}$$

00GA 4. *Invertibility.* We have

$$(R^\dagger)^\dagger = R.$$

00GB 5. *Identity.* We have

$$\chi_A^\dagger(-_1, -_2) = \chi_A(-_1, -_2).$$

Proof. **Item 1, Interaction With Ranges and Domains:** Clear.

Item 2, Interaction With Composition I: Clear.

Item 3, Interaction With Composition II: Clear.

Item 4, Invertibility: Clear.

Item 5, Identity: Clear. □

5.3.12 Composition of Relations

Let A , B , and C be sets and let $R \subset A \times B$ and $S \subset B \times C$ be relations.

00GD **Definition 5.3.12.1.1.** The **composition of R and S** is the relation $S \diamond R$ defined as follows:

- Viewing relations from A to C as subsets of $A \times C$, we define

$$S \diamond R \stackrel{\text{def}}{=} \left\{ (a, c) \in A \times C \mid \begin{array}{l} \text{there exists some } b \in B \text{ such} \\ \text{that } a \sim_R b \text{ and } b \sim_S c \end{array} \right\}.$$

- Viewing relations as functions $A \times B \rightarrow \{\text{true}, \text{false}\}$, we define

$$\begin{aligned} (S \diamond R)^{-1}_{-2} &\stackrel{\text{def}}{=} \int^{y \in B} S_y^{-1} \times R^y_{-2} \\ &= \bigvee_{y \in B} S_y^{-1} \times R^y_{-2}, \end{aligned}$$

where the join \bigvee is taken in the poset $(\{\text{true}, \text{false}\}, \leq)$ of ??.

- Viewing relations as functions $A \rightarrow \mathcal{P}(B)$, we define

$$\begin{array}{ccc} S \diamond R & \stackrel{\text{def}}{=} & \text{Lan}_{\chi_B}(S) \circ R, \\ & & \begin{array}{c} B \xrightarrow{S} \mathcal{P}(C), \\ \chi_B \downarrow \quad \nearrow \text{Lan}_{\chi_B}(S) \\ A \xrightarrow[R]{} \mathcal{P}(B) \end{array} \end{array}$$

where $\text{Lan}_{\chi_B}(S)$ is computed by the formula

$$\begin{aligned} [\text{Lan}_{\chi_B}(S)](V) &\cong \int^{y \in B} \chi_{\mathcal{P}(B)}(\chi_y, V) \odot S_y \\ &\cong \int^{y \in B} \chi_V(y) \odot S_y \\ &\cong \bigcup_{y \in B} \chi_V(y) \odot S_y \\ &\cong \bigcup_{y \in V} S_y \end{aligned}$$

for each $V \in \mathcal{P}(B)$. In other words, $S \diamond R$ is defined by³¹

$$\begin{aligned} [S \diamond R](a) &\stackrel{\text{def}}{=} S(R(a)) \\ &\stackrel{\text{def}}{=} \bigcup_{x \in R(a)} S(x). \end{aligned}$$

for each $a \in A$.

00GE Example 5.3.12.1.2. Here are some examples of composition of relations.

³¹That is: the relation R may send $a \in A$ to a number of elements $\{b_i\}_{i \in I}$ in B , and then the relation S may send the image of each of the b_i 's to a number of elements $\{S(b_i)\}_{i \in I} = \left\{ \{c_j\}_{j_i \in J_i} \right\}_{i \in I}$ in C .

1. *Composing Less/Greater Than Equal With Greater/Less Than Equal Signs.* We have

$$\begin{aligned}\leq \diamond \geq &= \sim_{\text{triv}}, \\ \geq \diamond \leq &= \sim_{\text{triv}}.\end{aligned}$$

2. *Composing Less/Greater Than Equal Signs With Less/Greater Than Equal Signs.* We have

$$\begin{aligned}\leq \diamond \leq &= \leq, \\ \geq \diamond \geq &= \geq.\end{aligned}$$

00GF **Proposition 5.3.12.1.3.** Let $R: A \rightarrow B$, $S: B \rightarrow C$, and $T: C \rightarrow D$ be relations.

00GG 1. *Interaction With Ranges and Domains.* We have

$$\begin{aligned}\text{dom}(S \diamond R) &\subset \text{dom}(R), \\ \text{range}(S \diamond R) &\subset \text{range}(S).\end{aligned}$$

00GH 2. *Associativity.* We have

$$(T \diamond S) \diamond R = T \diamond (S \diamond R).$$

00GJ 3. *Unitality.* We have

$$\begin{aligned}\chi_B \diamond R &= R, \\ R \diamond \chi_A &= R.\end{aligned}$$

00GK 4. *Interaction With Inverses.* We have

$$(S \diamond R)^\dagger = R^\dagger \diamond S^\dagger.$$

00GL 5. *Interaction With Composition.* We have

$$\begin{aligned}\chi_B(-_1, -_2) &\subset R \diamond R^\dagger, \\ \chi_A(-_1, -_2) &\subset R^\dagger \diamond R.\end{aligned}$$

Proof. **Item 1**, *Interaction With Ranges and Domains:* Clear.

Item 2, Associativity: Indeed, we have

$$\begin{aligned}
 (T \diamond S) \diamond R &\stackrel{\text{def}}{=} \left(\int^{y \in C} T_x^{-1} \times S_{-2}^x \right) \diamond R \\
 &\stackrel{\text{def}}{=} \int^{x \in B} \left(\int^{y \in C} T_x^{-1} \times S_y^x \right) \diamond R_{-2}^y \\
 &= \int^{x \in B} \int^{y \in C} \left(T_x^{-1} \times S_y^x \right) \diamond R_{-2}^y \\
 &= \int^{y \in C} \int^{x \in B} \left(T_x^{-1} \times S_y^x \right) \diamond R_{-2}^y \\
 &= \int^{y \in C} \int^{x \in B} T_x^{-1} \times \left(S_y^x \diamond R_{-2}^y \right) \\
 &= \int^{x \in B} T_x^{-1} \times \left(\int^{y \in C} S_y^x \diamond R_{-2}^y \right) \\
 &\stackrel{\text{def}}{=} \int^{x \in B} T_x^{-1} \times (S \diamond R)_{-2}^x \\
 &\stackrel{\text{def}}{=} T \diamond (S \diamond R).
 \end{aligned}$$

In the language of relations, given $a \in A$ and $d \in D$, the stated equality witnesses the equivalence of the following two statements:

1. We have $a \sim_{(T \diamond S) \diamond R} d$, i.e. there exists some $b \in B$ such that:
 - (a) We have $a \sim_R b$;
 - (b) We have $b \sim_{T \diamond S} d$, i.e. there exists some $c \in C$ such that:
 - i. We have $b \sim_S c$;
 - ii. We have $c \sim_T d$;
2. We have $a \sim_{T \diamond (S \diamond R)} d$, i.e. there exists some $c \in C$ such that:
 - (a) We have $a \sim_{S \diamond R} c$, i.e. there exists some $b \in B$ such that:
 - i. We have $a \sim_R b$;
 - ii. We have $b \sim_S c$;
 - (b) We have $c \sim_T d$;

both of which are equivalent to the statement

- There exist $b \in B$ and $c \in C$ such that $a \sim_R b \sim_S c \sim_T d$.

Item 3, Unitality: Indeed, we have

$$\begin{aligned}\chi_B \diamond R &\stackrel{\text{def}}{=} \int^{x \in B} (\chi_B)_x^{-1} \times R_{-2}^x \\ &= \bigvee_{x \in B} (\chi_B)_x^{-1} \times R_{-2}^x \\ &= \bigvee_{\substack{x \in B \\ x = -1}} R_{-2}^x \\ &= R_{-2}^{-1},\end{aligned}$$

and

$$\begin{aligned}R \diamond \chi_A &\stackrel{\text{def}}{=} \int^{x \in A} R_x^{-1} \times (\chi_A)_{-2}^x \\ &= \bigvee_{x \in B} R_x^{-1} \times (\chi_A)_{-2}^x \\ &= \bigvee_{\substack{x \in B \\ x = -2}} R_x^{-1} \\ &= R_{-2}^{-1}.\end{aligned}$$

In the language of relations, given $a \in A$ and $b \in B$:

- The equality

$$\chi_B \diamond R = R$$

witnesses the equivalence of the following two statements:

1. We have $a \sim_b B$.
2. There exists some $b' \in B$ such that:
 - (a) We have $a \sim_R b'$
 - (b) We have $b' \sim_{\chi_B} b$, i.e. $b' = b$.

- The equality

$$R \diamond \chi_A = R$$

witnesses the equivalence of the following two statements:

1. There exists some $a' \in A$ such that:
 - (a) We have $a \sim_{\chi_B} a'$, i.e. $a = a'$.
 - (b) We have $a' \sim_R b$
2. We have $a \sim_b B$.

Item 4, Interaction With Inverses: Clear.

Item 5, Interaction With Composition: Clear. □

5.3.13 The Collage of a Relation

Let A and B be sets and let $R: A \rightarrow B$ be a relation from A to B .

00GN **Definition 5.3.13.1.1.** The **collage** of R ³² is the poset $\mathbf{Coll}(R) \stackrel{\text{def}}{=} (\mathbf{Coll}(R), \leq_{\mathbf{Coll}(R)})$ consisting of

- *The Underlying Set.* The set $\mathbf{Coll}(R)$ defined by

$$\mathbf{Coll}(R) \stackrel{\text{def}}{=} A \coprod B.$$

- *The Partial Order.* The partial order

$$\leq_{\mathbf{Coll}(R)}: \mathbf{Coll}(R) \times \mathbf{Coll}(R) \rightarrow \{\text{true, false}\}$$

on $\mathbf{Coll}(R)$ defined by

$$\leq(a, b) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } a = b \text{ or } a \sim_R b, \\ \text{false} & \text{otherwise.} \end{cases}$$

00GP **Proposition 5.3.13.1.2.** Let A and B be sets and let $R: A \rightarrow B$ be a relation from A to B .

00GQ 1. *Functoriality I.* The assignment $R \mapsto \mathbf{Coll}(R)$ defines a functor³³

$$\mathbf{Coll}: \mathbf{Rel}(A, B) \rightarrow \mathbf{Pos}_{/\Delta^1}(A, B),$$

where

³²Further Terminology: Also called the **cograph** of R .

³³Here $\mathbf{Pos}_{/\Delta^1}(A, B)$ is the category defined as the pullback

$$\mathbf{Pos}_{/\Delta^1}(A, B) \stackrel{\text{def}}{=} \underset{[A], \mathbf{Pos}, \mathbf{fib}_0}{\mathbf{pt}} \times \underset{\mathbf{fib}_1, \mathbf{Pos}, [B]}{\mathbf{Pos}_{/\Delta^1}} \mathbf{pt},$$

as in the diagram

$$\begin{array}{ccccc} & & \mathbf{Pos}_{/\Delta^1}(A, B) & & \\ & \searrow & & \swarrow & \\ & & \mathbf{Pos}_{/\Delta^1} \times_{\mathbf{Pos}} \mathbf{pt} & & \mathbf{pt} \times_{\mathbf{Pos}} \mathbf{Pos}_{/\Delta^1} \\ & \swarrow & & \searrow & \\ \mathbf{pt} & & \mathbf{Pos}_{/\Delta^1} & & \mathbf{pt} \\ & \searrow & \swarrow & \searrow & \\ & [A] & & \mathbf{fib}_{[0]} & & [B] \\ & \mathbf{Pos} & & & \mathbf{Pos} & \end{array}$$

Explicitly, an object of $\mathbf{Pos}_{/\Delta^1}(A, B)$ is a pair (X, ϕ_X) consisting of

- A poset X ;
- A morphism $\phi_X: X \rightarrow \Delta^1$;

- *Action on Objects.* For each $R \in \text{Obj}(\mathbf{Rel}(A, B))$, we have

$$[\mathbf{Coll}](R) \stackrel{\text{def}}{=} (\mathbf{Coll}(R), \phi_R)$$

for each $R \in \mathbf{Rel}(A, B)$, where

- The poset $\mathbf{Coll}(R)$ is the collage of R of [Definition 5.3.13.1.1](#);
- The morphism $\phi_R: \mathbf{Coll}(R) \rightarrow \Delta^1$ is given by

$$\phi_R(x) \stackrel{\text{def}}{=} \begin{cases} 0 & \text{if } x \in A, \\ 1 & \text{if } x \in B \end{cases}$$

for each $x \in \mathbf{Coll}(R)$;

- *Action on Morphisms.* For each $R, S \in \text{Obj}(\mathbf{Rel}(A, B))$, the action on Hom-sets

$$\mathbf{Coll}_{R,S}: \text{Hom}_{\mathbf{Rel}(A,B)}(R, S) \rightarrow \text{Pos}(\mathbf{Coll}(R), \mathbf{Coll}(S))$$

of \mathbf{Coll} at (R, S) is given by sending an inclusion

$$\iota: R \subset S$$

to the morphism

$$\mathbf{Coll}(\iota): \mathbf{Coll}(R) \rightarrow \mathbf{Coll}(S)$$

of posets over Δ^1 defined by

$$[\mathbf{Coll}(\iota)](x) \stackrel{\text{def}}{=} x$$

for each $x \in \mathbf{Coll}(R)$.³⁴

00GR 2. *Equivalence.* The functor of [Item 1](#) is an equivalence of categories.

Proof. [Item 1](#), *Functionality:* Clear.

[Item 2](#), *Equivalence:* Omitted. □

5.4 Equivalence Relations

5.4.1 Reflexive Relations

5.4.1.1 Foundations

Let A be a set.

such that $\phi_X^{-1}(0) = A$ and $\phi_X^{-1}(1) = B$, with morphisms between such objects being morphisms of posets over Δ^1 .

³⁴Note that this is indeed a morphism of posets: if $x \leq_{\mathbf{Coll}(R)} y$, then $x = y$ or $x \sim_R y$, so we have either

00GV Definition 5.4.1.1.1. A **reflexive relation** is equivalently:³⁵

- An \mathbb{E}_0 -monoid in $(N_\bullet(\mathbf{Rel}(A, A)), \chi_A)$;
- A pointed object in $(\mathbf{Rel}(A, A), \chi_A)$.

00GW Remark 5.4.1.1.2. In detail, a relation R on A is **reflexive** if we have an inclusion

$$\eta_R: \chi_A \subset R$$

of relations in $\mathbf{Rel}(A, A)$, i.e. if, for each $a \in A$, we have $a \sim_R a$.

00GX Definition 5.4.1.1.3. Let A be a set.

1. The **set of reflexive relations on A** is the subset $\mathbf{Rel}^{\text{refl}}(A, A)$ of $\mathbf{Rel}(A, A)$ spanned by the reflexive relations.
00GY
2. The **poset of relations on A** is the subposet $\mathbf{Rel}^{\text{refl}}(A, A)$ of $\mathbf{Rel}(A, A)$ spanned by the reflexive relations.
00GZ

00H0 Proposition 5.4.1.1.4. Let R and S be relations on A .

00H1 1. *Interaction With Inverses.* If R is reflexive, then so is R^\dagger .

00H2 2. *Interaction With Composition.* If R and S are reflexive, then so is $S \diamond R$.

Proof. **Item 1, Interaction With Inverses:** Clear.

Item 2, Interaction With Composition: Clear. □

5.4.1.2 The Reflexive Closure of a Relation

Let R be a relation on A .

00H4 Definition 5.4.1.2.1. The **reflexive closure** of \sim_R is the relation \sim_R^{refl} ³⁶ satisfying the following universal property:³⁷

- (★) Given another reflexive relation \sim_S on A such that $R \subset S$, there exists an inclusion $\sim_R^{\text{refl}} \subset \sim_S$.

00H5 Construction 5.4.1.2.2. Concretely, \sim_R^{refl} is the free pointed object on R in $(\mathbf{Rel}(A, A), \chi_A)$ ³⁸, being given by

$$\begin{aligned} R^{\text{refl}} &\stackrel{\text{def}}{=} R \coprod^{\mathbf{Rel}(A, A)} \Delta_A \\ &= R \cup \Delta_A \\ &= \{(a, b) \in A \times A \mid \text{we have } a \sim_R b \text{ or } a = b\}. \end{aligned}$$

$x = y$ or $x \sim_S y$ (as $R \subset S$), and thus $x \leq_{\mathbf{Coll}(S)} y$.

³⁵Note that since $\mathbf{Rel}(A, A)$ is posetal, reflexivity is a property of a relation, rather than extra structure.

³⁶Further Notation: Also written R^{refl} .

³⁷Slogan: The reflexive closure of R is the smallest reflexive relation containing R .

³⁸Or, equivalently, the free \mathbb{E}_0 -monoid on R in $(N_\bullet(\mathbf{Rel}(A, A)), \chi_A)$.

Proof. Clear. \square

00H6 **Proposition 5.4.1.2.3.** Let R be a relation on A .

00H7 1. *Adjointness.* We have an adjunction

$$\left((-)^{\text{refl}} \dashv \text{忘} \right) : \mathbf{Rel}(A, A) \begin{array}{c} \xrightarrow{(-)^{\text{refl}}} \\[-1ex] \perp \\[-1ex] \text{忘} \end{array} \mathbf{Rel}^{\text{refl}}(A, A),$$

witnessed by a bijection of sets

$$\mathbf{Rel}^{\text{refl}}(R^{\text{refl}}, S) \cong \mathbf{Rel}(R, S),$$

natural in $R \in \text{Obj}(\mathbf{Rel}^{\text{refl}}(A, A))$ and $S \in \text{Obj}(\mathbf{Rel}(A, A))$.

00H8 2. *The Reflexive Closure of a Reflexive Relation.* If R is reflexive, then $R^{\text{refl}} = R$.

00H9 3. *Idempotency.* We have

$$(R^{\text{refl}})^{\text{refl}} = R^{\text{refl}}.$$

00HA 4. *Interaction With Inverses.* We have

$$\begin{array}{ccc} \mathbf{Rel}(A, A) & \xrightarrow{(-)^{\text{refl}}} & \mathbf{Rel}(A, A) \\ \left(R^{\dagger} \right)^{\text{refl}} = \left(R^{\text{refl}} \right)^{\dagger}, & \downarrow (-)^{\dagger} & \downarrow (-)^{\dagger} \\ \mathbf{Rel}(A, A) & \xrightarrow{(-)^{\text{refl}}} & \mathbf{Rel}(A, A). \end{array}$$

00HB 5. *Interaction With Composition.* We have

$$\begin{array}{ccc} \mathbf{Rel}(A, A) \times \mathbf{Rel}(A, A) & \xrightarrow{\diamond} & \mathbf{Rel}(A, A) \\ (S \diamond R)^{\text{refl}} = S^{\text{refl}} \diamond R^{\text{refl}}, & \downarrow (-)^{\text{refl}} \times (-)^{\text{refl}} & \downarrow (-)^{\text{refl}} \\ \mathbf{Rel}(A, A) \times \mathbf{Rel}(A, A) & \xrightarrow{\diamond} & \mathbf{Rel}(A, A). \end{array}$$

Proof. **Item 1, Adjointness:** This is a rephrasing of the universal property of the reflexive closure of a relation, stated in [Definition 5.4.1.2.1](#).

Item 2, The Reflexive Closure of a Reflexive Relation: Clear.

Item 3, Idempotency: This follows from [Item 2](#).

Item 4, Interaction With Inverses: Clear.

Item 5, Interaction With Composition: This follows from [Item 2](#) of [Proposition 5.4.1.1.4](#). \square

5.4.2 Symmetric Relations

5.4.2.1 Foundations

Let A be a set.

00HE Definition 5.4.2.1.1. A relation R on A is **symmetric** if, for each $a, b \in A$, the following conditions are equivalent:³⁹

1. We have $a \sim_R b$.
2. We have $b \sim_R a$.

00HF Definition 5.4.2.1.2. Let A be a set.

1. The **set of symmetric relations on A** is the subset $\text{Rel}^{\text{symm}}(A, A)$ of $\text{Rel}(A, A)$

00HG spanned by the symmetric relations.

2. The **poset of relations on A** is the subposet $\text{Rel}^{\text{symm}}(A, A)$ of $\text{Rel}(A, A)$

00HH spanned by the symmetric relations.

00HJ Proposition 5.4.2.1.3. Let R and S be relations on A .

00HK 1. *Interaction With Inverses.* If R is symmetric, then so is R^\dagger .

00HL 2. *Interaction With Composition.* If R and S are symmetric, then so is $S \diamond R$.

Proof. **Item 1, Interaction With Inverses:** Clear.

Item 2, Interaction With Composition: Clear. □

5.4.2.2 The Symmetric Closure of a Relation

Let R be a relation on A .

00HN Definition 5.4.2.2.1. The **symmetric closure** of \sim_R is the relation \sim_R^{symm} ⁴⁰ satisfying the following universal property:⁴¹

- (★) Given another symmetric relation \sim_S on A such that $R \subset S$, there exists an inclusion $\sim_R^{\text{symm}} \subset \sim_S$.

00HP Construction 5.4.2.2.2. Concretely, \sim_R^{symm} is the symmetric relation on A defined by

$$\begin{aligned} R^{\text{symm}} &\stackrel{\text{def}}{=} R \cup R^\dagger \\ &= \{(a, b) \in A \times A \mid \text{we have } a \sim_R b \text{ or } b \sim_R a\}. \end{aligned}$$

Proof. Clear. □

³⁹That is, R is symmetric if $R^\dagger = R$.

⁴⁰Further Notation: Also written R^{symm} .

⁴¹Slogan: The symmetric closure of R is the smallest symmetric relation containing R .

00HQ Proposition 5.4.2.2.3. Let R be a relation on A .

00HR 1. *Adjointness.* We have an adjunction

$$\left((-)^{\text{symm}} \dashv \overline{\text{Sym}} \right) : \mathbf{Rel}(A, A) \begin{array}{c} \xrightarrow{(-)^{\text{symm}}} \\ \perp \\ \xleftarrow{\overline{\text{Sym}}} \end{array} \mathbf{Rel}^{\text{symm}}(A, A),$$

witnessed by a bijection of sets

$$\mathbf{Rel}^{\text{symm}}(R^{\text{symm}}, S) \cong \mathbf{Rel}(R, S),$$

natural in $R \in \text{Obj}(\mathbf{Rel}^{\text{symm}}(A, A))$ and $S \in \text{Obj}(\mathbf{Rel}(A, A))$.

00HS 2. *The Symmetric Closure of a Symmetric Relation.* If R is symmetric, then $R^{\text{symm}} = R$.

00HT 3. *Idempotency.* We have

$$(R^{\text{symm}})^{\text{symm}} = R^{\text{symm}}.$$

00HU 4. *Interaction With Inverses.* We have

$$\begin{array}{ccc} \mathbf{Rel}(A, A) & \xrightarrow{(-)^{\text{symm}}} & \mathbf{Rel}(A, A) \\ \left(R^{\dagger} \right)^{\text{symm}} = \left(R^{\text{symm}} \right)^{\dagger}, & \downarrow (-)^{\dagger} & \downarrow (-)^{\dagger} \\ \mathbf{Rel}(A, A) & \xrightarrow{(-)^{\text{symm}}} & \mathbf{Rel}(A, A). \end{array}$$

00HV 5. *Interaction With Composition.* We have

$$\begin{array}{ccc} \mathbf{Rel}(A, A) \times \mathbf{Rel}(A, A) & \xrightarrow{\diamond} & \mathbf{Rel}(A, A) \\ (S \diamond R)^{\text{symm}} = S^{\text{symm}} \diamond R^{\text{symm}}, & \downarrow (-)^{\text{symm}} \times (-)^{\text{symm}} & \downarrow (-)^{\text{symm}} \\ \mathbf{Rel}(A, A) \times \mathbf{Rel}(A, A) & \xrightarrow{\diamond} & \mathbf{Rel}(A, A). \end{array}$$

Proof. **Item 1, Adjointness:** This is a rephrasing of the universal property of the symmetric closure of a relation, stated in [Definition 5.4.2.2.1](#).

Item 2, The Symmetric Closure of a Symmetric Relation: Clear.

Item 3, Idempotency: This follows from [Item 2](#).

Item 4, Interaction With Inverses: Clear.

Item 5, Interaction With Composition: This follows from [Item 2](#) of [Proposition 5.4.2.1.3](#).

□

5.4.3 Transitive Relations

5.4.3.1 Foundations

Let A be a set.

00HY Definition 5.4.3.1.1. A transitive relation is equivalently:⁴²

- A non-unital \mathbb{E}_1 -monoid in $(N_{\bullet}(\mathbf{Rel}(A, A)), \diamond)$;
- A non-unital monoid in $(\mathbf{Rel}(A, A), \diamond)$.

00HZ Remark 5.4.3.1.2. In detail, a relation R on A is **transitive** if we have an inclusion

$$\mu_R: R \diamond R \subset R$$

of relations in $\mathbf{Rel}(A, A)$, i.e. if, for each $a, c \in A$, the following condition is satisfied:

- (★) If there exists some $b \in A$ such that $a \sim_R b$ and $b \sim_R c$, then $a \sim_R c$.

00J0 Definition 5.4.3.1.3. Let A be a set.

1. The **set of transitive relations from A to B** is the subset $\mathbf{Rel}^{\text{trans}}(A)$ of $\mathbf{Rel}(A, A)$ spanned by the transitive relations.
2. The **poset of relations from A to B** is the subposet $\mathbf{Rel}^{\text{trans}}(A)$ of $\mathbf{Rel}(A, A)$ spanned by the transitive relations.

00J3 Proposition 5.4.3.1.4. Let R and S be relations on A .

00J4 1. Interaction With Inverses. If R is transitive, then so is R^\dagger .

00J5 2. Interaction With Composition. If R and S are transitive, then $S \diamond R$ **may fail to be transitive**.

Proof. **Item 1, Interaction With Inverses:** Clear.

Item 2, Interaction With Composition: See [MSE2096272].⁴³ □

⁴²Note that since $\mathbf{Rel}(A, A)$ is posetal, transitivity is a property of a relation, rather than extra structure.

⁴³*Intuition:* Transitivity for R and S fails to imply that of $S \diamond R$ because the composition operation for relations intertwines R and S in an incompatible way:

1. If $a \sim_{S \diamond R} c$ and $c \sim_{S \diamond R} e$, then:

- (a) There is some $b \in A$ such that:

- i. $a \sim_R b$;
- ii. $b \sim_S c$;

- (b) There is some $d \in A$ such that:

- i. $c \sim_R d$;
- ii. $d \sim_S e$.

5.4.3.2 The Transitive Closure of a Relation

Let R be a relation on A .

00J7 Definition 5.4.3.2.1. The **transitive closure** of \sim_R is the relation \sim_R^{trans} ⁴⁴ satisfying the following universal property:⁴⁵

- (★) Given another transitive relation \sim_S on A such that $R \subset S$, there exists an inclusion $\sim_R^{\text{trans}} \subset \sim_S$.

00J8 Construction 5.4.3.2.2. Concretely, \sim_R^{trans} is the free non-unital monoid on R in $(\mathbf{Rel}(A, A), \diamond)$ ⁴⁶, being given by

$$\begin{aligned} R^{\text{trans}} &\stackrel{\text{def}}{=} \bigcup_{n=1}^{\infty} R^{\diamond n} \\ &\stackrel{\text{def}}{=} \bigcup_{n=1}^{\infty} R^{\diamond n} \\ &\stackrel{\text{def}}{=} \left\{ (a, b) \in A \times B \mid \begin{array}{l} \text{there exists some } (x_1, \dots, x_n) \in R^{\times n} \\ \text{such that } a \sim_R x_1 \sim_R \dots \sim_R x_n \sim_R b \end{array} \right\}. \end{aligned}$$

Proof. Clear. □

00J9 Proposition 5.4.3.2.3. Let R be a relation on A .

00JA 1. *Adjointness.* We have an adjunction

$$((-)^{\text{trans}} \dashv \overline{\text{Forget}}) : \mathbf{Rel}(A, A) \begin{array}{c} \xrightarrow{(-)^{\text{trans}}} \\ \perp \\ \xleftarrow{\text{Forget}} \end{array} \mathbf{Rel}^{\text{trans}}(A, A),$$

witnessed by a bijection of sets

$$\mathbf{Rel}^{\text{trans}}(R^{\text{trans}}, S) \cong \mathbf{Rel}(R, S),$$

natural in $R \in \text{Obj}(\mathbf{Rel}^{\text{trans}}(A, A))$ and $S \in \text{Obj}(\mathbf{Rel}(A, B))$.

00JB 2. *The Transitive Closure of a Transitive Relation.* If R is transitive, then $R^{\text{trans}} = R$.

00JC 3. *Idempotency.* We have

$$(R^{\text{trans}})^{\text{trans}} = R^{\text{trans}}.$$

⁴⁴Further Notation: Also written R^{trans} .

⁴⁵Slogan: The transitive closure of R is the smallest transitive relation containing R .

⁴⁶Or, equivalently, the free non-unital \mathbb{E}_1 -monoid on R in $(\mathbf{N}_{\bullet}(\mathbf{Rel}(A, A)), \diamond)$.

00JD 4. *Interaction With Inverses.* We have

$$\left(R^\dagger\right)^{\text{trans}} = \left(R^{\text{trans}}\right)^\dagger, \quad \begin{array}{ccc} \text{Rel}(A, A) & \xrightarrow{(-)^{\text{trans}}} & \text{Rel}(A, A) \\ (-)^\dagger \downarrow & & \downarrow (-)^\dagger \\ \text{Rel}(A, A) & \xrightarrow{(-)^{\text{trans}}} & \text{Rel}(A, A). \end{array}$$

00JE 5. *Interaction With Composition.* We have

$$(S \diamond R)^{\text{trans}} \stackrel{\text{poss.}}{\neq} S^{\text{trans}} \diamond R^{\text{trans}}, \quad \begin{array}{ccc} \text{Rel}(A, A) \times \text{Rel}(A, A) & \xrightarrow{\diamond} & \text{Rel}(A, A) \\ (-)^{\text{trans}} \times (-)^{\text{trans}} \downarrow & \times & \downarrow (-)^{\text{trans}} \\ \text{Rel}(A, A) \times \text{Rel}(A, A) & \xrightarrow{\diamond} & \text{Rel}(A, A). \end{array}$$

Proof. **Item 1, Adjointness:** This is a rephrasing of the universal property of the transitive closure of a relation, stated in [Definition 5.4.3.2.1](#).

Item 2, The Transitive Closure of a Transitive Relation: Clear.

Item 3, Idempotency: This follows from [Item 2](#).

Item 4, Interaction With Inverses: We have

$$\begin{aligned} \left(R^\dagger\right)^{\text{trans}} &= \bigcup_{n=1}^{\infty} \left(R^\dagger\right)^{\diamond n} && \text{(by Construction 5.4.3.2.2)} \\ &= \bigcup_{n=1}^{\infty} (R^{\diamond n})^\dagger && \text{(by Item 4 of Proposition 5.3.12.1.3)} \\ &= \left(\bigcup_{n=1}^{\infty} R^{\diamond n}\right)^\dagger && \text{(by Item 1 of Proposition 5.3.6.1.2)} \\ &= \left(R^{\text{trans}}\right)^\dagger. && \text{(by Construction 5.4.3.2.2)} \end{aligned}$$

Item 5, Interaction With Composition: This follows from [Item 2 of Proposition 5.4.3.1.4](#). □

5.4.4 Equivalence Relations

5.4.4.1 Foundations

Let A be a set.

00JH Definition 5.4.4.1.1. A relation R is an **equivalence relation** if it is reflexive, symmetric, and transitive.⁴⁷

⁴⁷Further Terminology: If instead R is just symmetric and transitive, then it is called a **partial equivalence**

00JJ Example 5.4.4.1.2. The **kernel of a function** $f: A \rightarrow B$ is the equivalence $\sim_{\text{Ker}(f)}$ on A obtained by declaring $a \sim_{\text{Ker}(f)} b$ iff $f(a) = f(b)$.⁴⁸

00JK Definition 5.4.4.1.3. Let A and B be sets.

1. The **set of equivalence relations from A to B** is the subset $\text{Rel}^{\text{eq}}(A, B)$ of $\text{Rel}(A, B)$

00JL spanned by the equivalence relations.

2. The **poset of relations from A to B** is the subposet $\text{Rel}^{\text{eq}}(A, B)$ of $\text{Rel}(A, B)$

00JM spanned by the equivalence relations.

5.4.4.2 The Equivalence Closure of a Relation

Let R be a relation on A .

00JP Definition 5.4.4.2.1. The **equivalence closure**⁴⁹ of \sim_R is the relation \sim_R^{eq} ⁵⁰ satisfying the following universal property:⁵¹

- (★) Given another equivalence relation \sim_S on A such that $R \subset S$, there exists an inclusion $\sim_R^{\text{eq}} \subset \sim_S$.

00JQ Construction 5.4.4.2.2. Concretely, \sim_R^{eq} is the equivalence relation on A defined by

$$\begin{aligned} R^{\text{eq}} &\stackrel{\text{def}}{=} \left(\left(R^{\text{refl}} \right)^{\text{symm}} \right)^{\text{trans}} \\ &= \left(\left(R^{\text{symm}} \right)^{\text{trans}} \right)^{\text{refl}} \\ &= \left\{ (a, b) \in A \times B \mid \begin{array}{l} \text{there exists } (x_1, \dots, x_n) \in R^{\times n} \text{ satisfying at} \\ \text{least one of the following conditions:} \end{array} \right\} \\ &\quad \left. \begin{array}{l} 1. \text{ The following conditions are satisfied:} \\ \quad (a) \text{ We have } a \sim_R x_1 \text{ or } x_1 \sim_R a; \\ \quad (b) \text{ We have } x_i \sim_R x_{i+1} \text{ or } x_{i+1} \sim_R x_i \\ \quad \text{for each } 1 \leq i \leq n-1; \\ \quad (c) \text{ We have } b \sim_R x_n \text{ or } x_n \sim_R b; \\ 2. \text{ We have } a = b. \end{array} \right\}. \end{aligned}$$

Proof. From the universal properties of the reflexive, symmetric, and transitive closures of a relation (Definitions 5.4.1.2.1, 5.4.2.2.1 and 5.4.3.2.1), we see that it suffices to prove that:

relation.

⁴⁸The kernel $\text{Ker}(f): A \rightarrow A$ of f is the monad induced by the adjunction $\text{Gr}(f) \dashv f^{-1}: A \rightleftarrows B$ in Rel of Item 2 of Proposition 5.3.1.1.2.

⁴⁹Further Terminology: Also called the **equivalence relation associated to \sim_R** .

⁵⁰Further Notation: Also written R^{eq} .

⁵¹Slogan: The equivalence closure of R is the smallest equivalence relation containing R .

- 00JR** 1. The symmetric closure of a reflexive relation is still reflexive;
00JS 2. The transitive closure of a symmetric relation is still symmetric;

which are both clear. \square

00JT Proposition 5.4.4.2.3. Let R be a relation on A .

- 00JU** 1. *Adjointness.* We have an adjunction

$$((-)^{\text{eq}} \dashv \overline{\text{eq}}) : \mathbf{Rel}(A, B) \begin{array}{c} \xrightarrow{(-)^{\text{eq}}} \\ \perp \\ \xleftarrow{\overline{\text{eq}}} \end{array} \mathbf{Rel}^{\text{eq}}(A, B),$$

witnessed by a bijection of sets

$$\mathbf{Rel}^{\text{eq}}(R^{\text{eq}}, S) \cong \mathbf{Rel}(R, S),$$

natural in $R \in \text{Obj}(\mathbf{Rel}^{\text{eq}}(A, B))$ and $S \in \text{Obj}(\mathbf{Rel}(A, B))$.

- 00JV** 2. *The Equivalence Closure of an Equivalence Relation.* If R is an equivalence relation, then $R^{\text{eq}} = R$.
00JW 3. *Idempotency.* We have

$$(R^{\text{eq}})^{\text{eq}} = R^{\text{eq}}.$$

Proof. **Item 1, Adjointness:** This is a rephrasing of the universal property of the equivalence closure of a relation, stated in [Definition 5.4.4.2.1](#).

Item 2, The Equivalence Closure of an Equivalence Relation: Clear.

Item 3, Idempotency: This follows from [Item 2](#). \square

5.4.5 Quotients by Equivalence Relations

5.4.5.1 Equivalence Classes

Let A be a set, let R be a relation on A , and let $a \in A$.

00JZ Definition 5.4.5.1.1. The **equivalence class associated to a** is the set $[a]$ defined by

$$\begin{aligned} [a] &\stackrel{\text{def}}{=} \{x \in X \mid x \sim_R a\} \\ &= \{x \in X \mid a \sim_R x\}. \end{aligned} \quad (\text{since } R \text{ is symmetric})$$

5.4.5.2 Quotients of Sets by Equivalence Relations

Let A be a set and let R be a relation on A .

00K1 Definition 5.4.5.2.1. The **quotient of X by R** is the set X/\sim_R defined by

$$X/\sim_R \stackrel{\text{def}}{=} \{[a] \in \mathcal{P}(X) \mid a \in X\}.$$

00K2 Remark 5.4.5.2.2. The reason we define quotient sets for equivalence relations only is that each of the properties of being an equivalence relation—reflexivity, symmetry, and transitivity—ensures that the equivalences classes $[a]$ of X under R are well-behaved:

- *Reflexivity.* If R is reflexive, then, for each $a \in X$, we have $a \in [a]$.
- *Symmetry.* The equivalence class $[a]$ of an element a of X is defined by

$$[a] \stackrel{\text{def}}{=} \{x \in X \mid x \sim_R a\},$$

but we could equally well define

$$[a]' \stackrel{\text{def}}{=} \{x \in X \mid a \sim_R x\}$$

instead. This is not a problem when R is symmetric, as we then have $[a] = [a]'$.⁵²

- *Transitivity.* If R is transitive, then $[a]$ and $[b]$ are disjoint iff $a \not\sim_R b$, and equal otherwise.

00K3 Proposition 5.4.5.2.3. Let $f: X \rightarrow Y$ be a function and let R be a relation on X .

00K4 1. *As a Coequaliser.* We have an isomorphism of sets

$$X/\sim_R^{\text{eq}} \cong \text{CoEq}\left(R \hookrightarrow X \times X \xrightarrow{\text{pr}_1} X\right),$$

where \sim_R^{eq} is the equivalence relation generated by \sim_R .

00K5 2. *As a Pushout.* We have an isomorphism of sets⁵³

$$\begin{array}{ccc} X/\sim_R^{\text{eq}} & \xleftarrow{\quad} & X \\ \uparrow & \lrcorner & \uparrow \\ X/\sim_R^{\text{eq}} & \cong & X \coprod_{\text{Eq}(\text{pr}_1, \text{pr}_2)} X, \\ & & \uparrow \\ & & X \leftarrow \text{Eq}(\text{pr}_1, \text{pr}_2). \end{array}$$

where \sim_R^{eq} is the equivalence relation generated by \sim_R .

⁵²When categorifying equivalence relations, one finds that $[a]$ and $[a]'$ correspond to presheaves and copresheaves; see ??.

⁵³Dually, we also have an isomorphism of sets

$$\begin{array}{ccc} \text{Eq}(\text{pr}_1, \text{pr}_2) & \longrightarrow & X \\ \downarrow & \lrcorner & \downarrow \\ X & \longrightarrow & X/\sim_R^{\text{eq}}. \end{array}$$

$$\text{Eq}(\text{pr}_1, \text{pr}_2) \cong X \times_{X/\sim_R^{\text{eq}}} X,$$

00K6 3. *The First Isomorphism Theorem for Sets.* We have an isomorphism of sets^{54,55}

$$X/\sim_{\text{Ker}(f)} \cong \text{Im}(f).$$

00K7 4. *Descending Functions to Quotient Sets, I.* Let R be an equivalence relation on X . The following conditions are equivalent:

- (a) There exists a map

$$\bar{f}: X/\sim_R \rightarrow Y$$

making the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ q \downarrow & \exists \nearrow \bar{f} & \\ X/\sim_R & & \end{array}$$

commute.

- (b) For each $x, y \in X$, if $x \sim_R y$, then $f(x) = f(y)$.

00K8 5. *Descending Functions to Quotient Sets, II.* Let R be an equivalence relation on X . If the conditions of Item 4 hold, then \bar{f} is the *unique* map making the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ q \downarrow & \exists! \nearrow \bar{f} & \\ X/\sim_R & & \end{array}$$

commute.

00K9 6. *Descending Functions to Quotient Sets, III.* Let R be an equivalence relation on X . If the conditions of Item 4 hold, then the following conditions are equivalent:

⁵⁴Further Terminology: The set $X/\sim_{\text{Ker}(f)}$ is often called the **coimage** of f , and denoted by $\text{Coim}(f)$.

⁵⁵In a sense this is a result relating the monad in **Rel** induced by f with the comonad in **Rel** induced by f , as the kernel and image

$$\begin{aligned} \text{Ker}(f) : X &\dashrightarrow X, \\ \text{Im}(f) &\subset Y \end{aligned}$$

of f are respectively the induced monads and comonads of the adjunction

$$\left(\text{Gr}(f) \dashv f^{-1} \right) : A \begin{array}{c} \xrightarrow{\quad} \\ \perp \\ \xleftarrow{\quad} \end{array} B$$

of Item 2 of Proposition 5.3.1.1.2.

- (a) The map \bar{f} is an injection.
- (b) For each $x, y \in X$, we have $x \sim_R y$ iff $f(x) = f(y)$.

00KA 7. *Descending Functions to Quotient Sets, IV.* Let R be an equivalence relation on X . If the conditions of [Item 4](#) hold, then the following conditions are equivalent:

- (a) The map $f: X \rightarrow Y$ is surjective.
- (b) The map $\bar{f}: X/\sim_R \rightarrow Y$ is surjective.

00KB 8. *Descending Functions to Quotient Sets, V.* Let R be a relation on X and let \sim_R^{eq} be the equivalence relation associated to R . The following conditions are equivalent:

00KC (a) The map f satisfies the equivalent conditions of [Item 4](#):

- There exists a map

$$\bar{f}: X/\sim_R^{\text{eq}} \rightarrow Y$$

making the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ q \downarrow & \nearrow \exists \bar{f} & \\ X/\sim_R^{\text{eq}} & & \end{array}$$

commute.

- For each $x, y \in X$, if $x \sim_R^{\text{eq}} y$, then $f(x) = f(y)$.

00KD (b) For each $x, y \in X$, if $x \sim_R y$, then $f(x) = f(y)$.

Proof. [Item 1](#), As a Coequaliser: Omitted.

[Item 2](#), As a Pushout: Omitted.

[Item 3](#), The First Isomorphism Theorem for Sets: Clear.

[Item 4](#), Descending Functions to Quotient Sets, I: See [[Pro24o](#)].

[Item 5](#), Descending Functions to Quotient Sets, II: See [[Pro24y](#)].

[Item 6](#), Descending Functions to Quotient Sets, III: See [[Pro24n](#)].

[Item 7](#), Descending Functions to Quotient Sets, IV: See [[Pro24m](#)].

[Item 8](#), Descending Functions to Quotient Sets, V: The implication [Item 8a](#) \implies [Item 8b](#) is clear.

Conversely, suppose that, for each $x, y \in X$, if $x \sim_R y$, then $f(x) = f(y)$. Spelling out the definition of the equivalence closure of R , we see that the condition $x \sim_R^{\text{eq}} y$ unwinds to the following:

(★) There exist $(x_1, \dots, x_n) \in R^{\times n}$ satisfying at least one of the following conditions:

1. The following conditions are satisfied:
 - (a) We have $x \sim_R x_1$ or $x_1 \sim_R x$;

- (b) We have $x_i \sim_R x_{i+1}$ or $x_{i+1} \sim_R x_i$ for each $1 \leq i \leq n - 1$;
(c) We have $y \sim_R x_n$ or $x_n \sim_R y$;
2. We have $x = y$.

Now, if $x = y$, then $f(x) = f(y)$ trivially; otherwise, we have

$$\begin{aligned} f(x) &= f(x_1), \\ f(x_1) &= f(x_2), \\ &\vdots \\ f(x_{n-1}) &= f(x_n), \\ f(x_n) &= f(y), \end{aligned}$$

and $f(x) = f(y)$, as we wanted to show. \square

5.5 Functoriality of Powersets

5.5.1 Direct Images

Let A and B be sets and let $R: A \rightarrow B$ be a relation.

00KG Definition 5.5.1.1.1. The **direct image function associated to R** is the function⁵⁶

$$R_*: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by^{57,58}

$$\begin{aligned} R_*(U) &\stackrel{\text{def}}{=} R(U) \\ &\stackrel{\text{def}}{=} \bigcup_{a \in U} R(a) \\ &= \left\{ b \in B \middle| \begin{array}{l} \text{there exists some } a \in U \\ \text{such that } b \in R(a) \end{array} \right\} \end{aligned}$$

for each $U \in \mathcal{P}(A)$.

⁵⁶Further Notation: Also written $\exists_R: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$. This notation comes from the fact that the following statements are equivalent, where $b \in B$ and $U \in \mathcal{P}(A)$:

- We have $b \in \exists_R(U)$.
- There exists some $a \in U$ such that $b \in R(a)$.

⁵⁷Further Terminology: The set $R(U)$ is called the **direct image of U by R** .

⁵⁸We also have

$$R_*(U) = B \setminus R_!(A \setminus U);$$

see Item 7 of Proposition 5.5.1.1.3.

00KH Remark 5.5.1.1.2. Identifying subsets of A with relations from pt to A via ?? of ??, we see that the direct image function associated to R is equivalently the function

$$R_* : \underbrace{\mathcal{P}(A)}_{\cong \text{Rel}(\text{pt}, A)} \rightarrow \underbrace{\mathcal{P}(B)}_{\cong \text{Rel}(\text{pt}, B)}$$

defined by

$$R_*(U) \stackrel{\text{def}}{=} R \diamond U$$

for each $U \in \mathcal{P}(A)$, where $R \diamond U$ is the composition

$$\text{pt} \xrightarrow{U} A \xrightarrow{R} B.$$

00KJ Proposition 5.5.1.1.3. Let $R: A \rightarrow B$ be a relation.

00KK 1. Functoriality. The assignment $U \mapsto R_*(U)$ defines a functor

$$R_*: (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

where

- *Action on Objects.* For each $U \in \mathcal{P}(A)$, we have

$$[R_*](U) \stackrel{\text{def}}{=} R_*(U);$$

- *Action on Morphisms.* For each $U, V \in \mathcal{P}(A)$:

- If $U \subset V$, then $R_*(U) \subset R_*(V)$.

00KL 2. Adjointness. We have an adjunction

$$(R_* \dashv R_{-1}): \mathcal{P}(A) \begin{array}{c} \xrightarrow{R_*} \\ \perp \\ \xleftarrow{R_{-1}} \end{array} \mathcal{P}(B),$$

witnessed by a bijections of sets

$$\text{Hom}_{\mathcal{P}(A)}(R_*(U), V) \cong \text{Hom}_{\mathcal{P}(B)}(U, R_{-1}(V)),$$

natural in $U \in \mathcal{P}(A)$ and $V \in \mathcal{P}(B)$, i.e. such that:

- (★) The following conditions are equivalent:

- We have $R_*(U) \subset V$;
- We have $U \subset R_{-1}(V)$.

00KM 3. *Preservation of Colimits.* We have an equality of sets

$$R_* \left(\bigcup_{i \in I} U_i \right) = \bigcup_{i \in I} R_*(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$. In particular, we have equalities

$$\begin{aligned} R_*(U) \cup R_*(V) &= R_*(U \cup V), \\ R_*(\emptyset) &= \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

00KN 4. *Oplax Preservation of Limits.* We have an inclusion of sets

$$R_* \left(\bigcap_{i \in I} U_i \right) \subset \bigcap_{i \in I} R_*(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$. In particular, we have inclusions

$$\begin{aligned} R_*(U \cap V) &\subset R_*(U) \cap R_*(V), \\ R_*(A) &\subset B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

00KP 5. *Symmetric Strict Monoidality With Respect to Unions.* The direct image function of [Item 1](#) has a symmetric strict monoidal structure

$$\left(R_*, R_*^\otimes, R_{*\mid \mathbb{P}}^\otimes \right) : (\mathcal{P}(A), \cup, \emptyset) \rightarrow (\mathcal{P}(B), \cup, \emptyset),$$

being equipped with equalities

$$\begin{aligned} R_{*\mid U,V}^\otimes : R_*(U) \cup R_*(V) &\xrightarrow{=} R_*(U \cup V), \\ R_{*\mid \mathbb{P}}^\otimes : \emptyset &\xrightarrow{=} \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

00KQ 6. *Symmetric Oplax Monoidality With Respect to Intersections.* The direct image function of [Item 1](#) has a symmetric oplax monoidal structure

$$\left(R_*, R_*^\otimes, R_{*\mid \mathbb{P}}^\otimes \right) : (\mathcal{P}(A), \cap, A) \rightarrow (\mathcal{P}(B), \cap, B),$$

being equipped with inclusions

$$\begin{aligned} R_{*\mid U,V}^\otimes : R_*(U \cap V) &\subset R_*(U) \cap R_*(V), \\ R_{*\mid \mathbb{P}}^\otimes : R_*(A) &\subset B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

00KR 7. *Relation to Direct Images With Compact Support.* We have

$$R_*(U) = B \setminus R_!(A \setminus U)$$

for each $U \in \mathcal{P}(A)$.

Proof. **Item 1, Functoriality:** Clear.

Item 2, Adjointness: This follows from ?? of ??.

Item 3, Preservation of Colimits: This follows from Item 2 and ?? of ??.

Item 4, Oplax Preservation of Limits: Omitted.

Item 5, Symmetric Strict Monoidality With Respect to Unions: This follows from Item 3.

Item 6, Symmetric Oplax Monoidality With Respect to Intersections: This follows from Item 4.

Item 7, Relation to Direct Images With Compact Support: The proof proceeds in the same way as in the case of functions (?? of ??): applying Item 7 of Proposition 5.5.4.1.3 to $A \setminus U$, we have

$$\begin{aligned} R_!(A \setminus U) &= B \setminus R_*(A \setminus (A \setminus U)) \\ &= B \setminus R_*(U). \end{aligned}$$

Taking complements, we then obtain

$$\begin{aligned} R_*(U) &= B \setminus (B \setminus R_!(A \setminus U)), \\ &= B \setminus R_!(A \setminus U), \end{aligned}$$

which finishes the proof. □

00KS Proposition 5.5.1.1.4. Let $R: A \rightarrow B$ be a relation.

00KT 1. *Functionality I.* The assignment $R \mapsto R_*$ defines a function

$$(-)_*: \text{Rel}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B)).$$

00KU 2. *Functionality II.* The assignment $R \mapsto R_*$ defines a function

$$(-)_*: \text{Rel}(A, B) \rightarrow \text{Pos}((\mathcal{P}(A), \subset), (\mathcal{P}(B), \subset)).$$

00KV 3. *Interaction With Identities.* For each $A \in \text{Obj}(\text{Sets})$, we have⁵⁹

$$(\chi_A)_* = \text{id}_{\mathcal{P}(A)};$$

⁵⁹That is, the postcomposition function

$$(\chi_A)_*: \text{Rel}(\text{pt}, A) \rightarrow \text{Rel}(\text{pt}, A)$$

is equal to $\text{id}_{\text{Rel}(\text{pt}, A)}$.

00KW 4. *Interaction With Composition.* For each pair of composable relations $R: A \rightarrow B$ and $S: B \rightarrow C$, we have⁶⁰

$$\begin{array}{ccc} \mathcal{P}(A) & \xrightarrow{R_*} & \mathcal{P}(B) \\ (S \diamond R)_* = S_* \circ R_*, & \searrow & \downarrow S_* \\ & (S \diamond R)_* & \\ & & \mathcal{P}(C). \end{array}$$

Proof. **Item 1, Functionality I:** Clear.

Item 2, Functionality II: Clear.

Item 3, Interaction With Identities: Indeed, we have

$$\begin{aligned} (\chi_A)_*(U) &\stackrel{\text{def}}{=} \bigcup_{a \in U} \chi_A(a) \\ &\stackrel{\text{def}}{=} \bigcup_{a \in U} \{a\} \\ &= U \\ &\stackrel{\text{def}}{=} \text{id}_{\mathcal{P}(A)}(U) \end{aligned}$$

for each $U \in \mathcal{P}(A)$. Thus $(\chi_A)_* = \text{id}_{\mathcal{P}(A)}$.

Item 4, Interaction With Composition: Indeed, we have

$$\begin{aligned} (S \diamond R)_*(U) &\stackrel{\text{def}}{=} \bigcup_{a \in U} [S \diamond R](a) \\ &\stackrel{\text{def}}{=} \bigcup_{a \in U} S(R(a)) \\ &\stackrel{\text{def}}{=} \bigcup_{a \in U} S_*(R(a)) \\ &= S_* \left(\bigcup_{a \in U} R(a) \right) \\ &\stackrel{\text{def}}{=} S_*(R_*(U)) \\ &\stackrel{\text{def}}{=} [S_* \circ R_*](U) \end{aligned}$$

for each $U \in \mathcal{P}(A)$, where we used **Item 3** of **Proposition 5.5.1.1.3**. Thus $(S \diamond R)_* = S_* \circ R_*$. \square

⁶⁰That is, we have

$$\begin{array}{ccc} \text{Rel(pt, } A) & \xrightarrow{R_*} & \text{Rel(pt, } B) \\ (S \diamond R)_* = S_* \circ R_*, & \searrow & \downarrow S_* \\ & (S \diamond R)_* & \\ & & \text{Rel(pt, } C). \end{array}$$

5.5.2 Strong Inverse Images

Let A and B be sets and let $R: A \rightarrow B$ be a relation.

00KY Definition 5.5.2.1.1. The **strong inverse image function associated to R** is the function

$$R_{-1}: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

defined by⁶¹

$$R_{-1}(V) \stackrel{\text{def}}{=} \{a \in A \mid R(a) \subset V\}$$

for each $V \in \mathcal{P}(B)$.

00KZ Remark 5.5.2.1.2. Identifying subsets of B with relations from pt to B via ?? of ??, we see that the inverse image function associated to R is equivalently the function

$$R_{-1}: \underbrace{\mathcal{P}(B)}_{\cong \text{Rel}(\text{pt}, B)} \rightarrow \underbrace{\mathcal{P}(A)}_{\cong \text{Rel}(\text{pt}, A)}$$

defined by

$$R_{-1}(V) \stackrel{\text{def}}{=} \text{Rift}_R(V), \quad \begin{array}{ccc} & A & \\ \text{Rift}_R(V) & \nearrow \dashv \searrow & \downarrow R \\ \text{pt} & \xrightarrow[V]{+} & B, \end{array}$$

and being explicitly computed by

$$\begin{aligned} R_{-1}(V) &\stackrel{\text{def}}{=} \text{Rift}_R(V) \\ &\cong \int_{x \in B} \text{Hom}_{\{\text{t}, \text{f}\}}(R_{-1}^x, V_{-2}^x), \end{aligned}$$

where we have used **Item 12 of Proposition 5.2.5.1.1.**

⁶¹Further Terminology: The set $R_{-1}(V)$ is called the **strong inverse image of V by R** .

Proof. We have

$$\begin{aligned}
 \text{Rift}_R(V) &\cong \int_{x \in B} \text{Hom}_{\{\text{t,f}\}}(R_{-1}^x, V_{-2}^x) \\
 &= \left\{ a \in A \mid \int_{x \in B} \text{Hom}_{\{\text{t,f}\}}(R_a^x, V_\star^x) = \text{true} \right\} \\
 &= \left\{ a \in A \mid \begin{array}{l} \text{for each } x \in B, \text{ at least one of the} \\ \text{following conditions hold:} \end{array} \right. \\
 &\quad \left. \begin{array}{l} 1. \text{ We have } R_a^x = \text{false} \\ 2. \text{ The following conditions hold:} \end{array} \right. \\
 &\quad \left. \begin{array}{l} (a) \text{ We have } R_a^x = \text{true} \\ (b) \text{ We have } V_\star^x = \text{true} \end{array} \right\} \\
 &= \left\{ a \in A \mid \begin{array}{l} \text{for each } x \in B, \text{ at least one of the} \\ \text{following conditions hold:} \end{array} \right. \\
 &\quad \left. \begin{array}{l} 1. \text{ We have } x \notin R(a) \\ 2. \text{ The following conditions hold:} \end{array} \right. \\
 &\quad \left. \begin{array}{l} (a) \text{ We have } x \in R(a) \\ (b) \text{ We have } x \in V \end{array} \right\} \\
 &= \{a \in A \mid \text{for each } x \in R(a), \text{ we have } x \in V\} \\
 &= \{a \in A \mid R(a) \subset V\} \\
 &\stackrel{\text{def}}{=} R_{-1}(V).
 \end{aligned}$$

This finishes the proof. □

00L0 Proposition 5.5.2.1.3. Let $R: A \rightarrow B$ be a relation.

00L1 1. *Functoriality.* The assignment $V \mapsto R_{-1}(V)$ defines a functor

$$R_{-1}: (\mathcal{P}(B), \subset) \rightarrow (\mathcal{P}(A), \subset)$$

where

- *Action on Objects.* For each $V \in \mathcal{P}(B)$, we have

$$[R_{-1}](V) \stackrel{\text{def}}{=} R_{-1}(V);$$

- *Action on Morphisms.* For each $U, V \in \mathcal{P}(B)$:

– If $U \subset V$, then $R_{-1}(U) \subset R_{-1}(V)$.

00L2 2. *Adjointness.* We have an adjunction

$$(R_* \dashv R_{-1}): \mathcal{P}(A) \begin{array}{c} \xrightarrow{R_*} \\ \perp \\ \xleftarrow{R_{-1}} \end{array} \mathcal{P}(B),$$

witnessed by a bijections of sets

$$\text{Hom}_{\mathcal{P}(A)}(R_*(U), V) \cong \text{Hom}_{\mathcal{P}(A)}(U, R_{-1}(V)),$$

natural in $U \in \mathcal{P}(A)$ and $V \in \mathcal{P}(B)$, i.e. such that:

(★) The following conditions are equivalent:

- We have $R_*(U) \subset V$;
- We have $U \subset R_{-1}(V)$.

00L3 3. *Lax Preservation of Colimits.* We have an inclusion of sets

$$\bigcup_{i \in I} R_{-1}(U_i) \subset R_{-1}\left(\bigcup_{i \in I} U_i\right),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(B)^{\times I}$. In particular, we have inclusions

$$\begin{aligned} R_{-1}(U) \cup R_{-1}(V) &\subset R_{-1}(U \cup V), \\ \emptyset &\subset R_{-1}(\emptyset), \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

00L4 4. *Preservation of Limits.* We have an equality of sets

$$R_{-1}\left(\bigcap_{i \in I} U_i\right) = \bigcap_{i \in I} R_{-1}(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(B)^{\times I}$. In particular, we have equalities

$$\begin{aligned} R_{-1}(U \cap V) &= R_{-1}(U) \cap R_{-1}(V), \\ R_{-1}(B) &= B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

00L5 5. *Symmetric Lax Monoidality With Respect to Unions.* The direct image with compact support function of Item 1 has a symmetric lax monoidal structure

$$\left(R_{-1}, R_{-1}^{\otimes}, R_{-1|*}^{\otimes}\right): (\mathcal{P}(A), \cup, \emptyset) \rightarrow (\mathcal{P}(B), \cup, \emptyset),$$

being equipped with inclusions

$$\begin{aligned} R_{-1|U,V}^{\otimes}: R_{-1}(U) \cup R_{-1}(V) &\subset R_{-1}(U \cup V), \\ R_{-1|*}^{\otimes}: \emptyset &\subset R_{-1}(\emptyset), \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

- 00L6** 6. *Symmetric Strict Monoidality With Respect to Intersections.* The direct image function of **Item 1** has a symmetric strict monoidal structure

$$\left(R_{-1}, R_{-1}^{\otimes}, R_{-1|*}^{\otimes} \right) : (\mathcal{P}(A), \cap, A) \rightarrow (\mathcal{P}(B), \cap, B),$$

being equipped with equalities

$$R_{-1|U,V}^{\otimes} : R_{-1}(U \cap V) \xrightarrow{\cong} R_{-1}(U) \cap R_{-1}(V),$$

$$R_{-1|*}^{\otimes} : R_{-1}(A) \xrightarrow{\cong} B,$$

natural in $U, V \in \mathcal{P}(B)$.

- 00L7** 7. *Interaction With Weak Inverse Images I.* We have

$$R_{-1}(V) = A \setminus R^{-1}(B \setminus V)$$

for each $V \in \mathcal{P}(B)$.

- 00L8** 8. *Interaction With Weak Inverse Images II.* Let $R: A \nrightarrow B$ be a relation from A to B .

- 00L9** (a) If R is a total relation, then we have an inclusion of sets

$$R_{-1}(V) \subset R^{-1}(V)$$

natural in $V \in \mathcal{P}(B)$.

- 00LA** (b) If R is total and functional, then the above inclusion is in fact an equality.

- 00LB** (c) Conversely, if we have $R_{-1} = R^{-1}$, then R is total and functional.

Proof. **Item 1, Functoriality:** Clear.

Item 2, Adjointness: This follows from ?? of ??.

Item 3, Lax Preservation of Colimits: Omitted.

Item 4, Preservation of Limits: This follows from **Item 2** and ?? of ??.

Item 5, Symmetric Lax Monoidality With Respect to Unions: This follows from **Item 3**.

Item 6, Symmetric Strict Monoidality With Respect to Intersections: This follows from **Item 4**.

Item 7, Interaction With Weak Inverse Images I: We claim we have an equality

$$R_{-1}(B \setminus V) = A \setminus R^{-1}(V).$$

Indeed, we have

$$R_{-1}(B \setminus V) = \{a \in A \mid R(a) \subset B \setminus V\},$$

$$A \setminus R^{-1}(V) = \{a \in A \mid R(a) \cap V = \emptyset\}.$$

Taking $V = B \setminus V$ then implies the original statement.

Item 8, Interaction With Weak Inverse Images II: **Item 8a** is clear, while **Items 8b and 8c** follow from **Item 6** of Proposition 5.3.1.1.2. \square

00LC Proposition 5.5.2.1.4. Let $R: A \rightarrow B$ be a relation.

00LD 1. *Functionality I.* The assignment $R \mapsto R_{-1}$ defines a function

$$(-)_{-1}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B)).$$

00LE 2. *Functionality II.* The assignment $R \mapsto R_{-1}$ defines a function

$$(-)_{-1}: \text{Sets}(A, B) \rightarrow \text{Pos}((\mathcal{P}(A), \subset), (\mathcal{P}(B), \subset)).$$

00LF 3. *Interaction With Identities.* For each $A \in \text{Obj}(\text{Sets})$, we have

$$(\text{id}_A)_{-1} = \text{id}_{\mathcal{P}(A)};$$

00LG 4. *Interaction With Composition.* For each pair of composable relations $R: A \rightarrow B$ and $S: B \rightarrow C$, we have

$$\begin{array}{ccc} \mathcal{P}(C) & \xrightarrow{S_{-1}} & \mathcal{P}(B) \\ (S \diamond R)_{-1} = R_{-1} \circ S_{-1}, & \searrow & \downarrow R_{-1} \\ & & \mathcal{P}(A). \end{array}$$

Proof. **Item 1, Functionality I:** Clear.

Item 2, Functionality II: Clear.

Item 3, Interaction With Identities: Indeed, we have

$$\begin{aligned} (\chi_A)_{-1}(U) &\stackrel{\text{def}}{=} \{a \in A \mid \chi_A(a) \subset U\} \\ &\stackrel{\text{def}}{=} \{a \in A \mid \{a\} \subset U\} \\ &= U \end{aligned}$$

for each $U \in \mathcal{P}(A)$. Thus $(\chi_A)_{-1} = \text{id}_{\mathcal{P}(A)}$.

Item 4, Interaction With Composition: Indeed, we have

$$\begin{aligned} (S \diamond R)_{-1}(U) &\stackrel{\text{def}}{=} \{a \in A \mid [S \diamond R](a) \subset U\} \\ &\stackrel{\text{def}}{=} \{a \in A \mid S(R(a)) \subset U\} \\ &\stackrel{\text{def}}{=} \{a \in A \mid S_*(R(a)) \subset U\} \\ &= \{a \in A \mid R(a) \subset S_{-1}(U)\} \\ &\stackrel{\text{def}}{=} R_{-1}(S_{-1}(U)) \\ &\stackrel{\text{def}}{=} [R_{-1} \circ S_{-1}](U) \end{aligned}$$

for each $U \in \mathcal{P}(C)$, where we used **Item 2 of Proposition 5.5.2.1.3**, which implies that the conditions

- We have $S_*(R(a)) \subset U$;
- We have $R(a) \subset S_{-1}(U)$;

are equivalent. Thus $(S \diamond R)_{-1} = R_{-1} \circ S_{-1}$. □

5.5.3 Weak Inverse Images

Let A and B be sets and let $R: A \rightarrow B$ be a relation.

00LJ Definition 5.5.3.1.1. The **weak inverse image function associated to R** ⁶² is the function

$$R^{-1}: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

defined by⁶³

$$R^{-1}(V) \stackrel{\text{def}}{=} \{a \in A \mid R(a) \cap V \neq \emptyset\}$$

for each $V \in \mathcal{P}(B)$.

00LK Remark 5.5.3.1.2. Identifying subsets of B with relations from B to pt via ?? of ??, we see that the weak inverse image function associated to R is equivalently the function

$$R^{-1}: \underbrace{\mathcal{P}(B)}_{\cong \text{Rel}(B,\text{pt})} \rightarrow \underbrace{\mathcal{P}(A)}_{\cong \text{Rel}(A,\text{pt})}$$

defined by

$$R^{-1}(V) \stackrel{\text{def}}{=} V \diamond R$$

for each $V \in \mathcal{P}(A)$, where $R \diamond V$ is the composition

$$A \xrightarrow{R} B \xrightarrow{V} \text{pt}.$$

Explicitly, we have

$$\begin{aligned} R^{-1}(V) &\stackrel{\text{def}}{=} V \diamond R \\ &\stackrel{\text{def}}{=} \int^{x \in B} V_x^{-1} \times R_x^x. \end{aligned}$$

⁶²Further Terminology: Also called simply the **inverse image function associated to R** .

⁶³Further Terminology: The set $R^{-1}(V)$ is called the **weak inverse image of V by R** or simply the **inverse image of V by R** .

Proof. We have

$$\begin{aligned}
 V \diamond R &\stackrel{\text{def}}{=} \int^{x \in B} V_x^{-1} \times R_{-2}^x \\
 &= \left\{ a \in A \mid \int^{x \in B} V_x^* \times R_a^x = \text{true} \right\} \\
 &= \left\{ a \in A \mid \begin{array}{l} \text{there exists } x \in B \text{ such that the} \\ \text{following conditions hold:} \end{array} \right. \\
 &\quad \left. \begin{array}{l} 1. \text{ We have } V_x^* = \text{true} \\ 2. \text{ We have } R_a^x = \text{true} \end{array} \right\} \\
 &= \left\{ a \in A \mid \begin{array}{l} \text{there exists } x \in B \text{ such that the} \\ \text{following conditions hold:} \end{array} \right. \\
 &\quad \left. \begin{array}{l} 1. \text{ We have } x \in V \\ 2. \text{ We have } x \in R(a) \end{array} \right\} \\
 &= \{a \in A \mid \text{there exists } x \in V \text{ such that } x \in R(a)\} \\
 &= \{a \in A \mid R(a) \cap V \neq \emptyset\} \\
 &\stackrel{\text{def}}{=} R^{-1}(V)
 \end{aligned}$$

This finishes the proof. \square

00LL Proposition 5.5.3.1.3. Let $R: A \rightarrow B$ be a relation.

00LM 1. Functoriality. The assignment $V \mapsto R^{-1}(V)$ defines a functor

$$R^{-1}: (\mathcal{P}(B), \subset) \rightarrow (\mathcal{P}(A), \subset)$$

where

- *Action on Objects.* For each $V \in \mathcal{P}(B)$, we have

$$[R^{-1}](V) \stackrel{\text{def}}{=} R^{-1}(V);$$

- *Action on Morphisms.* For each $U, V \in \mathcal{P}(B)$:

– If $U \subset V$, then $R^{-1}(U) \subset R^{-1}(V)$.

00LN 2. Adjointness. We have an adjunction

$$(R^{-1} \dashv R_!): \mathcal{P}(B) \begin{array}{c} \xrightarrow{R^{-1}} \\ \perp \\ \xleftarrow{R_!} \end{array} \mathcal{P}(A),$$

witnessed by a bijections of sets

$$\text{Hom}_{\mathcal{P}(A)}(R^{-1}(U), V) \cong \text{Hom}_{\mathcal{P}(A)}(U, R_!(V)),$$

natural in $U \in \mathcal{P}(A)$ and $V \in \mathcal{P}(B)$, i.e. such that:

(★) The following conditions are equivalent:

- We have $R^{-1}(U) \subset V$;
- We have $U \subset R_!(V)$.

00LP 3. *Preservation of Colimits.* We have an equality of sets

$$R^{-1}\left(\bigcup_{i \in I} U_i\right) = \bigcup_{i \in I} R^{-1}(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(B)^{\times I}$. In particular, we have equalities

$$\begin{aligned} R^{-1}(U) \cup R^{-1}(V) &= R^{-1}(U \cup V), \\ R^{-1}(\emptyset) &= \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

00LQ 4. *Oplax Preservation of Limits.* We have an inclusion of sets

$$R^{-1}\left(\bigcap_{i \in I} U_i\right) \subset \bigcap_{i \in I} R^{-1}(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(B)^{\times I}$. In particular, we have inclusions

$$\begin{aligned} R^{-1}(U \cap V) &\subset R^{-1}(U) \cap R^{-1}(V), \\ R^{-1}(A) &\subset B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

00LR 5. *Symmetric Strict Monoidality With Respect to Unions.* The direct image function of [Item 1](#) has a symmetric strict monoidal structure

$$\left(R^{-1}, R^{-1,\otimes}, R_{\sharp}^{-1,\otimes}\right): (\mathcal{P}(A), \cup, \emptyset) \rightarrow (\mathcal{P}(B), \cup, \emptyset),$$

being equipped with equalities

$$\begin{aligned} R_{U,V}^{-1,\otimes}: R^{-1}(U) \cup R^{-1}(V) &\xrightarrow{=} R^{-1}(U \cup V), \\ R_{\sharp}^{-1,\otimes}: \emptyset &\xrightarrow{=} \emptyset, \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

00LS 6. *Symmetric Oplax Monoidality With Respect to Intersections.* The direct image function of [Item 1](#) has a symmetric oplax monoidal structure

$$\left(R^{-1}, R^{-1,\otimes}, R_{\sharp}^{-1,\otimes}\right): (\mathcal{P}(A), \cap, A) \rightarrow (\mathcal{P}(B), \cap, B),$$

being equipped with inclusions

$$\begin{aligned} R_{U,V}^{-1,\otimes}: R^{-1}(U \cap V) &\subset R^{-1}(U) \cap R^{-1}(V), \\ R_{\sharp}^{-1,\otimes}: R^{-1}(A) &\subset B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(B)$.

00LT 7. *Interaction With Strong Inverse Images I.* We have

$$R^{-1}(V) = A \setminus R_{-1}(B \setminus V)$$

for each $V \in \mathcal{P}(B)$.

00LU 8. *Interaction With Strong Inverse Images II.* Let $R: A \rightarrow B$ be a relation from A to B .

00LV (a) If R is a total relation, then we have an inclusion of sets

$$R_{-1}(V) \subset R^{-1}(V)$$

natural in $V \in \mathcal{P}(B)$.

(b) If R is total and functional, then the above inclusion is in fact an equality.

00LW (c) Conversely, if we have $R_{-1} = R^{-1}$, then R is total and functional.

Proof. **Item 1, Functoriality:** Clear.

Item 2, Adjointness: This follows from ?? of ??.

Item 3, Preservation of Colimits: This follows from Item 2 and ?? of ??.

Item 4, Oplax Preservation of Limits: Omitted.

Item 5, Symmetric Strict Monoidality With Respect to Unions: This follows from Item 3.

Item 6, Symmetric Oplax Monoidality With Respect to Intersections: This follows from Item 4.

Item 7, Interaction With Strong Inverse Images I: This follows from Item 7 of Proposition 5.5.2.1.3.

Item 8, Interaction With Strong Inverse Images II: This was proved in Item 8 of Proposition 5.5.2.1.3. \square

00LY **Proposition 5.5.3.1.4.** Let $R: A \rightarrow B$ be a relation.

00LZ 1. *Functionality I.* The assignment $R \mapsto R^{-1}$ defines a function

$$(-)^{-1}: \text{Rel}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B)).$$

00M0 2. *Functionality II.* The assignment $R \mapsto R_{-1}$ defines a function

$$(-)^{-1}: \text{Rel}(A, B) \rightarrow \text{Pos}((\mathcal{P}(A), \subset), (\mathcal{P}(B), \subset)).$$

00M1 3. *Interaction With Identities.* For each $A \in \text{Obj}(\text{Sets})$, we have⁶⁴

$$(\chi_A)^{-1} = \text{id}_{\mathcal{P}(A)};$$

⁶⁴That is, the postcomposition

$$(\chi_A)^{-1}: \text{Rel}(\text{pt}, A) \rightarrow \text{Rel}(\text{pt}, A)$$

is equal to $\text{id}_{\text{Rel}(\text{pt}, A)}$.

- 00M2** 4. *Interaction With Composition.* For each pair of composable relations $R: A \rightarrow B$ and $S: B \rightarrow C$, we have⁶⁵

$$(S \diamond R)^{-1} = R^{-1} \circ S^{-1}, \quad \begin{array}{ccc} \mathcal{P}(C) & \xrightarrow{S^{-1}} & \mathcal{P}(B) \\ & \searrow (S \diamond R)^{-1} & \downarrow R^{-1} \\ & & \mathcal{P}(A). \end{array}$$

Proof. Item 1, Functionality I: Clear.

Item 2, Functionality II: Clear.

Item 3, Interaction With Identities: This follows from ?? of ??.

Item 4, Interaction With Composition: This follows from ?? of ??.

□

5.5.4 Direct Images With Compact Support

Let A and B be sets and let $R: A \rightarrow B$ be a relation.

- 00M4** **Definition 5.5.4.1.1.** The **direct image with compact support function associated to R** is the function⁶⁶

$$R_!: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by^{67,68}

$$\begin{aligned} R_!(U) &\stackrel{\text{def}}{=} \left\{ b \in B \mid \begin{array}{l} \text{for each } a \in A, \text{ if we have} \\ b \in R(a), \text{ then } a \in U \end{array} \right\} \\ &= \left\{ b \in B \mid R^{-1}(b) \subset U \right\} \end{aligned}$$

for each $U \in \mathcal{P}(A)$.

⁶⁵That is, we have

$$(S \diamond R)^{-1} = R^{-1} \circ S^{-1}, \quad \begin{array}{ccc} \text{Rel(pt, } C) & \xrightarrow{R^{-1}} & \text{Rel(pt, } B) \\ & \searrow (S \diamond R)^{-1} & \downarrow S^{-1} \\ & & \text{Rel(pt, } A). \end{array}$$

⁶⁶Further Notation: Also written $\forall_R: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$. This notation comes from the fact that the following statements are equivalent, where $b \in B$ and $U \in \mathcal{P}(A)$:

- We have $b \in \forall_R(U)$.
- For each $a \in A$, if $b \in R(a)$, then $a \in U$.

⁶⁷Further Terminology: The set $R_!(U)$ is called the **direct image with compact support of U by R** .

⁶⁸We also have

$$R_!(U) = B \setminus R_*(A \setminus U);$$

see Item 7 of Proposition 5.5.4.1.3.

00M5 Remark 5.5.4.1.2. Identifying subsets of B with relations from pt to B via ?? of ??, we see that the direct image with compact support function associated to R is equivalently the function

$$R_! : \underbrace{\mathcal{P}(A)}_{\cong \text{Rel}(A, \text{pt})} \rightarrow \underbrace{\mathcal{P}(B)}_{\cong \text{Rel}(B, \text{pt})}$$

defined by

$$R_!(U) \stackrel{\text{def}}{=} \text{Ran}_R(U), \quad \begin{array}{ccc} & B & \\ & \nearrow R & \downarrow \\ A & \xrightarrow[U]{} & \text{pt}, \end{array} \quad \text{Ran}_R(U)$$

being explicitly computed by

$$\begin{aligned} R^*(U) &\stackrel{\text{def}}{=} \text{Ran}_R(U) \\ &\cong \int_{a \in A} \text{Hom}_{\{\text{t,f}\}}(R_a^{-2}, U_a^{-1}), \end{aligned}$$

where we have used Item 11 of Proposition 5.2.5.1.1.

Proof. We have

$$\begin{aligned}
 \text{Ran}_R(V) &\cong \int_{a \in A} \text{Hom}_{\{\text{t,f}\}}(R_a^{-2}, U_a^{-1}) \\
 &= \left\{ b \in B \mid \int_{a \in A} \text{Hom}_{\{\text{t,f}\}}(R_a^b, U_a^\star) = \text{true} \right\} \\
 &= \left\{ b \in B \mid \begin{array}{l} \text{for each } a \in A, \text{ at least one of the} \\ \text{following conditions hold:} \end{array} \right. \\
 &\quad \left. \begin{array}{l} 1. \text{ We have } R_a^b = \text{false} \\ 2. \text{ The following conditions hold:} \end{array} \right. \\
 &\quad \left. \begin{array}{l} (a) \text{ We have } R_a^b = \text{true} \\ (b) \text{ We have } U_a^\star = \text{true} \end{array} \right\} \\
 &= \left\{ b \in B \mid \begin{array}{l} \text{for each } a \in A, \text{ at least one of the} \\ \text{following conditions hold:} \end{array} \right. \\
 &\quad \left. \begin{array}{l} 1. \text{ We have } b \notin R(A) \\ 2. \text{ The following conditions hold:} \end{array} \right. \\
 &\quad \left. \begin{array}{l} (a) \text{ We have } b \in R(a) \\ (b) \text{ We have } a \in U \end{array} \right\} \\
 &= \left\{ b \in B \mid \begin{array}{l} \text{for each } a \in A, \text{ if we have} \\ b \in R(a), \text{ then } a \in U \end{array} \right\} \\
 &= \left\{ b \in B \mid R^{-1}(b) \subset U \right\} \\
 &\stackrel{\text{def}}{=} R^{-1}(U).
 \end{aligned}$$

This finishes the proof. □

00M6 Proposition 5.5.4.1.3. Let $R: A \rightarrow B$ be a relation.

00M7 1. *Functoriality.* The assignment $U \mapsto R_!(U)$ defines a functor

$$R_!: (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

where

- *Action on Objects.* For each $U \in \mathcal{P}(A)$, we have

$$[R_!](U) \stackrel{\text{def}}{=} R_!(U);$$

- *Action on Morphisms.* For each $U, V \in \mathcal{P}(A)$:
 - If $U \subset V$, then $R_!(U) \subset R_!(V)$.

00M8 2. *Adjointness.* We have an adjunction

$$(R^{-1} \dashv R_!): \mathcal{P}(B) \begin{array}{c} \xrightarrow{R^{-1}} \\ \perp \\ \xleftarrow{R_!} \end{array} \mathcal{P}(A),$$

witnessed by a bijections of sets

$$\text{Hom}_{\mathcal{P}(A)}(R^{-1}(U), V) \cong \text{Hom}_{\mathcal{P}(A)}(U, R_!(V)),$$

natural in $U \in \mathcal{P}(A)$ and $V \in \mathcal{P}(B)$, i.e. such that:

(★) The following conditions are equivalent:

- We have $R^{-1}(U) \subset V$;
- We have $U \subset R_!(V)$.

00M9 3. *Lax Preservation of Colimits.* We have an inclusion of sets

$$\bigcup_{i \in I} R_!(U_i) \subset R_!\left(\bigcup_{i \in I} U_i\right),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$. In particular, we have inclusions

$$\begin{aligned} R_!(U) \cup R_!(V) &\subset R_!(U \cup V), \\ \emptyset &\subset R_!(\emptyset), \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

00MA 4. *Preservation of Limits.* We have an equality of sets

$$R_!\left(\bigcap_{i \in I} U_i\right) = \bigcap_{i \in I} R_!(U_i),$$

natural in $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$. In particular, we have equalities

$$\begin{aligned} R_!(U \cap V) &= R_!(U) \cap R_!(V), \\ R_!(A) &= B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

00MB 5. *Symmetric Lax Monoidality With Respect to Unions.* The direct image with compact support function of **Item 1** has a symmetric lax monoidal structure

$$(R_!, R_!^\otimes, R_{!|k}^\otimes): (\mathcal{P}(A), \cup, \emptyset) \rightarrow (\mathcal{P}(B), \cup, \emptyset),$$

being equipped with inclusions

$$\begin{aligned} R_{!|U,V}^\otimes: R_!(U) \cup R_!(V) &\subset R_!(U \cup V), \\ R_{!|k}^\otimes: \emptyset &\subset R_!(\emptyset), \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

- 00MC** 6. *Symmetric Strict Monoidality With Respect to Intersections.* The direct image function of **Item 1** has a symmetric strict monoidal structure

$$\left(R_!, R_!^\otimes, R_{!|*}^\otimes \right) : (\mathcal{P}(A), \cap, A) \rightarrow (\mathcal{P}(B), \cap, B),$$

being equipped with equalities

$$\begin{aligned} R_{!|U,V}^\otimes : R_!(U \cap V) &\xrightarrow{=} R_!(U) \cap R_!(V), \\ R_{!|*}^\otimes : R_!(A) &\xrightarrow{=} B, \end{aligned}$$

natural in $U, V \in \mathcal{P}(A)$.

- 00MD** 7. *Relation to Direct Images.* We have

$$R_!(U) = B \setminus R_*(A \setminus U)$$

for each $U \in \mathcal{P}(A)$.

Proof. **Item 1, Functoriality:** Clear.

Item 2, Adjointness: This follows from ?? of ??.

Item 3, Lax Preservation of Colimits: Omitted.

Item 4, Preservation of Limits: This follows from **Item 2** and ?? of ??.

Item 5, Symmetric Lax Monoidality With Respect to Unions: This follows from **Item 3**.

Item 6, Symmetric Strict Monoidality With Respect to Intersections: This follows from **Item 4**.

Item 7, Relation to Direct Images: This follows from **Item 7** of **Proposition 5.5.1.1.3**.

Alternatively, we may prove it directly as follows, with the proof proceeding in the same way as in the case of functions (?? of ??).

We claim that $R_!(U) = B \setminus R_*(A \setminus U)$:

- *The First Implication.* We claim that

$$R_!(U) \subset B \setminus R_*(A \setminus U).$$

Let $b \in R_!(U)$. We need to show that $b \notin R_*(A \setminus U)$, i.e. that there is no $a \in A \setminus U$ such that $b \in R(a)$.

This is indeed the case, as otherwise we would have $a \in R^{-1}(b)$ and $a \notin U$, contradicting $R^{-1}(b) \subset U$ (which holds since $b \in R_!(U)$).

Thus $b \in B \setminus R_*(A \setminus U)$.

- *The Second Implication.* We claim that

$$B \setminus R_*(A \setminus U) \subset R_!(U).$$

Let $b \in B \setminus R_*(A \setminus U)$. We need to show that $b \in R_!(U)$, i.e. that $R^{-1}(b) \subset U$.

Since $b \notin R_*(A \setminus U)$, there exists no $a \in A \setminus U$ such that $b \in R(a)$, and hence $R^{-1}(b) \subset U$.

Thus $b \in R_!(U)$.

This finishes the proof. \square

00ME Proposition 5.5.4.1.4. Let $R: A \rightarrow B$ be a relation.

00MF 1. *Functionality I.* The assignment $R \mapsto R_!$ defines a function

$$(-)_!: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B)).$$

00MG 2. *Functionality II.* The assignment $R \mapsto R_!$ defines a function

$$(-)_!: \text{Sets}(A, B) \rightarrow \text{Hom}_{\text{Pos}}((\mathcal{P}(A), \subset), (\mathcal{P}(B), \subset)).$$

00MH 3. *Interaction With Identities.* For each $A \in \text{Obj}(\text{Sets})$, we have

$$(\text{id}_A)_! = \text{id}_{\mathcal{P}(A)};$$

00MJ 4. *Interaction With Composition.* For each pair of composable relations $R: A \rightarrow B$ and $S: B \rightarrow C$, we have

$$\begin{array}{ccc} \mathcal{P}(A) & \xrightarrow{R_!} & \mathcal{P}(B) \\ (S \diamond R)_! = S_! \circ R_!, & \searrow & \downarrow S_! \\ & (S \diamond R)_! & \\ & & \mathcal{P}(C). \end{array}$$

Proof. **Item 1, Functionality I:** Clear.

Item 2, Functionality II: Clear.

Item 3, Interaction With Identities: Indeed, we have

$$\begin{aligned} (\chi_A)_!(U) &\stackrel{\text{def}}{=} \{a \in A \mid \chi_A^{-1}(a) \subset U\} \\ &\stackrel{\text{def}}{=} \{a \in A \mid \{a\} \subset U\} \\ &= U \end{aligned}$$

for each $U \in \mathcal{P}(A)$. Thus $(\chi_A)_! = \text{id}_{\mathcal{P}(A)}$.

Item 4, Interaction With Composition: Indeed, we have

$$\begin{aligned} (S \diamond R)_!(U) &\stackrel{\text{def}}{=} \{c \in C \mid [S \diamond R]^{-1}(c) \subset U\} \\ &\stackrel{\text{def}}{=} \{c \in C \mid S^{-1}(R^{-1}(c)) \subset U\} \\ &= \{c \in C \mid R^{-1}(c) \subset S_!(U)\} \\ &\stackrel{\text{def}}{=} R_!(S_!(U)) \\ &\stackrel{\text{def}}{=} [R_! \circ S_!](U) \end{aligned}$$

for each $U \in \mathcal{P}(C)$, where we used **Item 2** of **Proposition 5.5.4.1.3**, which implies that the conditions

- We have $S^{-1}(R^{-1}(c)) \subset U$;
- We have $R^{-1}(c) \subset S_!(U)$;

are equivalent. Thus $(S \diamond R)_! = S_! \circ R_!$. \square

5.5.5 Functoriality of Powersets

00ML Proposition 5.5.5.1.1. The assignment $X \mapsto \mathcal{P}(X)$ defines functors⁶⁹

$$\begin{aligned}\mathcal{P}_*: \text{Rel} &\rightarrow \text{Sets}, \\ \mathcal{P}_{-1}: \text{Rel}^{\text{op}} &\rightarrow \text{Sets}, \\ \mathcal{P}^{-1}: \text{Rel}^{\text{op}} &\rightarrow \text{Sets}, \\ \mathcal{P}_!: \text{Rel} &\rightarrow \text{Sets}\end{aligned}$$

where

- *Action on Objects.* For each $A \in \text{Obj}(\text{Rel})$, we have

$$\begin{aligned}\mathcal{P}_*(A) &\stackrel{\text{def}}{=} \mathcal{P}(A), \\ \mathcal{P}_{-1}(A) &\stackrel{\text{def}}{=} \mathcal{P}(A), \\ \mathcal{P}^{-1}(A) &\stackrel{\text{def}}{=} \mathcal{P}(A), \\ \mathcal{P}_!(A) &\stackrel{\text{def}}{=} \mathcal{P}(A);\end{aligned}$$

- *Action on Morphisms.* For each morphism $R: A \rightarrow B$ of Rel, the images

$$\begin{aligned}\mathcal{P}_*(R): \mathcal{P}(A) &\rightarrow \mathcal{P}(B), \\ \mathcal{P}_{-1}(R): \mathcal{P}(B) &\rightarrow \mathcal{P}(A), \\ \mathcal{P}^{-1}(R): \mathcal{P}(B) &\rightarrow \mathcal{P}(A), \\ \mathcal{P}_!(R): \mathcal{P}(A) &\rightarrow \mathcal{P}(B)\end{aligned}$$

of R by \mathcal{P}_* , \mathcal{P}_{-1} , \mathcal{P}^{-1} , and $\mathcal{P}_!$ are defined by

$$\begin{aligned}\mathcal{P}_*(R) &\stackrel{\text{def}}{=} R_*, \\ \mathcal{P}_{-1}(R) &\stackrel{\text{def}}{=} R_{-1}, \\ \mathcal{P}^{-1}(R) &\stackrel{\text{def}}{=} R^{-1}, \\ \mathcal{P}_!(R) &\stackrel{\text{def}}{=} R_!,\end{aligned}$$

as in Definitions 5.5.1.1.1, 5.5.2.1.1, 5.5.3.1.1 and 5.5.4.1.1.

Proof. This follows from Items 3 and 4 of Proposition 5.5.1.1.4, Items 3 and 4 of Proposition 5.5.2.1.4, Items 3 and 4 of Proposition 5.5.3.1.4, and Items 3 and 4 of Proposition 5.5.4.1.4. \square

5.5.6 Functoriality of Powersets: Relations on Powersets

00MM Let A and B be sets and let $R: A \rightarrow B$ be a relation.

⁶⁹The functor $\mathcal{P}_*: \text{Rel} \rightarrow \text{Sets}$ admits a left adjoint; see Item 3 of Proposition 5.3.1.1.2.

00MN Definition 5.5.6.1.1. The **relation on powersets associated to R** is the relation

$$\mathcal{P}(R): \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by⁷⁰

$$\mathcal{P}(R)_U^V \stackrel{\text{def}}{=} \mathbf{Rel}(\chi_{\text{pt}}, V \diamond R \diamond U)$$

for each $U \in \mathcal{P}(A)$ and each $V \in \mathcal{P}(B)$.

00MP Remark 5.5.6.1.2. In detail, we have $U \sim_{\mathcal{P}(R)} V$ iff the following equivalent conditions hold:

- We have $\chi_{\text{pt}} \subset V \diamond R \diamond U$.
- We have $(V \diamond R \diamond U)_{\star}^{\star} = \text{true}$, i.e. we have

$$\int^{a \in A} \int^{b \in B} V_b^{\star} \times R_a^b \times U_{\star}^a = \text{true}.$$

- There exists some $a \in A$ and some $b \in B$ such that:
 - We have $U_{\star}^a = \text{true}$;
 - We have $R_a^b = \text{true}$;
 - We have $V_b^{\star} = \text{true}$.
- There exists some $a \in A$ and some $b \in B$ such that:
 - We have $a \in U$;
 - We have $a \sim_R b$;
 - We have $b \in V$.

00MQ Proposition 5.5.6.1.3. The assignment $R \mapsto \mathcal{P}(R)$ defines a functor

$$\mathcal{P}: \mathbf{Rel} \rightarrow \mathbf{Rel}.$$


Proof. Omitted. □

5.6 Relative Preorders

5.6.1 The Left Skew Monoidal Structure on $\mathbf{Rel}(A, B)$

Let A and B be sets and let $J: A \rightarrow B$ be a relation.

⁷⁰Illustration:



5.6.1.1 The Left Skew Monoidal Product

00MU Definition 5.6.1.1.1. The **left J -skew monoidal product** of $\mathbf{Rel}(A, B)$ is the functor

$$\triangleleft_J : \mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(A, B)$$

where

- *Action on Objects.* For each $R, S \in \text{Obj}(\mathbf{Rel}(A, B))$, we have

$$S \triangleleft_J R \stackrel{\text{def}}{=} S \diamond \text{Rift}_J(R), \quad \begin{array}{ccc} A & \xrightarrow{S} & B; \\ \text{Rift}_J(R) \swarrow & \nearrow & \downarrow J \\ A & \xrightarrow{R} & B \end{array}$$

- *Action on Morphisms.* For each $R, S, R', S' \in \text{Obj}(\mathbf{Rel}(A, B))$, the action on Hom-sets

$$(\triangleleft_J)_{(G,F),(G',F')} : \text{Hom}_{\mathbf{Rel}(A,B)}(S, S') \times \text{Hom}_{\mathbf{Rel}(A,B)}(R, R') \rightarrow \text{Hom}_{\mathbf{Rel}(A,B)}(S \triangleleft_J R, S' \triangleleft_J R')$$

of \triangleleft_J at $((R, S), (R', S'))$ is defined by⁷¹

$$\beta \triangleleft_J \alpha \stackrel{\text{def}}{=} \beta \diamond \text{Rift}_J(\alpha), \quad \begin{array}{ccc} & S & \\ \text{Rift}_J(R) \swarrow & \nearrow & \downarrow \beta \\ A & \xrightarrow{\alpha} & B \\ \text{Rift}_J(\alpha) \swarrow & \nearrow & \downarrow S' \\ A & \xrightarrow{R} & B \\ \text{Rift}_J(R') \swarrow & \nearrow & \downarrow \alpha \\ A & \xrightarrow{R'} & B \end{array}$$

for each $\beta \in \text{Hom}_{\mathbf{Rel}(A,B)}(S, S')$ and each $\alpha \in \text{Hom}_{\mathbf{Rel}(A,B)}(R, R')$.

5.6.1.2 The Left Skew Monoidal Unit

00MW Definition 5.6.1.2.1. The **left J -skew monoidal unit** of $\mathbf{Rel}(A, B)$ is the functor

$$\mathbb{M}_{\triangleleft}^{\mathbf{Rel}(A,B)} : \text{pt} \rightarrow \mathbf{Rel}(A, B)$$

picking the object

$$\mathbb{M}_{\mathbf{Rel}(A,B)}^{\triangleleft} \stackrel{\text{def}}{=} J$$

of $\mathbf{Rel}(A, B)$.

⁷¹Since $\mathbf{Rel}(A, B)$ is posetal, this is to say that if $S \subset S'$ and $R \subset R'$, then $S \triangleleft_J R \subset S' \triangleleft_J R'$.

5.6.1.3 The Left Skew Associators

00MY Definition 5.6.1.3.1. The **left J -skew associator of $\mathbf{Rel}(A, B)$** is the natural transformation

$$\alpha^{\mathbf{Rel}(A, B), \triangleleft} : \triangleleft_J \circ (\triangleleft_J \times \text{id}) \Longrightarrow \triangleleft_J \circ (\text{id} \times \triangleleft_J),$$

whose component

$$\alpha_{T, S, R}^{\mathbf{Rel}(A, B), \triangleleft} : \underbrace{(T \triangleleft_J S) \triangleleft_J R}_{\stackrel{\text{def}}{=} T \diamond \text{Rift}_J(S) \diamond \text{Rift}_J(R)} \hookrightarrow \underbrace{T \triangleleft_J (S \triangleleft_J R)}_{\stackrel{\text{def}}{=} T \diamond \text{Rift}_J(S \diamond \text{Rift}_J(R))}$$

at (T, S, R) is given by

$$\alpha_{T, S, R}^{\mathbf{Rel}(A, B), \triangleleft} \stackrel{\text{def}}{=} \text{id}_T \diamond \gamma,$$

where

$$\gamma : \text{Rift}_J(S) \diamond \text{Rift}_J(R) \hookrightarrow \text{Rift}_J(S \diamond \text{Rift}_J(R))$$

is the inclusion adjunct to the inclusion

$$\epsilon_S \star \text{id}_{\text{Rift}_J(R)} : \underbrace{J \diamond \text{Rift}_J(S) \diamond \text{Rift}_J(R)}_{\stackrel{\text{def}}{=} J_*(\text{Rift}_J(S) \diamond \text{Rift}_J(R))} \hookrightarrow S \diamond \text{Rift}_J(R)$$

under the adjunction $J_* \dashv \text{Rift}_J$, where $\epsilon : J \diamond \text{Rift}_J \Longrightarrow \text{id}_{\mathbf{Rel}(A, B)}$ is the counit of the adjunction $J_* \dashv \text{Rift}_J$.

5.6.1.4 The Left Skew Left Unitors

00NO Definition 5.6.1.4.1. The **left J -skew left unit of $\mathbf{Rel}(A, B)$** is the natural transformation

$$\lambda^{\mathbf{Rel}(A, B), \triangleleft} : \triangleleft_J \circ (\mathbb{1}_{\triangleleft}^{\mathbf{Rel}(A, B)} \times \text{id}) \Longrightarrow \text{id},$$

whose component

$$\lambda_R^{\mathbf{Rel}(A, B), \triangleleft} : \underbrace{J \triangleleft_J R}_{\stackrel{\text{def}}{=} J \diamond \text{Rift}_J(R)} \hookrightarrow R$$

at R is given by

$$\lambda_R^{\mathbf{Rel}(A, B), \triangleleft} \stackrel{\text{def}}{=} \epsilon_R,$$

where $\epsilon : J \diamond \text{Rift}_J \Longrightarrow \text{id}_{\mathbf{Rel}(A, B)}$ is the counit of the adjunction $J_* \dashv \text{Rift}_J$.

5.6.1.5 The Left Skew Right Unitors

00N2 Definition 5.6.1.5.1. The **left J -skew right unit of $\mathbf{Rel}(A, B)$** is the natural transformation

$$\rho^{\mathbf{Rel}(A, B), \triangleleft} : \text{id} \Longrightarrow \triangleleft_J \circ (\text{id} \times \mathbb{1}_{\triangleleft}^{\mathbf{Rel}(A, B)}),$$

whose component

$$\rho_R^{\mathbf{Rel}(A,B), \triangleleft} : R \hookrightarrow \underbrace{R \triangleleft_J J}_{\stackrel{\text{def}}{=} R \diamond \text{Rift}_J(J)}$$

at R is given by

$$\rho_R^{\mathbf{Rel}(A,B), \triangleleft} \stackrel{\text{def}}{=} \text{id}_R \star \sigma,$$

where $\sigma : \text{id}_A \implies \text{Rift}_J(J)$ is the universal transformation included in the data of the right Kan lift $\text{Rift}_J(J)$.

5.6.1.6 The Left Skew Monoidal Structure on $\mathbf{Rel}(A, B)$

00N4 Definition 5.6.1.6.1. The left J -skew monoidal category of relations from A to B is the left skew monoidal category

$$\left(\mathbf{Rel}(A, B), \triangleleft_J, \mu_{\triangleleft}^{\mathbf{Rel}(A, B)}, \alpha^{\mathbf{Rel}(A, B), \triangleleft}, \lambda^{\mathbf{Rel}(A, B), \triangleleft}, \rho^{\mathbf{Rel}(A, B), \triangleleft} \right)$$

consisting of

- *The Underlying Category.* The posetal category associated to the poset $\mathbf{Rel}(A, B)$ of relations from A to B of Item 2 of Definition 5.1.1.2;
- *The Skew Monoidal Product.* The functor \triangleleft_J of Definition 5.6.1.1.1;
- *The Skew Monoidal Unit.* The functor $\mu_{\triangleleft}^{\mathbf{Rel}(A, B)}$ of Definition 5.6.1.2.1;
- *The Skew Associators.* The natural transformation $\alpha^{\mathbf{Rel}(A, B), \triangleleft}$ of Definition 5.6.1.3.1;
- *The Skew Left Unitors.* The natural transformation $\lambda^{\mathbf{Rel}(A, B), \triangleleft}$ of Definition 5.6.1.4.1;
- *The Skew Right Unitors.* The natural transformation $\rho^{\mathbf{Rel}(A, B), \triangleleft}$ of Definition 5.6.1.5.1.

5.6.2 Left Relative Preorders

Let A and B be sets and let $J : A \nrightarrow B$ be a relation.

00N6 Definition 5.6.2.1.1. A left J -relative preorder from A to B is equivalently:

- An \mathbb{E}_1 -skew monoid in $(\mathbf{N}_*(\mathbf{Rel}(A, B)), \triangleleft_J, J)$;
- A skew monoid in $(\mathbf{Rel}(A, B), \triangleleft_J, J)$.

00N7 Remark 5.6.2.1.2. In detail, a left J -relative preorder (R, μ_R, η_R) from A to B consists of

- *The Underlying Relation.* A relation

$$R : A \nrightarrow B,$$

called the **underlying relation of** (R, μ_R, η_R) ;

- *The Multiplication Inclusion.* An inclusion of relations

$$\mu_R: R \triangleleft_J R \subset R,$$

called the **multiplication** of (R, μ_R, η_R) ;

- *The Unit Inclusion.* An inclusion of relations

$$\eta_R: J \subset R,$$

called the **unit** of (R, μ_R, η_R) .

00N8 Remark 5.6.2.1.3. In other words, a **left J -relative preorder from A to B** is a relation $R: A \rightarrow B$ from A to B satisfying the following conditions:

1. *J -Transitivity.* For each $a \in A$ and each $c \in B$, we have

$$a \sim_{R \circ \text{Rift}_J(R)} c$$

i.e. the following condition is satisfied:⁷²

- (★) If there exists some $b \in A$ such that:


- We have $a \sim_{\text{Rift}_J(R)} b$, i.e. for each $x \in B$, if $b \sim_J x$, then $a \sim_R x$;⁷³
- We have $b \sim_R c$;

then $a \sim_R c$.

2. *J -Unitality.* For each $a \in A$ and each $b \in B$, the following condition is satisfied:


- (★) If $a \sim_J b$, then $a \sim_R b$.

⁷²Illustration: If we have



then $a \sim_R c$.

⁷³Illustration:



5.6.3 The Right Skew Monoidal Structure on $\mathbf{Rel}(A, B)$

Let A and B be sets and let $J: A \rightarrow B$ be a relation.

5.6.3.1 The Right Skew Monoidal Product

00NB Definition 5.6.3.1.1. The **right J -skew monoidal product of $\mathbf{Rel}(A, B)$** is the functor

$$\triangleright_J: \mathbf{Rel}(A, B) \times \mathbf{Rel}(A, B) \rightarrow \mathbf{Rel}(A, B)$$

where

- *Action on Objects.* For each $R, S \in \text{Obj}(\mathbf{Rel}(A, B))$, we have

$$S \triangleright_J R \stackrel{\text{def}}{=} \text{Ran}_J(S) \diamond R,$$

- *Action on Morphisms.* For each $R, S, R', S' \in \text{Obj}(\mathbf{Rel}(A, B))$, the action on Hom-sets

$$(\triangleright_J)_{(S, R), (S', R')} : \text{Hom}_{\mathbf{Rel}(A, B)}(S, S') \times \text{Hom}_{\mathbf{Rel}(A, B)}(R, R') \rightarrow \text{Hom}_{\mathbf{Rel}(A, B)}(S \triangleright_J R, S' \triangleright_J R')$$

of \triangleright_J at $((S, R), (S', R'))$ is defined by⁷⁴

$$\beta \triangleright_J \alpha \stackrel{\text{def}}{=} \text{Ran}_J(\beta) \diamond \alpha,$$

for each $\beta \in \text{Hom}_{\mathbf{Rel}(A, B)}(S, S')$ and each $\alpha \in \text{Hom}_{\mathbf{Rel}(A, B)}(R, R')$.

5.6.3.2 The Right Skew Monoidal Unit

00ND Definition 5.6.3.2.1. The **right J -skew monoidal unit of $\mathbf{Rel}(A, B)$** is the functor

$$\mathbb{1}^{\mathbf{Rel}(A, B)}: \text{pt} \rightarrow \mathbf{Rel}(A, B)$$

picking the object

$$\mathbb{1}^{\triangleright}_{\mathbf{Rel}(A, B)} \stackrel{\text{def}}{=} J$$

of $\mathbf{Rel}(A, B)$.

⁷⁴Since $\mathbf{Rel}(A, B)$ is posetal, this is to say that if $S \subset S'$ and $R \subset R'$, then $S \triangleright_J R \subset S' \triangleright_J R'$.

5.6.3.3 The Right Skew Associators

00NF **Definition 5.6.3.3.1.** The **right J -skew associator of $\mathbf{Rel}(A, B)$** is the natural transformation

$$\alpha^{\mathbf{Rel}(A, B), \triangleright} : \triangleright_J \circ (\text{id} \times \triangleright_J) \Longrightarrow \triangleright_J \circ (\triangleright_J \times \text{id}),$$

whose component

$$\alpha_{T, S, R}^{\mathbf{Rel}(A, B), \triangleright} : \underbrace{T \triangleright_J (S \triangleright_J R)}_{\stackrel{\text{def}}{=} \text{Ran}_J(T) \diamond (\text{Ran}_J(S) \diamond R)} \hookrightarrow \underbrace{(\overline{T \triangleright_J S}) \triangleright_J R}_{\stackrel{\text{def}}{=} \text{Ran}_J(\text{Ran}_J(T) \diamond S) \diamond R}$$

at (T, S, R) is given by

$$\alpha_{T, S, R}^{\mathbf{Rel}(A, B), \triangleright} \stackrel{\text{def}}{=} \gamma \diamond \text{id}_R,$$

where

$$\gamma : \text{Ran}_J(T) \diamond \text{Ran}_J(S) \hookrightarrow \text{Ran}_J(\text{Ran}_J(T) \diamond S)$$

is the inclusion adjunct to the inclusion

$$\text{id}_{\text{Ran}_J(T)} \diamond \epsilon_S : \underbrace{\text{Ran}_J(T) \diamond \text{Ran}_J(S) \diamond J}_{\stackrel{\text{def}}{=} f^*(\text{Ran}_J(T) \diamond \text{Ran}_J(S))} \hookrightarrow \text{Ran}_J(T) \diamond S$$

under the adjunction $J^* \dashv \text{Ran}_J$, where $\epsilon : \text{Ran}_J \diamond J \Longrightarrow \text{id}_{\mathbf{Rel}(A, B)}$ is the counit of the adjunction $J^* \dashv \text{Ran}_J$.

5.6.3.4 The Right Skew Left Unitors

00NH **Definition 5.6.3.4.1.** The **right J -skew left unitor of $\mathbf{Rel}(A, B)$** is the natural transformation

$$\lambda^{\mathbf{Rel}(A, B), \triangleright} : \text{id} \Longrightarrow \triangleright_J \circ (\wp_{\triangleright}^{\mathbf{Rel}(A, B)} \times \text{id}),$$

whose component

$$\lambda_R^{\mathbf{Rel}(A, B), \triangleright} : R \hookrightarrow \underbrace{J \triangleright_J R}_{\stackrel{\text{def}}{=} \text{Ran}_J(J) \diamond R}$$

at R is given by

$$\lambda_R^{\mathbf{Rel}(A, B), \triangleright} \stackrel{\text{def}}{=} \sigma \diamond \text{id}_R,$$

where $\sigma : \text{id}_B \Longrightarrow \text{Ran}_J(J)$ is the universal transformation included in the data of the right Kan extension $\text{Ran}_J(J)$.

5.6.3.5 The Right Skew Right Unitors

00NK **Definition 5.6.3.5.1.** The **right J -skew right unitor of $\mathbf{Rel}(A, B)$** is the natural transformation

$$\rho^{\mathbf{Rel}(A, B), \triangleright} : \triangleright_J \circ (\text{id} \times \wp_{\triangleright}^{\mathbf{Rel}(A, B)}) \Longrightarrow \text{id},$$

whose component

$$\rho_S^{\mathbf{Rel}(A,B),\triangleright} : \underbrace{S \triangleright_J J}_{\stackrel{\text{def}}{=} \text{Ran}_J(S) \diamond J} \hookrightarrow S$$

at S is given by

$$\rho_S^{\mathbf{Rel}(A,B),\triangleright} \stackrel{\text{def}}{=} \epsilon_R,$$

where $\epsilon : \text{Ran}_J \diamond J \implies \text{id}_{\mathbf{Rel}(A,B)}$ is the counit of the adjunction $J^* \dashv \text{Ran}_J$.

5.6.3.6 The Right Skew Monoidal Structure on $\mathbf{Rel}(A, B)$

00NM Definition 5.6.3.6.1. The **right J -skew monoidal category of functors from A to B** is the right skew monoidal category

$$\left(\mathbf{Rel}(A, B), \triangleright_J, \mathbb{1}_\triangleright^{\mathbf{Rel}(A,B)}, \alpha^{\mathbf{Rel}(A,B),\triangleright}, \lambda^{\mathbf{Rel}(A,B),\triangleright}, \rho^{\mathbf{Rel}(A,B),\triangleright} \right)$$

consisting of

- *The Underlying Category.* The posetal category associated to the poset $\mathbf{Rel}(A, B)$ of relations from A to B of Item 2 of Definition 5.1.1.2;
- *The Skew Monoidal Product.* The functor \triangleright_J of Definition 5.6.3.1.1;
- *The Skew Monoidal Unit.* The functor $\mathbb{1}_\triangleright^{\mathbf{Rel}(A,B)}$ of Definition 5.6.3.2.1;
- *The Skew Associators.* The natural transformation $\alpha^{\mathbf{Rel}(A,B),\triangleright}$ of Definition 5.6.3.3.1;
- *The Skew Left Unitors.* The natural transformation $\lambda^{\mathbf{Rel}(A,B),\triangleright}$ of Definition 5.6.3.4.1;
- *The Skew Right Unitors.* The natural transformation $\rho^{\mathbf{Rel}(A,B),\triangleright}$ of Definition 5.6.3.5.1.

5.6.4 Right Relative Preorders

00NN Let A and B be sets and let $J : A \rightarrow B$ be a relation.

00NP Definition 5.6.4.1.1. A **right J -relative preorder from A to B** is equivalently:

- An \mathbb{E}_1 -skew monoid in $(N_\bullet(\mathbf{Rel}(A, B)), \triangleright_J, J)$;
- A skew monoid in $(\mathbf{Rel}(A, B), \triangleright_J, J)$.

00NQ Remark 5.6.4.1.2. In detail, a **right J -relative preorder** (R, μ_R, η_R) from A to B consists of

- *The Underlying Relation.* A relation

$$R : A \rightarrow B,$$

called the **underlying relation** of (R, μ_R, η_R) ;

- *The Multiplication Inclusion.* An inclusion of relations

$$\mu_R: R \triangleright_J R \subset R,$$

called the **multiplication** of (R, μ_R, η_R) ;

- *The Unit Inclusion.* An inclusion of relations

$$\eta_R: J \subset R,$$

called the **unit** of (R, μ_R, η_R) .

00NR Remark 5.6.4.1.3. In other words, a **right J -relative preorder from A to B** is a relation $R: A \rightarrow B$ from A to B satisfying the following conditions:

1. *J -Transitivity.* For each $a \in A$ and each $c \in B$, we have

$$a \sim_{\text{Ran}_J(R) \diamond R} c,$$

i.e. the following condition is satisfied:⁷⁵


- (★) If there exists some $b \in B$ such that:

- We have $a \sim_R b$;
 - We have $b \sim_{\text{Ran}_J(R)} c$, i.e. for each $x \in A$, if $x \sim_J b$, then $x \sim_R c$;⁷⁶
- then $a \sim_R c$.

2. *J -Unitarity.* For each $a \in A$ and each $b \in B$, the following condition is satisfied:


- (★) If $a \sim_J b$, then $a \sim_R b$.

⁷⁵Illustration: If we have



then $a \sim_R c$.

⁷⁶Illustration:



Appendices

5.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets
5. Relations
6. Spans
7. Posets

Indexed and Fibred Sets

7. Indexed Sets
8. Fibred Sets
9. Un/Straightening for Indexed and Fibred Sets

Category Theory

11. Categories
12. Types of Morphisms in Categories
13. Adjunctions and the Yoneda Lemma
14. Constructions With Categories
15. Profunctors
16. Cartesian Closed Categories
17. Kan Extensions

Bicategories

18. Bicategories
19. Internal Adjunctions

Internal Category Theory

- 20. Internal Categories
- 33. Hypersemirings and Hyperrings
- Cyclic Stuff**
- 21. The Cycle Category
- 34. Quantales
- Cubical Stuff**
- 22. The Cube Category
- 35. Near-Semirings
- Globular Stuff**
- 23. The Globe Category
- 36. Near-Rings
- Cellular Stuff**
- 24. The Cell Category
- Real Analysis**
- 37. Real Analysis in One Variable
- 38. Real Analysis in Several Variables
- Monoids**
- 25. Monoids
- 39. Measurable Spaces
- 26. Constructions With Monoids
- 40. Measures and Integration
- Monoids With Zero**
- 27. Monoids With Zero
- 40. Probability Theory
- 28. Constructions With Monoids With Zero
- Stochastic Analysis**
- 41. Stochastic Processes, Martingales, and Brownian Motion
- 42. Itô Calculus
- 43. Stochastic Differential Equations
- Groups**
- 29. Groups
- 30. Constructions With Groups
- Differential Geometry**
- 44. Topological and Smooth Manifolds
- Hyper Algebra**
- 31. Hypermonoids
- 32. Hypergroups
- Schemes**
- 45. Schemes

5.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
Category Theory	27. Monoids With Zero
11. Categories	28. Constructions With Monoids With Zero
12. Types of Morphisms in Categories	Groups
13. Adjunctions and the Yoneda Lemma	29. Groups
14. Constructions With Categories	30. Constructions With Groups
15. Profunctors	Hyper Algebra
16. Cartesian Closed Categories	31. Hypermonoids
17. Kan Extensions	32. Hypergroups
Bicategories	33. Hypersemirings and Hyperrings
18. Bicategories	34. Quantales
19. Internal Adjunctions	Near-Rings
Internal Category Theory	35. Near-Semirings
20. Internal Categories	36. Near-Rings
	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 6

Spans

00NS This chapter contains some material about spans. Notably, we discuss and explore:

1. The basic definitions around spans ([Section 6.1](#));
2. The relation between spans and functions ([Proposition 6.8.1.1.1](#));
3. The relation between spans and relations ([Propositions 6.8.2.2.1](#) and [6.8.3.1.1](#) and [Remark 6.8.5.1.1](#)).
4. “Hyperpointed sets” (??). I don’t know why I wrote this...

TODO:

1. https://www.sciencedirect.com/science/article/pii/0022404994900094?ref=pdf_download&fr=RR-2&rr=834107b75c906aa4
2. <https://arxiv.org/abs/1605.08100>
3. <https://arxiv.org/abs/1603.08181>
4. <https://arxiv.org/abs/1601.02307>
5. <https://arxiv.org/abs/1507.01460>
6. <https://arxiv.org/abs/1506.08870>
7. <https://arxiv.org/abs/1505.00048>
8. <https://arxiv.org/abs/1501.07592>
9. <https://arxiv.org/abs/1501.04664>
10. <https://arxiv.org/abs/1501.00792>
11. <https://arxiv.org/abs/1412.6560>
12. <https://arxiv.org/abs/1412.0212>

13. <https://arxiv.org/abs/1409.0837>
14. <https://arxiv.org/abs/1408.5220>
15. <https://arxiv.org/abs/1308.6548>
16. <https://arxiv.org/abs/1304.0219>
17. <https://arxiv.org/abs/1210.8192>
18. <https://arxiv.org/abs/1210.1433>
19. <https://arxiv.org/abs/1201.3789>
20. <https://arxiv.org/abs/1112.0560>
21. <https://arxiv.org/abs/1109.1598>
22. <https://arxiv.org/abs/1101.4594>
23. <https://arxiv.org/abs/1012.6001>
24. <https://arxiv.org/abs/1011.3243>
25. <https://arxiv.org/abs/0910.2996>
26. <https://arxiv.org/abs/0810.2361>
27. <https://arxiv.org/abs/0803.2429>
28. <https://arxiv.org/abs/0712.2525>
29. <https://arxiv.org/abs/0706.1286>
30. <https://arxiv.org/abs/math/0611930>
31. <https://arxiv.org/abs/2311.15342>
32. <https://arxiv.org/abs/2310.19428>
33. <https://arxiv.org/abs/2309.08084>
34. <https://arxiv.org/abs/2308.01662>
35. <https://arxiv.org/abs/2301.11860>
36. <https://arxiv.org/abs/2301.01199>
37. <https://arxiv.org/abs/2212.09060>
38. <https://arxiv.org/abs/2208.07183>
39. <https://arxiv.org/abs/2205.06892>

40. <https://arxiv.org/abs/2203.16179>
41. <https://arxiv.org/abs/2201.09551>
42. <https://arxiv.org/abs/2112.04599>
43. <https://arxiv.org/abs/2111.10968>
44. <https://arxiv.org/abs/2107.07621>
45. <https://arxiv.org/abs/2106.14743>
46. <https://arxiv.org/abs/2105.14654>
47. <https://arxiv.org/abs/2102.08051>
48. <https://arxiv.org/abs/2102.04386>
49. <https://arxiv.org/abs/2101.06734>
50. <https://arxiv.org/abs/2011.11042>
51. <https://arxiv.org/abs/2010.15722>
52. <https://arxiv.org/abs/2006.10375>
53. <https://arxiv.org/abs/2006.10375>
54. <https://arxiv.org/abs/2005.10496>
55. <https://arxiv.org/abs/2003.11541>
56. <https://arxiv.org/abs/2002.10334>
57. <https://arxiv.org/abs/1909.00069>
58. <https://arxiv.org/abs/1907.02695>
59. <https://arxiv.org/abs/1905.06671>
60. define a relational span
61. consider giving Ran and Rift their dedicated sections on the relations chapter,
perhaps together with the other sections on co/limits
62. <https://arxiv.org/abs/1710.02742>
63. <https://arxiv.org/search/math?searchtype=author&query=Walker,+Charles>
64. <https://arxiv.org/abs/1706.09575>
65. <https://arxiv.org/abs/1710.01465>

-
- 66. fibred categories: <https://arxiv.org/abs/1806.02376>
 - 67. <https://arxiv.org/abs/1806.10477v2>
 - 68. double categorical limits in $\mathbf{Rel}^{\mathrm{dbl}}$
 - 69. double categorical limits in $\mathbf{Span}^{\mathrm{dbl}}$
 - 70. internal adjoint equivalences in **Rel**
 - 71. internal adjoint equivalences in \mathbf{Span}
 - 72. 2-categorical limits in **Rel**;
 - 73. morphism of internal adjunctions in **Rel**;
 - 74. morphism of internal adjunctions in \mathbf{Span} ;
 - 75. morphism of co/monads in \mathbf{Span} ;
 - 76. What is $\mathrm{Adj}(\mathbf{Span}(A, B))$?
 - 77. monoids, comonoids, pseudomonoids, etc. in \mathbf{Span} .
 - 78. write down the dumb intuition about spans inducing morphisms $\mathrm{Sets}(S, A) \rightarrow \mathrm{Sets}(S, B)$ instead of $\mathcal{P}(A) \rightarrow \mathcal{P}(B)$ from the similarity between

$$S \rightarrow A \times B$$

and

$$A \times B \rightarrow \{t, f\}.$$

This intuition is justified by taking $A = \mathrm{pt}$ or $B = \mathrm{pt}$.

- 79. What about using the direct image with compact support in $g(f^{-1}(a))$?
- 80. Monads in \mathbf{Span} | develop this in the level of morphisms too
- 81. Comonads in \mathbf{Span} are spans whose legs are equal | develop this in the level of morphisms too
- 82. Does \mathbf{Span} have an internal **Hom**?
- 83. Examples of spans
- 84. Functional and total spans
- 85. closed symmetric monoidal category of spans
- 86. double category of relations
- 87. collage of a span

-
88. equivalence spans?
89. functoriality of powersets for spans
90. Is Span a closed bicategory?
91. skew monoidal structure on $\text{Span}(A, B)$
92. Adjunctions in Span
93. Isomorphisms in Span
94. Equivalences in Span
95. Interaction between the above notions in Span vs. in **Rel** via the comparison functors
96. $\text{Hom}_C(S, A) \times \text{Hom}_C(f^*(S), A)$.
97. Proof of non-existence of left Kan extensions/lifts in **Rel** (when do these exist btw?)
98. description of unitors and associators of span
99. add intuition for spans as relations with multiple witnesses:

(a) Given a span $A \xleftarrow{f} S \xrightarrow{g} B$, we have a functor

$$\text{St}(S) : (A \times B)_{\text{disc}} \rightarrow \text{Sets}$$

given by

$$\begin{aligned} [\text{St}(S)](a, b) &\stackrel{\text{def}}{=} \text{St}(S)_{ab} \\ &\stackrel{\text{def}}{=} \{s \in S \mid f(s) = a \text{ and } g(s) = b\}. \end{aligned}$$

(b) Given a functor

$$S : (A \times B)_{\text{disc}} \rightarrow \text{Sets},$$

we have a map of sets

$$\text{Un}(S) : \coprod_{(a,b) \in A \times B} S(a, b) \rightarrow A \times B,$$

determining a span from A to B .

- (c) How do these interact with left/right Kan extensions/lifts?
- (d) Un/straightening for spans of categories: assignment $(a, b) \mapsto \text{Wits}_S(a, b)$.
- (e) Fix the TODO below

Contents


6.1	Spans	222
6.1.1	The Walking Span	222
6.1.2	Spans	222
6.1.3	Morphisms of Spans	223
6.1.4	Functional Spans.....	223
6.1.5	Total Spans	225
6.2	Categories of Spans	225
6.2.1	The Category of Spans Between Two Sets	225
6.2.2	The Bicategory of Spans.....	227
6.2.3	The Monoidal Bicategory of Spans.....	229
6.2.4	The Double Category of Spans.....	229
6.2.5	Properties of The Bicategory of Spans	233
6.3	Limits of Spans.....	236
6.3.1	tmp2.....	236
6.3.2	tmp	236
6.3.3	Left Kan Extensions	236
6.3.4	Right Kan Extensions.....	237
6.3.5	Right Kan Lifts	238
6.4	Colimits of Spans	239
6.5	Constructions With Spans	239
6.5.1	Representable Spans.....	239
6.5.2	Composition of Spans.....	239
6.5.3	Horizontal Composition of Morphisms of Spans.....	240
6.5.4	Properties of Composition of Spans	240
6.5.5	The Inverse of a Span	241
6.6	Functionality of Spans	241
6.6.1	Direct Images.....	241
6.6.2	Functionality of Spans on Powersets	241
6.7	Un/Straightening for Spans.....	241
6.7.1	Straightening for Spans	241
6.7.2	Unstraightening for Spans.....	242
6.7.3	The Un/Straightening Equivalence for Spans.....	243
6.8	Comparison of Spans to Functions and Relations.....	243
6.8.1	Comparison to Functions	243
6.8.2	Comparison to Relations: From Span to Rel.....	245
6.8.3	Comparison to Relations: From Rel to Span.....	247
6.8.4	Comparison to Relations: The Wehrheim–Woodward Construction	249

6.8.5	Comparison to Multirelations	249
6.8.6	Comparison to Relations via Double Categories	249
6.A	Other Chapters	250
6.2	Other Chapters	251

6.1 Spans

6.1.1 The Walking Span

00NV Definition 6.1.1.1. The **walking span** is the category Λ that looks like this:



6.1.2 Spans

Let A and B be sets.

00NX Definition 6.1.2.1.1. A **span from A to B** ¹ is a functor $F: \Lambda \rightarrow \text{Sets}$ such that


$$\begin{aligned} F([-1]) &= A, \\ F([1]) &= B. \end{aligned}$$

00NY Remark 6.1.2.1.2. In detail, a **span from A to B** is a triple (S, f, g) consisting of^{2,3}

- *The Underlying Set.* A set S , called the **underlying set of (S, f, g)** ;
- *The Legs.* A pair of functions $f: S \rightarrow A$ and $g: S \rightarrow B$.

¹Further Terminology: Also called a **roof from A to B** or a **correspondence from A to B** .

²Picture:



³Every span (S, f, g) from A to B determines in particular a relation $R: A \nrightarrow B$ via

$$R \stackrel{\text{def}}{=} \{(f(a), g(a)) \mid a \in A\},$$

i.e. where $R(a) = g(f^{-1}(a))$ for each $a \in A$; see [Proposition 6.8.2.2.1](#).

00NZ Remark 6.1.2.1.3. A span

$$\begin{array}{ccc} & S & \\ f \swarrow & & \searrow g \\ A & & B. \end{array}$$

from A to B may be thought of as a relation from A to B which can relate an element $a \in A$ to an element $b \in B$ in multiple ways via f and g , with the “set of witnesses of $a \sim_S b$ ” being given by

$$\text{Wits}_S(a, b) \stackrel{\text{def}}{=} \{s \in S \mid a = f(s) \text{ and } g(s) = b\}.$$

This analogy is made precise by [Remark 6.8.5.1.1](#) and [Section 6.7](#).

6.1.3 Morphisms of Spans

00P1 Definition 6.1.3.1.1. A **morphism of spans** from (R, f_1, g_1) to (S, f_2, g_2) ⁴ is a natural transformation $(R, f_1, g_1) \Rightarrow (S, f_2, g_2)$.

00P2 Remark 6.1.3.1.2. In detail, a **morphism of spans** from (R, f_1, g_1) to (S, f_2, g_2) is a function $\phi: R \rightarrow S$ making the diagram⁵

$$\begin{array}{ccccc} & R & & S & \\ & \swarrow f_1 & \downarrow \phi & \searrow g_2 & \\ A & \xlongequal{\quad} & B & \xlongequal{\quad} & A \\ & \searrow f_2 & & \uparrow & \swarrow g_1 \\ & & B & & \end{array}$$

commute.

6.1.4 Functional Spans

Let $\lambda = \left(A \xleftarrow{f} S \xrightarrow{g} B \right)$ be a span.

00P4 Definition 6.1.4.1.1. The span λ is **functional** if it satisfies the following equivalent conditions:

⁴Further Terminology: Also called a **morphism of roofs** from (R, f_1, g_1) to (S, f_2, g_2) or a **morphism of correspondences** from (R, f_1, g_1) to (S, f_2, g_2) .

⁵Alternative Picture:

$$\begin{array}{ccccc} & R & & S & \\ & \swarrow f_1 & \downarrow \phi & \searrow g_1 & \\ A & & B & & \\ & \uparrow f_2 & & \downarrow g_2 & \\ & & S & & \end{array}$$

1. The associated relation $g \circ f^{-1}$ of λ is functional.
2. For each $s, t \in S$, if $f(s) = f(t)$, then $g(s) = g(t)$.
3. this “ f -relative injectivity” condition is the same as being a monomorphism/monoid/whatever in nice category | maybe this is the same as being a skew monoid in $\text{Span}(A, B)$ or something?
 1. a⁶

00P5 Definition 6.1.4.1.2. The span λ is **total** if f is surjective.

Let $\lambda = \left(A \xleftarrow{f} S \xrightarrow{g} B \right)$ be a span. A morphism of spans from id_A to $\lambda \diamond \lambda^\dagger$ is a morphism

$$s: A \rightarrow S \times_B S$$

making the diagram

$$\begin{array}{ccc} & A & \\ id_A \swarrow & \downarrow s & \searrow id_A \\ A & & A \\ \uparrow f \circ pr'_1 & \downarrow & \downarrow f \circ pr'_2 \\ S \times_B S & & \end{array}$$

commute, where $S \times_B S$ is the pullback

$$\begin{array}{ccc} S \times_B S & \longrightarrow & S \\ \downarrow \lrcorner & & \downarrow g \\ S & \xrightarrow{g} & B \end{array}$$

of S with itself along g . In particular, $pr_1 \circ s$ and $pr_2 \circ s$ are both left-inverses/retractions

⁶Here we could perhaps also use the direct image with compact support $g_!$ of g (see ??) instead of the usual direct image, although the expression for $g_!(f^{-1}(a))$ seems a bit weird. It can also actually be given as a right Kan extension (?? of ??):

$$\begin{aligned} g_!(f^{-1}(a)) &= g_!(\{s \in S \mid f(s) = a\}) \\ &= \{b \in B \mid g^{-1}(b) \subset \{s \in S \mid f(s) = a\}\} \\ &= \{b \in B \mid \text{for each } s \in S, \text{if } g(s) = b, \text{ then } f(s) = a\} \\ &= [\text{Ran}_g^\dagger(f)](a) \end{aligned}$$

for f , i.e. we have

$$\begin{aligned} (\text{pr}_1 \circ s) \circ f &\cong \text{id}_A, \\ (\text{pr}_2 \circ s) \circ f &\cong \text{id}_A. \end{aligned}$$

Thus, by ?? of ??, f is injective if $A \neq \emptyset$.

6.1.5 Total Spans

6.2 Categories of Spans


6.2.1 The Category of Spans Between Two Sets

Let A and B be sets.

00P9 Definition 6.2.1.1.1. The **category of spans from A to B** is the category $\text{Span}(A, B)$ defined by


$$\text{Span}(A, B) \stackrel{\text{def}}{=} \text{Fun}(\Lambda, \text{Sets}) \times_{\text{ev}_{[-1]}, \text{Sets}, [A]} \text{pt} \times_{[B], \text{Sets}, \text{ev}_{[1]}} \text{Fun}(\Lambda, \text{Sets}),$$

as in the diagram



00PA Remark 6.2.1.1.2. In detail, the **category of spans from A to B** is the category $\text{Span}(A, B)$ where

as in the diagram



- *Objects.* The objects of $\text{Span}(A, B)$ are spans from A to B ;
- *Morphisms.* The morphism of $\text{Span}(A, B)$ are morphisms of spans;
- *Identities.* The unit map

$$\mathbb{1}_{(S,f,g)}^{\text{Span}(A,B)} : \text{pt} \rightarrow \text{Hom}_{\text{Span}(A,B)}((S,f,g), (S,f,g))$$

of $\text{Span}(A, B)$ at (S, f, g) is defined by⁷

$$\text{id}_{(S,f,g)}^{\text{Span}(A,B)} \stackrel{\text{def}}{=} \text{id}_S;$$

- *Composition.* The composition map

$$\circ_{R,S,T}^{\text{Span}(A,B)} : \text{Hom}_{\text{Span}(A,B)}(S, T) \times \text{Hom}_{\text{Span}(A,B)}(R, S) \rightarrow \text{Hom}_{\text{Span}(A,B)}(R, T)$$

of $\text{Span}(A, B)$ at $((R, f_1, g_1), (S, f_2, g_2), (T, f_3, g_3))$ is defined by⁸

$$\psi \circ_{R,S,T}^{\text{Span}(A,B)} \phi \stackrel{\text{def}}{=} \psi \circ \phi.$$


00PB Proposition 6.2.1.1.3. Let A and B be sets.

00PC 1. *As a Pullback.* We have an isomorphism of categories


$$\begin{array}{ccc} \text{Span}(A, B) & \longrightarrow & \text{Sets}_{/B} \\ \downarrow & \lrcorner & \downarrow \text{忘} \\ \text{Sets}_{/A} & \longrightarrow & \text{Sets}. \end{array}$$

$\cong \text{Sets}_{/A} \times_{\text{Sets}} \text{Sets}_{/B}$

⁷Picture:



⁸Picture:



Proof. **Item 1, As a Pullback:** In detail, the pullback $\text{Sets}_{/A} \times_{\text{Sets}} \text{Sets}_{/B}$ is the category where

- *Objects.* The objects of $\text{Sets}_{/A} \times_{\text{Sets}} \text{Sets}_{/B}$ consist of pairs $((S, f), (S', g))$ of objects of Sets consisting of

- A pair (S, f) in $\text{Obj}(\text{Sets}_{/A})$ consisting of a set S and a map $f: S \rightarrow A$;
- A pair (S', g) in $\text{Obj}(\text{Sets}_{/B})$ consisting of a set S' and a map $g: S' \rightarrow B$;

such that

$$\underbrace{\mathfrak{F}(S, f)}_{\stackrel{\text{def}}{=} S} = \underbrace{\mathfrak{F}(S', g)}_{\stackrel{\text{def}}{=} S'}.$$

Thus the objects of $\text{Sets}_{/A} \times_{\text{Sets}} \text{Sets}_{/B}$ are the same as spans from A to B .

- *Morphisms.* A morphism of $\text{Sets}_{/A} \times_{\text{Sets}} \text{Sets}_{/B}$ from (S, f, g) to (S', f', g') consists of a pair of morphisms

$$\begin{aligned}\phi: S &\rightarrow S' \\ \psi: S &\rightarrow S'\end{aligned}$$

such that the diagrams

$$\begin{array}{ccc} S & \xrightarrow{\phi} & S' \\ f \searrow & & \swarrow f' \\ & A & \end{array} \quad \begin{array}{ccc} S & \xrightarrow{\psi} & S' \\ g \searrow & & \swarrow g' \\ & B & \end{array}$$

such that

$$\underbrace{\mathfrak{F}(\phi)}_{\stackrel{\text{def}}{=} \phi} = \underbrace{\mathfrak{F}(\psi)}_{\stackrel{\text{def}}{=} \psi}.$$

Thus the morphisms of $\text{Sets}_{/A} \times_{\text{Sets}} \text{Sets}_{/B}$ are also the same as morphisms of spans from (S, f, g) to (S', f', g') .

- *Identities and Composition.* The identities and composition of $\text{Sets}_{/A} \times_{\text{Sets}} \text{Sets}_{/B}$ are also the same as those in $\text{Span}(A, B)$.

This finishes the proof. □

6.2.2 The Bicategory of Spans

00PE Definition 6.2.2.1.1. The **bicategory of spans** is the bicategory Span where

- *Objects.* The objects of Span are sets;

- *Hom-Categories.* For each $A, B \in \text{Obj}(\text{Span})$, we have

$$\text{Hom}_{\text{Span}}(A, B) \stackrel{\text{def}}{=} \text{Span}(A, B);$$

- *Identities.* For each $A \in \text{Obj}(\text{Span})$, the unit functor

$$\mathbb{1}_A^{\text{Span}} : \text{pt} \rightarrow \text{Span}(A, A)$$

of Span at A is the functor picking the span $(A, \text{id}_A, \text{id}_A)$:

$$\begin{array}{ccc} & A & \\ \text{id}_A \swarrow & & \searrow \text{id}_A \\ A & & A. \end{array}$$

- *Composition.* For each $A, B, C \in \text{Obj}(\text{Span})$, the composition bifunctor

$$\circ_{A,B,C}^{\text{Span}} : \text{Span}(B, C) \times \text{Span}(A, B) \rightarrow \text{Span}(A, C)$$

of Span at (A, B, C) is the bifunctor where

- *Action on Objects.* The composition of two spans

$$\begin{array}{ccc} & R & \\ f_1 \swarrow & & \searrow g_1 \\ A & & B \end{array} \quad \text{and} \quad \begin{array}{ccc} & S & \\ f_2 \swarrow & & \searrow g_2 \\ B & & C \end{array}$$

is the span $(R \times_B S, f_1 \circ \text{pr}_1, g_2 \circ \text{pr}_2)$, constructed as in the diagram

$$\begin{array}{ccccc} & & R \times_B S & & \\ & \swarrow f_1 \circ \text{pr}_1 & \downarrow \text{pr}_1 & \searrow \text{pr}_2 & \\ R & & & & S \\ \downarrow f_1 & \swarrow g_1 & & \searrow f_2 & \downarrow g_2 \\ A & & B & & C \end{array}$$

- *Action on Morphisms.* The horizontal composition of 2-morphisms is defined via functoriality of pullbacks: given morphisms of spans

$$\begin{array}{ccc} & R & \\ f \swarrow & \downarrow \phi & \searrow g \\ A & & B \end{array} \quad \text{and} \quad \begin{array}{ccc} & S & \\ h \swarrow & \downarrow \psi & \searrow k \\ B & & C \\ \downarrow h' & \swarrow g' & \searrow k' \\ S' & & \end{array}$$

their horizontal composition is the morphism of spans

$$\begin{array}{ccc}
 & R \times_B S & \\
 f \circ \text{pr}_1 \swarrow & \downarrow \exists! & \searrow k \circ \text{pr}_2 \\
 A & & C, \\
 h' \circ \text{pr}'_1 \swarrow & \downarrow & \searrow k' \circ \text{pr}'_2 \\
 R' \times_B S' & &
 \end{array}$$

constructed as in the diagram

$$\begin{array}{ccccc}
 & R \times_B S & & & \\
 f \circ \text{pr}_1 \swarrow & \downarrow \exists! & \searrow \text{pr}_2 & & k \circ \text{pr}_2 \\
 R & & S & & C; \\
 f \swarrow & \downarrow g & \searrow h & \searrow k & \\
 A & & B & & C \\
 f' \swarrow & \downarrow g' & \searrow h' & \searrow k' & \\
 R' & & S' & & \\
 f' \circ \text{pr}'_1 \swarrow & \downarrow \text{pr}'_1 & \searrow \text{pr}'_2 & & k' \circ \text{pr}'_2 \\
 R' \times_B S' & & & &
 \end{array}$$

- *Associators and Unitors.* The associator and unitors are defined using the universal property of the pullback.

6.2.3 The Monoidal Bicategory of Spans

6.2.4 The Double Category of Spans

00PH Definition 6.2.4.1.1. The **double category of spans** is the double category Span^{dbl} where

- *Objects.* The objects of Span^{dbl} are sets;
- *Vertical Morphisms.* The vertical morphisms of Span^{dbl} are functions $f: A \rightarrow B$;
- *Horizontal Morphisms.* The horizontal morphisms of Span^{dbl} are spans $(S, \phi, \psi): A \dashrightarrow X$;

- *2-Morphisms.* A 2-cell

$$\begin{array}{ccc} A & \xrightarrow{(R,\phi_R,\psi_R)} & B \\ f \downarrow & \Downarrow \alpha & \downarrow g \\ X & \xrightarrow{(S,\phi_S,\psi_S)} & Y \end{array}$$

of Span^{dbl} is a morphism of spans from the span

$$\begin{array}{ccccc} & & R & & \\ & \swarrow \phi_R & & \searrow \psi_R & \\ A & & B & & Y \\ & \searrow g & & & \end{array}$$

to the span

$$\begin{array}{ccccc} & & A \times_X S & & \\ & \swarrow & \downarrow & \searrow & \\ & A & & S & \\ f \swarrow & \searrow f & \swarrow \phi_S & \searrow \psi_S & \\ X & & X & & Y \end{array};$$

- *Horizontal Identities.* The horizontal unit functor

$$\mathbb{1}_{\text{Span}^{\text{dbl}}} : (\text{Span}^{\text{dbl}})_0 \rightarrow (\text{Span}^{\text{dbl}})_1$$

of Span^{dbl} is the functor where

- *Action on Objects.* For each $A \in \text{Obj}((\text{Span}^{\text{dbl}})_0)$, we have

$$\mathbb{1}_A \stackrel{\text{def}}{=} (A, \text{id}_A, \text{id}_A),$$

as in the diagram

$$\begin{array}{ccc} & A & \\ \text{id}_A \swarrow & \Downarrow & \searrow \text{id}_A \\ A & & A \end{array};$$

- *Action on Morphisms.* For each vertical morphism $f: A \rightarrow B$ of Span^{dbl} , i.e.

each map of sets f from A to B , the identity 2-morphism

$$\begin{array}{ccc} A & \xrightarrow{\quad \sharp_A \quad} & A \\ f \downarrow & \parallel \sharp_f \downarrow & \downarrow f \\ B & \xrightarrow{\quad \sharp_B \quad} & B \end{array}$$

of f is the morphism of spans from

$$\begin{array}{ccc} & A & \\ & \swarrow \text{id}_A & \searrow \text{id}_A \\ A & & A \\ & \swarrow & \searrow f \\ & & B \end{array}$$

to

$$\begin{array}{ccccc} & & A \times_B B & & \\ & \swarrow & \downarrow & \searrow & \\ & A & & B & \\ f \swarrow & \searrow f & \swarrow \text{id}_B & \searrow & \text{id}_B \\ B & & B & & B \end{array}$$

given by the isomorphism $A \xrightarrow{\cong} A \times_B B$;

- *Vertical Identities.* For each $A \in \text{Obj}(\text{Span}^{\text{dbl}})$, we have

$$\text{id}_A^{\text{Span}^{\text{dbl}}} \stackrel{\text{def}}{=} \text{id}_A;$$

- *Identity 2-Morphisms.* For each horizontal morphism $R: A \rightarrow B$ of Span^{dbl} , the identity 2-morphism

$$\begin{array}{ccc} A & \xrightarrow{\quad S \quad} & B \\ \text{id}_A \downarrow & \parallel \text{id}_S \downarrow & \downarrow \text{id}_B \\ A & \xrightarrow{\quad S \quad} & B \end{array}$$

of R is the morphism of spans from

$$\begin{array}{ccc} & S & \\ & \swarrow \phi_S & \searrow \psi_S \\ A & & B \\ & \swarrow & \searrow \text{id}_B \\ & & B \end{array}$$

to

$$\begin{array}{ccccc}
 & & A \times_A S & & \\
 & \swarrow & \downarrow & \searrow & \\
 A & \xrightarrow{\quad id_A \quad} & A & \xrightarrow{\quad id_A \quad} & A \\
 & \parallel & \parallel & \parallel & \\
 & & S & & B \\
 & & \phi_S \swarrow & \psi_S \searrow & \\
 & & & & B
 \end{array}$$

given by the isomorphism $S \xrightarrow{\cong} A \times_A S$;

- *Horizontal Composition.* The horizontal composition functor

$$\odot^{\text{Span}^{\text{dbl}}} : (\text{Span}^{\text{dbl}})_1 \times_{(\text{Span}^{\text{dbl}})_0} (\text{Span}^{\text{dbl}})_1 \rightarrow (\text{Span}^{\text{dbl}})_1$$

of Span^{dbl} is the functor where

- *Action on Objects.* For each composable pair

$$\begin{array}{ccc}
 (R, \phi_R, \psi_R) & & (S, \phi_S, \psi_S) \\
 A \xrightarrow{\quad} B \xrightarrow{\quad} C
 \end{array}$$

of horizontal morphisms of Span^{dbl} , we have

$$(S, \phi_S, \psi_S) \odot (R, \phi_R, \psi_R) \stackrel{\text{def}}{=} S \circ_{A,B,C}^{\text{Span}} R,$$

where $S \circ_{A,B,C}^{\text{Span}} R$ is the composition of (R, ϕ_R, ψ_R) and (S, ϕ_S, ψ_S) defined as in [Definition 6.2.2.1.1](#);

- *Action on Morphisms.* For each horizontally composable pair

$$\begin{array}{ccc}
 A & \xrightarrow{(R, \phi_R, \psi_R)} & B \\
 f \downarrow & \parallel \alpha \Downarrow & \downarrow g \\
 X & \xrightarrow{(T, \phi_T, \psi_T)} & Y
 \end{array}
 \qquad
 \begin{array}{ccc}
 B & \xrightarrow{(S, \phi_S, \psi_S)} & C \\
 g \downarrow & \parallel \beta \Downarrow & \downarrow h \\
 Y & \xrightarrow{(U, \phi_U, \psi_U)} & Z
 \end{array}$$

of 2-morphisms of Span^{dbl} ,

- *Vertical Composition of 1-Morphisms.* For each composable pair $A \xrightarrow{F} B \xrightarrow{G} C$ of vertical morphisms of Span^{dbl} , i.e. maps of sets, we have

$$g \circ^{\text{Span}^{\text{dbl}}} f \stackrel{\text{def}}{=} g \circ f;$$

- *Vertical Composition of 2-Morphisms.* For each vertically composable pair

$$\begin{array}{ccc}
 A & \xrightarrow{(R,\phi_R,\psi_R)} & X \\
 f \downarrow & \Downarrow \alpha & \downarrow g \\
 B & \xrightarrow{(S,\phi_S,\psi_S)} & Y
 \end{array}
 \quad
 \begin{array}{ccc}
 B & \xrightarrow{(S,\phi_S,\psi_S)} & Y \\
 h \downarrow & \Downarrow \beta & \downarrow k \\
 C & \xrightarrow{(T,\phi_T,\psi_T)} & Z
 \end{array}$$

of 2-morphisms of Span^{dbl} ,

- *Associators and Unitors.* The associator and unitors of Span^{dbl} are defined using the universal property of the pullback.

6.2.5 Properties of The Bicategory of Spans

00PK Proposition 6.2.5.1.1. Let $\lambda = (A \xleftarrow{f} S \xrightarrow{g} B)$ be a span.

- 00PL 1. *Self-Duality.*
- 00PM 2. *Isomorphisms in Span.*
- 00PN 3. *Equivalences in Span.*
- 00PP 4. *Adjunctions in Span.* Let A and B be sets.⁹
- 00PQ (a) We have a natural bijection

$$\left\{ \begin{array}{c} \text{Adjunctions in Span} \\ \text{from } A \text{ to } B \end{array} \right\} \cong \left\{ \begin{array}{c} \text{Spans } A \xleftarrow{f} S \xrightarrow{g} B \\ \text{from } A \text{ to } B \text{ with} \\ f \text{ an isomorphism} \end{array} \right\}.$$

- 00PR (b) We have an equivalence of categories

$$\text{MapSpan}(A, B) \stackrel{\text{eq.}}{\cong} \text{Sets}(A, B)_{\text{disc}},$$

where $\text{MapSpan}(A, B)$ is the full subcategory of $\text{Span}(A, B)$ spanned by the spans $A \xleftarrow{f} S \xrightarrow{g} B$ from A to B with f an isomorphism.

- 00PS (c) We have a biequivalence of bicategories

$$\text{MapSpan} \stackrel{\text{eq.}}{\cong} \text{Sets}_{\text{bidisc}},$$

where MapSpan is the sub-bicategory of Span whose Hom-categories are given by $\text{MapSpan}(A, B)$.

⁹In the literature (e.g. [ref]),... are called maps and denoted by $\text{MapSpan}(A, B)$

- 00PT 5. *Monads in Span.*
- 00PU 6. *Comonads in Span.*
- 00PV 7. *Monomorphisms in Span.*
- 00PW 8. *Epimorphisms in Span.*
- 00PX 9. *Existence of Right Kan Extensions.*
- 00PY 10. *Existence of Right Kan Lifts.*
- 00PZ 11. *Closedness.*

Proof. Item 1, SelfDuality:

Item 2, Isomorphisms in Span:

Item 3, Equivalences in Span:

Item 4, Adjunctions in Span: We first prove Item 4a.

We proceed step by step:

1. *From Adjunctions in Span to Functions.* An adjunction in Span from A to B consists of a pair of spans

$$\begin{array}{ccc} & R & \\ f \swarrow & & \searrow g \\ A & & B \end{array} \quad \text{and} \quad \begin{array}{ccc} & S' & \\ h \swarrow & & \searrow k \\ B & & A \end{array}$$

together with maps

$$\begin{array}{ccc} & A & \\ id_A \swarrow & \downarrow & \searrow id_A \\ A & \phi & A \\ \uparrow & \downarrow & \uparrow \\ f \circ pr'_1 & \nearrow & \searrow k \circ pr'_2 \\ S \times_B S' & & \end{array} \quad \text{and} \quad \begin{array}{ccc} & S' \times_A S & \\ h \circ pr_1 \swarrow & \downarrow & \searrow g \circ pr_2 \\ B & \psi & B \\ \uparrow & \downarrow & \uparrow \\ id_B \swarrow & \downarrow & \searrow id_B \\ B & & B \end{array}$$

We claim that these conditions

2. *From Functions to Adjunctions in Rel.*
3. *Invertibility: From Functions to Adjunctions Back to Functions.*
4. *Invertibility: From Adjunctions to Functions Back to Adjunctions.*

We now proceed to the proof of Item 4b. For this, we will construct a functor

$$F: \text{Sets}(A, B)_{\text{disc}} \rightarrow \text{MapSpan}(A, B)$$

and prove it to be essentially surjective and fully faithful, and thus an equivalence by ?? of ???. Indeed, given a map $f: A \rightarrow B$, let $F(f)$ be the representable span associated to f of **Definition 6.5.1.1.1**, and let F send the unique (identity) morphism from f to itself to the identity morphism of $F(f)$ in $\text{MapSpan}(A, B)$. We now prove that F is fully faithful and essentially surjective:

1. *F Is Fully Faithful:* Given maps $f, g: A \rightrightarrows B$, we need to show that

$$\text{Hom}_{\text{MapSpan}(A, B)}(F(f), F(g)) = \begin{cases} \text{pt} & \text{if } f = g, \\ \emptyset & \text{otherwise.} \end{cases}$$

Indeed, a morphism from $F(f)$ to $F(g)$ takes the form

$$\begin{array}{ccc} & A & \\ \swarrow & & \searrow \\ A & \phi & B \\ \swarrow & & \searrow \\ & A & \end{array}$$

From the relations $\text{id}_A = \text{id}_A \circ \phi$ and $f = g \circ \phi$, we see that $\phi = \text{id}_A$, and thus from the relation $f = g \circ \phi$ there is such a morphism iff $f = g$.

2. *F Is Essentially Surjective:* Let λ be a span of the form

$$\begin{array}{ccc} & S & \\ \phi \nearrow & \searrow f \\ A & & B \end{array}$$

we claim that $\lambda \cong F(f \circ \phi^{-1})$. Indeed, we have morphisms

$$\begin{array}{ccc} & S & \\ \phi \nearrow & \searrow f & \\ A & \phi \downarrow & B \\ \swarrow & \nearrow f \circ \phi^{-1} & \\ & A & \end{array} \quad \text{and} \quad \begin{array}{ccc} & A & \\ \swarrow & \nearrow f \circ \phi^{-1} & \\ A & \phi^{-1} \downarrow & B \\ \nearrow & \swarrow f & \\ & S & \end{array}$$

inverse to each other in $\text{MapSpan}(A, B)$, and thus $\lambda \cong F(f \circ \phi^{-1})$.

Finally, we prove **Item 4c**.

Item 5, Monads in Span:

Item 6, Comonads in Span:

Item 7, Monomorphisms in Span:

Item 8, Epimorphisms in Span:

Item 9, Existence of Right Kan Extensions:

Item 10, Existence of Right Kan Lifts:

Item 11, Closedness:

□

6.3 Limits of Spans

6.3.1 tmp2

$$\text{Hom}_{\text{Rel}(A,X)}(\text{Lan}_S(R), T) \cong \text{Hom}_{\text{Rel}(B,X)}(R, T \diamond S)$$

1. $\text{Lan}_S(R) \subset T$, i.e. if $a \sim_{\text{Lan}_S(R)} x$, then $a \sim_T x$.
2. $R \subset T \diamond S$, i.e. if $b \sim_R x$, then there exists some $a \in A$ such that $a \sim_S b$ and $b \sim_T x$.

6.3.2 tmp

$$\begin{array}{ccc}
 & \begin{array}{c} S \\ f \swarrow \quad \searrow g \\ A \qquad B \end{array} & \\
 & \Downarrow & \\
 & \begin{array}{ccc} S' & & S' \times_B S' \\ \phi_{S'} \swarrow \quad \searrow \psi_{S'} & \mapsto & f \circ \text{pr}_1 \swarrow \quad \searrow \psi_{S'} \circ \text{pr}_2 \\ B \qquad X & & A \qquad X \end{array} & \\
 & \Downarrow & \\
 & \begin{array}{ccc} S'' & & ? \\ \phi_{S''} \swarrow \quad \searrow \psi_{S''} & \mapsto & ? \swarrow \quad \searrow ? \\ A \qquad X & & B \qquad X \end{array} & \\
 & \Downarrow & \\
 & \begin{array}{ccc} S' \times_B S' & & S' \\ f \circ \text{pr}_1 \swarrow \quad \downarrow \xi \quad \searrow \psi_{S'} \circ \text{pr}_2 & & \phi_{S'} \swarrow \quad \downarrow \xi^\dagger \quad \searrow \psi_{S'} \\ A \qquad S'' & \xrightarrow{\quad \psi_{S''} \quad} & B \qquad \Pi_g(S'') \\ \phi_{S''} \swarrow \quad \uparrow \quad \searrow \psi_{S''} & & ? \swarrow \quad \downarrow \xi^\dagger \quad \searrow ? \\ A \qquad X & & B \qquad X \end{array} &
 \end{array}$$

$\text{Hom}_{\text{Span}(A,X)}(S \times_B S', K) \cong \text{Hom}_{\text{Span}(B,X)}(S', R(K))$

6.3.3 Left Kan Extensions

Let $\lambda = \left(A \xleftarrow{f} S \xrightarrow{g} B \right)$ be a span.

00Q2 Proposition 6.3.3.1.1. The left Kan extension

$$\text{Lan}_\lambda: \text{Span}(A, X) \rightarrow \text{Span}(B, X)$$

along λ in Span exists and is the functor given on objects by sending a span λ' in $\text{Span}(A, X)$ as in

$$\begin{array}{ccc} & S' & \\ \phi \swarrow & & \searrow \psi \\ A & & X \end{array}$$

to the span

$$\text{Lan}_\lambda(\lambda') \stackrel{\text{def}}{=} (\text{Lan}_\lambda(S'), \text{Lan}_\lambda(\phi), \text{Lan}_\lambda(\psi)),$$

in $\text{Span}(B, X)$ where

- The set $\text{Lan}_\lambda(S')$ is given by

$$\begin{aligned} \text{Lan}_\lambda(S') &\stackrel{\text{def}}{=} \Sigma_g(S') \\ &\stackrel{\text{def}}{=} S' \end{aligned}$$

where $\Sigma_g(S')$ is the dependent sum of $\phi: S' \rightarrow A$ along g of ??;

- The map $\text{Lan}_\lambda(\phi): \text{Lan}_\lambda(S') \rightarrow B$ is given by $\Sigma_g(\phi)$;
- The map $\text{Lan}_\lambda(\psi): \text{Lan}_\lambda(S') \rightarrow X$ is given by ψ .

Proof.

□

6.3.4 Right Kan Extensions

Let $\lambda = (A \xleftarrow{f} S \xrightarrow{g} B)$ be a span.

00Q4 **Proposition 6.3.4.1.1.** The right Kan extension

$$\text{Ran}_\lambda: \text{Span}(A, X) \rightarrow \text{Span}(B, X)$$

along λ in Span exists and is the functor given on objects by sending a span λ' in $\text{Span}(A, X)$ as in

$$\begin{array}{ccc} & S' & \\ \phi_{S'} \swarrow & & \searrow \psi_{S'} \\ A & & X \end{array}$$

to the span

$$\text{Ran}_\lambda(\lambda') \stackrel{\text{def}}{=} (\text{Ran}_\lambda(S'), \text{Ran}_\lambda(\phi_{S'}), \text{Ran}_\lambda(\psi_{S'})),$$

in $\text{Span}(B, X)$ where

- The set $\text{Ran}_\lambda(S')$ is given by

$$\text{Ran}_\lambda(S') \stackrel{\text{def}}{=} \coprod_{b \in B} \prod_{s \in g^{-1}(b)} \phi_{S'}^{-1}(f(s));$$

- The map $\text{Ran}_\lambda(\phi_{S'}) : \text{Ran}_\lambda(S') \rightarrow B$ is given by

$$[\text{Ran}_\lambda(\phi_{S'})](b, (s'_s)_{s \in g^{-1}(b)}) \stackrel{\text{def}}{=} b;$$

for each $(b, (s'_s)_{s \in g^{-1}(b)}) \in \text{Ran}_\lambda(S')$;

- The map $\text{Ran}_\lambda(\psi_{S'}) : \text{Ran}_\lambda(S') \rightarrow X$ is given by

$$[\text{Ran}_\lambda(\psi_{S'})](b, (s'_s)_{s \in g^{-1}(b)}) \stackrel{\text{def}}{=} \psi_{S'}(s'_i)$$

for each $(b, (s'_s)_{s \in g^{-1}(b)}) \in \text{Ran}_\lambda(S')$, where the i in s'_i denotes any $s \in g^{-1}(b)$, as we have $\psi_{S'}(s'_i) = \psi_{S'}(s'_j)$ for all $s \in g^{-1}(b)$.¹⁰

Proof.

□

6.3.5 Right Kan Lifts

(Although right Kan lifts aren't really limits, this is probably the most appropriate to place this section.)

Let $\lambda = \left(A \xleftarrow{f} S \xrightarrow{g} B \right)$ be a span.

00Q6 Proposition 6.3.5.1.1. The right Kan lift

$$\text{Rift}_\lambda : \text{Span}(X, B) \rightarrow \text{Span}(X, A)$$

along λ in Span exists and is the functor given on objects by sending a span λ' in $\text{Span}(X, B)$ as in

$$\begin{array}{ccc} & S' & \\ \phi \swarrow & & \searrow \psi \\ X & & B \end{array}$$

to the span

$$\text{Rift}_\lambda(\lambda') \stackrel{\text{def}}{=} (\text{Rift}_\lambda(S'), \text{Rift}_\lambda(\phi), \text{Rift}_\lambda(\psi)),$$

in $\text{Span}(X, A)$ where

- The set $\text{Rift}_\lambda(S')$ is given by

$$\text{Rift}_\lambda(S') \stackrel{\text{def}}{=} \Pi_f(S'),$$

where $\Pi_f(S')$ is the dependent product of $\psi : S' \rightarrow A$ along f of ??;

- The map $\text{Rift}_\lambda(\phi) : \text{Rift}_\lambda(S') \rightarrow X$ is given by ϕ ;
- The map $\text{Rift}_\lambda(\psi) : \text{Rift}_\lambda(S') \rightarrow A$ is given by $\Pi_f(\psi)$.

Proof.

□

¹⁰Indeed

6.4 Colimits of Spans

6.5 Constructions With Spans

6.5.1 Representable Spans

00QA Definition 6.5.1.1.1. Let $f: A \rightarrow B$ be a function.

- The **representable span associated to f** is the span

$$\begin{array}{ccc} & A & \\ id_A & \swarrow \searrow & \\ A & & B \end{array}$$

from A to B .

- The **corepresentable span associated to f** is the span

$$\begin{array}{ccc} & A & \\ f & \swarrow \searrow & id_A \\ B & & A \end{array}$$

from B to A .

6.5.2 Composition of Spans

00QC Definition 6.5.2.1.1. The **composition** of two spans

$$\begin{array}{ccc} & R & \\ f_1 & \swarrow \searrow & g_1 \\ A & & B \end{array} \quad \text{and} \quad \begin{array}{ccc} & S & \\ f_2 & \swarrow \searrow & g_2 \\ B & & C \end{array}$$

is the span $(R \times_B S, f_1 \circ \text{pr}_1, g_2 \circ \text{pr}_2)$, constructed as in the diagram

$$\begin{array}{ccccc} & R \times_B S & & & \\ & \text{pr}_1 \swarrow \searrow \text{pr}_2 & & & \\ & R & & S & \\ f_1 \curvearrowleft & \swarrow \searrow & & \swarrow \searrow & g_2 \circ \text{pr}_2 \\ A & & B & & C \end{array}$$

6.5.3 Horizontal Composition of Morphisms of Spans

00QE Definition 6.5.3.1.1. The **horizontal composition** of a pair of 2-morphisms of spans

$$\begin{array}{ccc} & R & \\ f \swarrow & \downarrow \phi & \searrow g \\ A & & B \\ \uparrow f' & \downarrow g' & \uparrow h \\ R' & & S' \end{array} \quad \text{and} \quad \begin{array}{ccc} & S & \\ h \swarrow & \downarrow \psi & \searrow k \\ B & & C \\ \uparrow h' & \downarrow k' & \uparrow \\ S' & & \end{array}$$

is the morphism of spans

$$\begin{array}{ccc} & R \times_B S & \\ f \circ \text{pr}_1 \swarrow & \downarrow \exists! & \searrow k \circ \text{pr}_2 \\ A & & C \\ \uparrow h' \circ \text{pr}'_1 & \downarrow & \uparrow k' \circ \text{pr}'_2 \\ R' \times_B S' & & \end{array}$$

constructed as in the diagram

$$\begin{array}{ccccc} & R \times_B S & & & \\ & \swarrow \text{pr}_1 & \downarrow \exists! & \searrow \text{pr}_2 & \\ & R & & S & \\ f \swarrow & \downarrow g & \swarrow h & \searrow k & \\ A & & B & & C \\ \uparrow f' & \uparrow g' & \uparrow h' & \uparrow k' & \\ R' & & S' & & \\ \uparrow f' \circ \text{pr}'_1 & \uparrow \text{pr}'_1 & \uparrow \text{pr}'_2 & \uparrow k' \circ \text{pr}'_2 & \\ R' \times_B S' & & & & \end{array}$$

6.5.4 Properties of Composition of Spans

00QG Proposition 6.5.4.1.1. Let $\lambda = \left(A \xleftarrow{f} S \xrightarrow{g} B \right)$ be a span.

00QH 1. *Functoriality.*

Proof.

□

6.5.5 The Inverse of a Span

6.6 Functoriality of Spans

6.6.1 Direct Images

6.6.2 Functoriality of Spans on Powersets

6.7 Un/Straightening for Spans

6.7.1 Straightening for Spans

Let A and B be sets and let (S, f, g) be a span from A to B .

00QQ **Definition 6.7.1.1.1.** The **straightening** of (S, f, g) is the $(A \times B)$ -indexed set

$$\text{St}_{A,B}(S) : (A \times B)_{\text{disc}} \rightarrow \text{Sets}$$

defined as the straightening of S , viewed as an $(A \times B)$ -fibred set, as in ??.

00QR **Remark 6.7.1.1.2.** In detail, $\text{St}_{A,B}(S)$ is the $(A \times B)$ -indexed set defined by¹¹

$$\begin{aligned} [\text{St}_{A,B}(S)](a, b) &\stackrel{\text{def}}{=} \text{Wits}_S(a, b) \\ &\stackrel{\text{def}}{=} S_{ab} \\ &\stackrel{\text{def}}{=} \{s \in S \mid f(s) = a \text{ and } g(s) = b\}. \end{aligned}$$

00QS **Proposition 6.7.1.1.3.** Let A and B be sets and let (S, f, g) be a span.

00QT 1. *Functoriality.* The assignment $(S, f, g) \mapsto \text{St}_{A,B}(S)$ defines a functor

$$\text{St}_{A,B} : \text{Span}(A, B) \rightarrow \text{ISets}(A \times B)$$

- *Action on Objects.* For each $(S, f, g) \in \text{Obj}(\text{Span}(A, B))$, we have

$$[\text{St}_{A,B}](S, f, g) \stackrel{\text{def}}{=} \text{St}_{A,B}(S);$$

- *Action on Morphisms.* For each $(S_1, f_1, g_1), (S_2, f_2, g_2) \in \text{Obj}(\text{Span}(A, B))$, the action on Hom-sets

$$\text{St}_{A,B|S_1, S_2} : \text{Hom}_{\text{Span}(A, B)}(S_1, S_2) \rightarrow \text{Hom}_{\text{ISets}(A \times B)}(\text{St}_{A,B}(S_1), \text{St}_{A,B}(S_2))$$

of $\text{St}_{A,B}$ at (S_1, S_2) is given by sending a morphism

$$\phi : (S_1, f_1, g_1) \rightarrow (S_2, f_2, g_2)$$

¹¹Here we may think of $\text{Wits}_S(a, b)$ as the “set of witnesses in S that $a \sim b$ holds”; see Remark 6.1.2.1.3.

of spans from A to B to the morphism

$$\text{St}_{A,B}(\phi) : \text{St}_{A,B}(S_1) \rightarrow \text{St}_{A,B}(S_2)$$

of $(A \times B)$ -indexed sets defined by

$$\text{St}_{A,B}(\phi) \stackrel{\text{def}}{=} \{\phi_{ab}^*\}_{(a,b) \in A \times B},$$

where ϕ_{ab}^* is the transport map associated to ϕ at $(a, b) \in A \times B$ of ??.

Proof. Item 1, Functoriality: This is the special case of ?? of ?? where $K = A \times B$. \square

6.7.2 Unstraightening for Spans

Let A and B be sets and let $S : (A \times B)_{\text{disc}} \rightarrow \text{Sets}$ be an $(A \times B)$ -indexed set.

00QV **Definition 6.7.2.1.1.** The **unstraightening** of S is the span

$$\begin{array}{ccc} & \text{Un}_{A,B}(S) & \\ f_{\text{Un}_{A,B}(S)} \swarrow & & \searrow g_{\text{Un}_{A,B}(S)} \\ A & & B \end{array}$$

from A to B where

$$\text{Un}_{A,B}(S) \stackrel{\text{def}}{=} \coprod_{(a,b) \in A \times B} S(a, b)$$

and where the maps $f_{\text{Un}_{A,B}(S)}$ and $g_{\text{Un}_{A,B}(S)}$ are given by

$$\begin{aligned} f_{\text{Un}_{A,B}(S)}((a, b), s) &\stackrel{\text{def}}{=} a, \\ g_{\text{Un}_{A,B}(S)}((a, b), s) &\stackrel{\text{def}}{=} b \end{aligned}$$

for each $((a, b), s) \in \text{Un}_{A,B}(S)$.

00QW **Proposition 6.7.2.1.2.** Let A and B be sets.

00QX 1. *Functoriality.* The assignment $S \mapsto \text{Un}_{A,B}(S)$ defines a functor

$$\text{Un}_{A,B} : \text{ISets}(A \times B) \rightarrow \text{Span}(A, B)$$

- *Action on Objects.* For each $S \in \text{Obj}(\text{ISets}(A \times B))$, we have

$$[\text{Un}_{A,B}](S) \stackrel{\text{def}}{=} \text{Un}_{A,B}(S);$$

- *Action on Morphisms.* For each $S, S' \in \text{Obj}(\text{ISets}(A \times B))$, the action on Hom-sets

$$\text{Un}_{A,B|S,S'} : \text{Hom}_{\text{ISets}(A \times B)}(S, S') \rightarrow \text{Hom}_{\text{Span}(A, B)}(\text{Un}_{A,B}(S), \text{Un}_{A,B}(S'))$$

of $\text{Un}_{A,B}$ at (S, S') is defined by

$$\text{Un}_{A,B|S,S'}(f) \stackrel{\text{def}}{=} \coprod_{(a,b) \in A \times B} f_{ab}.$$

- 00QY** 2. *Interaction With Fibres.* Viewing the legs of $\text{Un}_{A,B}(S)$ as a morphism $(f, g): \text{Un}_{A,B}(S) \rightarrow A \times B$, we have a bijection of sets

$$(f, g)^{-1}_{\text{Un}_{A,B}(S)}(a, b) \cong S(a, b)$$

for each $(a, b) \in A \times B$.

- 00QZ** 3. *As a Pullback.* We have a bijection of sets

$$\begin{array}{ccc} \text{Un}_{A,B}(S) & \longrightarrow & \text{Sets}_* \\ \downarrow & \dashv & \downarrow \text{忘} \\ \text{Un}_{A,B}(S) \cong (A \times B)_{\text{disc}} \times_{\text{Sets}} \text{Sets}_* & & \\ \downarrow & & \downarrow \\ (A \times B)_{\text{disc}} & \xrightarrow[S]{} & \text{Sets}. \end{array}$$

- 00R0** 4. *As a Colimit.* We have a bijection of sets

$$\text{Un}_{A,B}(S) \cong \text{colim}(S).$$

Proof. **Item 1, Functoriality:** This is the special case of ?? of ?? where $K = A \times B$.

Item 2, Interaction With Fibres: This is the special case of ?? of ?? where $K = A \times B$.

Item 3, As a Pullback: This is the special case of ?? of ?? where $K = A \times B$.

Item 4, As a Colimit: This is the special case of ?? of ?? where $K = A \times B$. \square

6.7.3 The Un/Straightening Equivalence for Spans

- 00R2** **Theorem 6.7.3.1.1.** We have an isomorphism of categories

$$(\text{St}_{A,B} \dashv \text{Un}_{A,B}): \text{Span}(A, B) \begin{array}{c} \xrightarrow[\text{St}_{A,B}]{} \\ \perp \\ \xleftarrow[\text{Un}_{A,B}]{} \end{array} \text{ISets}(A \times B).$$

Proof. This is the special case of ?? where $K = A \times B$. \square

6.8 Comparison of Spans to Functions and Relations

6.8.1 Comparison to Functions

- 00R5** **Proposition 6.8.1.1.1.** We have a pseudofunctor

$$\iota: \text{Sets}_{\text{bidisc}} \rightarrow \text{Span}$$

from $\text{Sets}_{\text{bidisc}}$ to Span where

- *Action on Objects.* For each $A \in \text{Obj}(\text{Sets}_{\text{bidisc}})$, we have

$$\iota(A) \stackrel{\text{def}}{=} A;$$

- *Action on Hom-Categories.* For each $A, B \in \text{Obj}(\text{Sets}_{\text{bidisc}})$, the action on Hom-categories

$$\iota_{A,B}: \text{Sets}(A, B)_{\text{disc}} \rightarrow \text{Span}(A, B)$$

of ι at (A, B) is the functor defined on objects by sending a function $f: A \rightarrow B$ to the span

$$\begin{array}{ccc} & A & \\ id_A \swarrow & & \searrow f \\ A & & B \end{array}$$

from A to B .

- *Strict Unity Constraints.* For each $A \in \text{Obj}(\text{Sets}_{\text{bidisc}})$, the strict unity constraint

$$\iota_A^0: id_{\iota(A)} \Rightarrow \iota(id_A)$$

of ι at A is given by the identity morphism of spans

$$\begin{array}{ccc} & A & \\ id_A \swarrow & \parallel & \searrow id_A \\ A & id & A, \\ id_A \swarrow & \parallel & \searrow id_A \\ A & & A \end{array}$$

as indeed $id_{\iota(A)} = \iota(id_A)$;

- *Pseudofunctoriality Constraints.* For each $A, B, C \in \text{Obj}(\text{Sets}_{\text{bidisc}})$, each $f \in \text{Hom}_{\text{Sets}_{\text{bidisc}}}(A, B)$, and each $g \in \text{Hom}_{\text{Sets}_{\text{bidisc}}}(B, C)$, the pseudofunctoriality constraint

$$\iota_{g,f}^2: \iota(g) \circ \iota(f) \Rightarrow \iota(g \circ f)$$

of ι at (f, g) is the morphism of spans from the span

$$\begin{array}{ccccc} & & A \times_B B & & \\ & \swarrow id_A \circ pr_1 & \downarrow pr_1 & \searrow pr_2 & \\ A & & B & & C \\ & \searrow id_A & f & \swarrow id_B & g \\ & & B & & \end{array}$$

to the span

$$\begin{array}{ccc} & A & \\ id_A \swarrow & & \searrow g \circ f \\ A & & C \end{array}$$

given by the isomorphism $A \times_B B \cong A$.

Proof. Omitted. □

6.8.2 Comparison to Relations: From Span to Rel

6.8.2.1 Relations Associated to Spans

Let $\lambda = \left(A \xleftarrow{f} S \xrightarrow{g} B \right)$ be a span.

00R8 Definition 6.8.2.1.1. The **relation associated to λ** is the relation

$$S(\lambda) : A \nrightarrow B$$

from A to B defined as follows:

- Viewing relations from A to B as functions $A \times B \rightarrow \{\text{true}, \text{false}\}$, we define

$$\iota_{A,B}(S)_b^a \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if there exists } x \in S \text{ such} \\ & \text{that } a = f(x) \text{ and } b = g(x), \\ \text{false} & \text{otherwise} \end{cases}$$

for each $(a, b) \in A \times B$.

- Viewing relations from A to B as functions $A \rightarrow \mathcal{P}(B)$, we define

$$[\iota_{A,B}(S)](a) \stackrel{\text{def}}{=} g(f^{-1}(a))$$

for each $a \in A$.

- Viewing relations from A to B as subsets of $A \times B$, we define

$$\iota_{A,B}(S) \stackrel{\text{def}}{=} \{(f(x), g(x)) \mid x \in S\}.$$

00R9 Proposition 6.8.2.1.2. Let $\lambda = \left(A \xleftarrow{f} S \xrightarrow{g} B \right)$ be a span.

00RA 1. *Interaction With Identities.*

00RB 2. *Interaction With Composition.*

00RC 3. *Interaction With Inverses.*

Proof.

□

6.8.2.2 The Comparison Functor from Span to Rel

00RE Proposition 6.8.2.2.1. We have a pseudofunctor

$$\iota : \text{Span} \rightarrow \text{Rel}$$

from **Span** to **Rel** where

- *Action on Objects.* For each $A \in \text{Obj}(\text{Span})$, we have

$$\iota(A) \stackrel{\text{def}}{=} A;$$

- *Action on Hom-Categories.* For each $A, B \in \text{Obj}(\text{Span})$, the action on Hom-categories

$$\iota_{A,B} : \text{Span}(A, B) \rightarrow \mathbf{Rel}(A, B)$$

of ι at (A, B) is the functor where

- *Action on Objects.* Given a span

$$\begin{array}{ccc} & S & \\ f \swarrow & & \searrow g \\ A & & B \end{array}$$

from A to B , the image

$$\iota_{A,B}(S) : A \dashrightarrow B$$

of S by ι is the relation from A to B defined as follows:

- * Viewing relations as functions $A \times B \rightarrow \{\text{true}, \text{false}\}$, we define

$$\iota_{A,B}(S)_b^a \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if there exists } x \in S \\ & \text{such that } a = f(x) \\ & \text{and } b = g(x), \\ \text{false} & \text{otherwise} \end{cases}$$

for each $(a, b) \in A \times B$;

- * Viewing relations as functions $A \rightarrow \mathcal{P}(B)$, we define

$$[\iota_{A,B}(S)](a) \stackrel{\text{def}}{=} g(f^{-1}(a))$$

for each $a \in A$;

- * Viewing relations as subsets of $A \times B$, we define

$$\iota_{A,B}(S) \stackrel{\text{def}}{=} \{(f(x), g(x)) \mid x \in S\}.$$

- *Action on Morphisms.* Given a morphism of spans

$$\begin{array}{ccc} & R & \\ f_R \swarrow & \downarrow \phi & \searrow g_R \\ A & & B \\ \uparrow f_S & \downarrow g_S & \nearrow \\ & S & \end{array}$$

we have a corresponding inclusion of relations

$$\iota_{A,B}(\phi) : \iota_{A,B}(R) \subset \iota_{A,B}(S),$$

since we have $a \sim_{\iota_{A,B}(R)} b$ iff there exists $x \in R$ such that $a = f_R(x)$ and $b = g_R(x)$, in which case we then have

$$\begin{aligned} a &= f_R(x) \\ &= f_S(\phi(x)), \\ b &= g_R(x) \\ &= g_S(\phi(x)), \end{aligned}$$

so that $a \sim_{\iota_{A,B}(S)} b$, and thus $\iota_{A,B}(R) \subset \iota_{A,B}(S)$.

Proof. Omitted. □

6.8.3 Comparison to Relations: From **Rel** to **Span**

00RG Proposition 6.8.3.1.1. We have a lax functor

$$(\iota, \iota^2, \iota^0) : \mathbf{Rel} \rightarrow \mathbf{Span}$$

from **Rel** to **Span** where

- *Action on Objects.* For each $A \in \text{Obj}(\mathbf{Span})$, we have

$$\iota(A) \stackrel{\text{def}}{=} A;$$

- *Action on Hom-Categories.* For each $A, B \in \text{Obj}(\mathbf{Span})$, the action on Hom-categories

$$\iota_{A,B} : \mathbf{Rel}(A, B) \rightarrow \mathbf{Span}(A, B)$$

of ι at (A, B) is the functor where

- *Action on Objects.* Given a relation $R : A \rightarrow B$ from A to B , we define a span

$$\iota_{A,B}(R) : A \dashrightarrow B$$

from A to B by

$$\iota_{A,B}(R) \stackrel{\text{def}}{=} (R, \uparrow \text{pr}_1 R, \uparrow \text{pr}_2 R),$$

where $R \subset A \times B$ and $\uparrow \text{pr}_1 R$ and $\uparrow \text{pr}_2 R$ are the restriction of the projections

$$\begin{aligned} \text{pr}_1 &: A \times B \rightarrow A, \\ \text{pr}_2 &: A \times B \rightarrow B \end{aligned}$$

to R ;

- *Action on Morphisms.* Given an inclusion $\phi: R \subset S$ of relations, we have a corresponding morphism of spans

$$\iota_{A,B}(\phi): \iota_{A,B}(R) \rightarrow \iota_{A,B}(S)$$

as in the diagram

$$\begin{array}{ccc} & R & \\ \lceil \text{pr}_1 R & \swarrow & \searrow \lceil \text{pr}_2 R \\ A & & B \\ \lceil \text{pr}_1 S & \nearrow & \searrow \lceil \text{pr}_2 S \\ & S & \end{array}$$

- *The Lax Functoriality Constraints.* The lax functoriality constraint

$$\iota_{R,S}^2: \iota(S) \circ \iota(R) \Longrightarrow \iota(S \diamond R)$$

of ι at (R, S) is given by the morphism of spans from

$$\begin{array}{ccccc} & R \times_B S & & & \\ & \swarrow \text{pr}_1 \quad \searrow \text{pr}_2 & & & \\ R & & S & & \\ \lceil \text{pr}_1 R \circ \text{pr}_1 & & \lceil \text{pr}_2 S \circ \text{pr}_2 & & \\ \lceil \text{pr}_1 R & \swarrow \text{pr}_2 R & \nearrow \lceil \text{pr}_1 S & \searrow \text{pr}_2 S & \lceil \text{pr}_2 S \\ A & & B & & C \end{array}$$

to

$$\begin{array}{ccc} & S \diamond R & \\ & \swarrow \lceil \text{pr}_1 S \diamond R \quad \searrow \lceil \text{pr}_2 S \diamond R & \\ A & & C \end{array}$$

given by the natural inclusion $R \times_B S \hookrightarrow S \diamond R$, since we have

$$\begin{aligned} R \times_B S &= \{((a_R, b_R), (b_S, c_S)) \in R \times S \mid b_R = b_S\}; \\ S \diamond R &= \left\{ (a, c) \in A \times C \mid \begin{array}{l} \text{there exists some } b \in B \text{ such that} \\ (a, b) \in R \text{ and } (b, c) \in S \end{array} \right\}; \end{aligned}$$

- *The Lax Unity Constraints.* The lax unity constraint¹²

$$\iota_A^0: \underbrace{\text{id}_{\iota(A)}}_{(A, \text{id}_A, \text{id}_A)} \Longrightarrow \underbrace{\iota(\chi_A)}_{(\Delta_A, \lceil \text{pr}_1 \Delta_A, \lceil \text{pr}_2 \Delta_A)}$$

¹²Which is in fact strong, as δ_A is an isomorphism.

of ι at A is given by the diagonal morphism of A , as in the diagram

$$\begin{array}{ccc}
 & A & \\
 id_A \swarrow & \downarrow & \searrow id_A \\
 A & \delta_A & A. \\
 \uparrow pr_1 \Delta_A & \downarrow & \uparrow pr_2 \Delta_A \\
 & \Delta_A &
 \end{array}$$

Proof. Omitted. □

6.8.4 Comparison to Relations: The Wehrheim–Woodward Construction

6.8.5 Comparison to Multirelations

00RK Remark 6.8.5.1.1. The pseudofunctor of [Proposition 6.8.2.2.1](#) and the lax functor of [Proposition 6.8.3.1.1](#) fail to be equivalences of bicategories. This happens essentially because a span $(S, f, g): A \rightarrow B$ from A to B may relate elements $a \in A$ and $b \in B$ by more than one element, e.g. there could be $s \neq s' \in S$ such that $a = f(s) = f(s')$ and $b = g(s) = g(s')$.

Thus, in a sense, spans may be thought of as “relations with multiplicity”. And indeed, if instead of considering relations from A to B , i.e. functions

$$R: A \times B \rightarrow \{\text{true, false}\}$$

from $A \times B$ to $\{\text{true, false}\} \cong \{0, 1\}$, we consider functions

$$R: A \times B \rightarrow \mathbb{N} \cup \{\infty\}$$

from $A \times B$ to $\mathbb{N} \cup \{\infty\}$, then we obtain the notion of a **multirelation from A to B** , and these turn out to assemble together with sets into a bicategory MRel that is biequivalent to Span ; see [[some-algebraic-laws-for-spans-and-their-connections-with-multirelations](#)].

6.8.6 Comparison to Relations via Double Categories

00RM Remark 6.8.6.1.1. There are double functors between the double categories Rel^{dbl} and Span^{dbl} analogous to the functors of [Propositions 6.8.2.2.1](#) and [6.8.3.1.1](#), assembling moreover into a strict-lax adjunction of double functors; see [[higher-dimensional-categories](#)].

Appendices

6.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets
5. Relations
6. Spans
7. Posets

Indexed and Fibred Sets

7. Indexed Sets
8. Fibred Sets
9. Un/Straightening for Indexed and Fibred Sets

Category Theory

11. Categories
12. Types of Morphisms in Categories
13. Adjunctions and the Yoneda Lemma
14. Constructions With Categories
15. Profunctors
16. Cartesian Closed Categories
17. Kan Extensions

Bicategories

18. Bicategories
19. Internal Adjunctions

Internal Category Theory

20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

41. Stochastic Processes, Martingales, and Brownian Motion

Measure Theory

39. Measurable Spaces

42. Itô Calculus

40. Measures and Integration

43. Stochastic Differential Equations

Probability Theory

40. Probability Theory

Differential Geometry

44. Topological and Smooth Manifolds

Stochastic Analysis

Schemes

45. Schemes

6.2 Other Chapters

Sets

1. Sets

15. Profunctors

2. Constructions With Sets

16. Cartesian Closed Categories

3. Pointed Sets

17. Kan Extensions

4. Tensor Products of Pointed Sets

Bicategories

5. Relations

18. Bicategories

6. Spans

19. Internal Adjunctions

7. Posets

Internal Category Theory

Indexed and Fibred Sets

20. Internal Categories

7. Indexed Sets

Cyclic Stuff

8. Fibred Sets

21. The Cycle Category

9. Un/Straightening for Indexed and Fibred Sets

Cubical Stuff

Category Theory

22. The Cube Category

11. Categories

Globular Stuff

12. Types of Morphisms in Categories

23. The Globe Category

13. Adjunctions and the Yoneda Lemma

Cellular Stuff

14. Constructions With Categories

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero	37. Real Analysis in One Variable
27. Monoids With Zero	38. Real Analysis in Several Variables
28. Constructions With Monoids With Zero	Measure Theory
Groups	39. Measurable Spaces
29. Groups	40. Measures and Integration
30. Constructions With Groups	Probability Theory
Hyper Algebra	40. Probability Theory
31. Hypermonoids	Stochastic Analysis
32. Hypergroups	41. Stochastic Processes, Martingales, and Brownian Motion
33. Hypersemirings and Hyperrings	42. Itô Calculus
34. Quantales	43. Stochastic Differential Equations
Near-Rings	Differential Geometry
35. Near-Semirings	44. Topological and Smooth Manifolds
36. Near-Rings	Schemes
Real Analysis	45. Schemes

Chapter 7

Posets

00RN Rename this to “Preorders, Partial Orders, and Posets”.

Contents

7.1	Section	253
7.1.1	Section	253
7.A	Other Chapters	253
7.2	Other Chapters	255

7.1 Section

7.1.1 Section

7.1.1.1 Section

Appendices

7.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets
5. Relations
6. Spans

7. Posets

Indexed and Fibred Sets

7. Indexed Sets
8. Fibred Sets
9. Un/Straightening for Indexed and Fibred Sets

Category Theory

11. Categories

- 12. Types of Morphisms in Categories
- 13. Adjunctions and the Yoneda Lemma
- 14. Constructions With Categories
- 15. Profunctors
- 16. Cartesian Closed Categories
- 17. Kan Extensions
- Bicategories**
- 18. Bicategories
- 19. Internal Adjunctions
- Internal Category Theory**
- 20. Internal Categories
- Cyclic Stuff**
- 21. The Cycle Category
- Cubical Stuff**
- 22. The Cube Category
- Globular Stuff**
- 23. The Globe Category
- Cellular Stuff**
- 24. The Cell Category
- Monoids**
- 25. Monoids
- 26. Constructions With Monoids
- Monoids With Zero**
- 27. Monoids With Zero
- 28. Constructions With Monoids With Zero
- Groups**
- 29. Groups
- 30. Constructions With Groups
- Hyper Algebra**
- 31. Hypermonoids
- 32. Hypergroups
- 33. Hypersemirings and Hyperrings
- 34. Quantales
- Near-Rings**
- 35. Near-Semirings
- 36. Near-Rings
- Real Analysis**
- 37. Real Analysis in One Variable
- 38. Real Analysis in Several Variables
- Measure Theory**
- 39. Measurable Spaces
- 40. Measures and Integration
- Probability Theory**
- 40. Probability Theory
- Stochastic Analysis**
- 41. Stochastic Processes, Martingales, and Brownian Motion
- 42. Itô Calculus
- 43. Stochastic Differential Equations
- Differential Geometry**
- 44. Topological and Smooth Manifolds
- Schemes**
- 45. Schemes

7.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
Category Theory	27. Monoids With Zero
11. Categories	28. Constructions With Monoids With Zero
12. Types of Morphisms in Categories	Groups
13. Adjunctions and the Yoneda Lemma	29. Groups
14. Constructions With Categories	30. Constructions With Groups
15. Profunctors	Hyper Algebra
16. Cartesian Closed Categories	31. Hypermonoids
17. Kan Extensions	32. Hypergroups
Bicategories	33. Hypersemirings and Hyperrings
18. Bicategories	34. Quantales
19. Internal Adjunctions	Near-Rings
Internal Category Theory	35. Near-Semirings
20. Internal Categories	36. Near-Rings
	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part II

Indexed and Fibred Sets

Chapter 8

Indexed Sets

00RS This chapter contains a discussion of indexed sets, the set-theoretical counterpart to indexed categories. In particular, here we explore:

1. Indexed sets, i.e. functors $K_{\text{disc}} \rightarrow \text{Sets}$ with K a set;
2. The limits and colimits in the category of K -indexed sets;
3. Constructions with indexed sets like dependent sums, dependent products, and internal Hom.

Contents

8.1	Indexed Sets.....	259
8.1.1	Foundations	259
8.1.2	Morphisms of Indexed Sets	259
8.1.3	The Category of Sets Indexed by a Fixed Set.....	259
8.1.4	The Category of Indexed Sets	260
8.2	Limits of Indexed Sets	261
8.2.1	Products of K -Indexed Sets	261
8.2.2	Pullbacks of K -Indexed Sets	262
8.2.3	Equalisers of K -Indexed Sets.....	262
8.2.4	Products in ISets	262
8.2.5	Pullbacks in ISets	262
8.2.6	Equalisers in ISets	263
8.3	Colimits of Indexed Sets.....	263
8.3.1	Coproducts of K -Indexed Sets.....	263
8.3.2	Pushouts of K -Indexed Sets.....	263
8.3.3	Coequalisers of K -Indexed Sets	264
8.4	Constructions With Indexed Sets.....	264
8.4.1	Change of Indexing.....	264
8.4.2	Dependent Sums.....	265

8.4.3	Dependent Products.....	266
8.4.4	Internal Homs.....	267
8.4.5	Adjointness of Indexed Sets.....	267
8.A	Other Chapters	267
8.2	Other Chapters	269

8.1 Indexed Sets

8.1.1 Foundations

Let K be a set.

00RV Definition 8.1.1.1. A **K -indexed set** is a functor $X: K_{\text{disc}} \rightarrow \text{Sets}$.

00RW Remark 8.1.1.2. By ??, a **K -indexed set** consists of a K -indexed collection

$$X^\dagger: K \rightarrow \text{Obj}(\text{Sets}),$$

of sets, assigning a set $X_x^\dagger \stackrel{\text{def}}{=} X_x$ to each element x of K .

8.1.2 Morphisms of Indexed Sets

Let $X: K_{\text{disc}} \rightarrow \text{Sets}$ and $Y: K_{\text{disc}} \rightarrow \text{Sets}$ be indexed sets.

00RY Definition 8.1.2.1.1. A **morphism of K -indexed sets from X to Y** ¹ is a natural transformation

$$f: X \Longrightarrow Y, \quad K_{\text{disc}} \begin{array}{c} X \\ f \Downarrow \\ Y \end{array} \rightarrow \text{Sets}$$

from X to Y .

00RZ Remark 8.1.2.1.2. In detail, a **morphism of K -indexed sets** consists of a K -indexed collection

$$\{f_x: X_x \rightarrow Y_x\}_{x \in K}$$

of maps of sets.

8.1.3 The Category of Sets Indexed by a Fixed Set

Let K be a set.

¹Further Terminology: Also called a **K -indexed map of sets from X to Y** .

00S1 Definition 8.1.3.1.1. The **category of K -indexed sets** is the category $\text{ISets}(K)$ defined by

$$\text{ISets}(K) \stackrel{\text{def}}{=} \text{Fun}(K_{\text{disc}}, \text{Sets}).$$

00S2 Remark 8.1.3.1.2. In detail, the **category of K -indexed sets** is the category $\text{ISets}(K)$ where

- *Objects.* The objects of $\text{ISets}(K)$ are K -indexed sets as in [Definition 8.1.1.1.1](#);
- *Morphisms.* The morphisms of $\text{ISets}(K)$ are morphisms of K -indexed sets as in [Definition 8.1.2.1.1](#);
- *Identities.* For each $X \in \text{Obj}(\text{ISets}(K))$, the unit map

$$\mathbb{1}_X^{\text{ISets}(K)} : \text{pt} \rightarrow \text{Hom}_{\text{ISets}(K)}(X, X)$$

of $\text{ISets}(K)$ at X is defined by

$$\text{id}_X^{\text{ISets}(K)} \stackrel{\text{def}}{=} \{\text{id}_{X_x}\}_{x \in K};$$

- *Composition.* For each $X, Y, Z \in \text{Obj}(\text{ISets}(K))$, the composition map

$$\circ^{\text{ISets}(K)}_{X, Y, Z} : \text{Hom}_{\text{ISets}(K)}(Y, Z) \times \text{Hom}_{\text{ISets}(K)}(X, Y) \rightarrow \text{Hom}_{\text{ISets}(K)}(X, Z)$$

of $\text{ISets}(K)$ at (X, Y, Z) is defined by

$$\{g_x\}_{x \in K} \circ^{\text{ISets}(K)}_{X, Y, Z} \{f_x\}_{x \in K} \stackrel{\text{def}}{=} \{g_x \circ f_x\}_{x \in K}.$$

8.1.4 The Category of Indexed Sets

00S4 Definition 8.1.4.1.1. The **category of indexed sets** is the category ISets defined as the Grothendieck construction of the functor $\text{ISets} : \text{Sets}^{\text{op}} \rightarrow \text{Cats}$ of [Proposition 8.4.1.4](#):

$$\text{ISets} \stackrel{\text{def}}{=} \int^{\text{Sets}} \text{ISets}.$$

00S5 Remark 8.1.4.1.2. In detail, the **category of indexed sets** is the category ISets where

- *Objects.* The objects of ISets are pairs (K, X) consisting of
 - *The Indexing Set.* A set K ;
 - *The Indexed Set.* A K -indexed set $X : K_{\text{disc}} \rightarrow \text{Sets}$;
- *Morphisms.* A morphism of ISets from (K, X) to (K', Y) is a pair (ϕ, f) consisting of
 - *The Reindexing Map.* A map of sets $\phi : K \rightarrow K'$;
 - *The Morphism of Indexed Sets.* A morphism of K -indexed sets $f : X \rightarrow$

$\phi_*(Y)$ as in the diagram

$$f: X \rightarrow \phi_*(Y), \quad \begin{array}{ccc} K_{\text{disc}} & \xrightarrow{\phi} & K'_{\text{disc}} \\ & \searrow f \quad \nearrow Y & \\ X & & Y \\ & & \text{Sets;} \end{array}$$

- *Identities.* For each $(K, X) \in \text{Obj}(\text{ISets})$, the unit map

$$\mathbb{1}_{(K,X)}^{\text{ISets}}: \text{pt} \rightarrow \text{ISets}((K,X), (K,X))$$

of ISets at (K, X) is defined by

$$\text{id}_{(K,X)}^{\text{ISets}} \stackrel{\text{def}}{=} (\text{id}_K, \text{id}_X).$$

- *Composition.* For each $\mathbf{X} = (K, X), \mathbf{Y} = (K', Y), \mathbf{Z} = (K'', Z) \in \text{Obj}(\text{ISets})$, the composition map

$$\circ_{\mathbf{X}, \mathbf{Y}, \mathbf{Z}}^{\text{ISets}}: \text{ISets}(\mathbf{Y}, \mathbf{Z}) \times \text{ISets}(\mathbf{X}, \mathbf{Y}) \rightarrow \text{ISets}(\mathbf{X}, \mathbf{Z})$$

of ISets at $(\mathbf{X}, \mathbf{Y}, \mathbf{Z})$ is defined by

$$(\psi, g) \circ (\phi, f) \stackrel{\text{def}}{=} (\psi \circ \phi, (g \star \text{id}_\phi) \circ f),$$

as in the diagram

$$\begin{array}{ccccc} K_{\text{disc}} & \xrightarrow{\phi} & K'_{\text{disc}} & \xrightarrow{\psi} & K''_{\text{disc}} \\ & \searrow f \quad \nearrow Y & \downarrow & \nearrow g \quad \searrow Z & \\ X & & Y & & Z \\ & & & & \text{Sets} \end{array}$$

for each $(\phi, f) \in \text{ISets}(\mathbf{X}, \mathbf{Y})$ and each $(\psi, g) \in \text{ISets}(\mathbf{Y}, \mathbf{Z})$.

8.2 Limits of Indexed Sets

8.2.1 Products of K -Indexed Sets

Let $X: K_{\text{disc}} \rightarrow \text{Sets}$ and $Y: K_{\text{disc}} \rightarrow \text{Sets}$ be indexed sets.

0058 Definition 8.2.1.1.1. The **product of X and Y** is the K -indexed set $X \times Y: K_{\text{disc}} \rightarrow \text{Sets}$ defined by

$$(X \times Y)_k \stackrel{\text{def}}{=} X_k \times Y_k$$

for each $k \in K$.

Proof. That this agrees with the categorical product in $\text{ISets}(K)$ follows from ?? of ?? \square

8.2.2 Pullbacks of K -Indexed Sets

Let $X, Y, Z: K_{\text{disc}} \rightarrow \text{Sets}$ be K -indexed sets and let $f: X \rightarrow Z$ and $g: Y \rightarrow Z$ be morphisms of K -indexed sets.

00SA Definition 8.2.2.1.1. The **pullback of X and Y over Z** is the K -indexed set $X \times_Z Y: K_{\text{disc}} \rightarrow \text{Sets}$ defined by

$$(X \times_Z Y)_k \stackrel{\text{def}}{=} X_k \times_{Z_k} Y_k$$

for each $k \in K$.

Proof. That this agrees with the categorical pullback in $\text{ISets}(K)$ follows from ?? of ??.

□

8.2.3 Equalisers of K -Indexed Sets

Let $X, Y: K_{\text{disc}} \rightarrow \text{Sets}$ be K -indexed sets and let $f, g: X \rightrightarrows Y$ be morphisms of K -indexed sets.

00SC Definition 8.2.3.1.1. The **equaliser of f and g** is the K -indexed set $\text{Eq}(f, g): K_{\text{disc}} \rightarrow \text{Sets}$ defined by

$$(\text{Eq}(f, g))_k \stackrel{\text{def}}{=} \text{Eq}(f_k, g_k)$$

for each $k \in K$.

Proof. That this agrees with the categorical equaliser in $\text{ISets}(K)$ follows from ?? of ??.

□

8.2.4 Products in ISets

Let $X: K_{\text{disc}} \rightarrow \text{Sets}$ and $Y: K'_{\text{disc}} \rightarrow \text{Sets}$ be indexed sets.

00SE Definition 8.2.4.1.1. The **product of X and Y** is the $(K \times K')$ -indexed set

$$X \times Y: (K \times K')_{\text{disc}} \rightarrow \text{Sets}$$

defined by

$$(X \times Y)_{(k, k')} \stackrel{\text{def}}{=} X_k \times Y_{k'}$$

for each $(k, k') \in K \times K'$.

Proof. We claim that this agrees with the categorical product in ISets .

□

8.2.5 Pullbacks in ISets

Let $X: K_{\text{disc}} \rightarrow \text{Sets}$ be a K -indexed set, let $Y: K'_{\text{disc}} \rightarrow \text{Sets}$ be a K' -indexed set, let $Z: K''_{\text{disc}} \rightarrow \text{Sets}$ be a K'' -indexed set, and let $(\phi, f): X \rightarrow Z$ and $(\psi, g): Y \rightarrow Z$ be morphisms of indexed sets (as in Remark 8.1.4.1.2).

00SG Definition 8.2.5.1.1. The **pullback of X and Y over Z** is the $(K \times_{K''} K)$ -indexed set

$$X \times_Z Y: (K \times_{K''} K)_{\text{disc}} \rightarrow \text{Sets}$$

defined by

$$\begin{aligned} (X \times_Z Y)_{(k,k')} &\stackrel{\text{def}}{=} X_k \times_{Z_{\phi(k)}} Y_{k'} \\ &\stackrel{\text{def}}{=} X_k \times_{Z_{\psi(k)}} Y_{k'} \end{aligned}$$

for each $(k, k') \in K \times_{K''} K'$.

Proof. We claim that this agrees with the categorical pullback in ISets . \square

8.2.6 Equalisers in ISets

Let $X: K_{\text{disc}} \rightarrow \text{Sets}$ be a K -indexed set, let $Y: K'_{\text{disc}} \rightarrow \text{Sets}$ be a K' -indexed set, and let $(\phi, f), (\psi, g): X \rightarrow Y$ be morphisms of indexed sets (as in Remark 8.1.4.1.2).

00SJ Definition 8.2.6.1.1. The **equaliser of (ϕ, f) and (ψ, g)** is the $\text{Eq}(\phi, \psi)$ -indexed set $\text{Eq}(f, g): \text{Eq}(\phi, \psi) \rightarrow \text{Sets}$ defined by

$$(\text{Eq}(f, g))_k \stackrel{\text{def}}{=} \text{Eq}(f_k, g_k)$$

for each $k \in \text{Eq}(\phi, \psi)$.

Proof. We claim that this agrees with the categorical equaliser in ISets . \square

8.3 Colimits of Indexed Sets

8.3.1 Coproducts of K -Indexed Sets

Let $X: K_{\text{disc}} \rightarrow \text{Sets}$ and $Y: K_{\text{disc}} \rightarrow \text{Sets}$ be indexed sets.

00SM Definition 8.3.1.1.1. The **coproduct** of X and Y is the K -indexed set $X \coprod Y: K_{\text{disc}} \rightarrow \text{Sets}$ defined by

$$(X \coprod Y)_k \stackrel{\text{def}}{=} X_k \coprod Y_k$$

for each $k \in K$.

Proof. That this agrees with the categorical coproduct in $\text{ISets}(K)$ follows from ?? of ??.

8.3.2 Pushouts of K -Indexed Sets

Let $X, Y, Z: K_{\text{disc}} \rightarrow \text{Sets}$ be K -indexed sets and let $f: Z \rightarrow X$ and $g: Z \rightarrow Y$ be morphisms of K -indexed sets.

00SP Definition 8.3.2.1.1. The **pushout** of X and Y is the K -indexed set $X \coprod_Z Y: K_{\text{disc}} \rightarrow \text{Sets}$ defined by

$$(X \coprod_Z Y)_k \stackrel{\text{def}}{=} X_k \coprod_{Z_k} Y_k$$

for each $k \in K$.

Proof. That this agrees with the categorical pushout in $\text{ISets}(K)$ follows from ?? of ??.

8.3.3 Coequalisers of K -Indexed Sets

Let $X, Y: K_{\text{disc}} \rightarrow \text{Sets}$ be K -indexed sets and let $f, g: X \rightrightarrows Y$ be morphisms of K -indexed sets.

00SR Definition 8.3.3.1.1. The **coequaliser** of X and Y is the K -indexed set $\text{CoEq}(f, g): K_{\text{disc}} \rightarrow \text{Sets}$ defined by

$$(\text{CoEq}(f, g))_k \stackrel{\text{def}}{=} \text{CoEq}(f_k, g_k)$$

for each $k \in K$.

Proof. That this agrees with the categorical coequaliser in $\text{ISets}(K)$ follows from ?? of ??.

□

8.4 Constructions With Indexed Sets

8.4.1 Change of Indexing

Let $\phi: K \rightarrow K'$ be a function and let X be a K' -indexed set.

00SU Definition 8.4.1.1.1. The **change of indexing of X to K** is the K -indexed set $\phi^*(X)$ defined by

$$\phi^*(X) \stackrel{\text{def}}{=} X \circ \phi_{\text{disc}}.$$

00SV Remark 8.4.1.1.2. In detail, the **change of indexing of X to K** is the K -indexed set $\phi^*(X)$ defined by

$$\phi^*(X)_x \stackrel{\text{def}}{=} X_{\phi(x)}$$

for each $x \in K$.

00SW Proposition 8.4.1.1.3. The assignment $X \mapsto \phi^*(X)$ defines a functor

$$\phi^*: \text{ISets}(K') \rightarrow \text{ISets}(K),$$

where

- *Action on Objects.* For each $X \in \text{Obj}(\text{ISets}(K'))$, we have

$$[\phi^*](X) \stackrel{\text{def}}{=} \phi^*(X);$$

- *Action on Morphisms.* For each $X, Y \in \text{Obj}(\text{ISets}(K'))$, the action on Hom-sets

$$\phi_{X,Y}^*: \text{Hom}_{\text{ISets}(K')}(X, Y) \rightarrow \text{Hom}_{\text{ISets}(K)}(\phi^*(X), \phi^*(Y))$$

of ϕ^* at (X, Y) is the map sending a morphism of K' -indexed sets

$$f = \{f_x: X_x \rightarrow Y_x\}_{x \in K'}$$

from X to Y to the morphism of K -indexed sets defined by

$$\phi^*(f) \stackrel{\text{def}}{=} \{f_{\phi(x)}: X_{\phi(x)} \rightarrow Y_{\phi(x)}\}_{x \in K}.$$

Proof. Omitted. □

00SX Proposition 8.4.1.1.4. The assignment $K \mapsto \text{ISets}(K)$ defines a functor

$$\text{ISets}: \text{Sets}^{\text{op}} \rightarrow \text{Cats},$$

where

- *Action on Objects.* For each $K \in \text{Obj}(\text{Sets})$, we have

$$[\text{ISets}](K) \stackrel{\text{def}}{=} \text{ISets}(K);$$

- *Action on Morphisms.* For each $K, K' \in \text{Obj}(\text{Sets})$, the action on Hom-sets

$$\text{ISets}_{K,K'}: \text{Sets}^{\text{op}}(K, K') \rightarrow \text{Fun}(\text{ISets}(K), \text{ISets}(K'))$$

of ISets at (K, K') is the map defined by

$$\text{ISets}_{K,K'}(\phi) \stackrel{\text{def}}{=} \phi^*$$

for each $\phi \in \text{Sets}^{\text{op}}(K, K')$.

Proof. Omitted. □

8.4.2 Dependent Sums

Let $\phi: K \rightarrow K'$ be a function and let X be a K -indexed set.

00SZ Definition 8.4.2.1.1. The **dependent sum** of X is the K' -indexed set $\Sigma_\phi(X)$ ² defined by

$$\Sigma_\phi(X) \stackrel{\text{def}}{=} \text{Lan}_\phi(X),$$

and hence given by

$$\Sigma_\phi(X)_x \cong \coprod_{y \in \phi^{-1}(x)} X_y$$

for each $x \in K'$.

00T0 Proposition 8.4.2.1.2. The assignment $X \mapsto \Sigma_\phi(X)$ defines a functor

$$\Sigma_\phi: \text{ISets}(K) \rightarrow \text{ISets}(K'),$$

where

- *Action on Objects.* For each $X \in \text{Obj}(\text{ISets}(K))$, we have

$$[\Sigma_\phi](X) \stackrel{\text{def}}{=} \Sigma_\phi(X);$$

²Further Notation: Also written $\phi_*(X)$.

- *Action on Morphisms.* For each $X, Y \in \text{Obj}(\text{ISets}(K))$, the action on Hom-sets

$$\Sigma_{\phi|X,Y}: \text{Hom}_{\text{ISets}(K)}(X, Y) \rightarrow \text{Hom}_{\text{ISets}(K')}\left(\Sigma_{\phi}(X), \Sigma_{\phi}(Y)\right)$$

of Σ_{ϕ} at (X, Y) is the map sending a morphism of K -indexed sets

$$f: X \rightarrow Y$$

to the morphism of K' -indexed sets defined by

$$\begin{aligned} \Sigma_{\phi}(f) &\stackrel{\text{def}}{=} \text{Lan}_{\phi}(f); \\ &\cong \coprod_{y \in \phi^{-1}(X)} f_y. \end{aligned}$$

Proof. Omitted. □

8.4.3 Dependent Products

Let $\phi: K \rightarrow K'$ be a function and let X be a K -indexed set.

00T2 Definition 8.4.3.1.1. The **dependent product** of X is the K' -indexed set $\Pi_{\phi}(X)$ ³ defined by

$$\Pi_{\phi}(X) \stackrel{\text{def}}{=} \text{Ran}_{\phi}(X),$$

and hence given by

$$\Pi_{\phi}(X)_x \cong \prod_{y \in \phi^{-1}(x)} X_y$$

for each $x \in K'$.

00T3 Proposition 8.4.3.1.2. The assignment $X \mapsto \Pi_{\phi}(X)$ defines a functor

$$\Pi_{\phi}: \text{ISets}(K) \rightarrow \text{ISets}(K'),$$

where

- *Action on Objects.* For each $X \in \text{Obj}(\text{ISets}(K))$, we have

$$[\Pi_{\phi}](X) \stackrel{\text{def}}{=} \Pi_{\phi}(X);$$

- *Action on Morphisms.* For each $X, Y \in \text{Obj}(\text{ISets}(K))$, the action on Hom-sets

$$\Pi_{\phi|X,Y}: \text{Hom}_{\text{ISets}(K)}(X, Y) \rightarrow \text{Hom}_{\text{ISets}(K')}\left(\Pi_{\phi}(X), \Pi_{\phi}(Y)\right)$$

of Π_{ϕ} at (X, Y) is the map sending a morphism of K -indexed sets

$$f: X \rightarrow Y$$

³Further Notation: Also written $\phi_!(X)$.

to the morphism of K' -indexed sets defined by

$$\begin{aligned}\Pi_\phi(f) &\stackrel{\text{def}}{=} \text{Ran}_\phi(f); \\ &\cong \prod_{y \in \phi^{-1}(x)} f_y.\end{aligned}$$

Proof. Omitted. \square

8.4.4 Internal Hom

Let K be a set and let X and Y be K -indexed sets.

00T5 Definition 8.4.4.1.1. The **internal Hom of indexed sets from X to Y** is the indexed set $\mathbf{Hom}_{\text{ISets}(K)}(X, Y)$ defined by

$$\mathbf{Hom}_{\text{ISets}(K)}(X, Y) \stackrel{\text{def}}{=} \text{Sets}(X_x, Y_x)$$

for each $x \in K$.

8.4.5 Adjointness of Indexed Sets

Let $\phi: K \rightarrow K'$ be a map of sets.

00T7 Proposition 8.4.5.1.1. We have a triple adjunction

$$(\Sigma_\phi \dashv \phi^* \dashv \Pi_\phi): \quad \text{ISets}(K) \begin{array}{c} \xleftarrow{\perp} \\[-1ex] \xleftarrow{\phi^*} \\[-1ex] \xleftarrow{\perp} \end{array} \text{ISets}(K').$$

Σ_ϕ
 Π_ϕ

Proof. This follows from ?? of ??.

\square

Appendices

8.A Other Chapters

Sets

- 1. Sets
- 2. Constructions With Sets
- 3. Pointed Sets
- 4. Tensor Products of Pointed Sets

Relations

- 6. Spans
- 7. Posets
- 7. Indexed Sets

Indexed and Fibred Sets

8. Fibred Sets

9. Un/Straightening for Indexed and Fibred Sets

Category Theory

11. Categories

12. Types of Morphisms in Categories

13. Adjunctions and the Yoneda Lemma

14. Constructions With Categories

15. Profunctors

16. Cartesian Closed Categories

17. Kan Extensions

Bicategories

18. Bicategories

19. Internal Adjunctions

Internal Category Theory

20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales, and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

8.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 9

Fibred Sets

00T8 This chapter contains a discussion of the un/straightening equivalence in the context of sets, as well as a general discussion of indexed and fibred sets. In particular, it contains:

1. A discussion of indexed sets (i.e. functors $K_{\text{disc}} \rightarrow \text{Sets}$ with K a set), constructions with them like dependent sums and dependent products, and their properties (????);
2. A discussion of fibred sets (i.e. maps of sets $X \rightarrow K$), constructions with them like dependent sums and dependent products, and their properties ([Sections 9.1](#) and [9.2](#));
3. A discussion of the un/straightening equivalence for indexed and fibred sets (??).

Contents

9.1	Fibred Sets	272
9.1.1	Foundations	272
9.1.2	Morphisms of Fibred Sets	272
9.1.3	The Category of Fibred Sets Over a Fixed Base	273
9.1.4	The Category of Fibred Sets	274
9.2	Constructions With Fibred Sets	275
9.2.1	Change of Base	275
9.2.2	Dependent Sums	277
9.2.3	Dependent Products	278
9.2.4	Internal Homs	282
9.2.5	Adjointness for Fibred Sets	283
9.A	Other Chapters	294
9.2	Other Chapters	296

9.1 Fibred Sets

9.1.1 Foundations

Let K be a set.

00TB Definition 9.1.1.1. A **K -fibred set** is a pair (X, ϕ) consisting of¹

- *The Underlying Set.* A set X , called the **underlying set of** (X, ϕ) ;
- *The Fibration.* A map of sets $\phi: X \rightarrow K$.

9.1.2 Morphisms of Fibred Sets

00TD Definition 9.1.2.1.1. A **morphism of K -fibred sets from** (X, ϕ) **to** (Y, ψ) is a function $f: X \rightarrow Y$ such that the diagram²

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \phi \searrow & & \swarrow \psi \\ & K & \end{array}$$

commutes.

¹Further Terminology: The **fibre of** (X, ϕ) **over** $x \in K$ is the set $\phi^{-1}(x)$ (also written ϕ_x) defined by

$$\begin{array}{ccc} \phi^{-1}(x) & \xrightarrow{\quad} & X \\ \downarrow & \lrcorner & \downarrow \phi \\ \text{pt} & \xrightarrow{[x]} & K. \end{array}$$

$\phi^{-1}(x) \stackrel{\text{def}}{=} \text{pt} \times_{[x], K, \phi} X,$

²Further Terminology: The **transport map associated to** f **at** $x \in K$ is the function

$$f_x^*: \phi^{-1}(x) \rightarrow \psi^{-1}(x)$$

given by the dashed map in the diagram

$$\begin{array}{ccccc} \phi^{-1}(x) & \xrightarrow{\quad} & X & \xrightarrow{f} & Y \\ \downarrow & \lrcorner & \downarrow \phi & \downarrow & \downarrow \psi \\ \text{pt} & \xrightarrow{[x]} & K & \xrightarrow{\quad} & K \\ \text{pt} & \xrightarrow{[x]} & K & \xrightarrow{\quad} & K \end{array}$$

$\psi^{-1}(x) \xrightarrow{\quad} \text{pt} \times_{[x], K, \phi} X \xrightarrow{\quad} Y$

9.1.3 The Category of Fibred Sets Over a Fixed Base

00TF Definition 9.1.3.1.1. The **category of K -fibred sets** is the category $\text{FibSets}(K)$ defined as the slice category $\text{Sets}_{/K}$ of Sets over K :

$$\text{FibSets}(K) \stackrel{\text{def}}{=} \text{Sets}_{/K}.$$

00TG Remark 9.1.3.1.2. In detail $\text{FibSets}(K)$ is the category where

- *Objects.* The objects of $\text{FibSets}(K)$ are pairs (X, ϕ) consisting of
 - *The Fibred Set.* A set X ;
 - *The Fibration.* A function $\phi: X \rightarrow K$;
- *Morphisms.* A morphism of $\text{FibSets}(K)$ from (X, ϕ) to (Y, ψ) is a function $f: X \rightarrow Y$ making the diagram

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \phi \searrow & & \swarrow \psi \\ & X & \end{array}$$

commute;

- *Identities.* For each $(X, \phi) \in \text{Obj}(\text{FibSets}(K))$, the unit map

$$\text{id}_{(X, \phi)}^{\text{FibSets}(K)}: \text{pt} \rightarrow \text{Hom}_{\text{FibSets}(K)}((X, \phi), (X, \phi))$$

of $\text{FibSets}(K)$ at (X, ϕ) is given by

$$\text{id}_{(X, \phi)}^{\text{FibSets}(K)} \stackrel{\text{def}}{=} \text{id}_X,$$

as witnessed by the commutativity of the diagram

$$\begin{array}{ccc} X & \xrightarrow{\text{id}_X} & X \\ \phi \searrow & & \swarrow \phi \\ & K & \end{array}$$

in Sets ;

- *Composition.* For each $\mathbf{X} = (X, \phi), \mathbf{Y} = (Y, \psi), \mathbf{Z} = (Z, \chi) \in \text{Obj}(\text{FibSets}(K))$, the composition map

$$\circ_{\mathbf{X}, \mathbf{Y}, \mathbf{Z}}^{\text{FibSets}(K)}: \text{Hom}_{\text{FibSets}(K)}(\mathbf{Y}, \mathbf{Z}) \times \text{Hom}_{\text{FibSets}(K)}(\mathbf{X}, \mathbf{Y}) \rightarrow \text{Hom}_{\text{FibSets}(K)}(\mathbf{X}, \mathbf{Z})$$

of $\text{FibSets}(K)$ at $(\mathbf{X}, \mathbf{Y}, \mathbf{Z})$ is defined by

$$\circ_{\mathbf{X}, \mathbf{Y}, \mathbf{Z}}^{\text{FibSets}(K)} \stackrel{\text{def}}{=} \circ_{X, Y, Z}^{\text{Sets}},$$

as witnessed by the commutativity of the diagram

$$\begin{array}{ccccc} X & \xrightarrow{f} & Y & \xrightarrow{g} & Z \\ & \searrow \phi & \downarrow \psi & \swarrow \chi & \\ & & K & & \end{array}$$

in Sets.

9.1.4 The Category of Fibred Sets

00TJ Definition 9.1.4.1.1. The **category of fibred sets** is the category FibSets defined as the Grothendieck construction of the functor $\text{FibSets}: \text{Sets}^{\text{op}} \rightarrow \text{Cats}$ of [Proposition 9.2.1.1.3](#):

$$\text{FibSets} \stackrel{\text{def}}{=} \int^{\text{Sets}} \text{FibSets}.$$

00TK Remark 9.1.4.1.2. In detail, the **category of fibred sets** is the category FibSets where

- **Objects.** The objects of FibSets are pairs $(K, (X, \phi_X))$ consisting of
 - *The Base Set.* A set K ;
 - *The Fibred Set.* A K -fibred set $\phi_X: X \rightarrow K$;
- **Morphisms.** A morphism of FibSets from $(K, (X, \phi_X))$ to $(K', (Y, \phi_Y))$ is a pair (ϕ, f) consisting of
 - *The Base Map.* A map of sets $\phi: K \rightarrow K'$;
 - *The Morphism of Fibred Sets.* A morphism of K -fibred sets

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \times_{K'} K \\ f: (X, \phi_X) \rightarrow \phi_Y^*(Y), & \phi_X \searrow & \swarrow \text{pr}_2 \\ & & K; \end{array}$$

- **Identities.** For each $(K, X) \in \text{Obj}(\text{FibSets})$, the unit map

$$\mu_{(K,X)}^{\text{FibSets}}: \text{pt} \rightarrow \text{FibSets}((K, X), (K, X))$$

of FibSets at (K, X) is defined by

$$\text{id}_{(K,X)}^{\text{FibSets}} \stackrel{\text{def}}{=} (\text{id}_K, \sim),$$

where \sim is the isomorphism $X \rightarrow X \times_K K$ as in the diagram

$$\begin{array}{ccc} X & \xrightarrow{\sim} & X \times_K K \\ \phi_X \searrow & & \swarrow \text{pr}_2 \\ & & K; \end{array}$$

- *Composition.* For each $\mathbf{X} = (K, X), \mathbf{Y} = (K', Y), \mathbf{Z} = (K'', Z) \in \text{Obj}(\text{FibSets})$, the composition map

$$\circ^{\text{FibSets}}_{\mathbf{X}, \mathbf{Y}, \mathbf{Z}} : \text{FibSets}(\mathbf{Y}, \mathbf{Z}) \times \text{FibSets}(\mathbf{X}, \mathbf{Y}) \rightarrow \text{FibSets}(\mathbf{X}, \mathbf{Z})$$

of FibSets at $(\mathbf{X}, \mathbf{Y}, \mathbf{Z})$ is defined by

$$g \circ^{\text{FibSets}}_{\mathbf{X}, \mathbf{Y}, \mathbf{Z}} f \stackrel{\text{def}}{=} (g \times_{K'} \text{id}_K) \circ f$$

as in the diagram

$$\begin{array}{ccccc} & & & \cong Z \times_{K''} K & \\ & & & \nearrow & \\ X & \xrightarrow{f} & Y \times_{K'} K & \xrightarrow{g \times_{K'} \text{id}_K} & (Z \times_{K''} K') \times_{K'} K \\ & \searrow \phi_X & \downarrow \text{pr}_2 & & \swarrow \text{pr}_2 \\ & & K; & & \end{array}$$

for each $f \in \text{Obj}(\text{FibSets}(\mathbf{X}, \mathbf{Y}))$ and each $g \in \text{Obj}(\text{FibSets}(\mathbf{Y}, \mathbf{Z}))$.

9.2 Constructions With Fibred Sets

9.2.1 Change of Base

Let $f : K \rightarrow K'$ be a function and let (X, ϕ_X) be a K' -fibred set.

00TN Definition 9.2.1.1.1. The **change of base of** (X, ϕ_X) **to** K is the K -fibred set $f^*(X)$ defined by

$$f^*(X) \stackrel{\text{def}}{=} (K \times_{K'} X, \text{pr}_1), \quad \begin{array}{ccc} f^*(X) & \xrightarrow{\text{pr}_2} & X \\ \text{pr}_1 \downarrow & \lrcorner & \downarrow \phi_X \\ K & \xrightarrow{f} & K'. \end{array}$$

00TP Proposition 9.2.1.1.2. The assignment $X \mapsto f^*(X)$ defines a functor

$$f^* : \text{FibSets}(K') \rightarrow \text{FibSets}(K),$$

where

- *Action on Objects.* For each $(X, \phi_X) \in \text{Obj}(\text{FibSets}(K'))$, we have

$$f^*(X, \phi_X) \stackrel{\text{def}}{=} f^*(X);$$

- *Action on Morphisms.* For each $(X, \phi_X), (Y, \phi_Y) \in \text{Obj}(\text{FibSets}(K'))$, the action on Hom-sets

$$f_{X,Y}^*: \text{Hom}_{\text{FibSets}(K')}(X, Y) \rightarrow \text{Hom}_{\text{FibSets}(K)}(f^*(X), f^*(Y))$$

of f^* at $((X, \phi_X), (Y, \phi_Y))$ is the map sending a morphism of K' -fibred sets

$$g: (X, \phi_X) \rightarrow (Y, \phi_Y)$$

to the morphism of K -fibred sets given by the dashed morphism in the diagram

$$\begin{array}{ccccc} f^*(X) & \xrightarrow{\quad} & X & \xrightarrow{\quad g \quad} & \\ \downarrow & \lrcorner & \downarrow \phi_X & \lrcorner & \downarrow \\ f^*(Y) & \xrightarrow{\quad} & Y & & \\ \downarrow & \lrcorner & \downarrow \phi_Y & & \\ K & \xrightarrow{f} & K' & & \\ \parallel & & \parallel & & \\ K & \xrightarrow{f} & K' & & \end{array}$$

Proof. Omitted. □

00TQ Proposition 9.2.1.1.3. The assignment $K \mapsto \text{FibSets}(K)$ defines a functor

$$\text{FibSets}: \text{Sets}^{\text{op}} \rightarrow \text{Cats},$$

where

- *Action on Objects.* For each $K \in \text{Obj}(\text{Sets})$, we have

$$[\text{FibSets}](K) \stackrel{\text{def}}{=} \text{FibSets}(K);$$

- *Action on Morphisms.* For each $K, K' \in \text{Obj}(\text{Sets})$, the action on Hom-sets

$$\text{Sets}_{/(-)|K,K'}: \text{Sets}^{\text{op}}(K, K') \rightarrow \text{Fun}(\text{FibSets}(K), \text{FibSets}(K'))$$

of $\text{Sets}_{/(-)}$ at (K, K') is the map sending a map of sets $f: K \rightarrow K'$ to the functor

$$\text{Sets}_{/f}: \text{FibSets}(K') \rightarrow \text{FibSets}(K)$$

defined by

$$\text{Sets}_{/f} \stackrel{\text{def}}{=} f^*.$$

Proof. Omitted. □

9.2.2 Dependent Sums

Let $f: K \rightarrow K'$ be a function and let (X, ϕ_X) be a K -fibred set.

00TS Definition 9.2.2.1.1. The **dependent sum**³ of (X, ϕ_X) is the K' -fibred set $\Sigma_f(X)$ ⁴ defined by

$$\begin{aligned}\Sigma_f(X) &\stackrel{\text{def}}{=} (\Sigma_f(X), \Sigma_f(\phi_X)) \\ &\stackrel{\text{def}}{=} (X, f \circ \phi_X).\end{aligned}$$

00TT Proposition 9.2.2.1.2. Let $f: K \rightarrow K'$ be a function.

00TU 1. *Functoriality.* The assignment $X \mapsto \Sigma_f(X)$ defines a functor

$$\Sigma_f: \text{FibSets}(K) \rightarrow \text{FibSets}(K'),$$

where

- *Action on Objects.* For each $(X, \phi_X) \in \text{Obj}(\text{FibSets}(K))$, we have

$$\Sigma_f(X, \phi_X) \stackrel{\text{def}}{=} (\Sigma_f(X), \Sigma_f(\phi_X));$$

- *Action on Morphisms.* For each $(X, \phi_X), (Y, \phi_Y) \in \text{Obj}(\text{FibSets}(K))$, the action on Hom-sets

$$\Sigma_{f|X,Y}: \text{Hom}_{\text{FibSets}(K)}(X, Y) \rightarrow \text{Hom}_{\text{FibSets}(K)}(\Sigma_f(X), \Sigma_f(Y))$$

of Σ_f at $((X, \phi_X), (Y, \phi_Y))$ is the map sending a morphism of K -fibred sets

$$g: (X, \phi_X) \rightarrow (Y, \phi_Y)$$

to the morphism of K' -fibred sets defined by

$$\Sigma_f(g) \stackrel{\text{def}}{=} g.$$

00TV 2. *Interaction With Fibres.* We have a bijection of sets

$$\Sigma_f(\phi_X)^{-1}(k') \cong \coprod_{k \in f^{-1}(k')} \phi_X^{-1}(k)$$

for each $k' \in K'$.

³The name “dependent sum” comes from the fact that the fibre $\Sigma_f(\phi_X)^{-1}(x)$ of $\Sigma_f(X)$ at $x \in K'$ is given by

$$\Sigma_f(\phi_X)^{-1}(x) \cong \coprod_{y \in f^{-1}(x)} \phi_X^{-1}(y);$$

see [Item 2 of Proposition 9.2.2.1.2](#).

⁴Further Notation: Also written $f_*(X)$.

Proof. **Item 1, Functoriality:** Omitted.

Item 2, Interaction With Fibres: Indeed, we have

$$\begin{aligned}\Sigma_f(\phi_X)^{-1}(k') &\stackrel{\text{def}}{=} \text{pt} \times_{[k'], K', f \circ \phi_X} X \\ &\cong \{x \in X \mid f(\phi_X(x)) = k'\} \\ &\cong \coprod_{k \in f^{-1}(k')} \{x \in X \mid \phi_X(x) = k\} \\ &\cong \coprod_{k \in f^{-1}(k')} \phi_X^{-1}(k)\end{aligned}$$

for each $k' \in K'$. □

9.2.3 Dependent Products

Let $f: K \rightarrow K'$ be a function and let (X, ϕ_X) be a K -fibred set.

00TX Definition 9.2.3.1.1. The **dependent product**⁵ of (X, ϕ_X) is the K' -fibred set $\Pi_f(X)$ ⁶ consisting of⁷

- *The Underlying Set.* The set $\Pi_f(X)$ defined by

$$\Pi_f(X) \stackrel{\text{def}}{=} \coprod_{k' \in K'} \prod_{k \in f^{-1}(k')} \phi_X^{-1}(k);$$

- *The Fibration.* The map of sets

$$\Pi_f(\phi_X): \Pi_f(X) \rightarrow K'$$

defined by sending an element of

$$\Pi_f(X) \stackrel{\text{def}}{=} \coprod_{k' \in K'} \prod_{k \in f^{-1}(k')} \phi_X^{-1}(k)$$

to its index k' in K' .

00TY Example 9.2.3.1.2. Here are some examples of dependent products of sets.

⁵The name “dependent product” comes from the fact that the fibre $\Pi_f(\phi_X)^{-1}(k')$ of $\Pi_f(X)$ at $k' \in K'$ is given by

$$\Pi_f(\phi_X)^{-1}(k') \cong \prod_{k \in f^{-1}(k')} \phi_X^{-1}(k);$$

see **Item 2 of Proposition 9.2.3.1.3.**

⁶Further Notation: Also written $f_!(X)$.

⁷We can also define dependent products via the internal **Hom** in **FibSets**(K'); see **Item 3 of Proposition 9.2.3.1.3.**

- 00TZ** 1. *Spaces of Sections.* Let $K = X, K' = \text{pt}$, let $\phi: E \rightarrow X$ be a map of sets, and write $!_X: X \rightarrow \text{pt}$ for the terminal map from X to pt . We have a bijection of sets

$$\begin{aligned}\Pi_{!_X}((E, \phi)) &\cong \Gamma_X(\phi) \\ &\stackrel{\text{def}}{=} \{h \in \text{Sets}(X, E) \mid \phi \circ h = \text{id}_X\}.\end{aligned}$$

- 00U0** 2. *Function Spaces.* Let $K = K' = \text{pt}$ and write $!_X: X \rightarrow \text{pt}$ and $!_Y: Y \rightarrow \text{pt}$ for the terminal maps from X and Y to pt . We have a bijection of sets

$$\text{Sets}(X, Y) \cong \Pi_{!_X}(!^*_X(Y, !_Y)).$$

Proof. **Item 1, Spaces of Sections:** Indeed, we have

$$\begin{aligned}\Pi_{!_X}((E, \phi)) &\stackrel{\text{def}}{=} \coprod_{\star \in \text{pt}} \prod_{k \in !_X^{-1}(\star)} \phi_X^{-1}(k) \\ &= \prod_{x \in X} \phi_X^{-1}(x) \\ &\cong \{h \in \text{Sets}(X, E) \mid \phi_X \circ h = \text{id}_X\} \\ &\stackrel{\text{def}}{=} \Gamma_X(\phi).\end{aligned}$$

Item 2, Function Spaces: Indeed, we have

$$\begin{aligned}\Pi_{!_X}(!^*_X(Y, !_Y)) &\stackrel{\text{def}}{=} \Pi_{!_X}(X \times_{!_X, \text{pt}, !_Y} Y) \\ &\stackrel{\text{def}}{=} \coprod_{\star \in \text{pt}} \prod_{x \in !_X^{-1}(\star)} \text{pr}_1^{-1}(x) \\ &= \prod_{x \in X} Y \\ &\cong \text{Sets}(X, Y).\end{aligned}$$

This finishes the proof. □

- 00U1** **Proposition 9.2.3.1.3.** Let $f: K \rightarrow K'$ be a function.

- 00U2** 1. *Functoriality.* The assignment $X \mapsto \Pi_f(X)$ defines a functor

$$\Pi_f: \text{FibSets}(K) \rightarrow \text{FibSets}(K'),$$

where

- *Action on Objects.* For each $(X, \phi_X) \in \text{Obj}(\text{FibSets}(K))$, we have

$$\Pi_f(X, \phi_X) \stackrel{\text{def}}{=} \Pi_f(X);$$

- *Action on Morphisms.* For each $(X, \phi_X), (Y, \phi_Y) \in \text{Obj}(\text{FibSets}(K))$, the

action on Hom-sets

$$\Pi_f|_{X,Y} : \text{Hom}_{\text{FibSets}(K)}(X, Y) \rightarrow \text{Hom}_{\text{FibSets}(K')}(\Pi_f(X), \Pi_f(Y))$$

of Π_f at $((X, \phi_X), (Y, \phi_Y))$ is the map sending a morphism of K -fibred sets

$$\xi : (X, \phi_X) \rightarrow (Y, \phi_Y), \quad \begin{array}{ccc} X & \xrightarrow{\xi} & Y \\ \phi_X \searrow & & \swarrow \phi_Y \\ K & & \end{array}$$

to the morphism

$$\begin{array}{ccc} \Pi_f(X) & \xrightarrow{\Pi_f(\xi)} & \Pi_f(Y) \\ \Pi_f(\phi_X) \searrow & & \swarrow \Pi_f(\phi_Y) \\ K & & \end{array}$$

of K' -fibred sets given by⁸

$$[\Pi_f(\xi)]((x_k)_{k \in f^{-1}(k')}) \stackrel{\text{def}}{=} (\xi(x_k))_{k \in f^{-1}(k')}$$

for each $(x_k)_{k \in f^{-1}(k')} \in \Pi_f(X)$.

00U3 2. *Interaction With Fibres.* We have a bijection of sets

$$\Pi_f(\phi_X)^{-1}(k') \cong \prod_{k \in f^{-1}(k')} \phi_X^{-1}(k);$$

for each $k' \in K'$.

00U4 3. *Construction Using the Internal Hom.* We have

$$\Pi_f(X, \phi_X) = \left(K' \times_{\text{Hom}_{\text{FibSets}(K')}((K, f), (K, f))} \text{Hom}_{\text{FibSets}(K')}((K, f), (X, f \circ \phi_X)), \text{pr}_1 \right),$$

forming a pullback diagram

$$\begin{array}{ccc} \Pi_f(X, \phi_X) & \xrightarrow{\text{pr}_2} & \text{Hom}_{\text{FibSets}(K')}((K, f), (X, f \circ \phi_X)) \\ \text{pr}_1 \downarrow & \lrcorner & \downarrow (\phi_X)_* \\ K' & \xrightarrow{I} & \text{Hom}_{\text{FibSets}(K')}((K, f), (K, f)), \end{array}$$

⁸Note that we indeed have $\xi(x_k) \in \phi_Y^{-1}(k)$, since

$$\begin{aligned} \phi_Y(\xi(x_k)) &= [\phi_Y \circ \xi](x_k) \\ &= \phi_X(x_k) \\ &= k, \end{aligned}$$

where the bottom map is given by

$$I(k') \stackrel{\text{def}}{=} \text{id}_{f^{-1}(k')}$$

for each $k' \in K'$ and where $\mathbf{Hom}_{\text{FibSets}(K')}$ denotes the internal Hom of $\text{FibSets}(K')$ of [Definition 9.2.4.1.1](#).

00U5 4. *Internal Hom via Dependent Products.* We have

$$\mathbf{Hom}_{\text{FibSets}(K)}(X, Y) \cong \Pi_{\phi_X}(\phi_X^*(Y)).$$

Proof. [Item 1, Functoriality:](#) Omitted.

[Item 2, Interaction With Fibres:](#) Clear.

[Item 3, Construction Using the Internal Hom:](#) Using the explicit formula for pullbacks of sets given in [??](#), we see that the pullback

$$K' \times_{\mathbf{Hom}_{\text{FibSets}(K')}}((K, f), (K, f)) \mathbf{Hom}_{\text{FibSets}(K')}((K, f), (X, f \circ \phi_X))$$

is given by

$$\left\{ (k', h) \in \coprod_{k' \in K'} \text{Sets}(f^{-1}(k'), \phi_X^{-1}(f^{-1}(k'))) \mid \phi_X \circ h = \text{id}_{f^{-1}(k')} \right\},$$

which is isomorphic to

$$\coprod_{k' \in K'} \{ h \in \text{Sets}(f^{-1}(k'), \phi_X^{-1}(f^{-1}(k'))) \mid \phi_X \circ h = \text{id}_{f^{-1}(k')} \}.$$

We claim that

$$\{ h \in \text{Sets}(f^{-1}(k'), \phi_X^{-1}(f^{-1}(k'))) \mid \phi_X \circ h = \text{id}_{f^{-1}(k')} \} \cong \prod_{k \in f^{-1}(k')} \phi_X^{-1}(k),$$

so that the pullback is indeed given by $\Pi_f(X)$. There are two cases:

1. If $f^{-1}(k') = \emptyset$, then there is only one map from $f^{-1}(k')$ to $\phi_X^{-1}(f^{-1}(k'))$ (the inclusion), so $\text{Sets}(f^{-1}(k'), \phi_X^{-1}(f^{-1}(k')) \cong \text{pt}$. Since products indexed by the empty set are isomorphic to pt , the isomorphism follows.
2. Otherwise, by the condition $\phi_X \circ h = \text{id}_{f^{-1}(k')}$, it follows that, for each $k \in f^{-1}(k')$, we must have

$$\phi_X(h(k)) = k,$$

and thus $h(k) \in \phi_X^{-1}(k)$. Therefore, a map from $f^{-1}(k')$ to $\phi_X^{-1}(f^{-1}(k'))$ consists of a choice of an element from $\phi_X^{-1}(k)$ for each $k \in f^{-1}(k')$, which is precisely given by an element of the product $\prod_{k \in f^{-1}(k')} \phi_X^{-1}(k)$, showing the bijection to be true.

Item 4, Internal Homs via Dependent Products: Indeed we have

$$\begin{aligned}
\Pi_{\phi_X}(\phi_X^*(Y)) &\stackrel{\text{def}}{=} \Pi_{\phi_X}(X \times_K Y) \\
&\stackrel{\text{def}}{=} \coprod_{k \in K} \prod_{x \in \phi_X^{-1}(k)} \text{pr}_1^{-1}(x) \\
&\cong \coprod_{k \in K} \prod_{x \in \phi_X^{-1}(k)} \{y \in Y \mid \phi_X(x) = \phi_Y(y)\} \\
&\cong \coprod_{k \in K} \prod_{x \in \phi_X^{-1}(k)} \{y \in Y \mid k = \phi_Y(y)\} \\
&\cong \coprod_{k \in K} \prod_{x \in \phi_X^{-1}(k)} \phi_Y^{-1}(k) \\
&\cong \coprod_{k \in K} \text{Sets}(\phi_X^{-1}(k), \phi_Y^{-1}(k)) \\
&\stackrel{\text{def}}{=} \mathbf{Hom}_{\mathbf{FibSets}(K)}(X, Y).
\end{aligned}$$

This finishes the proof. \square

9.2.4 Internal Hom

Let K be a set and let (X, ϕ_X) and (Y, ϕ_Y) be K -fibred sets.

00U7 Definition 9.2.4.1.1. The **internal Hom of fibred sets from** (X, ϕ_X) **to** (Y, ϕ_Y) is the fibred set $\mathbf{Hom}_{\mathbf{FibSets}(K)}(X, Y)$ consisting of

- *The Underlying Set.* The set $\mathbf{Hom}_{\mathbf{FibSets}(K)}(X, Y)$ defined by

$$\mathbf{Hom}_{\mathbf{FibSets}(K)}(X, Y) \stackrel{\text{def}}{=} \coprod_{k \in K} \text{Sets}(\phi_X^{-1}(k), \phi_Y^{-1}(k));$$

- *The Fibration.* The map of sets⁹

$$\begin{aligned}
\phi_{\mathbf{Hom}_{\mathbf{FibSets}(K)}(X, Y)} : & \underbrace{\mathbf{Hom}_{\mathbf{FibSets}(K)}(X, Y)}_{\coprod_{k \in K} \text{Sets}(\phi_X^{-1}(k), \phi_Y^{-1}(k))} \rightarrow K
\end{aligned}$$

defined by sending a map $f : \phi_X^{-1}(k) \rightarrow \phi_Y^{-1}(k)$ to its index $k \in K$.

where we have used that ξ is a morphism of K -fibred sets for the second equality.

⁹The fibres of the internal **Hom** of $\mathbf{FibSets}(K)$ are precisely the sets $\text{Sets}(\phi_X^{-1}(k), \phi_Y^{-1}(k))$, i.e. we have

$$\phi_{\mathbf{Hom}_{\mathbf{FibSets}(K)}(X, Y)|k} \cong \text{Sets}(\phi_X^{-1}(k), \phi_Y^{-1}(k))$$

for each $k \in K$.

Proof. Omitted. \square

00U8 **Proposition 9.2.4.1.2.** Let K be a set and let (X, ϕ_X) and (Y, ϕ_Y) be K -fibred sets.

00U9 1. *Functoriality.* Let (X, ϕ_X) and (Y, ϕ_Y) be K -fibred sets.

(a) The assignment $X \mapsto \mathbf{Hom}_{\mathbf{FibSets}(K)}(X, Y)$ defines a functor

$$\mathbf{Hom}_{\mathbf{FibSets}(K)}(X, -) : \mathbf{FibSets}(K) \rightarrow \mathbf{FibSets}(K).$$

(b) The assignment $Y \mapsto \mathbf{Hom}_{\mathbf{FibSets}(K)}(X, Y)$ defines a functor

$$\mathbf{Hom}_{\mathbf{FibSets}(K)}(-, Y) : \mathbf{FibSets}(K)^{\text{op}} \rightarrow \mathbf{FibSets}(K).$$

(c) The assignment $(X, Y) \mapsto \mathbf{Hom}_{\mathbf{FibSets}(K)}(X, Y)$ defines a functor

$$\mathbf{Hom}_{\mathbf{FibSets}(K)}(-_1, -_2) : \mathbf{FibSets}(K)^{\text{op}} \times \mathbf{FibSets}(K) \rightarrow \mathbf{FibSets}(K).$$

00UA 2. *Internal Homs via Dependent Products.* We have

$$\mathbf{Hom}_{\mathbf{FibSets}(K)}(X, Y) \cong \Pi_{\phi_X}(\phi_X^*(Y)).$$

Proof. Item 1, Functoriality: Omitted.

Item 2, Internal Homs via Dependent Products: This was proved in Item 4 of Proposition 9.2.3.1.3. \square

9.2.5 Adjointness for Fibred Sets

Let $f : K \rightarrow K'$ be a map of sets.

00UC **Proposition 9.2.5.1.1.** We have a triple adjunction

$$(\Sigma_f \dashv f^* \dashv \Pi_f) : \mathbf{FibSets}(K) \begin{array}{c} \xleftarrow{\perp} \\[-1ex] f^* \\[-1ex] \xrightarrow{\perp} \end{array} \mathbf{FibSets}(K').$$

We offer two proofs. The first uses the corresponding adjunction for indexed sets (??) and the un/straightening equivalence together with its compatibility with dependent sums and products to “transfer” the adjunction to fibred sets, while the second is a direct proof.

Proof. The Adjunction $\Sigma_f \dashv f^*$: The adjunction

$$(\Sigma_f \dashv f^*) : \mathbf{ISets}(K) \begin{array}{c} \xleftarrow{\perp} \\[-1ex] f^* \\[-1ex] \xrightarrow{\perp} \end{array} \mathbf{ISets}(K')$$

of ?? gives a unit and counit of the form

$$\begin{aligned}\eta: \text{id}_{\text{ISets}(K)} &\Rightarrow \Sigma_f \circ f^*, \\ \epsilon: f^* \circ \Sigma_f &\Rightarrow \text{id}_{\text{ISets}(K')}.\end{aligned}$$

With these in hand, we construct natural transformations

$$\begin{aligned}\eta': \text{id}_{\text{FibSets}(K)} &\Rightarrow \Sigma_f \circ f^*, \\ \epsilon': f^* \circ \Sigma_f &\Rightarrow \text{id}_{\text{FibSets}(K')}\end{aligned}$$

as follows:

1. *The Unit.* We define $\eta': \text{id}_{\text{FibSets}(K)} \Rightarrow \Sigma_f \circ f^*$ as the pasting of the diagram

$$\begin{array}{ccccc} & & \text{FibSets}(K') & & \\ & \nearrow \Sigma_f & \uparrow \text{St}_{K'} & \searrow f^* & \\ \text{FibSets}(K) & (1) & \text{ISets}(K') & (2) & \text{FibSets}(K) \\ \uparrow \text{id}_{\text{FibSets}(K)} & \swarrow \text{St}_K & \uparrow \eta \parallel f^* & \swarrow \text{St}_K & \uparrow \text{id}_{\text{FibSets}(K)} \\ \text{FibSets}(K) & (3) & \text{ISets}(K) & (4) & \text{FibSets}(K) \\ \uparrow \text{Un}_K & \nearrow \Sigma_f & \text{id}_{\text{ISets}(K)} \longrightarrow \text{id}_{\text{ISets}(K)} & \swarrow \text{Un}_K & \\ \text{FibSets}(K) & (5) & & & \text{FibSets}(K), \end{array}$$

where:

- (a) Subdiagram (1) commutes by ?? of ??.
- (b) Subdiagram (2) commutes by ?? of ??.
- (c) Subdiagram (3) commutes by ??.
- (d) Subdiagram (4) commutes by ??.
- (e) Subdiagram (5) commutes by unitality of composition.

2. *The Counit.* We define $\epsilon': f^* \circ \Sigma_f \Rightarrow \text{id}_{\text{FibSets}(K')}$ as the pasting of the diagram

$$\begin{array}{ccccc} \text{FibSets}(K') & \xrightarrow{\text{id}_{\text{FibSets}(K')}} & \text{FibSets}(K') & & \\ \downarrow \text{id}_{\text{FibSets}(K')} & \swarrow \text{Un}_K & & \searrow \text{Un}_K & \downarrow \text{id}_{\text{FibSets}(K')} \\ (2) & \text{ISets}(K') & \xrightarrow{\text{id}_{\text{ISets}(K)}} \text{ISets}(K') & (3) & \\ \downarrow \text{id}_{\text{FibSets}(K')} & \swarrow \text{St}_{K'} & \uparrow \epsilon \parallel \Sigma_f & \swarrow \text{St}_{K'} & \downarrow \text{id}_{\text{FibSets}(K')} \\ \text{FibSets}(K') & (4) & \text{ISets}(K) & (5) & \text{FibSets}(K') \\ \downarrow f^* & \nearrow \Sigma_f & \downarrow \text{St}_K & \nearrow \Sigma_f & \\ & & \text{FibSets}(K) & & \end{array}$$

where:

- (a) Subdiagram (1) commutes by unitality of composition.
 - (b) Subdiagram (2) commutes by ??.
 - (c) Subdiagram (3) commutes by ??.
 - (d) Subdiagram (4) commutes by ?? of ??.
 - (e) Subdiagram (5) commutes by ?? of ??.

Next, we prove the left triangle identity,

$$\begin{array}{ccc}
 \text{FibSets}(K') & \xrightarrow{\text{id}_{\text{FibSets}(K')}} & \text{FibSets}(K') \\
 \Sigma_f \nearrow \quad \uparrow \eta \quad \searrow f^* & \quad \uparrow \epsilon \quad \uparrow & \quad \nearrow \Sigma_f \\
 \text{FibSets}(K) & \xrightarrow{\text{id}_{\text{FibSets}(K)}} & \text{FibSets}(K)
 \end{array} = \begin{array}{ccc}
 \text{FibSets}(K') & \xrightarrow{\text{id}_{\text{FibSets}(K')}} & \text{FibSets}(K') \\
 \Sigma_f \nearrow \quad \uparrow \text{id}_{\Sigma_f} \quad \searrow \Sigma_f & & \\
 \text{FibSets}(K) & \xrightarrow{\text{id}_{\text{FibSets}(K)}} & \text{FibSets}(K)
 \end{array}$$

whose left side in our case looks like this:

It can be rearranged into

where:

1. Subdiagram (1) commutes by ??.
 2. Subdiagram (2) commutes by unitality of composition.
 3. Subdiagram (3) commutes by ??.

And then, it can be rearranged into

$$\begin{array}{ccccc}
 & & \text{FibSets}(K') & & \\
 & \nearrow \Sigma_f & \uparrow \text{id}_{\text{FibSets}(K')} & & \\
 \text{FibSets}(K) & & \text{ISets}(K') & & \text{FibSets}(K') \\
 & \downarrow \text{St}_K & \downarrow \text{id}_{\text{Sets}(K)} & \downarrow \text{id}_{\text{FibSets}(K')} & \\
 & \text{ISets}(K) & \xrightarrow{\eta} \text{ISets}(K) & \xrightarrow{\epsilon} \text{ISets}(K') & \\
 & \uparrow \Sigma_f & \uparrow f^* & \uparrow \Sigma_f & \\
 & \text{FibSets}(K) & \xrightarrow{\text{id}_{\text{FibSets}(K)}} \text{FibSets}(K) & \xrightarrow{\text{id}_{\text{FibSets}(K')}} \text{FibSets}(K') & \\
 & \downarrow \text{Un}_K & & \downarrow \text{St}_K & \\
 & & & \downarrow \Sigma_f & \\
 & & & \text{FibSets}(K') &
 \end{array}$$

which by the left triangle identity for (η, ϵ) , becomes

$$\begin{array}{ccccc}
 & & \text{FibSets}(K') & & \\
 & \nearrow \Sigma_f & \uparrow \text{id}_{\text{FibSets}(K')} & & \\
 \text{FibSets}(K) & & \text{ISets}(K') & & \text{FibSets}(K') \\
 & \downarrow \text{St}_K & \downarrow \text{id}_{\text{Sets}(K)} & \downarrow \text{id}_{\text{FibSets}(K')} & \\
 & \text{ISets}(K) & \xrightarrow{\text{id}} \text{ISets}(K) & \xrightarrow{\text{id}} \text{ISets}(K') & \\
 & \uparrow \Sigma_f & \uparrow \text{id}_{\text{Sets}(K)} & \uparrow \Sigma_f & \\
 & \text{FibSets}(K) & \xrightarrow{\text{id}_{\text{FibSets}(K)}} \text{FibSets}(K) & \xrightarrow{\text{id}_{\text{FibSets}(K')}} \text{FibSets}(K') & \\
 & \downarrow \text{Un}_K & & \downarrow \text{St}_K & \\
 & & & \downarrow \Sigma_f & \\
 & & & \text{FibSets}(K') &
 \end{array}$$

finishing the proof of the left triangle identity. The proof of the right triangle identity is similar, and is thus omitted.

The Adjunction $f^ \dashv \Pi_f$:* This proof is similar to the proof of the adjunction $\Sigma_f \dashv f^*$, and is thus omitted. \square

We proceed to the direct proof of [Proposition 9.2.5.1.1](#).

Proof. *The Adjunction $\Sigma_f \dashv f^*$:* We claim there's a bijection

$$\text{Hom}_{\text{FibSets}(K')}(\Sigma_f(X), Y) \cong \text{Hom}_{\text{FibSets}(K)}(X, f^*(Y)),$$

natural in $(X, \phi_X) \in \text{Obj}(\text{FibSets}(K))$ and $(Y, \phi_Y) \in \text{Obj}(\text{FibSets}(K'))$:

- *Map I.* We define a map

$$\Phi_{X,Y}: \text{Hom}_{\text{FibSets}(K')}(\Sigma_f(X), Y) \rightarrow \text{Hom}_{\text{FibSets}(K)}(X, f^*(Y)),$$

by sending a morphism

$$\begin{array}{ccc}
 X & \xrightarrow{\xi} & Y \\
 \xi: \Sigma_f(X) \rightarrow Y, & \downarrow \phi_X & \swarrow \phi_Y \\
 & K & \\
 & \downarrow f & \nearrow K' \\
 & K' &
 \end{array}$$

of K' -fibred sets to the morphism

$$\xi^\dagger: X \rightarrow f^*(Y), \quad \begin{array}{ccc} X & \xrightarrow{\xi^\dagger} & K \times_{K'} Y \\ \phi_X \searrow & & \swarrow \text{pr}_1 \\ & K' & \end{array}$$

of K -fibred sets given by the dashed morphism in the diagram

$$\begin{array}{ccccc} X & \xrightarrow{\quad \exists! \quad} & K \times_{K'} Y & \xrightarrow{\text{pr}_2} & Y \\ \phi_X \curvearrowleft & & \downarrow \text{pr}_1 & & \downarrow \phi_Y \\ & & K & \xrightarrow{f} & K'. \end{array}$$

- *Map II.* We define a map

$$\Psi_{X,Y}: \text{Hom}_{\text{FibSets}(K)}(X, f^*(Y)) \rightarrow \text{Hom}_{\text{FibSets}(K')}\left(\Sigma_f(X), Y\right),$$

given by sending a map

$$\xi: X \rightarrow f^*(Y), \quad \begin{array}{ccc} X & \xrightarrow{\xi} & K \times_{K'} Y \\ \phi_X \searrow & & \swarrow \text{pr}_1 \\ & K' & \end{array}$$

of K' -fibred sets to the map

$$\xi^\dagger: \Sigma_f(X) \rightarrow Y, \quad \begin{array}{ccc} X & \xrightarrow{\xi^\dagger} & Y \\ \phi_X \searrow & & \swarrow \phi_Y \\ K & \xrightarrow{f} & K' \end{array}$$

of K -fibred sets given by

$$\xi^\dagger \stackrel{\text{def}}{=} \text{pr}_2 \circ \xi,$$

where we indeed have

$$\begin{aligned} \phi_Y \circ (\text{pr}_2 \circ \xi) &= (\phi_Y \circ \text{pr}_2) \circ \xi \\ &= (f \circ \text{pr}_1) \circ \xi && \text{(by the pullback square of } K \times_{K'} Y\text{)} \\ &= f \circ (\text{pr}_1 \circ \xi) \\ &= f \circ \phi_X. && \text{(since } \xi \text{ is a morphism of } K'\text{-fibred sets)} \end{aligned}$$

- *Naturality I.* We need to show that, given a morphism

$$\alpha: (X, \phi_X) \rightarrow (X', \phi_{X'})$$

of K -fibred sets, the diagram

$$\begin{array}{ccc} \text{Hom}_{\text{FibSets}(K')}(\Sigma_f(X'), Y) & \xrightarrow{\Phi_{X',Y}} & \text{Hom}_{\text{FibSets}(K)}(X', f^*(Y)), \\ \downarrow \Sigma_f(\alpha)^* & & \downarrow \alpha^* \\ \text{Hom}_{\text{FibSets}(K')}(\Sigma_f(X), Y) & \xrightarrow{\Phi_{X,Y}} & \text{Hom}_{\text{FibSets}(K)}(X, f^*(Y)) \end{array}$$

commutes. Indeed, given a morphism

$$\begin{array}{ccc} X' & \xrightarrow{\xi} & Y \\ \phi_{X'} \searrow & K & \swarrow \phi_Y \\ & f \downarrow & \\ & K' & \end{array}$$

of K' -fibred-sets, the map $\Phi_{X',Y}(\xi) \circ \alpha$ is the composition, coloured in **vermillion**, of the dashed arrow with α in the diagram

$$\begin{array}{ccccc} X & \xrightarrow{\quad \quad} & X' & \xrightarrow{\quad \quad} & Y \\ \alpha \searrow & \nearrow \exists! & \xi \searrow & \nearrow \exists! & \phi_Y \downarrow \\ & X' & \xrightarrow{\quad \quad} & K \times_{K'} Y & \xrightarrow{\quad \quad} Y \\ \phi_{X'} \circ \alpha \searrow & \nearrow \phi_{X'} & \downarrow \text{pr}_1 & \downarrow \text{pr}_2 & \downarrow \phi_Y \\ & K & \xrightarrow{\quad f \quad} & K' & \end{array}$$

while $\Phi_{X,Y}(\xi \circ \Sigma_f(\alpha))$ is given by the dashed arrow, coloured in **blue**. Since both the **blue arrow** and the **vermillion arrow** make the outer pullback diagram commute, it follows from the universal property of the pullback that they must be equal, i.e. that

$$\Phi_{X',Y}(\xi) \circ \alpha = \Phi_{X,Y}(\xi \circ \Sigma_f(\alpha)),$$

showing that the naturality diagram above indeed commutes.

- *Naturality II.* We need to show that, given a morphism

$$\beta: (Y, \phi_Y) \rightarrow (Y', \phi_{Y'})$$

of K -fibred sets, the diagram

$$\begin{array}{ccc} \mathrm{Hom}_{\mathsf{FibSets}(K')}(\Sigma_f(X), Y) & \xrightarrow{\Phi_{X,Y}} & \mathrm{Hom}_{\mathsf{FibSets}(K)}(X, f^*(Y)), \\ \beta_* \downarrow & & \downarrow f^*(\beta)_* \\ \mathrm{Hom}_{\mathsf{FibSets}(K')}(\Sigma_f(X), Y') & \xrightarrow{\Phi_{X,Y'}} & \mathrm{Hom}_{\mathsf{FibSets}(K)}(X, f^*(Y')) \end{array}$$

commutes. Indeed, given a morphism

$$\begin{array}{ccc} X' & \xrightarrow{\xi} & Y \\ \phi_{X'} \searrow & K & \swarrow \phi_Y \\ f \searrow & & K' \end{array}$$

of K' -fibred-sets, the map $f^*(\beta) \circ \Phi_{X,Y}(\xi)$ is the composition, coloured in **vermillion**, of the dashed arrow from X to $K \times_{K'} Y$ with the dashed arrow from $K \times_{K'} Y$ to $K \times_{K'} Y'$ in the diagram

$$\begin{array}{ccccc}
X & \xrightarrow{\hspace{1cm}} & \xi & \xrightarrow{\hspace{1cm}} & Y \\
\exists! \searrow & & \exists! \downarrow & & \beta \searrow \\
K \times_{K'} Y & \xrightarrow{\hspace{1cm}} & & \xrightarrow{\hspace{1cm}} & Y' \\
\downarrow & \lrcorner \quad \lrcorner & \downarrow & \lrcorner & \downarrow \\
K \times_{K'} Y' & \xrightarrow{\hspace{1cm}} & & \xrightarrow{\hspace{1cm}} & Y' \\
\downarrow & \lrcorner & \downarrow & \lrcorner & \downarrow \\
K & \xrightarrow{f} & K & \xrightarrow{f} & K' \\
\parallel & \parallel & \parallel & \parallel & \parallel \\
& & & & \phi_{Y'} \\
& & & & \phi_Y
\end{array}$$

while $\Phi_{X,Y'}(\beta \circ \xi)$ is given by the dashed arrow from X to $K \times_{K'} Y'$, coloured in blue. Since both the blue arrow and the vermillion arrow make the outer pullback diagram for $K \times_{K'} Y'$ commute, it follows from the universal property of the pullback that they must be equal, i.e. that

$$f^*(\beta) \circ \Phi_{X,Y}(\xi) = \Phi_{X,Y'}(\beta \circ \xi),$$

showing that the naturality diagram above indeed commutes.

- *Invertibility I.* We claim that

$$\Psi_{X,Y} \circ \Phi_{X,Y} = \text{id}_{\text{Hom}_{\text{FibSets}(K')}(\Sigma_f(X), Y)}.$$

Indeed, $\Phi_{X,Y}$ sends a map

$$\xi: \Sigma_f(X) \rightarrow Y, \quad \begin{array}{ccc} X & \xrightarrow{\xi} & Y \\ \phi_X \searrow & K & \swarrow \phi_Y \\ f \downarrow & K' & \end{array}$$

of K' -fibred sets to the dashed morphism in the diagram

$$\begin{array}{ccccc} X & \xrightarrow{\quad \exists! \quad} & K \times_{K'} Y & \xrightarrow{\text{pr}_2} & Y \\ \phi_X \searrow & \text{pr}_1 \downarrow & \downarrow & & \downarrow \phi_Y \\ & K & \xrightarrow{f} & K' & \end{array}$$

and $\Psi_{X,Y}$ then postcomposes that map with pr_2 , which, by the commutativity of the diagram above, is ξ again, showing the claimed equality to be true.

- *Invertibility II.* We claim that

$$\Phi_{X,Y} \circ \Psi_{X,Y} = \text{id}_{\text{Hom}_{\text{FibSets}(K)}(X, f^*(Y))}.$$

Indeed, $\Psi_{X,Y}$ sends a map

$$\xi: X \rightarrow f^*(Y), \quad \begin{array}{ccc} X & \xrightarrow{\xi} & K \times_{K'} Y \\ \phi_X \searrow & \text{pr}_1 \swarrow & \\ & K' & \end{array}$$

of K' -fibred sets to $\text{pr}_2 \circ \xi$, which is then sent by $\Phi_{X,Y}$ to the dashed morphism in the diagram

$$\begin{array}{ccccc} X & \xrightarrow{\quad \exists! \quad} & K \times_{K'} Y & \xrightarrow{\text{pr}_2 \circ \xi} & Y \\ \phi_X \searrow & \text{pr}_1 \downarrow & \downarrow & & \downarrow \phi_Y \\ & K & \xrightarrow{f} & K' & \end{array}$$

which, by the commutativity of the subdiagram marked with (\dagger) , is given by ξ again, showing the claimed equality to be true.

The Adjunction $f^ \dashv \Pi_f$:* We claim there's a bijection

$$\text{Hom}_{\text{FibSets}(K)}(f^*(X), Y) \cong \text{Hom}_{\text{FibSets}(K')}(X, \Pi_f(Y))$$

natural in $(X, \phi_X) \in \text{Obj}(\text{FibSets}(K))$ and $(Y, \phi_Y) \in \text{Obj}(\text{FibSets}(K'))$:

1. *Map I.* We define a map

$$\Phi_{X,Y}: \text{Hom}_{\text{FibSets}(K)}(f^*(X), Y) \rightarrow \text{Hom}_{\text{FibSets}(K')}(X, \Pi_f(Y))$$

defined as follows. Given a morphism

$$\begin{array}{c} K \times_{K'} X \xrightarrow{\xi} Y \\ \xi: f^*(X) \rightarrow Y, \quad \text{pr}_1 \searrow \quad \swarrow \phi_Y \\ K \end{array}$$

of K -fibred sets, where

$$\begin{aligned} f^*(X) &\stackrel{\text{def}}{=} K \times_{K'} X \\ &\stackrel{\text{def}}{=} \{(k, x) \in K \times X \mid f(k) = \phi_X(x)\}, \end{aligned}$$

we construct a morphism

$$\begin{array}{c} X \xrightarrow{\xi^\dagger} \Pi_f(Y) \\ \xi^\dagger: X \rightarrow \Pi_f(Y), \quad \phi_X \searrow \quad \swarrow \Pi_f(\phi_Y) \\ K' \end{array}$$

of K' -fibred sets, where

$$\Pi_f(Y) \stackrel{\text{def}}{=} \coprod_{k' \in K'} \prod_{k \in f^{-1}(k')} \phi_Y^{-1}(k),$$

by defining

$$\xi^\dagger(x) \stackrel{\text{def}}{=} (\xi(k, x))_{k \in f^{-1}(\phi_X(x))}$$

for each $x \in X$. There are two things to be checked here:

- We have $\xi(k, x) \in \phi_Y^{-1}(\phi_X(x))$ since $\phi_Y(\xi(k, x)) = \phi_X(x)$ as ξ is a morphism of K -fibred sets.
- The map ξ^\dagger is indeed a morphism of K' -fibred sets, i.e. we have

$$\Pi_f(\phi_Y) \circ \xi^\dagger = \phi_X,$$

since

$$[\Pi_f(\phi_Y)]((\xi(k, x))_{k \in f^{-1}(\phi_X(x))}) = \phi_X(x)$$

for each $x \in X$.

2. *Map II.* We define a map

$$\Psi_{X,Y}: \text{Hom}_{\text{FibSets}(K')}(X, \Pi_f(Y)) \rightarrow \text{Hom}_{\text{FibSets}(K)}(f^*(X), Y)$$

as follows. Given a morphism

$$\begin{array}{ccc} X & \xrightarrow{\xi} & \Pi_f(Y) \\ \xi: X \rightarrow \Pi_f(Y), & \phi_X \searrow & \swarrow \Pi_f(\phi_Y) \\ & K' & \end{array}$$

of K' -fibred sets, where

$$\Pi_f(Y) \stackrel{\text{def}}{=} \coprod_{k' \in K'} \prod_{k \in f^{-1}(k')} \phi_Y^{-1}(k),$$

we construct a morphism

$$\begin{array}{ccc} \xi^\dagger: f^*(X) \rightarrow Y, & K \times_{K'} X & \xrightarrow{\xi^\dagger} Y \\ & \text{pr}_1 \searrow & \swarrow \phi_Y \\ & K & \end{array}$$

of K -fibred sets, where

$$\begin{aligned} f^*(X) &\stackrel{\text{def}}{=} K \times_{K'} X \\ &\stackrel{\text{def}}{=} \{(k, x) \in K \times X \mid f(k) = \phi_X(x)\}, \end{aligned}$$

by defining

$$\xi^\dagger(k, x) \stackrel{\text{def}}{=} \xi(x)_k$$

for each $(k, x) \in f^*(X)$, where $\xi(x)_k$ is the k th component of $\xi(x) = (y_k)_{k \in f^{-1}(k')}$.

We also need to check that ξ^\dagger is a morphism of K -fibred sets, i.e. that

$$\phi_Y \circ \xi^\dagger = \text{pr}_1,$$

or

$$\phi_Y(\xi^\dagger(k, x)) = k,$$

for each $(k, x) \in f^*(X)$, which is clear, since $\xi^\dagger(k, x) \in \phi_Y^{-1}(k)$ by definition.

3. *Naturality I.* We need to show that, given a morphism

$$\alpha: (X, \phi_X) \rightarrow (X', \phi_{X'})$$

of K' -fibred sets, the diagram

$$\begin{array}{ccc} \text{Hom}_{\text{FibSets}(K')}(f^*(X'), Y) & \xrightarrow{\Phi_{X', Y}} & \text{Hom}_{\text{FibSets}(K)}(X', \Pi_f(Y)) \\ f^*(\alpha)^* \downarrow & & \downarrow \alpha^* \\ \text{Hom}_{\text{FibSets}(K')}(f^*(X), Y) & \xrightarrow{\Phi_{X, Y}} & \text{Hom}_{\text{FibSets}(K)}(X, \Pi_f(Y)) \end{array}$$

commutes. Indeed, given a morphism $\xi: f^*(X') \rightarrow Y$ of K' -fibred sets, we have

$$\begin{aligned} [[\Phi_{X,Y} \circ f^*(\alpha)](\xi)](x) &\stackrel{\text{def}}{=} [\Phi_{X,Y}(\xi \circ f^*(\alpha))](x) \\ &\stackrel{\text{def}}{=} ([\xi \circ f^*(\alpha)](k, x))_{k \in f^{-1}(\phi_X(x))} \\ &\stackrel{\text{def}}{=} (\xi(k, \alpha(x)))_{k \in f^{-1}(\phi_X(x))} \\ &\stackrel{\text{def}}{=} \alpha^*\left((\xi(k, x))_{k \in f^{-1}(\phi_X(x))}\right) \\ &\stackrel{\text{def}}{=} \alpha^*\left(\xi^\dagger(x)\right) \\ &\stackrel{\text{def}}{=} [[\alpha^* \circ \Phi_{X,Y}](\xi)](x) \end{aligned}$$

for each $x \in X$.

4. *Naturality II.* We need to show that, given a morphism

$$\beta: (Y, \phi_Y) \rightarrow (Y', \phi_{Y'})$$

of K -fibred sets, the diagram

$$\begin{array}{ccc} \text{Hom}_{\text{FibSets}(K')}(f^*(X), Y) & \xrightarrow{\Phi_{X,Y}} & \text{Hom}_{\text{FibSets}(K)}(X, \Pi_f(Y)) \\ \beta_* \downarrow & & \downarrow \Pi_f(\beta)_* \\ \text{Hom}_{\text{FibSets}(K')}(f^*(X), Y') & \xrightarrow{\Phi_{X,Y'}} & \text{Hom}_{\text{FibSets}(K)}(X, \Pi_f(Y')) \end{array}$$

commutes. Indeed, given a morphism $\xi: X \rightarrow \Pi_f(Y)$ of K -fibred sets, we have

$$\begin{aligned} [[\Phi_{X,Y'} \circ \beta_*](\xi)](x) &\stackrel{\text{def}}{=} [\Phi_{X,Y'}(\beta \circ \xi)](x) \\ &\stackrel{\text{def}}{=} [\Phi_{X,Y'}(\beta \circ \xi)](x) \\ &\stackrel{\text{def}}{=} ([\beta \circ \xi](k, x))_{k \in f^{-1}(\phi_X(x))} \\ &\stackrel{\text{def}}{=} (\beta(\xi(k, x)))_{k \in f^{-1}(\phi_X(x))} \\ &\stackrel{\text{def}}{=} \Pi_f(\beta)_*\left((\xi(k, x))_{k \in f^{-1}(\phi_X(x))}\right) \\ &\stackrel{\text{def}}{=} [\Pi_f(\beta)_* \circ \xi^\dagger](x) \\ &\stackrel{\text{def}}{=} [\Pi_f(\beta)_* \circ \Phi_{X,Y'}(\xi)](x) \end{aligned}$$

for each $x \in X$.

5. *Invertibility I.* We claim that

$$\Psi_{X,Y} \circ \Phi_{X,Y} = \text{id}_{\text{Hom}_{\text{FibSets}(K)}(f^*(X), Y)}.$$

Indeed, given a morphism $\xi: f^*(X') \rightarrow Y$ of K' -fibred sets, we have

$$\begin{aligned} [[\Psi_{X,Y} \circ \Phi_{X,Y}](\xi)](k, x) &\stackrel{\text{def}}{=} [\Psi_{X,Y}(\Phi_{X,Y}(\xi))](k, x) \\ &\stackrel{\text{def}}{=} ([\Phi_{X,Y}(\xi)](x))_k \\ &\stackrel{\text{def}}{=} \left((\xi(k_1, x))_{k_1 \in f^{-1}(\phi_X(x))}\right)_k \\ &\stackrel{\text{def}}{=} \xi(k, x) \end{aligned}$$

for each $(k, x) \in f^*(X)$, and thus the stated equality follows.

6. *Invertibility II.* We claim that

$$\Phi_{X,Y} \circ \Psi_{X,Y} = \text{id}_{\text{Hom}_{\text{FibSets}(K')}(X, \Pi_f(Y))}.$$

Indeed, given a morphism $\xi: X \rightarrow \Pi_f(Y)$ of K -fibred sets, write

$$\xi(x) = (y_k)_{k \in f^{-1}(k'_x)}.$$

We then have

$$\begin{aligned} [[\Phi_{X,Y} \circ \Psi_{X,Y}](\xi)](x) &\stackrel{\text{def}}{=} [\Phi_{X,Y}(\Psi_{X,Y}(\xi))](x) \\ &\stackrel{\text{def}}{=} ([\Psi_{X,Y}(\xi)](k_1, x))_{k_1 \in f^{-1}(\phi_X(x))} \\ &\stackrel{\text{def}}{=} ((\xi(x))_{k_1})_{k_1 \in f^{-1}(\phi_X(x))} \\ &\stackrel{\text{def}}{=} \left(((y_k)_{k \in f^{-1}(k'_x)})_{k_1} \right)_{k_1 \in f^{-1}(\phi_X(x))} \\ &\stackrel{\text{def}}{=} (y_{k_1})_{k_1 \in f^{-1}(\phi_X(x))} \\ &= (y_{k_1})_{k_1 \in f^{-1}(k'_x)} \\ &= (y_k)_{k \in f^{-1}(k'_x)} \\ &\stackrel{\text{def}}{=} \xi(x) \end{aligned}$$

for each $x \in X$, where the equality $\phi_X(x) = k'_x$ follows from the fact that ξ is a morphism of K' -fibred sets. Thus the stated equality follows.

This finishes the proof. \square

Appendices

9.A Other Chapters

Sets

- 1. Sets
- 2. Constructions With Sets
- 3. Pointed Sets
- 4. Tensor Products of Pointed Sets
- 5. Relations
- 6. Spans

Posets

Indexed and Fibred Sets

- 7. Indexed Sets
- 8. Fibred Sets
- 9. Un/Straightening for Indexed and Fibred Sets

Category Theory

- 11. Categories

- 12. Types of Morphisms in Categories
- 13. Adjunctions and the Yoneda Lemma
- 14. Constructions With Categories
- 15. Profunctors
- 16. Cartesian Closed Categories
- 17. Kan Extensions
- Bicategories**
- 18. Bicategories
- 19. Internal Adjunctions
- Internal Category Theory**
- 20. Internal Categories
- Cyclic Stuff**
- 21. The Cycle Category
- Cubical Stuff**
- 22. The Cube Category
- Globular Stuff**
- 23. The Globe Category
- Cellular Stuff**
- 24. The Cell Category
- Monoids**
- 25. Monoids
- 26. Constructions With Monoids
- Monoids With Zero**
- 27. Monoids With Zero
- 28. Constructions With Monoids With Zero
- Groups**
- 29. Groups
- 30. Constructions With Groups
- Hyper Algebra**
- 31. Hypermonoids
- 32. Hypergroups
- 33. Hypersemirings and Hyperrings
- 34. Quantales
- Near-Rings**
- 35. Near-Semirings
- 36. Near-Rings
- Real Analysis**
- 37. Real Analysis in One Variable
- 38. Real Analysis in Several Variables
- Measure Theory**
- 39. Measurable Spaces
- 40. Measures and Integration
- Probability Theory**
- 40. Probability Theory
- Stochastic Analysis**
- 41. Stochastic Processes, Martingales, and Brownian Motion
- 42. Itô Calculus
- 43. Stochastic Differential Equations
- Differential Geometry**
- 44. Topological and Smooth Manifolds
- Schemes**
- 45. Schemes

9.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
Category Theory	27. Monoids With Zero
11. Categories	28. Constructions With Monoids With Zero
12. Types of Morphisms in Categories	Groups
13. Adjunctions and the Yoneda Lemma	29. Groups
14. Constructions With Categories	30. Constructions With Groups
15. Profunctors	Hyper Algebra
16. Cartesian Closed Categories	31. Hypermonoids
17. Kan Extensions	32. Hypergroups
Bicategories	33. Hypersemirings and Hyperrings
18. Bicategories	34. Quantales
19. Internal Adjunctions	Near-Rings
Internal Category Theory	35. Near-Semirings
20. Internal Categories	36. Near-Rings
	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 10

Un/Straightening for Indexed and Fibred Sets

00UD This chapter contains a discussion of the un/sstraightening equivalence in the context of sets, as well as a general discussion of indexed and fibred sets. In particular, it contains:

1. A discussion of indexed sets (i.e. functors $K_{\text{disc}} \rightarrow \text{Sets}$ with K a set), constructions with them like dependent sums and dependent products, and their properties (????);
2. A discussion of fibred sets (i.e. maps of sets $X \rightarrow K$), constructions with them like dependent sums and dependent products, and their properties (????);
3. A discussion of the un/sstraightening equivalence for indexed and fibred sets ([Section 10.1](#)).

Contents

10.1	Un/Straightening for Indexed and Fibred Sets	298
10.1.1	Straightening for Fibred Sets	298
10.1.2	Unstraightening for Indexed Sets	301
10.1.3	The Un/Straightening Equivalence	304
10.2	Miscellany	304
10.2.1	Other Kinds of Un/Straightening	304
10.A	Other Chapters	305
10.2	Other Chapters	306

10.1 Un/Straightening for Indexed and Fibred Sets

10.1.1 Straightening for Fibred Sets

Let K be a set and let (X, ϕ) be a K -fibred set.

00UG Definition 10.1.1.1.1. The **straightening** of (X, ϕ) is the K -indexed set

$$\text{St}_K(X, \phi) : K_{\text{disc}} \rightarrow \text{Sets}$$

defined by

$$\text{St}_K(X, \phi)_x \stackrel{\text{def}}{=} \phi^{-1}(x)$$

for each $x \in K$.

00UH Proposition 10.1.1.1.2. Let K be a set.

00UJ 1. Functoriality. The assignment $(X, \phi) \mapsto \text{St}_K(X, \phi)$ defines a functor

$$\text{St}_K : \text{FibSets}(K) \rightarrow \text{ISets}(K)$$

- *Action on Objects.* For each $(X, \phi) \in \text{Obj}(\text{FibSets}(K))$, we have

$$[\text{St}_K](X, \phi) \stackrel{\text{def}}{=} \text{St}_K(X, \phi);$$

- *Action on Morphisms.* For each $(X, \phi), (Y, \psi) \in \text{Obj}(\text{FibSets}(K))$, the action on Hom-sets

$$\text{St}_{K|X,Y} : \text{Hom}_{\text{FibSets}(K)}(X, Y) \rightarrow \text{Hom}_{\text{ISets}(K)}(\text{St}_K(X), \text{St}_K(Y))$$

of St_K at (X, Y) is given by sending a morphism

$$f : (X, \phi) \rightarrow (Y, \psi)$$

of K -fibred sets to the morphism

$$\text{St}_K(f) : \text{St}_K(X, \phi) \rightarrow \text{St}_K(Y, \psi)$$

of K -indexed sets defined by

$$\text{St}_K(f) \stackrel{\text{def}}{=} \{f_x^*\}_{x \in K},$$

where f_x^* is the transport map associated to f at $x \in K$ of ??.

00UK 2. Interaction With Change of Base/Indexing. Let $f : K \rightarrow K'$ be a map of sets. The diagram

$$\begin{array}{ccc} \text{FibSets}(K') & \xrightarrow{f^*} & \text{FibSets}(K) \\ \text{St}_{K'} \downarrow & & \downarrow \text{St}_K \\ \text{ISets}(K') & \xrightarrow{f^*} & \text{ISets}(K) \end{array}$$

commutes.

00UL 3. *Interaction With Dependent Sums.* Let $f: K \rightarrow K'$ be a map of sets. The diagram

$$\begin{array}{ccc} \text{FibSets}(K) & \xrightarrow{\Sigma_f} & \text{FibSets}(K') \\ \text{St}_K \downarrow & & \downarrow \text{St}_{K'} \\ \text{ISets}(K) & \xrightarrow{\Sigma_f} & \text{ISets}(K') \end{array}$$

commutes.

00UM 4. *Interaction With Dependent Products.* Let $f: K \rightarrow K'$ be a map of sets. The diagram

$$\begin{array}{ccc} \text{Sets}_{/K} & \xrightarrow{\Pi_f} & \text{FibSets}(K') \\ \text{St}_K \downarrow & & \downarrow \text{St}_{K'} \\ \text{ISets}(K) & \xrightarrow{\Pi_f} & \text{ISets}(K') \end{array}$$

commutes.

Proof. **Item 1, Functoriality:** Omitted.

Item 2, Interaction With Change of Base/Indexing: Indeed, we have

$$\begin{aligned} \text{St}_K(f^*(X, \phi))_x &\stackrel{\text{def}}{=} \text{St}_K(K \times_{K'} X)_x \\ &\stackrel{\text{def}}{=} \left(\text{pr}_1^{K \times_{K'} X} \right)^{-1}(x) \\ &= \{(k, y) \in K \times_{K'} X \mid \text{pr}_1^{K \times_{K'} X}(k, y) = x\} \\ &= \{(k, y) \in K \times_{K'} X \mid k = x\} \\ &= \{(k, y) \in K \times X \mid k = x \text{ and } f(k) = \phi(y)\} \\ &\cong \{y \in X \mid \phi(y) = f(x)\} \\ &= \phi^{-1}(f(x)) \\ &\stackrel{\text{def}}{=} f^*(\phi^{-1}(x)) \\ &\stackrel{\text{def}}{=} f^*(\text{St}_{K'}(X, \phi)_x) \end{aligned}$$

for each $(X, \phi) \in \text{Obj}(\text{FibSets}(K'))$ and each $x \in K$, and similarly for morphisms.

Item 3, Interaction With Dependent Sums: Indeed, we have

$$\begin{aligned} \text{St}_{K'}(\Sigma_f(X, \phi))_x &\stackrel{\text{def}}{=} \Sigma_f(\phi)^{-1}(x) \\ &\cong \coprod_{\substack{y \in X \\ f(y)=x}} \phi^{-1}(y) \\ &\cong \Sigma_f(\phi^{-1}(x)) \\ &\stackrel{\text{def}}{=} \Sigma_f(\text{St}_K(X, \phi)_x) \end{aligned}$$

for each $(X, \phi) \in \text{Obj}(\text{FibSets}(K))$ and each $x \in K'$, where we have used ?? of ?? for the first bijection, and similarly for morphisms.

Item 4, Interaction With Dependent Products: Indeed, we have

$$\begin{aligned} \text{St}_{K'}(\Pi_f(X, \phi))_x &\stackrel{\text{def}}{=} \Pi_f(\phi)^{-1}(x) \\ &\cong \prod_{\substack{y \in X \\ f(y)=x}} \phi^{-1}(y) \\ &\cong \Pi_f(\phi^{-1}(x)) \\ &\stackrel{\text{def}}{=} \Pi_f(\text{St}_K(X, \phi)_x) \end{aligned}$$

for each $(X, \phi) \in \text{Obj}(\text{FibSets}(K))$ and each $x \in K'$, where we have used ?? of ?? for the first bijection, and similarly for morphisms. \square

10.1.2 Unstraightening for Indexed Sets

Let K be a set and let X be a K -indexed set.

00UP Definition 10.1.2.1.1. The **unstraightening** of X is the K -fibred set

$$\phi_{\text{Un}_K} : \text{Un}_K(X) \rightarrow K$$

consisting of

- *The Underlying Set.* The set $\text{Un}_K(X)$ defined by

$$\text{Un}_K(X) \stackrel{\text{def}}{=} \coprod_{x \in K} X_x;$$

- *The Fibration.* The map of sets

$$\phi_{\text{Un}_K} : \text{Un}_K(X) \rightarrow K$$

defined by sending an element of $\coprod_{x \in K} X_x$ to its index in K .

00UQ Proposition 10.1.2.1.2. Let K be a set.

00UR 1. *Functoriality.* The assignment $X \mapsto \text{Un}_K(X)$ defines a functor

$$\text{Un}_K : \text{ISets}(K) \rightarrow \text{FibSets}(K)$$

- *Action on Objects.* For each $X \in \text{Obj}(\text{ISets}(K))$, we have

$$[\text{Un}_K](X) \stackrel{\text{def}}{=} \text{Un}_K(X);$$

- *Action on Morphisms.* For each $X, Y \in \text{Obj}(\text{ISets}(K))$, the action on Hom-sets

$$\text{Un}_{K|X,Y} : \text{Hom}_{\text{ISets}(K)}(X, Y) \rightarrow \text{Hom}_{\text{FibSets}(K)}(\text{Un}_K(X), \text{Un}_K(Y))$$

of Un_K at (X, Y) is defined by

$$\text{Un}_{K|X,Y}(f) \stackrel{\text{def}}{=} \coprod_{x \in K} f_x^*.$$

00US 2. *Interaction With Fibres.* We have a bijection of sets

$$\phi_{\text{Un}_K}^{-1}(x) \cong X_x$$

for each $x \in K$.

00UT 3. *As a Pullback.* We have a bijection of sets

$$\begin{array}{ccc} \text{Un}_K(X) & \rightarrow & \text{Sets}_* \\ \downarrow & \lrcorner & \downarrow \text{忘} \\ \text{Un}_K(X) \cong K_{\text{disc}} \times_{\text{Sets}} \text{Sets}_*, & & \\ \downarrow & & \\ K_{\text{disc}} & \xrightarrow[X]{} & \text{Sets}. \end{array}$$

00UU 4. *As a Colimit.* We have a bijection of sets

$$\text{Un}_K(X) \cong \text{colim}(X).$$

00UV 5. *Interaction With Change of Indexing/Base.* Let $f: K \rightarrow K'$ be a map of sets. The diagram

$$\begin{array}{ccc} \text{ISets}(K') & \xrightarrow{f^*} & \text{ISets}(K) \\ \text{Un}_{K'} \downarrow & & \downarrow \text{Un}_K \\ \text{FibSets}(K') & \xrightarrow[f^*]{} & \text{FibSets}(K) \end{array}$$

commutes.

00UW 6. *Interaction With Dependent Sums.* Let $f: K \rightarrow K'$ be a map of sets. The diagram

$$\begin{array}{ccc} \text{ISets}(K) & \xrightarrow{\Sigma_f} & \text{ISets}(K') \\ \text{Un}_K \downarrow & & \downarrow \text{Un}_{K'} \\ \text{FibSets}(K) & \xrightarrow[\Sigma_f]{} & \text{FibSets}(K') \end{array}$$

commutes.

00UX 7. *Interaction With Dependent Products.* Let $f: K \rightarrow K'$ be a map of sets. The diagram

$$\begin{array}{ccc} \text{ISets}(K) & \xrightarrow{\Pi_f} & \text{ISets}(K') \\ \text{Un}_K \downarrow & & \downarrow \text{Un}_{K'} \\ \text{FibSets}(K) & \xrightarrow[\Pi_f]{} & \text{FibSets}(K') \end{array}$$

commutes.

Proof. **Item 1**, Functoriality: Omitted.

Item 2, Interaction With Fibres: Omitted.

Item 3, As a Pullback: Omitted.

Item 4, As a Colimit: Clear.

Item 5, Interaction With Change of Indexing/Base: Indeed, we have

$$\begin{aligned}
 \text{Un}_K(f^*(X)) &\stackrel{\text{def}}{=} \text{Un}_K(X \circ f) \\
 &\stackrel{\text{def}}{=} \coprod_{x \in K} X_{f(x)} \\
 &\cong \left\{ (x, (y, a)) \in K \times \coprod_{y \in K'} X_y \mid f(x) = y \right\} \\
 &\cong K \times_{K'} \coprod_{y \in K'} X_y \\
 &\stackrel{\text{def}}{=} K \times_{K'} \text{Un}_{K'}(X) \\
 &\stackrel{\text{def}}{=} f^*(\text{Un}_{K'}(X))
 \end{aligned}$$

for each $X \in \text{Obj}(\text{!Sets}(K'))$. Similarly, it can be shown that we also have $\text{Un}_K(f^*(\phi)) = f^*(\text{Un}_{K'}(\phi))$ and that $\text{Un}_K \circ f^* = f^* \circ \text{Un}_{K'}$ also holds on morphisms.

Item 6, Interaction With Dependent Sums: Indeed, we have

$$\begin{aligned}
 \text{Un}_{K'}(\Sigma_f(X)) &\stackrel{\text{def}}{=} \coprod_{x \in K'} \Sigma_f(X)_x \\
 &\cong \coprod_{x \in K'} \coprod_{y \in f^{-1}(x)} X_y \\
 &\cong \coprod_{y \in K} X_y \\
 &\cong \text{Un}_K(X) \\
 &\stackrel{\text{def}}{=} \Sigma_f(\text{Un}_K(X))
 \end{aligned}$$

for each $X \in \text{Obj}(\text{!Sets}(K))$, where we have used ?? of ?? for the first bijection. Similarly, it can be shown that we also have $\text{Un}_{K'}(\Sigma_f(\phi)) = \Sigma_f(\phi_{\text{Un}_K})$ and that $\text{Un}_{K'} \circ \Sigma_f = \Sigma_f \circ \text{Un}_K$ also holds on morphisms.

Item 7, Interaction With Dependent Products: Indeed, we have

$$\begin{aligned} \text{Un}_{K'}(\Pi_f(X)) &\stackrel{\text{def}}{=} \coprod_{x \in K'} \Pi_f(X)_x \\ &\cong \coprod_{x \in K'} \prod_{y \in f^{-1}(x)} X_y \\ &\cong \left\{ (x, h) \in \coprod_{x \in K'} \text{Sets}\left(f^{-1}(x), \phi_{\text{Un}_K}^{-1}(f^{-1}(x))\right) \mid \phi \circ h = \text{id}_{f^{-1}(x)} \right\} \\ &\stackrel{\text{def}}{=} \Pi_f\left(\coprod_{y \in K} X_y\right) \\ &\stackrel{\text{def}}{=} \Pi_f(\text{Un}_K(X)) \end{aligned}$$

for each $X \in \text{Obj}(\text{ISets}(K))$, where we have used ?? of ?? for the first bijection. Similarly, it can be shown that we also have $\text{Un}_{K'}(\Pi_f(\phi)) = \Pi_f(\phi_{\text{Un}_K})$ and that $\text{Un}_{K'} \circ \Pi_f = \Pi_f \circ \text{Un}_K$ also holds on morphisms. \square

10.1.3 The Un/Straightening Equivalence

00UZ Theorem 10.1.3.1.1. We have an isomorphism of categories

$$(\text{St}_K \dashv \text{Un}_K): \quad \text{FibSets}(K) \begin{array}{c} \xrightarrow{\text{St}_K} \\ \perp \\ \xleftarrow{\text{Un}_K} \end{array} \text{ISets}(K).$$

Proof. Omitted. \square

10.2 Miscellany

10.2.1 Other Kinds of Un/Straightening

00V2 Remark 10.2.1.1.1. There are also other kinds of un/straightening for sets, where Sets is replaced by **Rel** or **Span**:

- *Un/Straightening With **Rel**, I.* We have an isomorphism of sets

$$\text{Rel}(A, B) \cong \text{Sets}(B \times A, \{\text{true}, \text{false}\}).$$

by the definition of a relation from A to B , ??.

- *Un/Straightening With **Rel**, II.* We have an equivalence of categories

$$\text{LaxFun}(K_{\text{disc}}, \mathbf{Rel}) \stackrel{\text{eq.}}{\cong} \text{Cats}_{/K_{\text{disc}}}^{\text{fth}},$$

where $\text{Cats}_{/K_{\text{disc}}}^{\text{fth}}$ is the full subcategory of $\text{Cats}_{/K_{\text{disc}}}$ spanned by the faithful functors; see [Nie04, Theorem 3.1].

- *Un/Straightening With Span, I.* For each $A, B \in \text{Obj}(\text{Sets})$, we have a morphism of sets

$$\text{Span}(A, B) \rightarrow \text{Sets}(A \times B, \mathbb{N} \cup \{\infty\})$$

which assemble into an equivalence of categories between $\text{Span}(\text{Sets})$ and the category MRel of “multirelations”; see ??.

- *Un/Straightening With Span, II.* We have an equivalence of categories

$$\text{LaxFun}(K_{\text{disc}}, \text{Span}) \xrightarrow{\text{eq}} \text{Cats}_{/K_{\text{disc}}};$$

see [nLa24a, Section 3].

Appendices

10.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets
5. Relations
6. Spans
7. Posets

Indexed and Fibred Sets

7. Indexed Sets
8. Fibred Sets
9. Un/Straightening for Indexed and Fibred Sets

Category Theory

11. Categories
12. Types of Morphisms in Categories
13. Adjunctions and the Yoneda Lemma

14. Constructions With Categories

15. Profunctors
16. Cartesian Closed Categories
17. Kan Extensions

Bicategories

18. Bicategories
19. Internal Adjunctions

Internal Category Theory

20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

<p>25. Monoids</p> <p>26. Constructions With Monoids</p> <p>Monoids With Zero</p> <p>27. Monoids With Zero</p> <p>28. Constructions With Monoids With Zero</p> <p>Groups</p> <p>29. Groups</p> <p>30. Constructions With Groups</p> <p>Hyper Algebra</p> <p>31. Hypermonoids</p> <p>32. Hypergroups</p> <p>33. Hypersemirings and Hyperrings</p> <p>34. Quantales</p> <p>Near-Rings</p> <p>35. Near-Semirings</p> <p>36. Near-Rings</p>	<p>Real Analysis</p> <p>37. Real Analysis in One Variable</p> <p>38. Real Analysis in Several Variables</p> <p>Measure Theory</p> <p>39. Measurable Spaces</p> <p>40. Measures and Integration</p> <p>Probability Theory</p> <p>40. Probability Theory</p> <p>Stochastic Analysis</p> <p>41. Stochastic Processes, Martingales, and Brownian Motion</p> <p>42. Itô Calculus</p> <p>43. Stochastic Differential Equations</p> <p>Differential Geometry</p> <p>44. Topological and Smooth Manifolds</p> <p>Schemes</p> <p>45. Schemes</p>
---	---

10.2 Other Chapters

<p>Sets</p> <p>1. Sets</p> <p>2. Constructions With Sets</p> <p>3. Pointed Sets</p> <p>4. Tensor Products of Pointed Sets</p> <p>5. Relations</p> <p>6. Spans</p> <p>7. Posets</p> <p>Indexed and Fibred Sets</p>	<p>7. Indexed Sets</p> <p>8. Fibred Sets</p> <p>9. Un/Straightening for Indexed and Fibred Sets</p> <p>Category Theory</p> <p>11. Categories</p> <p>12. Types of Morphisms in Categories</p> <p>13. Adjunctions and the Yoneda Lemma</p> <p>14. Constructions With Categories</p> <p>15. Profunctors</p>
---	--

16. Cartesian Closed Categories	Hyper Algebra
17. Kan Extensions	31. Hypermonoids
Bicategories	32. Hypergroups
18. Bicategories	33. Hypersemirings and Hyperrings
19. Internal Adjunctions	34. Quantales
Internal Category Theory	Near-Rings
20. Internal Categories	35. Near-Semirings
Cyclic Stuff	36. Near-Rings
21. The Cycle Category	Real Analysis
Cubical Stuff	37. Real Analysis in One Variable
22. The Cube Category	38. Real Analysis in Several Variables
Globular Stuff	Measure Theory
23. The Globe Category	39. Measurable Spaces
Cellular Stuff	40. Measures and Integration
24. The Cell Category	Probability Theory
Monoids	40. Probability Theory
25. Monoids	Stochastic Analysis
26. Constructions With Monoids	41. Stochastic Processes, Martingales, and Brownian Motion
Monoids With Zero	42. Itô Calculus
27. Monoids With Zero	43. Stochastic Differential Equations
28. Constructions With Monoids With Zero	Differential Geometry
Groups	44. Topological and Smooth Manifolds
29. Groups	Schemes
30. Constructions With Groups	45. Schemes

Part III

Category Theory

Chapter 11

Categories

00V3 Create tags (see [MSE 350788] for some of these):

1. define bicategory $\text{Adj}(C)$
2. internal **Hom** in categories of co/Cartesian fibrations
3. <https://mathoverflow.net/questions/460146/universal-property-of-isbell-duality>
4. <http://www.tac.mta.ca/tac/volumes/36/12/36-12abs.html>
5. justify adjunctions via homs
6. <https://ncatlab.org/nlab/show/enrichment+versus+internalisation>
7. <https://arxiv.org/pdf/2004.08964.pdf>

Contents

11.1 Categories	311
11.1.1 Foundations	311
11.1.2 Examples of Categories	313
11.1.3 Subcategories	315
11.1.4 Skeletons of Categories	316
11.1.5 Precomposition and Postcomposition	317
11.2 The Quadruple Adjunction With Sets.....	318
11.2.1 Statement	318
11.2.2 Connected Components of Categories	319
11.2.3 Sets of Connected Components of Categories	320
11.2.4 Connected Categories	321
11.2.5 Discrete Categories	321

11.2.6	Indiscrete Categories	323
11.3	Groupoids	324
11.3.1	Foundations	324
11.3.2	The Groupoid Completion of a Category	324
11.3.3	The Core of a Category	327
11.4	Functors.....	330
11.4.1	Foundations	330
11.4.2	Faithful Functors.....	332
11.4.3	Full Functors	333
11.4.4	Fully Faithful Functors	334
11.4.5	Essentially Surjective Functors	334
11.4.6	Conservative Functors	334
11.4.7	Equivalences of Categories.....	335
11.4.8	Isomorphisms of Categories	337
11.4.9	The Natural Transformation Associated to a Functor	338
11.5	Natural Transformations	339
11.5.1	Foundations	339
11.5.2	Vertical Composition of Natural Transformations.....	341
11.5.3	Horizontal Composition of Natural Transformations	344
11.5.4	Properties of Natural Transformations.....	347
11.5.5	Natural Isomorphisms.....	348
11.6	Categories of Categories.....	349
11.6.1	Functor Categories	349
11.6.2	The Category of Categories and Functors	352
11.6.3	The 2-Category of Categories, Functors, and Natural Transformations	353
11.6.4	The Category of Groupoids.....	354
11.6.5	The 2-Category of Groupoids	354
11.7	Miscellany	354
11.7.1	Concrete Categories	354
11.7.2	Balanced Categories	354
11.7.3	Monoid Actions on Objects of Categories	354
11.7.4	Group Actions on Objects of Categories	355
11.A	Other Chapters	355
11.2	Other Chapters	357

11.1 Categories

11.1.1 Foundations

00V6 Definition 11.1.1.1.1. A category $(C, \circ^C, \mathbb{M}^C)$ consists of^{1,2}

- *Objects.* A class $\text{Obj}(C)$ of **objects**;
- *Morphisms.* For each $A, B \in \text{Obj}(C)$, a class $\text{Hom}_C(A, B)$, called the **class of morphisms of C from A to B** ;
- *Identities.* For each $A \in \text{Obj}(C)$, a map of sets

$$\mathbb{M}_A^C : \text{pt} \rightarrow \text{Hom}_C(A, A),$$

called the **unit map of C at A** , determining a morphism

$$\text{id}_A : A \rightarrow A$$

of C , called the **identity morphism of A** ;

- *Composition.* For each $A, B, C \in \text{Obj}(C)$, a map of sets

$$\circ_{A,B,C}^C : \text{Hom}_C(B, C) \times \text{Hom}_C(A, B) \rightarrow \text{Hom}_C(A, C),$$

called the **composition map of C at (A, B, C)** ;

such that the following conditions are satisfied:

1. *Associativity.* The diagram

$$\begin{array}{ccc}
& \text{Hom}_C(C, D) \times (\text{Hom}_C(B, C) \times \text{Hom}_C(A, B)) & \\
& \swarrow \alpha_{\text{Hom}_C(C,D), \text{Hom}_C(B,C), \text{Hom}_C(A,B)}^{\text{Sets}} \quad \searrow \text{id}_{\text{Hom}_C(C,D)} \times \circ_{A,B,C}^C & \\
(\text{Hom}_C(C, D) \times \text{Hom}_C(B, C)) \times \text{Hom}_C(A, B) & & \text{Hom}_C(C, D) \times \text{Hom}_C(A, C) \\
& \downarrow \circ_{B,C,D}^C \times \text{id}_{\text{Hom}_C(A,B)} & \downarrow \circ_{A,C,D}^C \\
& \text{Hom}_C(B, D) \times \text{Hom}_C(A, B) & \xrightarrow{\circ_{A,B,D}^C} \text{Hom}_C(A, D)
\end{array}$$

¹Further Notation: We also write $C(A, B)$ for $\text{Hom}_C(A, B)$.

²Further Notation: We write $\text{Mor}(C)$ for the class of all morphisms of C .

commutes, i.e. for each composable triple (f, g, h) of morphisms of C , we have

$$(f \circ g) \circ h = f \circ (g \circ h).$$

2. *Left Unitality.* The diagram

$$\begin{array}{ccc} \text{pt} \times \text{Hom}_C(A, B) & & \\ \downarrow \mathbb{1}_B^C \times \text{id}_{\text{Hom}_C(A, B)} & \nearrow \lambda_{\text{Hom}_C(A, B)}^{\text{Sets}} & \\ \text{Hom}_C(B, B) \times \text{Hom}_C(A, B) & \xrightarrow{\circ_{A, B, B}^C} & \text{Hom}_C(A, B) \end{array}$$

commutes, i.e. for each morphism $f: A \rightarrow B$ of C , we have

$$\text{id}_B \circ f = f.$$

3. *Right Unitality.* The diagram

$$\begin{array}{ccc} \text{Hom}_C(A, B) \times \text{pt} & & \\ \downarrow \text{id}_{\text{Hom}_C(A, B)} \times \mathbb{1}_A^C & \nearrow \rho_{\text{Hom}_C(A, B)}^{\text{Sets}} & \\ \text{Hom}_C(A, B) \times \text{Hom}_C(A, A) & \xrightarrow{\circ_{A, A, B}^C} & \text{Hom}_C(A, B) \end{array}$$

commutes, i.e. for each morphism $f: A \rightarrow B$ of C , we have

$$f \circ \text{id}_A = f.$$

00V7 Definition 11.1.1.2. Let κ be a regular cardinal. A category C is

1. **Locally small** if, for each $A, B \in \text{Obj}(C)$, the class $\text{Hom}_C(A, B)$ is a set.
2. **Locally essentially small** if, for each $A, B \in \text{Obj}(C)$, the class

$$\text{Hom}_C(A, B) / \{\text{isomorphisms}\}$$

is a set.

3. **Small** if C is locally small and $\text{Obj}(C)$ is a set.
4. **κ -Small** if C is locally small, $\text{Obj}(C)$ is a set, and we have $\#\text{Obj}(C) < \kappa$.

11.1.2 Examples of Categories

00V9 Example 11.1.2.1.1. The **punctual category**³ is the category pt where

- *Objects.* We have

$$\text{Obj}(\text{pt}) \stackrel{\text{def}}{=} \{\star\};$$

- *Morphisms.* The unique Hom-set of pt is defined by

$$\text{Hom}_{\text{pt}}(\star, \star) \stackrel{\text{def}}{=} \{\text{id}_\star\};$$

- *Identities.* The unit map

$$\mu_\star^{\text{pt}} : \text{pt} \rightarrow \text{Hom}_{\text{pt}}(\star, \star)$$

of pt at \star is defined by

$$\text{id}_\star^{\text{pt}} \stackrel{\text{def}}{=} \text{id}_\star;$$

- *Composition.* The composition map

$$\circ_{\star, \star, \star}^{\text{pt}} : \text{Hom}_{\text{pt}}(\star, \star) \times \text{Hom}_{\text{pt}}(\star, \star) \rightarrow \text{Hom}_{\text{pt}}(\star, \star)$$

of pt at (\star, \star, \star) is given by the bijection $\text{pt} \times \text{pt} \cong \text{pt}$.

00VA Example 11.1.2.1.2. We have an isomorphism of categories⁴

$$\begin{array}{ccc} \text{Mon} & \longrightarrow & \text{Cats} \\ \downarrow \lrcorner & & \downarrow \text{Obj} \\ \text{Mon} \cong \text{pt} \times_{\text{Sets}} \text{Cats}, & & \\ \downarrow & & \downarrow \\ \text{pt} & \xrightarrow{[\text{pt}]} & \text{Sets} \end{array}$$

via the delooping functor $B : \text{Mon} \rightarrow \text{Cats}$ of ?? of ??.

Proof. Omitted. □

00VB Example 11.1.2.1.3. The **empty category** is the category \emptyset_{cat} where

³Further Terminology: Also called the **singleton category**.

⁴This can be enhanced to an isomorphism of 2-categories

$$\begin{array}{ccc} \text{Mon}_{2-\text{disc}} & \rightarrow & \text{Cats}_{2,*} \\ \downarrow \lrcorner & & \downarrow \text{Obj} \\ \text{Mon}_{2-\text{disc}} \cong \text{pt}_{\text{bi}} \times_{\text{Sets}_{2-\text{disc}}} \text{Cats}_{2,*}, & & \\ \downarrow & & \downarrow \\ \text{pt}_{\text{bi}} & \xrightarrow{[\text{pt}]} & \text{Sets}_{2-\text{disc}} \end{array}$$

between the discrete 2-category $\text{Mon}_{2-\text{disc}}$ on Mon and the 2-category of pointed categories with one object.

- *Objects.* We have

$$\text{Obj}(\emptyset_{\text{cat}}) \stackrel{\text{def}}{=} \emptyset;$$

- *Morphisms.* We have

$$\text{Mor}(\emptyset_{\text{cat}}) \stackrel{\text{def}}{=} \emptyset;$$

- *Identities and Composition.* Having no objects, \emptyset_{cat} has no unit nor composition maps.

00VC Example 11.1.2.1.4. The n th ordinal category is the category \bowtie where⁵

- *Objects.* We have

$$\text{Obj}(\bowtie) \stackrel{\text{def}}{=} \{[0], \dots, [n]\};$$

- *Morphisms.* For each $[i], [j] \in \text{Obj}(\bowtie)$, we have

$$\text{Hom}_{\bowtie}([i], [j]) \stackrel{\text{def}}{=} \begin{cases} \{\text{id}_{[i]}\} & \text{if } [i] = [j], \\ \{[i] \rightarrow [j]\} & \text{if } [j] < [i], \\ \emptyset & \text{if } [j] > [i]; \end{cases}$$

- *Identities.* For each $[i] \in \text{Obj}(\bowtie)$, the unit map

$$\mathbb{1}_{[i]}^{\bowtie} : \text{pt} \rightarrow \text{Hom}_{\bowtie}([i], [i])$$

of \bowtie at $[i]$ is defined by

$$\text{id}_{[i]}^{\bowtie} \stackrel{\text{def}}{=} \text{id}_{[i]};$$

⁵In other words, \bowtie is the category associated to the poset

$$[0] \rightarrow [1] \rightarrow \dots \rightarrow [n-1] \rightarrow [n].$$

The category \bowtie for $n \geq 2$ may also be defined in terms of \star and joins: we have isomorphisms of categories

$$\begin{aligned} \mathbb{1} &\cong \mathbb{1} \star \mathbb{1}, \\ \mathbb{1} &\cong \mathbb{1} \star \mathbb{1} \\ &\cong (\mathbb{1} \star \mathbb{1}) \star \mathbb{1}, \\ \mathbb{1} &\cong \mathbb{1} \star \mathbb{1} \\ &\cong (\mathbb{1} \star \mathbb{1}) \star \mathbb{1} \\ &\cong ((\mathbb{1} \star \mathbb{1}) \star \mathbb{1}) \star \mathbb{1}, \\ \mathbb{1} &\cong \mathbb{1} \star \mathbb{1} \\ &\cong (\mathbb{1} \star \mathbb{1}) \star \mathbb{1} \\ &\cong ((\mathbb{1} \star \mathbb{1}) \star \mathbb{1}) \star \mathbb{1} \\ &\cong (((\mathbb{1} \star \mathbb{1}) \star \mathbb{1}) \star \mathbb{1}) \star \mathbb{1}, \end{aligned}$$

and so on.

- *Composition.* For each $[i], [j], [k] \in \text{Obj}(\bowtie)$, the composition map

$$\circ_{[i],[j],[k]}^\bowtie : \text{Hom}_\bowtie([j], [k]) \times \text{Hom}_\bowtie([i], [j]) \rightarrow \text{Hom}_\bowtie([i], [k])$$

of \bowtie at $([i], [j], [k])$ is defined by

$$\begin{aligned} \text{id}_{[i]} \circ \text{id}_{[i]} &= \text{id}_{[i]}, \\ ([j] \rightarrow [k]) \circ ([i] \rightarrow [j]) &= ([i] \rightarrow [k]). \end{aligned}$$

00VD Example 11.1.2.1.5. Here we list all the other categories that appear throughout this work.

- The category Sets_* of pointed sets of ??.
- The category Rel of sets and relations of ??.
- The category $\text{Span}(A, B)$ of spans from a set A to a set B of ??.
- The category $\text{ISets}(K)$ of K -indexed sets of ??.
- The category ISets of indexed sets of ??.
- The category $\text{FibSets}(K)$ of K -fibred sets of ??.
- The category FibSets of fibred sets of ??.

11.1.3 Subcategories

Let C be a category.

00VF Definition 11.1.3.1.1. A **subcategory** of C is a category \mathcal{A} satisfying the following conditions:

1. *Objects.* We have $\text{Obj}(\mathcal{A}) \subset \text{Obj}(C)$.
2. *Morphisms.* For each $A, B \in \text{Obj}(\mathcal{A})$, we have

$$\text{Hom}_{\mathcal{A}}(A, B) \subset \text{Hom}_C(A, B).$$

3. *Identities.* For each $A \in \text{Obj}(\mathcal{A})$, we have

$$\mathbb{1}_A^{\mathcal{A}} = \mathbb{1}_A^C.$$

4. *Composition.* For each $A, B, C \in \text{Obj}(\mathcal{A})$, we have

$$\circ_{A,B,C}^{\mathcal{A}} = \circ_{A,B,C}^C.$$

00VG Definition 11.1.3.1.2. A subcategory \mathcal{A} of C is **full** if the canonical inclusion functor $\mathcal{A} \rightarrow C$ is full, i.e. if, for each $A, B \in \text{Obj}(\mathcal{A})$, the inclusion

$$\iota_{A,B}: \text{Hom}_{\mathcal{A}}(A, B) \hookrightarrow \text{Hom}_C(A, B)$$

is surjective (and thus bijective).

00VH Definition 11.1.3.1.3. A subcategory \mathcal{A} of a category C is **strictly full** if it satisfies the following conditions:

1. *Fullness.* The subcategory \mathcal{A} is full.
2. *Closedness Under Isomorphisms.* The class $\text{Obj}(\mathcal{A})$ is closed under isomorphisms.⁶

00VJ Definition 11.1.3.1.4. A subcategory \mathcal{A} of C is **wide**⁷ if $\text{Obj}(\mathcal{A}) = \text{Obj}(C)$.

11.1.4 Skeletons of Categories

00VL Definition 11.1.4.1.1. A⁸ **skeleton** of a category C is a full subcategory $\text{Sk}(C)$ with one object from each isomorphism class of objects of C .

00VM Definition 11.1.4.1.2. A category C is **skeletal** if $C \cong \text{Sk}(C)$.⁹

00VN Proposition 11.1.4.1.3. Let C be a category.

00VP 1. *Existence.* Assuming the axiom of choice, $\text{Sk}(C)$ always exists.

00VQ 2. *Pseudofunctoriality.* The assignment $C \mapsto \text{Sk}(C)$ defines a pseudofunctor

$$\text{Sk}: \text{Cats}_2 \rightarrow \text{Cats}_2.$$

00VR 3. *Uniqueness Up to Equivalence.* Any two skeletons of C are equivalent.

00VS 4. *Inclusions of Skeletons Are Equivalences.* The inclusion

$$\iota_C: \text{Sk}(C) \hookrightarrow C$$

of a skeleton of C into C is an equivalence of categories.

Proof. **Item 1, Existence:** See [nLab23, Section “Existence of Skeletons of Categories”].

Item 2, Pseudofunctoriality: See [nLab23, Section “Skeletons as an Endo-Pseudofunctor on \mathbf{Cat} ”].

Item 3, Uniqueness Up to Equivalence: Clear.

Item 4, Inclusions of Skeletons Are Equivalences: Clear. □

⁶That is, given $A \in \text{Obj}(\mathcal{A})$ and $C \in \text{Obj}(C)$, if $C \cong A$, then $C \in \text{Obj}(\mathcal{A})$.

⁷Further Terminology: Also called **lluf**.

⁸Due to Item 3 of Proposition 11.1.4.1.3, we often refer to any such full subcategory $\text{Sk}(C)$ of C as *the* skeleton of C .

⁹That is, C is **skeletal** if isomorphic objects of C are equal.

11.1.5 Precomposition and Postcomposition

Let C be a category and let $A, B, C \in \text{Obj}(C)$.

00VU Definition 11.1.5.1.1. Let $f: A \rightarrow B$ and $g: B \rightarrow C$ be morphisms of C .

- The **precomposition function associated to f** is the function

$$f^*: \text{Hom}_C(B, C) \rightarrow \text{Hom}_C(A, C)$$

defined by

$$f^*(\phi) \stackrel{\text{def}}{=} \phi \circ f$$

for each $\phi \in \text{Hom}_C(B, C)$.

- The **postcomposition function associated to g** is the function

$$g_*: \text{Hom}_C(A, B) \rightarrow \text{Hom}_C(A, C)$$

defined by

$$g_*(\phi) \stackrel{\text{def}}{=} g \circ \phi$$

for each $\phi \in \text{Hom}_C(A, B)$.

00VV Proposition 11.1.5.1.2. Let $A, B, C, D \in \text{Obj}(C)$ and let $f: A \rightarrow B$ and $g: B \rightarrow C$ be morphisms of C .

00VW 1. *Interaction Between Precomposition and Postcomposition.* We have

$$\begin{array}{ccc} \text{Hom}_C(B, C) & \xrightarrow{g_*} & \text{Hom}_C(B, D) \\ g_* \circ f^* = f^* \circ g_*, & f^* \downarrow & \downarrow f^* \\ \text{Hom}_C(A, C) & \xrightarrow{g_*} & \text{Hom}_C(A, D). \end{array}$$

00VX 2. *Interaction With Composition I.* We have

$$\begin{array}{ccc} \text{Hom}_C(X, A) & \xrightarrow{f_*} & \text{Hom}_C(X, B) \\ (g \circ f)^* = f^* \circ g^*, & \searrow (g \circ f)_* & \downarrow g_* \\ & & \text{Hom}_C(X, C), \\ \text{Hom}_C(C, X) & \xrightarrow{g^*} & \text{Hom}_C(B, X) \\ (g \circ f)_* = g_* \circ f_*, & \searrow (g \circ f)^* & \downarrow f^* \\ & & \text{Hom}_C(A, X). \end{array}$$

00VY 3. *Interaction With Composition II.* We have

$$\begin{array}{ccc} \text{pt} & \xrightarrow{[f]} & \text{Hom}_C(A, B) \\ & \searrow [g \circ f] & \downarrow g^* \\ & & \text{Hom}_C(A, C) \end{array} \quad \begin{array}{c} [g \circ f] = g_* \circ [f], \\ [g \circ f] = f^* \circ [g], \end{array} \quad \begin{array}{ccc} \text{pt} & \xrightarrow{[g]} & \text{Hom}_C(B, C) \\ & \searrow [g \circ f] & \downarrow f^* \\ & & \text{Hom}_C(A, C). \end{array}$$

00VZ 4. *Interaction With Composition III.* We have

$$\begin{array}{ccc} \text{Hom}_C(B, C) \times \text{Hom}_C(A, B) & \xrightarrow{\circ_{A,B,C}^C} & \text{Hom}_C(A, C) \\ f^* \circ \circ_{A,B,C}^C = \circ_{X,B,C}^C \circ (f^* \times \text{id}), & \text{id} \times f^* \downarrow & \downarrow f^* \\ \text{Hom}_C(B, C) \times \text{Hom}_C(X, B) & \xrightarrow{\circ_{X,B,C}^C} & \text{Hom}_C(X, C), \\ \text{Hom}_C(B, C) \times \text{Hom}_C(A, B) & \xrightarrow{\circ_{A,B,C}^C} & \text{Hom}_C(A, C) \\ g_* \circ \circ_{A,B,C}^C = \circ_{A,B,D}^C \circ (\text{id} \times g_*), & g_* \times \text{id} \downarrow & \downarrow g^* \\ \text{Hom}_C(B, D) \times \text{Hom}_C(A, B) & \xrightarrow{\circ_{A,B,D}^C} & \text{Hom}_C(A, D). \end{array}$$

00W0 5. *Interaction With Identities.* We have

$$\begin{aligned} (\text{id}_A)^* &= \text{id}_{\text{Hom}_C(A, B)}, \\ (\text{id}_B)_* &= \text{id}_{\text{Hom}_C(A, B)}. \end{aligned}$$

Proof. Item 1, Interaction Between Precomposition and Postcomposition: Clear.

Item 2, Interaction With Composition I: Clear.

Item 3, Interaction With Composition II: Clear.

Item 4, Interaction With Composition III: Clear.

Item 5, Interaction With Identities: Clear. □

11.2 The Quadruple Adjunction With Sets

11.2.1 Statement

Let C be a category.

00W3 **Proposition 11.2.1.1.1.** We have a quadruple adjunction

$$(\pi_0 \dashv (-)_{\text{disc}} \dashv \text{Obj} \dashv (-)_{\text{indisc}}): \quad \text{Sets} \leftrightarrow \text{Cats},$$

witnessed by bijections of sets

$$\begin{aligned}\text{Hom}_{\text{Sets}}(\pi_0(C), X) &\cong \text{Hom}_{\text{Cats}}(C, X_{\text{disc}}), \\ \text{Hom}_{\text{Cats}}(X_{\text{disc}}, C) &\cong \text{Hom}_{\text{Sets}}(X, \text{Obj}(C)), \\ \text{Hom}_{\text{Sets}}(\text{Obj}(C), X) &\cong \text{Hom}_{\text{Cats}}(C, X_{\text{indisc}}),\end{aligned}$$

natural in $C \in \text{Obj}(\text{Cats})$ and $X \in \text{Obj}(\text{Sets})$, where

- The functor

$$\pi_0: \text{Cats} \rightarrow \text{Sets},$$

the **connected components functor**, is the functor sending a category to its set of connected components of [Definition 11.2.3.1.1](#).

- The functor

$$(-)_{\text{disc}}: \text{Sets} \rightarrow \text{Cats},$$

the **discrete category functor**, is the functor sending a set to its associated discrete category of [Definition 11.2.5.1.1](#).

- The functor

$$\text{Obj}: \text{Cats} \rightarrow \text{Sets},$$

the **object functor**, is the functor sending a category to its set of objects.

- The functor

$$(-)_{\text{indisc}}: \text{Sets} \rightarrow \text{Cats},$$

the **indiscrete category functor**, is the functor sending a set to its associated indiscrete category of [Definition 11.2.6.1.1](#).

Proof. Omitted. □

11.2.2 Connected Components of Categories

Let C be a category.

00W5 Definition 11.2.2.1.1. A **connected component** of C is a full subcategory \mathcal{I} of C satisfying the following conditions:¹⁰

1. *Non-Emptiness.* We have $\text{Obj}(\mathcal{I}) \neq \emptyset$.
2. *Connectedness.* There exists a zigzag of arrows between any two objects of \mathcal{I} .

¹⁰In other words, a **connected component** of C is an element of the set $\text{Obj}(C)/\sim$ with \sim the equivalence relation generated by the relation \sim' obtained by declaring $A \sim' B$ iff there exists a morphism of C from A to B .

11.2.3 Sets of Connected Components of Categories

Let C be a category.

00W7 Definition 11.2.3.1.1. The **set of connected components** of C is the set $\pi_0(C)$ whose elements are the connected components of C .

00W8 Proposition 11.2.3.1.2. Let C be a category.

00W9 1. *Functoriality.* The assignment $C \mapsto \pi_0(C)$ defines a functor

$$\pi_0 : \text{Cats} \rightarrow \text{Sets}.$$

00WA 2. *Adjointness.* We have a quadruple adjunction

$$(\pi_0 \dashv (-)_{\text{disc}} \dashv \text{Obj} \dashv (-)_{\text{indisc}}) : \text{Sets} \rightleftarrows \text{Cats.}$$

00WB 3. *Interaction With Groupoids.* If C is a groupoid, then we have an isomorphism of categories

$$\pi_0(C) \cong K(C),$$

where $K(C)$ is the set of isomorphism classes of C of ??.

00WC 4. *Preservation of Colimits.* The functor π_0 of Item 1 preserves colimits. In particular, we have bijections of sets

$$\begin{aligned} \pi_0(C \coprod \mathcal{D}) &\cong \pi_0(C) \coprod \pi_0(\mathcal{D}), \\ \pi_0(C \coprod_{\mathcal{E}} \mathcal{D}) &\cong \pi_0(C) \coprod_{\pi_0(\mathcal{E})} \pi_0(\mathcal{D}), \\ \pi_0\left(\text{CoEq}\left(C \xrightarrow[G]{F} \mathcal{D}\right)\right) &\cong \text{CoEq}\left(\pi_0(C) \xrightarrow[\pi_0(G)]{\pi_0(F)} \pi_0(\mathcal{D})\right), \end{aligned}$$

natural in $C, \mathcal{D}, \mathcal{E} \in \text{Obj}(\text{Cats})$.

00WD 5. *Symmetric Strong Monoidality With Respect to Coproducts.* The connected components functor of Item 1 has a symmetric strong monoidal structure

$$\left(\pi_0, \pi_0^{\coprod}, \pi_{0|k}^{\coprod}\right) : (\text{Cats}, \coprod, \emptyset_{\text{cat}}) \rightarrow (\text{Sets}, \coprod, \emptyset),$$

being equipped with isomorphisms

$$\begin{aligned} \pi_{0|C,\mathcal{D}}^{\coprod} : \pi_0(C) \coprod \pi_0(\mathcal{D}) &\xrightarrow{\cong} \pi_0(C \coprod \mathcal{D}), \\ \pi_{0|k}^{\coprod} : \emptyset &\xrightarrow{\cong} \pi_0(\emptyset_{\text{cat}}), \end{aligned}$$

natural in $C, \mathcal{D} \in \text{Obj}(\text{Cats})$.

- 00WE** 6. *Symmetric Strong Monoidality With Respect to Products.* The connected components functor of [Item 1](#) has a symmetric strong monoidal structure

$$\left(\pi_0, \pi_0^\otimes, \pi_{0|*}^\otimes \right) : (\text{Cats}, \times, \text{pt}) \rightarrow (\text{Sets}, \times, \text{pt}),$$

being equipped with isomorphisms

$$\begin{aligned} \pi_{0|C,D}^\otimes : \pi_0(C) \times \pi_0(D) &\xrightarrow{\cong} \pi_0(C \times D), \\ \pi_{0|*}^\otimes : \text{pt} &\xrightarrow{\cong} \pi_0(\text{pt}), \end{aligned}$$

natural in $C, D \in \text{Obj}(\text{Cats})$.

Proof. [Item 1](#), *Functoriality:* Clear.

[Item 2](#), *Adjointness:* This is proved in [Proposition 11.2.1.1.1](#).

[Item 3](#), *Interaction With Groupoids:* Clear.

[Item 4](#), *Preservation of Colimits:* This follows from [Item 2](#) and ?? of ??.

[Item 5](#), *Symmetric Strong Monoidality With Respect to Coproducts:* Omitted.

[Item 6](#), *Symmetric Strong Monoidality With Respect to Products:* Omitted. \square

11.2.4 Connected Categories

- 00WG** **Definition 11.2.4.1.1.** A category C is **connected** if $\pi_0(C) \cong \text{pt}$.^{11,12}

11.2.5 Discrete Categories

Let X be a set.

- 00WJ** **Definition 11.2.5.1.1.** The **discrete category on a set** X is the category X_{disc} where

- *Objects.* We have

$$\text{Obj}(X_{\text{disc}}) \stackrel{\text{def}}{=} X;$$

- *Morphisms.* For each $A, B \in \text{Obj}(X_{\text{disc}})$, we have

$$\text{Hom}_{X_{\text{disc}}}(A, B) \stackrel{\text{def}}{=} \begin{cases} \text{id}_A & \text{if } A = B, \\ \emptyset & \text{if } A \neq B; \end{cases}$$

- *Identities.* For each $A \in \text{Obj}(X_{\text{disc}})$, the unit map

$$\mathbb{1}_A^{X_{\text{disc}}} : \text{pt} \rightarrow \text{Hom}_{X_{\text{disc}}}(A, A)$$

of X_{disc} at A is defined by

$$\text{id}_A^{X_{\text{disc}}} \stackrel{\text{def}}{=} \text{id}_A;$$

¹¹*Further Terminology:* A category is **disconnected** if it is not connected.

¹²*Example:* A groupoid is connected iff any two of its objects are isomorphic.

- *Composition.* For each $A, B, C \in \text{Obj}(X_{\text{disc}})$, the composition map

$$\circ_{A,B,C}^{X_{\text{disc}}} : \text{Hom}_{X_{\text{disc}}}(B, C) \times \text{Hom}_{X_{\text{disc}}}(A, B) \rightarrow \text{Hom}_{X_{\text{disc}}}(A, C)$$

of X_{disc} at (A, B, C) is defined by

$$\text{id}_A \circ \text{id}_A \stackrel{\text{def}}{=} \text{id}_A.$$

00WK Proposition 11.2.5.1.2. Let X be a set.

00WL 1. *Functoriality.* The assignment $X \mapsto X_{\text{disc}}$ defines a functor

$$(-)_{\text{disc}} : \text{Sets} \rightarrow \text{Cats}.$$

00WM 2. *Adjointness.* We have a quadruple adjunction

$$(\pi_0 \dashv (-)_{\text{disc}} \dashv \text{Obj} \dashv (-)_{\text{indisc}}) : \text{Sets} \rightleftarrows \text{Cats.}$$

```

    \begin{array}{ccc}
    (\pi_0 \dashv (-)_{\text{disc}} \dashv \text{Obj} \dashv (-)_{\text{indisc}}) : & \text{Sets} & \rightleftarrows \text{Cats.} \\
    & \begin{array}{c} \xrightarrow{\pi_0} \\ \downarrow \\ \xrightarrow{(-)_{\text{disc}}} \\ \downarrow \\ \xrightarrow{\text{Obj}} \\ \downarrow \\ \xrightarrow{(-)_{\text{indisc}}} \end{array} &
    \end{array}
  
```

00WN 3. *Symmetric Strong Monoidality With Respect to Coproducts.* The functor of **Item 1** has a symmetric strong monoidal structure

$$((-)_{\text{disc}}, (-)_{\text{disc}}^{\coprod}, (-)_{\text{disc}|\wp}^{\coprod}) : (\text{Sets}, \coprod, \emptyset) \rightarrow (\text{Cats}, \coprod, \emptyset_{\text{cat}}),$$

being equipped with isomorphisms

$$(-)_{\text{disc}|X,Y}^{\coprod} : X_{\text{disc}} \coprod Y_{\text{disc}} \xrightarrow{\cong} (X \coprod Y)_{\text{disc}},$$

$$(-)_{\text{disc}|\wp}^{\coprod} : \emptyset_{\text{cat}} \xrightarrow{\cong} \emptyset_{\text{disc}},$$

natural in $X, Y \in \text{Obj}(\text{Sets})$.

00WP 4. *Symmetric Strong Monoidality With Respect to Products.* The functor of **Item 1** has a symmetric strong monoidal structure

$$((-)_{\text{disc}}, (-)_{\text{disc}}^{\otimes}, (-)_{\text{disc}|\wp}^{\otimes}) : (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Cats}, \times, \text{pt}),$$

being equipped with isomorphisms

$$(-)_{\text{disc}|X,Y}^{\otimes} : X_{\text{disc}} \times Y_{\text{disc}} \xrightarrow{\cong} (X \times Y)_{\text{disc}},$$

$$(-)_{\text{disc}|\wp}^{\otimes} : \text{pt} \xrightarrow{\cong} \text{pt}_{\text{disc}},$$

natural in $X, Y \in \text{Obj}(\text{Sets})$.

Proof. **Item 1, Functoriality:** Clear.

Item 2, Adjointness: This is proved in [Proposition 11.2.1.1.1](#).

Item 3, Symmetric Strong Monoidality With Respect to Coproducts: Omitted.

Item 4, Symmetric Strong Monoidality With Respect to Products: Omitted. \square

11.2.6 Indiscrete Categories

00WR Definition 11.2.6.1.1. The **indiscrete category on a set** X^{13} is the category X_{indisc} where

- *Objects.* We have

$$\text{Obj}(X_{\text{indisc}}) \stackrel{\text{def}}{=} X;$$

- *Morphisms.* For each $A, B \in \text{Obj}(X_{\text{indisc}})$, we have

$$\text{Hom}_{X_{\text{disc}}}(A, B) \stackrel{\text{def}}{=} \{[A] \rightarrow [B]\};$$

- *Identities.* For each $A \in \text{Obj}(X_{\text{indisc}})$, the unit map

$$\text{id}_A^{X_{\text{indisc}}} : \text{pt} \rightarrow \text{Hom}_{X_{\text{indisc}}}(A, A)$$

of X_{indisc} at A is defined by

$$\text{id}_A^{X_{\text{indisc}}} \stackrel{\text{def}}{=} \{[A] \rightarrow [A]\};$$

- *Composition.* For each $A, B, C \in \text{Obj}(X_{\text{indisc}})$, the composition map

$$\circ_{A,B,C}^{X_{\text{indisc}}} : \text{Hom}_{X_{\text{indisc}}}(B, C) \times \text{Hom}_{X_{\text{indisc}}}(A, B) \rightarrow \text{Hom}_{X_{\text{indisc}}}(A, C)$$

of X_{disc} at (A, B, C) is defined by


$$([B] \rightarrow [C]) \circ ([A] \rightarrow [B]) \stackrel{\text{def}}{=} ([A] \rightarrow [C]).$$

00WS Proposition 11.2.6.1.2. Let X be a set.

00WT 1. Functoriality. The assignment $X \mapsto X_{\text{indisc}}$ defines a functor

$$(-)_{\text{indisc}} : \text{Sets} \rightarrow \text{Cats}.$$

00WU 2. Adjointness. We have a quadruple adjunction



00WV 3. Symmetric Strong Monoidality With Respect to Products. The functor of Item 1 has

¹³Further Terminology: Also called the **chaotic category** on X .

a symmetric strong monoidal structure

$$\left((-)_{\text{indisc}}, (-)_{\text{indisc}}^{\otimes}, (-)_{\text{indisc}|k}^{\otimes} \right): (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Cats}, \times, \text{pt}),$$

being equipped with isomorphisms

$$\begin{aligned} (-)_{\text{indisc}|X,Y}^{\otimes}: X_{\text{indisc}} \times Y_{\text{indisc}} &\xrightarrow{\cong} (X \times Y)_{\text{indisc}}, \\ (-)_{\text{indisc}|k}^{\otimes}: \text{pt} &\xrightarrow{\cong} \text{pt}_{\text{indisc}}, \end{aligned}$$

natural in $X, Y \in \text{Obj}(\text{Sets})$.

Proof. **Item 1, Functoriality:** Clear.

Item 2, Adjointness: This is proved in [Proposition 11.2.1.1.1](#).

Item 3, Symmetric Strong Monoidality With Respect to Products: Omitted. \square

11.3 Groupoids

11.3.1 Foundations

Let C be a category.

00WY Definition 11.3.1.1.1. A morphism $f: A \rightarrow B$ of C is an **isomorphism** if there exists a morphism $f^{-1}: B \rightarrow A$ of C such that

$$\begin{aligned} f \circ f^{-1} &= \text{id}_B, \\ f^{-1} \circ f &= \text{id}_A. \end{aligned}$$

00WZ Definition 11.3.1.1.2. A **groupoid** is a category in which every morphism is an isomorphism.

11.3.2 The Groupoid Completion of a Category

Let C be a category.

00X1 Definition 11.3.2.1.1. The **groupoid completion of C** ¹⁴ is the pair $(K_0(C), \iota_C)$ consisting of

- A groupoid $K_0(C)$;
- A functor $\iota_C: C \rightarrow K_0(C)$;

satisfying the following universal property:¹⁵

¹⁴Further Terminology: Also called the **Grothendieck groupoid** of C or the **Grothendieck groupoid completion** of C .

¹⁵See [Item 5 of Proposition 11.3.2.1.2](#) for an explicit construction.

(UP) Given another such pair (\mathcal{G}, i) , there exists a unique functor $K_0(C) \xrightarrow{\exists!} \mathcal{G}$ making the diagram

$$\begin{array}{ccc} & K_0(C) & \\ \iota_C \nearrow & \downarrow \exists! & \\ C & \xrightarrow{i} & \mathcal{G} \end{array}$$

commute.

00X2 Proposition 11.3.2.1.2. Let C be a category.

00X3 1. *Functoriality.* The assignment $C \mapsto K_0(C)$ defines a functor

$$K_0 : \text{Cats} \rightarrow \text{Grpd}.$$

00X4 2. *2-Functoriality.* The assignment $C \mapsto K_0(C)$ defines a 2-functor

$$K_0 : \text{Cats}_2 \rightarrow \text{Grpd}_2.$$

00X5 3. *Adjointness.* We have an adjunction

$$(K_0 \dashv \iota) : \text{Cats} \begin{array}{c} \xrightarrow{K_0} \\ \perp \\ \iota \end{array} \text{Grpd},$$

witnessed by a bijection of sets

$$\text{Hom}_{\text{Grpd}}(K_0(C), \mathcal{G}) \cong \text{Hom}_{\text{Cats}}(C, \mathcal{G}),$$

natural in $C \in \text{Obj}(\text{Cats})$ and $\mathcal{G} \in \text{Obj}(\text{Grpd})$, forming, together with the functor **Core** of Item 1 of [Proposition 11.3.3.1.3](#), a triple adjunction

$$(K_0 \dashv \iota \dashv \text{Core}) : \text{Cats} \begin{array}{c} \xrightarrow{K_0} \\ \perp \\ \iota \\ \perp \\ \text{Core} \end{array} \text{Grpd},$$

witnessed by bijections of sets

$$\text{Hom}_{\text{Grpd}}(K_0(C), \mathcal{G}) \cong \text{Hom}_{\text{Cats}}(C, \mathcal{G}),$$

$$\text{Hom}_{\text{Cats}}(\mathcal{G}, \mathcal{D}) \cong \text{Hom}_{\text{Grpd}}(\mathcal{G}, \text{Core}(\mathcal{D})),$$

natural in $C, \mathcal{D} \in \text{Obj}(\text{Cats})$ and $\mathcal{G} \in \text{Obj}(\text{Grpd})$.

00X6 4. *2-Adjointness.* We have a 2-adjunction

$$(K_0 \dashv \iota) : \text{Cats} \begin{array}{c} \xrightarrow{K_0} \\ \perp_2 \\ \iota \end{array} \text{Grpd},$$

witnessed by an isomorphism of categories

$$\text{Fun}(K_0(C), \mathcal{G}) \cong \text{Fun}(C, \mathcal{G}),$$

natural in $C \in \text{Obj}(\text{Cats})$ and $\mathcal{G} \in \text{Obj}(\text{Grpd})$, forming, together with the 2-functor Core of [Item 2 of Proposition 11.3.3.1.3](#), a triple 2-adjunction

$$(K_0 \dashv \iota \dashv \text{Core}): \text{Cats} \begin{array}{c} \xleftarrow{\quad K_0 \quad} \\[-1ex] \xleftarrow{\quad \iota \quad} \\[-1ex] \xrightarrow{\quad \text{Core} \quad} \end{array} \text{Grpd},$$

witnessed by isomorphisms of categories

$$\begin{aligned} \text{Fun}(K_0(C), \mathcal{G}) &\cong \text{Fun}(C, \mathcal{G}), \\ \text{Fun}(\mathcal{G}, \mathcal{D}) &\cong \text{Fun}(\mathcal{G}, \text{Core}(\mathcal{D})), \end{aligned}$$

natural in $C, \mathcal{D} \in \text{Obj}(\text{Cats})$ and $\mathcal{G} \in \text{Obj}(\text{Grpd})$.

- [00X7](#) 5. *Interaction With Classifying Spaces.* We have an isomorphism of groupoids

$$K_0(C) \cong \Pi_{\leq 1}(|N_\bullet(C)|),$$

natural in $C \in \text{Obj}(\text{Cats})$; i.e. the diagram

$$\begin{array}{ccc} \text{Cats} & \xrightarrow{K_0} & \text{Grp} \\ N_\bullet \downarrow & \Downarrow \gamma & \uparrow \Pi_{\leq 1} \\ s\text{Sets} & \xrightarrow{|-|} & \text{Top} \end{array}$$

commutes up to natural isomorphism.

- [00X8](#) 6. *Symmetric Strong Monoidality With Respect to Coproducts.* The groupoid completion functor of [Item 1](#) has a symmetric strong monoidal structure

$$\left(K_0, K_0^{\coprod}, K_{0|\mathbb{1}}^{\coprod} \right): (\text{Cats}, \coprod, \emptyset_{\text{cat}}) \rightarrow (\text{Grpd}, \coprod, \emptyset_{\text{cat}})$$

being equipped with isomorphisms

$$\begin{aligned} K_{0|C,D}^{\coprod}: K_0(C) \coprod K_0(D) &\xrightarrow{\cong} K_0(C \coprod D), \\ K_{0|\mathbb{1}}^{\coprod}: \emptyset_{\text{cat}} &\xrightarrow{\cong} K_0(\emptyset_{\text{cat}}), \end{aligned}$$

natural in $C, D \in \text{Obj}(\text{Cats})$.

- 00X9** 7. *Symmetric Strong Monoidality With Respect to Products.* The groupoid completion functor of **Item 1** has a symmetric strong monoidal structure

$$\left(K_0, K_0^\times, K_{0|*}^\times \right) : (\text{Cats}, \times, \text{pt}) \rightarrow (\text{Grpd}, \times, \text{pt})$$

being equipped with isomorphisms

$$\begin{aligned} K_{0|C,D}^\times : K_0(C) \times K_0(D) &\xrightarrow{\cong} K_0(C \times D), \\ K_{0|*}^\times : \text{pt} &\xrightarrow{\cong} K_0(\text{pt}), \end{aligned}$$

natural in $C, D \in \text{Obj}(\text{Cats})$.

Proof. **Item 1**, Functoriality: Omitted.

Item 2, 2-Functoriality: Omitted.

Item 3, Adjointness: Omitted.

Item 4, 2-Adjointness: Omitted.

Item 5, Interaction With Classifying Spaces: See Corollary 18.33 of <https://web.ma.utexas.edu/users/dafra/M392C-2012/Notes/lecture18.pdf>.

Item 6, Symmetric Strong Monoidality With Respect to Coproducts: Omitted.

Item 7, Symmetric Strong Monoidality With Respect to Products: Omitted. \square

11.3.3 The Core of a Category

Let C be a category.

- 00XB** **Definition 11.3.3.1.1.** The **core** of C is the pair $(\text{Core}(C), \iota_C)$ ¹⁶ consisting of

1. A groupoid $\text{Core}(C)$;
2. A functor $\iota_C : \text{Core}(C) \hookrightarrow C$;

satisfying the following universal property:

(UP) Given another such pair (\mathcal{G}, i) , there exists a unique functor $\mathcal{G} \xrightarrow{\exists!} \text{Core}(C)$ making the diagram

$$\begin{array}{ccc} & \text{Core}(C) & \\ & \nearrow \exists! & \downarrow \iota_C \\ \mathcal{G} & \xrightarrow{i} & C \end{array}$$

commute.

¹⁶Further Notation: Also written C^\simeq .

00XC Construction 11.3.3.1.2. The core of C is the wide subcategory of C spanned by the isomorphisms of C , i.e. the category $\text{Core}(C)$ where¹⁷

1. *Objects.* We have

$$\text{Obj}(\text{Core}(C)) \stackrel{\text{def}}{=} \text{Obj}(C);$$

2. *Morphisms.* The morphisms of $\text{Core}(C)$ are the isomorphisms of C .

Proof. This follows from the fact that functors preserve isomorphisms. \square

00XD Proposition 11.3.3.1.3. Let C be a category.

00XE 1. *Functoriality.* The assignment $C \mapsto \text{Core}(C)$ defines a functor

$$\text{Core}: \text{Cats} \rightarrow \text{Grpd}.$$

00XF 2. *2-Functoriality.* The assignment $C \mapsto \text{Core}(C)$ defines a 2-functor

$$\text{Core}: \text{Cats}_2 \rightarrow \text{Grpd}_2.$$

00XG 3. *Adjointness.* We have an adjunction

$$(\iota \dashv \text{Core}): \quad \text{Grpd} \begin{array}{c} \xrightarrow{\iota} \\ \perp \\ \xleftarrow{\text{Core}} \end{array} \text{Cats},$$

witnessed by a bijection of sets

$$\text{Hom}_{\text{Cats}}(\mathcal{G}, \mathcal{D}) \cong \text{Hom}_{\text{Grpd}}(\mathcal{G}, \text{Core}(\mathcal{D})),$$

natural in $\mathcal{G} \in \text{Obj}(\text{Grpd})$ and $\mathcal{D} \in \text{Obj}(\text{Cats})$, forming, together with the functor K_0 of Item 1 of Proposition 11.3.2.1.2, a triple adjunction

$$(K_0 \dashv \iota \dashv \text{Core}): \quad \text{Cats} \begin{array}{c} \xleftarrow{\iota} \\ \perp \\ \xrightarrow{\text{Core}} \\ \perp \\ \xleftarrow{K_0} \end{array} \text{Grpd},$$

witnessed by bijections of sets

$$\text{Hom}_{\text{Grpd}}(K_0(C), \mathcal{G}) \cong \text{Hom}_{\text{Cats}}(C, \mathcal{G}),$$

$$\text{Hom}_{\text{Cats}}(\mathcal{G}, \mathcal{D}) \cong \text{Hom}_{\text{Grpd}}(\mathcal{G}, \text{Core}(\mathcal{D})),$$

natural in $C, \mathcal{D} \in \text{Obj}(\text{Cats})$ and $\mathcal{G} \in \text{Obj}(\text{Grpd})$.

¹⁷*Slogan:* The groupoid $\text{Core}(C)$ is the maximal subgroupoid of C .

00XH 4. *2-Adjointness.* We have an adjunction

$$(\iota \dashv \text{Core}): \text{Grpd} \begin{array}{c} \xrightarrow{\iota} \\ \perp_2 \\ \xleftarrow{\text{Core}} \end{array} \text{Cats},$$

witnessed by an isomorphism of categories

$$\text{Fun}(\mathcal{G}, \mathcal{D}) \cong \text{Fun}(\mathcal{G}, \text{Core}(\mathcal{D})),$$

natural in $\mathcal{G} \in \text{Obj}(\text{Grpd})$ and $\mathcal{D} \in \text{Obj}(\text{Cats})$, forming, together with the 2-functor K_0 of [Item 2 of Proposition 11.3.2.1.2](#), a triple 2-adjunction

$$(K_0 \dashv \iota \dashv \text{Core}): \text{Cats} \begin{array}{c} \xrightarrow{K_0} \\ \perp_2 \\ \xleftarrow{\iota} \\ \perp_2 \\ \xleftarrow{\text{Core}} \end{array} \text{Grpd},$$

witnessed by isomorphisms of categories

$$\text{Fun}(K_0(C), \mathcal{G}) \cong \text{Fun}(C, \mathcal{G}),$$

$$\text{Fun}(\mathcal{G}, \mathcal{D}) \cong \text{Fun}(\mathcal{G}, \text{Core}(\mathcal{D})),$$

natural in $C, \mathcal{D} \in \text{Obj}(\text{Cats})$ and $\mathcal{G} \in \text{Obj}(\text{Grpd})$.

00XJ 5. *Symmetric Strong Monoidality With Respect to Products.* The core functor of [Item 1](#) has a symmetric strong monoidal structure

$$(\text{Core}, \text{Core}^\times, \text{Core}_\times^\times): (\text{Cats}, \times, \text{pt}) \rightarrow (\text{Grpd}, \times, \text{pt})$$

being equipped with isomorphisms

$$\begin{aligned} \text{Core}_{C,D}^\times: \text{Core}(C) \times \text{Core}(D) &\xrightarrow{\cong} \text{Core}(C \times D), \\ \text{Core}_\times^\times: \text{pt} &\xrightarrow{\cong} \text{Core}(\text{pt}), \end{aligned}$$

natural in $C, D \in \text{Obj}(\text{Cats})$.

00XK 6. *Symmetric Strong Monoidality With Respect to Coproducts.* The core functor of [Item 1](#) has a symmetric strong monoidal structure

$$(\text{Core}, \text{Core}^{\coprod}, \text{Core}_\times^{\coprod}): (\text{Cats}, \coprod, \emptyset_{\text{cat}}) \rightarrow (\text{Grpd}, \coprod, \emptyset_{\text{cat}})$$

being equipped with isomorphisms

$$\begin{aligned} \text{Core}_{C,D}^{\coprod}: \text{Core}(C) \coprod \text{Core}(D) &\xrightarrow{\cong} \text{Core}(C \coprod D), \\ \text{Core}_\times^{\coprod}: \emptyset_{\text{cat}} &\xrightarrow{\cong} \text{Core}(\emptyset_{\text{cat}}), \end{aligned}$$

natural in $C, D \in \text{Obj}(\text{Cats})$.

Proof. **Item 1**, Functoriality: Omitted.

Item 2, 2-Functoriality: Omitted.

Item 3, Adjointness: The adjunction $(K_0 \dashv \iota)$ follows from the universal property of the Gabriel–Zisman localisation of a category with respect to a class of morphisms (??), while the adjunction $(\iota \dashv \text{Core})$ is a reformulation of the universal property of the core of a category (Definition 11.3.3.1.1).¹⁸

Item 4, 2-Adjointness: Omitted.

Item 5, Symmetric Strong Monoidality With Respect to Products: Omitted.

Item 6, Symmetric Strong Monoidality With Respect to Coproducts: Omitted. \square

11.4 Functors

11.4.1 Foundations

Let C and \mathcal{D} be categories.

00XN **Definition 11.4.1.1.1.** A functor $F: C \rightarrow \mathcal{D}$ from C to \mathcal{D} ¹⁹ consists of²⁰

1. Action on Objects. A map of sets

$$F: \text{Obj}(C) \rightarrow \text{Obj}(\mathcal{D}),$$

called the **action on objects of F** ;

2. Action on Morphisms. For each $A, B \in \text{Obj}(C)$, a map

$$F_{A,B}: \text{Hom}_C(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F_A, F_B),$$

called the **action on morphisms of F at (A, B)** ²¹;

satisfying the following conditions:

1. Preservation of Identities. For each $A \in \text{Obj}(C)$, the diagram

$$\begin{array}{ccc} & \text{pt} & \\ & \downarrow \varepsilon_A^C & \searrow \varepsilon_{F_A}^{\mathcal{D}} \\ \text{Hom}_C(A, A) & \xrightarrow{F_{A,A}} & \text{Hom}_{\mathcal{D}}(F_A, F_A) \end{array}$$

commutes, i.e. we have

$$F(\text{id}_A) = \text{id}_{F_A}.$$

¹⁸Reference: [Rie17, Example 4.1.15]

¹⁹Further Terminology: Also called a **covariant functor**.

²⁰Further Notation: Given functors $F: C \rightarrow \mathcal{D}$ and $G: C^{\text{op}} \rightarrow \mathcal{D}$, we will sometimes write F_A for $F(A)$ (resp. G^A for $G(A)$) and F_f for $F(f)$ (resp. G^f for $G(f)$). This has been called Einstein notation in the literature.

²¹Further Terminology: Also called **action on Hom-sets of F at (A, B)** .

2. *Preservation of Composition.* For each $A, B, C \in \text{Obj}(C)$, the diagram

$$\begin{array}{ccc} \text{Hom}_C(B, C) \times \text{Hom}_C(A, B) & \xrightarrow{\circ_{A,B,C}^C} & \text{Hom}_C(A, C) \\ F_{B,C} \times F_{A,B} \downarrow & & \downarrow F_{A,C} \\ \text{Hom}_{\mathcal{D}}(F_B, F_C) \times \text{Hom}_{\mathcal{D}}(F_A, F_B) & \xrightarrow{\circ_{F_A, F_B, F_C}^{\mathcal{D}}} & \text{Hom}_{\mathcal{D}}(F_A, F_C) \end{array}$$

commutes, i.e. for each composable pair (g, f) of morphisms of C , we have

$$F(g \circ f) = F(g) \circ F(f).$$

00XP Example 11.4.1.1.2. The **identity functor** of a category C is the functor $\text{id}_C: C \rightarrow C$ where

1. *Action on Objects.* For each $A \in \text{Obj}(C)$, we have

$$\text{id}_C(A) \stackrel{\text{def}}{=} A;$$

2. *Action on Morphisms.* For each $A, B \in \text{Obj}(C)$, the action on morphisms map

$$(\text{id}_C)_{A,B}: \text{Hom}_C(A, B) \rightarrow \underbrace{\text{Hom}_C(\text{id}_C(A), \text{id}_C(B))}_{\stackrel{\text{def}}{=} \text{Hom}_C(A, B)}$$

of id_C at (A, B) is defined by

$$(\text{id}_C)_{A,B} \stackrel{\text{def}}{=} \text{id}_{\text{Hom}_C(A, B)}.$$

Proof. Preservation of Identities: We have $\text{id}_C(\text{id}_A) \stackrel{\text{def}}{=} \text{id}_A$ for each $A \in \text{Obj}(C)$ by definition.

Preservation of Compositions: For each composable pair $A \xrightarrow{f} B \xrightarrow{g} B$ of morphisms of C , we have

$$\begin{aligned} \text{id}_C(g \circ f) &\stackrel{\text{def}}{=} g \circ f \\ &\stackrel{\text{def}}{=} \text{id}_C(g) \circ \text{id}_C(f). \end{aligned}$$

This finishes the proof. □

00XQ Definition 11.4.1.1.3. The **composition** of two functors $F: C \rightarrow \mathcal{D}$ and $G: \mathcal{D} \rightarrow \mathcal{E}$ is the functor $G \circ F$ where

• *Action on Objects.* For each $A \in \text{Obj}(C)$, we have

$$[G \circ F](A) \stackrel{\text{def}}{=} G(F(A));$$

- *Action on Morphisms.* For each $A, B \in \text{Obj}(C)$, the action on morphisms map

$$(G \circ F)_{A,B} : \text{Hom}_C(A, B) \rightarrow \text{Hom}_{\mathcal{E}}(G_{F_A}, G_{F_B})$$

of $G \circ F$ at (A, B) is defined by

$$[G \circ F](f) \stackrel{\text{def}}{=} G(F(f)).$$

Proof. Preservation of Identities: For each $A \in \text{Obj}(C)$, we have

$$\begin{aligned} G_{F_{\text{id}_A}} &= G_{\text{id}_{F_A}} && (\text{functoriality of } F) \\ &= \text{id}_{G_{F_A}}. && (\text{functoriality of } G) \end{aligned}$$

Preservation of Composition: For each composable pair (g, f) of morphisms of C , we have

$$\begin{aligned} G_{F_{g \circ f}} &= G_{F_g \circ F_f} && (\text{functoriality of } F) \\ &= G_{F_g} \circ G_{F_f}. && (\text{functoriality of } G) \end{aligned}$$

This finishes the proof. □

00XR Proposition 11.4.1.1.4. Let $F: C \rightarrow \mathcal{D}$ be a functor.

00XS 1. *Preservation of Isomorphisms.* If f is an isomorphism in C , then $F(f)$ is an isomorphism in \mathcal{D} .²²

Proof. Item 1, Preservation of Isomorphisms: Indeed, we have

$$\begin{aligned} F(f)^{-1} \circ F(f) &= F(f^{-1} \circ f) \\ &= F(\text{id}_A) \\ &= \text{id}_{F(A)} \end{aligned}$$

and

$$\begin{aligned} F(f) \circ F(f)^{-1} &= F(f \circ f^{-1}) \\ &= F(\text{id}_A) \\ &= \text{id}_{F(A)}, \end{aligned}$$

showing $F(f)$ to be an isomorphism. □

11.4.2 Faithful Functors

Let C and \mathcal{D} be categories.

²²When the converse holds, we call F *conservative*, see Definition 11.4.6.1.1.

00XU Definition 11.4.2.1.1. A functor $F: C \rightarrow \mathcal{D}$ is **faithful** if, for each $A, B \in \text{Obj}(C)$, the action on morphisms map

$$F_{A,B}: \text{Hom}_C(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F_A, F_B)$$

of F at (A, B) is injective.

00XV Proposition 11.4.2.1.2. Let $F: C \rightarrow \mathcal{D}$ be a functor

00XW 1. Characterisations. The following conditions are equivalent:

- (a) The functor $F: C \rightarrow \mathcal{D}$ is faithful.
- (b) For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(C, \mathcal{X})$$

is faithful.

- (c) For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, C) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is faithful.

Proof. Item 1, Characterisations: Omitted. □

11.4.3 Full Functors

Let C and \mathcal{D} be categories.

00XY Definition 11.4.3.1.1. A functor $F: C \rightarrow \mathcal{D}$ is **full** if, for each $A, B \in \text{Obj}(C)$, the action on morphisms map

$$F_{A,B}: \text{Hom}_C(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F_A, F_B)$$

of F at (A, B) is surjective.

00XZ Proposition 11.4.3.1.2. Let $F: C \rightarrow \mathcal{D}$ be a functor

00Y0 1. Characterisations. The following conditions are equivalent:

- (a) The functor $F: C \rightarrow \mathcal{D}$ is full.
- (b) For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(C, \mathcal{X})$$

is full.

- (c) For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, C) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is full.

Proof. Item 1, Characterisations: Omitted. □

11.4.4 Fully Faithful Functors

Let C and \mathcal{D} be categories.

00Y2 Definition 11.4.4.1.1. A functor $F: C \rightarrow \mathcal{D}$ is **fully faithful** if F is full and faithful, i.e. if, for each $A, B \in \text{Obj}(C)$, the action on morphisms map

$$F_{A,B}: \text{Hom}_C(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F_A, F_B)$$

of F at (A, B) is bijective.

00Y3 Proposition 11.4.4.1.2. Let $F: C \rightarrow \mathcal{D}$ be a functor

00Y4 1. *Characterisations.* The following conditions are equivalent:

- (a) The functor $F: C \rightarrow \mathcal{D}$ is fully faithful.
- (b) For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the precomposition functor

$$F^*: \text{Fun}(\mathcal{D}, \mathcal{X}) \rightarrow \text{Fun}(C, \mathcal{X})$$

is fully faithful.

- (c) For each $\mathcal{X} \in \text{Obj}(\text{Cats})$, the postcomposition functor

$$F_*: \text{Fun}(\mathcal{X}, C) \rightarrow \text{Fun}(\mathcal{X}, \mathcal{D})$$

is fully faithful.

00Y5 2. *Conservativity.* If F is fully faithful, then F is conservative.

Proof. **Item 1, Characterisations:** Omitted.

Item 2, Conservativity: This is proved in **Item 2 of Proposition 11.4.6.1.2.**

□

11.4.5 Essentially Surjective Functors

Let C and \mathcal{D} be categories.

00Y7 Definition 11.4.5.1.1. A functor $F: C \rightarrow \mathcal{D}$ is **essentially surjective** if, for each $D \in \text{Obj}(\mathcal{D})$, there exists some object A of C with $F(A) \cong D$.

11.4.6 Conservative Functors

Let C and \mathcal{D} be categories.

00Y9 Definition 11.4.6.1.1. A functor $F: C \rightarrow \mathcal{D}$ is **conservative** if it satisfies the following condition:

- (★) For each $f \in \text{Mor}(C)$, if $F(f)$ is an isomorphism in \mathcal{D} , then f is an isomorphism in C .

00YA Proposition 11.4.6.1.2. Let $F: C \rightarrow \mathcal{D}$ be a functor.

00YB 1. *Characterisations.* The following conditions are equivalent:

- (a) The functor F is conservative.
- (b) For each $f \in \text{Mor}(C)$, the morphism $F(f)$ is an isomorphism in \mathcal{D} iff f is an isomorphism in C .

00YC 2. *Interaction With Fully Faithfulness.* Every fully faithful functor is conservative.

Proof. **Item 1, Characterisations:** This follows from **Item 1** of [Proposition 11.4.1.1.4](#).

Item 2, Interaction With Fully Faithfulness: Let $F: C \rightarrow \mathcal{D}$ be a fully faithful functor, let $f: A \rightarrow B$ be a morphism of C , and suppose that F_f is an isomorphism. We have

$$\begin{aligned} F(\text{id}_B) &= \text{id}_{F(B)} \\ &= F(f) \circ F(f)^{-1} \\ &= F(f \circ f^{-1}). \end{aligned}$$

Similarly, $F(\text{id}_A) = F(f^{-1} \circ f)$. But since F is fully faithful, we must have

$$\begin{aligned} f \circ f^{-1} &= \text{id}_B, \\ f^{-1} \circ f &= \text{id}_A, \end{aligned}$$

showing f to be an isomorphism. Thus F is conservative. \square

11.4.7 Equivalences of Categories

00YE **Definition 11.4.7.1.1.** Let C and \mathcal{D} be categories.

- An **equivalence of categories** between C and \mathcal{D} consists of a pair of functors

$$\begin{aligned} F: C &\rightarrow \mathcal{D}, \\ G: \mathcal{D} &\rightarrow C \end{aligned}$$

together with natural isomorphisms

$$\begin{aligned} \eta: \text{id}_C &\xrightarrow{\cong} G \circ F, \\ \epsilon: F \circ G &\xrightarrow{\cong} \text{id}_{\mathcal{D}}. \end{aligned}$$

- An **adjoint equivalence of categories** between C and \mathcal{D} is an equivalence (F, G, η, ϵ) between C and \mathcal{D} which is also an adjunction.

00YF **Proposition 11.4.7.1.2.** Let $F: C \rightarrow \mathcal{D}$ be a functor.

00YG 1. *Characterisations.* If C and \mathcal{D} are small²³, then the following conditions are

²³Otherwise there will be size issues. One can also work with large categories and universes, or require F

equivalent:²⁴

- 00YH** (a) The functor F is an equivalence of categories.
- 00YJ** (b) The functor F is fully faithful and essentially surjective.
- 00YK** (c) The induced functor

$$\uparrow F\text{Sk}(C): \text{Sk}(C) \rightarrow \text{Sk}(\mathcal{D})$$

is an *isomorphism* of categories.

- 00YL** 2. *Two-Out-of-Three*. Let

$$\begin{array}{ccc} C & \xrightarrow{G \circ F} & \mathcal{E} \\ F \searrow & \nearrow G & \\ \mathcal{D} & & \end{array}$$

be a diagram in Cats . If two out of the three functors among F , G , and $G \circ F$ are equivalences of categories, then so is the third.

- 00YM** 3. *Stability Under Composition*. Let

$$C \xrightleftharpoons[G]{F} \mathcal{D} \xrightleftharpoons[G']{F'} \mathcal{E}$$

be a diagram in Cats . If (F, G) and (F', G') are equivalences of categories, then so is their composite $(F' \circ F, G' \circ G)$.

- 00YN** 4. *Equivalences vs. Adjoint Equivalences*. Every equivalence of categories can be promoted to an adjoint equivalence.²⁵

- 00YP** 5. *Interaction With Groupoids*. If C and \mathcal{D} are groupoids, then the following conditions are equivalent:

- (a) The functor F is an equivalence of groupoids.
- (b) The following conditions are satisfied:
 - i. The functor F induces a bijection

$$\pi_0(F): \pi_0(C) \rightarrow \pi_0(\mathcal{D})$$

of sets.

to be *constructively* essentially surjective; see [MSE1465107].

²⁴In ZFC, the equivalence between Item 1a and Item 1b is equivalent to the axiom of choice; see [MO 119454].

In Univalent Foundations, this is true without requiring neither the axiom of choice nor the law of the excluded middle.

²⁵More precisely, we can promote an equivalence of categories (F, G, η, ϵ) to adjoint equivalences (F, G, η', ϵ) and (F, G, η, ϵ') .

ii. For each $A \in \text{Obj}(C)$, the induced map

$$F_{x,x} : \text{Aut}_C(A) \rightarrow \text{Aut}_{\mathcal{D}}(F_A)$$

is an isomorphism of groups.

Proof. **Item 1, Characterisations:** We claim that **Items 1a** to **1c** are indeed equivalent:

1. **Item 1a** \implies **Item 1b**. Clear.

2. **Item 1b** \implies **Item 1a**. Since F is essentially surjective and C and \mathcal{D} are small, we can choose, using the axiom of choice, for each $B \in \text{Obj}(\mathcal{D})$, an object j_B of C and an isomorphism $i_B : B \rightarrow F_{j_B}$ of \mathcal{D} .

Since F is fully faithful, we can extend the assignment $B \mapsto j_B$ to a *unique* functor $j : \mathcal{D} \rightarrow C$ such that the isomorphisms $i_B : B \rightarrow F_{j_B}$ assemble into a natural isomorphism $\eta : \text{id}_{\mathcal{D}} \xrightarrow{\cong} F \circ j$, with a similar natural isomorphism $\epsilon : \text{id}_C \xrightarrow{\cong} j \circ F$. Hence F is an equivalence.

3. **Item 1a** \implies **Item 1c**. This follows from ??.

Item 2, Two-Out-of-Three: Omitted.

Item 3, Stability Under Composition: Clear.

Item 4, Equivalences vs. Adjoint Equivalences: See [Rie17, Proposition 4.4.5].

Item 5, Interaction With Groupoids: See [nLa24b, Proposition 4.4]. □

11.4.8 Isomorphisms of Categories

00YR Definition 11.4.8.1.1. An **isomorphism of categories** is a pair of functors

$$\begin{aligned} F &: C \rightarrow \mathcal{D}, \\ G &: \mathcal{D} \rightarrow C \end{aligned}$$

such that we have

$$G \circ F = \text{id}_C,$$

$$F \circ G = \text{id}_{\mathcal{D}}.$$

00YS Example 11.4.8.1.2. Categories can be equivalent but non-isomorphic. For example, the category consisting of two isomorphic objects is equivalent to pt, but not isomorphic to it.

00YT Proposition 11.4.8.1.3. Let $F : C \rightarrow \mathcal{D}$ be a functor.

00YU 1. **Characterisations.** If C and \mathcal{D} are small, then the following conditions are equivalent:

- (a) The functor F is an isomorphism of categories.
- (b) The functor F is fully faithful and a bijection on objects.

Proof. **Item 1, Characterisations:** Omitted, but similar to **Item 1 of Proposition 11.4.7.1.2.** □

11.4.9 The Natural Transformation Associated to a Functor

00YW Definition 11.4.9.1.1. Every functor $F: \mathcal{C} \rightarrow \mathcal{D}$ defines a natural transformation²⁶

$$\begin{array}{ccc} \mathcal{C}^{\text{op}} \times \mathcal{C} & \xrightarrow{F^{\text{op}} \times F} & \mathcal{D}^{\text{op}} \times \mathcal{D} \\ F^{\dagger}: \text{Hom}_{\mathcal{C}} \Rightarrow \text{Hom}_{\mathcal{D}} \circ (F^{\text{op}} \times F), & \swarrow \quad \searrow & \swarrow \quad \searrow \\ \text{Hom}_{\mathcal{C}} & \begin{array}{c} \cong \\ F^{\dagger} \\ \cong \end{array} & \text{Hom}_{\mathcal{D}} \\ & \downarrow & \downarrow \\ & \text{Sets}, & \end{array}$$

called the **natural transformation associated to F** , consisting of the collection

$$\left\{ F_{A,B}^{\dagger}: \text{Hom}_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(F_A, F_B) \right\}_{(A,B) \in \text{Obj}(\mathcal{C}^{\text{op}} \times \mathcal{C})}$$

with

$$F_{A,B}^{\dagger} \stackrel{\text{def}}{=} F_{A,B}.$$

Proof. The naturality condition for F^{\dagger} is the requirement that for each morphism

$$(\phi, \psi): (X, Y) \rightarrow (A, B)$$

of $\mathcal{C}^{\text{op}} \times \mathcal{C}$, the diagram

$$\begin{array}{ccc} \text{Hom}_{\mathcal{C}}(X, Y) & \xrightarrow{\phi^* \circ \psi_* = \psi_* \circ \phi^*} & \text{Hom}_{\mathcal{C}}(A, B) \\ \downarrow F_{X,Y} & & \downarrow F_{A,B} \\ \text{Hom}_{\mathcal{D}}(F_X, F_Y) & \xrightarrow{F(\phi)^* \circ F(\psi)_* = F(\psi)_* \circ F(\phi)^*} & \text{Hom}_{\mathcal{D}}(F_A, F_B), \end{array}$$

acting on elements as

$$\begin{array}{ccc} f & \longmapsto & \psi \circ f \circ \phi \\ \downarrow & & \downarrow \\ F(f) & \longmapsto & F(\psi) \circ F(f) \circ F(\phi) = F(\psi \circ f \circ \phi) \end{array}$$

commutes, which follows from the functoriality of F . □

00YX Proposition 11.4.9.1.2. Let $F: \mathcal{C} \rightarrow \mathcal{D}$ and $G: \mathcal{D} \rightarrow \mathcal{E}$ be functors.

00YY 1. *Interaction With Natural Isomorphisms.* The following conditions are equivalent:

- (a) The natural transformation $F^{\dagger}: \text{Hom}_{\mathcal{C}} \Rightarrow \text{Hom}_{\mathcal{D}} \circ (F^{\text{op}} \times F)$ associated to F is a natural isomorphism.

²⁶This is the 1-categorical version of ?? of ??.

(b) The functor F is fully faithful.

00YZ 2. *Interaction With Composition.* We have an equality of pasting diagrams

$$\begin{array}{ccc} C^{\text{op}} \times C & \xrightarrow{F^{\text{op}} \times F} & \mathcal{D}^{\text{op}} \times \mathcal{D} \xrightarrow{G^{\text{op}} \times G} \mathcal{E}^{\text{op}} \times \mathcal{E} \\ \searrow \text{Hom}_C \quad \swarrow F^\dagger \quad \downarrow \text{Hom}_{\mathcal{D}} \quad \nearrow G^\dagger \quad \searrow \text{Hom}_{\mathcal{E}} \\ & \text{Sets} & \end{array} = \begin{array}{ccc} C^{\text{op}} \times C & \xrightarrow{(G \circ F)^{\text{op}} \times (G \circ F)} & \mathcal{E}^{\text{op}} \times \mathcal{E}, \\ \searrow \text{Hom}_C \quad \swarrow (G \circ F)^\dagger \quad \searrow \text{Hom}_{\mathcal{E}} \\ & \text{Sets} & \end{array}$$

in Cats_2 , i.e. we have

$$(G \circ F)^\dagger = (G^\dagger \star \text{id}_{F^{\text{op}} \times F}) \circ F^\dagger.$$

00Z0 3. *Interaction With Identities.* We have

$$\text{id}_C^\dagger = \text{id}_{\text{Hom}_C(-1, -2)},$$

i.e. the natural transformation associated to id_C is the identity natural transformation of the functor $\text{Hom}_C(-1, -2)$.

Proof. **Item 1, Interaction With Natural Isomorphisms:** Clear.

Item 2, Interaction With Composition: Clear.

Item 3, Interaction With Identities: Clear. □

11.5 Natural Transformations

11.5.1 Foundations

Let C and \mathcal{D} be categories and $F, G: C \Rightarrow \mathcal{D}$ be functors.

00Z3 **Definition 11.5.1.1.1.** A **transformation**^{27,28} $\alpha: F \xrightarrow{\text{unnat}} G$ from F to G is a collection

$$\{\alpha_A: F(A) \rightarrow G(A)\}_{A \in \text{Obj}(C)}$$

of morphisms of \mathcal{D} .

²⁷Further Terminology: Also called an **unnatural transformation** for emphasis.

²⁸Further Notation: We write $\text{UnNat}(F, G)$ for the set of unnatural transformations from F to G .

00Z4 Definition 11.5.1.1.2. A **natural transformation**²⁹ $\alpha: F \Rightarrow G$ from F to G is a transformation

$$\{\alpha_A: F(A) \rightarrow G(A)\}_{A \in \text{Obj}(C)}$$

from F to G such that, for each morphism $f: A \rightarrow B$ of C , the diagram

$$\begin{array}{ccc} F(A) & \xrightarrow{F(f)} & F(B) \\ \alpha_A \downarrow & & \downarrow \alpha_B \\ G(A) & \xrightarrow{G(f)} & G(B) \end{array}$$

commutes.^{30,31}

00Z5 Example 11.5.1.1.3. The **identity natural transformation** $\text{id}_F: F \Rightarrow F$ of F is the natural transformation consisting of the collection

$$\{\text{id}_{F(A)}: F(A) \rightarrow F(A)\}_{A \in \text{Obj}(C)}.$$

Proof. The naturality condition for id_F is the requirement that, for each morphism $f: A \rightarrow B$ of C , the diagram

$$\begin{array}{ccc} F(A) & \xrightarrow{F(f)} & F(B) \\ \text{id}_{F(A)} \downarrow & & \downarrow \text{id}_{F(B)} \\ F(A) & \xrightarrow{F(f)} & F(B) \end{array}$$


commutes, which follows from unitality of the composition of C . □

00Z6 Definition 11.5.1.1.4. Two natural transformations $\alpha, \beta: F \Rightarrow G$ are **equal** if we have

$$\alpha_A = \beta_A$$

for each $A \in \text{Obj}(C)$.

²⁹Pictured in diagrams as



³⁰Further Terminology: The morphism $\alpha_A: F_A \rightarrow G_A$ is called the **component of α at A** .

³¹Further Notation: We write $\text{Nat}(F, G)$ for the set of natural transformations from F to G .

11.5.2 Vertical Composition of Natural Transformations

00Z8 Definition 11.5.2.1.1. The **vertical composition** of two natural transformations $\alpha: F \Rightarrow G$ and $\beta: G \Rightarrow H$ as in the diagram

$$\begin{array}{ccccc} & & F & & \\ & \swarrow \alpha & \downarrow & \searrow & \\ C & \xrightarrow{G} & \mathcal{D} & & \\ & \beta \downarrow & & & \\ & & H & & \end{array}$$

is the natural transformation $\beta \circ \alpha: F \Rightarrow H$ consisting of the collection

$$\{(\beta \circ \alpha)_A: F(A) \rightarrow H(A)\}_{A \in \text{Obj}(C)}$$

with

$$(\beta \circ \alpha)_A \stackrel{\text{def}}{=} \beta_A \circ \alpha_A$$

for each $A \in \text{Obj}(C)$.

Proof. The naturality condition for $\beta \circ \alpha$ is the requirement that the boundary of the diagram

$$\begin{array}{ccc} F(A) & \xrightarrow{F(f)} & F(B) \\ \alpha_A \downarrow & (1) & \downarrow \alpha_B \\ G(A) & \xrightarrow{G(f)} & G(B) \\ \beta_A \downarrow & (2) & \downarrow \beta_B \\ H(A) & \xrightarrow{H(f)} & H(B) \end{array}$$

commutes. Since

1. Subdiagram (1) commutes by the naturality of α ;
2. Subdiagram (2) commutes by the naturality of β ;

so does the boundary diagram. Hence $\beta \circ \alpha$ is a natural transformation. \square

00Z9 Proposition 11.5.2.1.2. Let C , \mathcal{D} , and \mathcal{E} be categories.

00ZA 1. *Functionality.* The assignment $(\beta, \alpha) \mapsto \beta \circ \alpha$ defines a function

$$\circ_{F,G,H}: \text{Nat}(G, H) \times \text{Nat}(F, G) \rightarrow \text{Nat}(F, H).$$

00ZB 2. *Associativity.* Let $F, G, H, K: \mathcal{C} \rightrightarrows \mathcal{D}$ be functors. The diagram

$$\begin{array}{ccc}
 & \text{Nat}(H, K) \times (\text{Nat}(G, H) \times \text{Nat}(F, G)) & \\
 & \swarrow \alpha_{\text{Nat}(H, K), \text{Nat}(G, H), \text{Nat}(F, G)}^{\text{Sets}} \quad \searrow \text{id}_{\text{Nat}(H, K) \times \circ_{F, G, H}} & \\
 (\text{Nat}(H, K) \times \text{Nat}(G, H)) \times \text{Nat}(F, G) & & \text{Nat}(H, K) \times \text{Nat}(F, H) \\
 \downarrow \circ_{G, H, K} \times \text{id}_{\text{Nat}(F, G)} & & \downarrow \circ_{F, H, K} \\
 \text{Nat}(G, K) \times \text{Nat}(F, G) & \xrightarrow{\circ_{F, G, K}} & \text{Nat}(F, K)
 \end{array}$$

commutes, i.e. given natural transformations

$$\begin{aligned}
 \alpha: F &\Longrightarrow G, \\
 \beta: G &\Longrightarrow H, \\
 \gamma: H &\Longrightarrow K,
 \end{aligned}$$

we have

$$(\gamma \circ \beta) \circ \alpha = \gamma \circ (\beta \circ \alpha).$$

00ZC 3. *Unitality.* Let $F, G: \mathcal{C} \rightrightarrows \mathcal{D}$ be functors.

(a) *Left Unitality.* The diagram

$$\begin{array}{ccc}
 \text{pt} \times \text{Nat}(F, G) & & \\
 \downarrow [\text{id}_G] \times \text{id}_{\text{Nat}(F, G)} & \searrow \lambda_{\text{Nat}(F, G)}^{\text{Sets}} & \\
 \text{Nat}(G, G) \times \text{Nat}(F, G) & \xrightarrow{\circ_{F, G, G}} & \text{Nat}(F, G)
 \end{array}$$

commutes, i.e. given a natural transformation $\alpha: F \Longrightarrow G$, we have

$$\text{id}_G \circ \alpha = \alpha.$$

(b) *Right Unitality.* The diagram

$$\begin{array}{ccc}
 \text{Nat}(F, G) \times \text{pt} & & \\
 \downarrow \text{id}_{\text{Nat}(F, G)} \times [\text{id}_F] & \searrow \rho_{\text{Nat}(F, G)}^{\text{Sets}} & \\
 \text{Nat}(F, G) \times \text{Nat}(F, F) & \xrightarrow{\circ_{F, F, G}^{\mathcal{C}}} & \text{Nat}(F, G)
 \end{array}$$

commutes, i.e. given a natural transformation $\alpha: F \Rightarrow G$, we have

$$\alpha \circ \text{id}_F = \alpha.$$

- 00ZD** 4. *Middle Four Exchange.* Let $F_1, F_2, F_3: C \rightarrow D$ and $G_1, G_2, G_3: D \rightarrow E$ be functors. The diagram

$$\begin{array}{ccc}
(\text{Nat}(G_2, G_3) \times \text{Nat}(G_1, G_2)) \times (\text{Nat}(F_2, F_3) \times \text{Nat}(F_1, F_2)) & \xleftarrow[\sim]{\mu_4} & (\text{Nat}(G_2, G_3) \times \text{Nat}(F_2, F_3)) \times (\text{Nat}(G_1, G_2) \times \text{Nat}(F_1, F_2)) \\
\downarrow \circ_{G_1, G_2, G_3} \times \circ_{F_1, F_2, F_3} & & \downarrow \star_{F_2, F_3, G_2, G_3} \times \star_{F_1, F_2, G_1, G_2} \\
\text{Nat}(G_1, G_3) \times \text{Nat}(F_1, F_3) & & \text{Nat}(G_2 \circ F_2, G_3 \circ F_3) \times \text{Nat}(G_1 \circ F_1, G_2 \circ F_2) \\
\searrow \star_{F_1, F_3, G_1, G_3} & & \swarrow \circ_{G_1 \circ F_1, G_2 \circ F_2, G_3 \circ F_3} \\
& \text{Nat}(G_1 \circ F_1, G_3 \circ F_3) &
\end{array}$$

commutes, i.e. given a diagram

$$\begin{array}{ccccc}
& F_1 & & G_1 & \\
& \alpha \Downarrow & & \beta \Downarrow & \\
C & \xrightarrow{F_2} & D & \xrightarrow{G_2} & E \\
\alpha' \Downarrow & \nearrow & \nearrow & \nearrow & \nearrow \\
& F_3 & & G_3 &
\end{array}$$

in Cats_2 , we have

$$(\beta' \star \alpha') \circ (\beta \star \alpha) = (\beta' \circ \beta) \star (\alpha' \circ \alpha).$$

Proof. **Item 1, Functionality:** Clear.

Item 2, Associativity: Indeed, we have

$$\begin{aligned}
((\gamma \circ \beta) \circ \alpha)_A &= (\gamma_A \circ \beta_A) \circ \alpha_A \\
&= \gamma_A \circ (\beta_A \circ \alpha_A) \\
&= (\gamma \circ (\beta \circ \alpha))_A
\end{aligned}$$

for each $A \in \text{Obj}(C)$, showing the desired equality.

Item 3, Unitality: We have

$$\begin{aligned}
(\text{id}_G \circ \alpha)_A &= \text{id}_G \circ \alpha_A \\
&= \alpha_A, \\
(\alpha \circ \text{id}_F)_A &= \alpha_A \circ \text{id}_F \\
&= \alpha_A
\end{aligned}$$

for each $A \in \text{Obj}(C)$, showing the desired equality.

Item 4, Middle Four Exchange: This is proved in [Item 4 of Proposition 11.5.3.1.2](#). \square

11.5.3 Horizontal Composition of Natural Transformations

00ZF Definition 11.5.3.1.1. The **horizontal composition**^{32,33} of two natural transformations $\alpha: F \Rightarrow G$ and $\beta: H \Rightarrow K$ as in the diagram

$$\begin{array}{ccc} C & \xrightarrow[F]{\alpha \parallel} & \mathcal{D} \\ & \downarrow G & \\ & & K \end{array} \quad \begin{array}{ccc} \mathcal{D} & \xrightarrow[H]{\beta \parallel} & \mathcal{E} \\ & \downarrow & \\ & & K \end{array}$$

of α and β is the natural transformation

$$\beta \star \alpha: (H \circ F) \Rightarrow (K \circ G),$$

as in the diagram

$$\begin{array}{ccc} C & \xrightarrow[H \circ F]{\parallel} & \mathcal{E}, \\ & \beta \star \alpha \downarrow & \\ & & K \circ G \end{array}$$

consisting of the collection

$$\{(\beta \star \alpha)_A: H(F(A)) \rightarrow K(G(A))\}_{A \in \text{Obj}(C)},$$

of morphisms of \mathcal{E} with

$$\begin{aligned} (\beta \star \alpha)_A &\stackrel{\text{def}}{=} \beta_{G(A)} \circ H(\alpha_A) \\ &= K(\alpha_A) \circ \beta_{F(A)}, \end{aligned} \quad \begin{array}{ccc} H(F(A)) & \xrightarrow{H(\alpha_A)} & H(G(A)) \\ \beta_{F(A)} \downarrow & & \downarrow \beta_{G(A)} \\ K(F(A)) & \xrightarrow{K(\alpha_A)} & K(G(A)). \end{array}$$

Proof. First, we claim that we indeed have

$$\begin{array}{ccc} H(F(A)) & \xrightarrow{H(\alpha_A)} & H(G(A)) \\ \beta_{G(A)} \circ H(\alpha_A) = K(\alpha_A) \circ \beta_{F(A)} \downarrow & & \downarrow \beta_{G(A)} \\ K(F(A)) & \xrightarrow{K(\alpha_A)} & K(G(A)). \end{array}$$

This is, however, simply the naturality square for β applied to the morphism $\alpha_A: F(A) \rightarrow G(A)$. Next, we check the naturality condition for $\beta \star \alpha$, which is the requirement that

³²Further Terminology: Also called the **Godement product** of α and β .

³³Horizontal composition forms a map

$$\star_{(F,H),(G,K)}: \text{Nat}(H,K) \times \text{Nat}(F,G) \rightarrow \text{Nat}(H \circ F, K \circ G).$$

the boundary of the diagram

$$\begin{array}{ccc}
 H(F(A)) & \xrightarrow{H(F(f))} & H(F(B)) \\
 H(\alpha_A) \downarrow & (1) & \downarrow H(\alpha_B) \\
 H(G(A)) & \xrightarrow{H(G(f))} & H(G(B)) \\
 \beta_{G(A)} \downarrow & (2) & \downarrow \beta_{G(B)} \\
 K(G(A)) & \xrightarrow{K(G(f))} & K(G(B))
 \end{array}$$

commutes. Since

1. Subdiagram (1) commutes by the naturality of α ;
2. Subdiagram (2) commutes by the naturality of β ;

so does the boundary diagram. Hence $\beta \circ \alpha$ is a natural transformation.³⁴

□

00ZG Proposition 11.5.3.1.2. Let C, \mathcal{D} , and \mathcal{E} be categories.

00ZH 1. *Functionality.* The assignment $(\beta, \alpha) \mapsto \beta \star \alpha$ defines a function

$$\star_{(F,G),(H,K)} : \text{Nat}(H, K) \times \text{Nat}(F, G) \rightarrow \text{Nat}(H \circ F, K \circ G).$$

00ZJ 2. *Associativity.* Let

$$C \xrightarrow[G_1]{F_1} \mathcal{D} \xrightarrow[G_2]{F_2} \mathcal{E} \xrightarrow[G_3]{F_3} \mathcal{F}$$

be a diagram in Cats_2 . The diagram

$$\begin{array}{ccc}
 \text{Nat}(F_3, G_3) \times \text{Nat}(F_2, G_2) \times \text{Nat}(F_1, G_1) & \xrightarrow{\star_{(F_2, G_2), (F_3, G_3)} \times \text{id}} & \text{Nat}(F_3 \circ F_2, G_3 \circ G_2) \times \text{Nat}(F_1, G_1) \\
 \downarrow \text{id} \times \star_{(F_1, G_1), (F_2, G_2)} & & \downarrow \star_{(F_3 \circ F_2), (G_3 \circ G_2, F_1, G_1)} \\
 \text{Nat}(F_3, G_3) \times \text{Nat}(F_2 \circ F_1, G_2 \circ G_1) & \xrightarrow{\star_{(F_2 \circ F_1), (G_2 \circ G_1, F_3, G_3)}} & \text{Nat}(F_3 \circ F_2 \circ F_1, G_3 \circ G_2 \circ G_1)
 \end{array}$$

commutes, i.e. given natural transformations

$$C \xrightarrow[G_1]{F_1} \mathcal{D} \xrightarrow[G_2]{F_2} \mathcal{E} \xrightarrow[G_3]{F_3} \mathcal{F},$$

we have

$$(\gamma \star \beta) \star \alpha = \gamma \star (\beta \star \alpha).$$

³⁴Reference: [Bor94, Proposition 1.3.4].

- 00ZK** 3. *Interaction With Identities.* Let $F: \mathcal{C} \rightarrow \mathcal{D}$ and $G: \mathcal{D} \rightarrow \mathcal{E}$ be functors. The diagram

$$\begin{array}{ccc} \text{pt} \times \text{pt} & \xrightarrow{[\text{id}_G] \times [\text{id}_F]} & \text{Nat}(G, G) \times \text{Nat}(F, F) \\ \uparrow \ddots & & \downarrow \star_{(F,F),(G,G)} \\ \text{pt} & \xrightarrow{[\text{id}_{G \circ F}]} & \text{Nat}(G \circ F, G \circ F) \end{array}$$

commutes, i.e. we have

$$\text{id}_G \star \text{id}_F = \text{id}_{G \circ F}.$$

- 00ZL** 4. *Middle Four Exchange.* Let $F_1, F_2, F_3: \mathcal{C} \rightarrow \mathcal{D}$ and $G_1, G_2, G_3: \mathcal{D} \rightarrow \mathcal{E}$ be functors. The diagram

$$\begin{array}{ccc} (\text{Nat}(G_2, G_3) \times \text{Nat}(G_1, G_2)) \times (\text{Nat}(F_2, F_3) \times \text{Nat}(F_1, F_2)) & \xleftarrow{\mu_4} & (\text{Nat}(G_2, G_3) \times \text{Nat}(F_2, F_3)) \times (\text{Nat}(G_1, G_2) \times \text{Nat}(F_1, F_2)) \\ \downarrow \circ_{G_1, G_2, G_3} \times \circ_{F_1, F_2, F_3} & & \downarrow \star_{F_2, F_3, G_2, G_3} \times \star_{F_1, F_2, G_1, G_2} \\ \text{Nat}(G_1, G_3) \times \text{Nat}(F_1, F_3) & & \text{Nat}(G_2 \circ F_2, G_3 \circ F_3) \times \text{Nat}(G_1 \circ F_1, G_2 \circ F_2) \\ & \searrow \star_{F_1, F_3, G_1, G_3} & \swarrow \circ_{G_1 \circ F_1, G_2 \circ F_2, G_3 \circ F_3} \\ & \text{Nat}(G_1 \circ F_1, G_3 \circ F_3) & \end{array}$$

commutes, i.e. given a diagram

$$\begin{array}{ccccc} & F_1 & & G_1 & \\ C & \xrightarrow{F_2} & \mathcal{D} & \xrightarrow{G_2} & \mathcal{E} \\ \alpha \Downarrow & & \beta \Downarrow & & \\ \alpha' \Downarrow & \nearrow & \nearrow & \nearrow & \\ & F_3 & & G_3 & \end{array}$$

in Cats_2 , we have

$$(\beta' \star \alpha') \circ (\beta \star \alpha) = (\beta' \circ \beta) \star (\alpha' \circ \alpha).$$

Proof. **Item 1, Functionality:** Clear.

Item 2, Associativity: Omitted.

Item 3, Interaction With Identities: We have

$$\begin{aligned} (\text{id}_G \star \text{id}_F)_A &\stackrel{\text{def}}{=} (\text{id}_G)_{F_A} \circ G_{(\text{id}_F)_A} \\ &\stackrel{\text{def}}{=} \text{id}_{G_{F_A}} \circ G_{\text{id}_{F_A}} \\ &= \text{id}_{G_{F_A}} \circ \text{id}_{G_{F_A}} \\ &= \text{id}_{G_{F_A}} \\ &\stackrel{\text{def}}{=} (\text{id}_{G \circ F})_A \end{aligned}$$

for each $A \in \text{Obj}(C)$, showing the desired equality.

Item 4, Middle Four Exchange: Let $A \in \text{Obj}(C)$ and consider the diagram

$$\begin{array}{ccccc}
 & & G_{F''_A} & & \\
 & \swarrow G_{\alpha'_A} & & \searrow \beta_{F''_A} & \\
 G_{F_A} \xrightarrow{G_{\alpha_A}} G_{F'_A} & (1) & G''_{F''_A} & \xrightarrow{\beta'_{F''_A}} G''_{F_A''} \\
 & \searrow \beta_{F'_A} & \nearrow G'_{\alpha'_A} & &
 \end{array}$$

The top composition is $((\beta' \circ \beta) \star (\alpha' \circ \alpha))_A$ and the bottom composition is $((\beta' \star \alpha') \circ (\beta \star \alpha))_A$. Since Subdiagram (1) commutes, they are equal. \square

11.5.4 Properties of Natural Transformations

00ZN Proposition 11.5.4.1.1. Let $F, G: C \Rightarrow \mathcal{D}$ be functors. The following data are equivalent:³⁵

- 00ZP** 1. A natural transformation $\alpha: F \Rightarrow G$.
- 00ZQ** 2. A functor $[\alpha]: C \rightarrow \mathcal{D}^*$ filling the diagram

$$\begin{array}{ccc}
 & \mathcal{D} & \\
 F \nearrow & \uparrow \text{ev}_0 & \\
 C & \xrightarrow{[\alpha]} & \mathcal{D}^*. \\
 G \searrow & \downarrow \text{ev}_1 & \\
 & \mathcal{D} &
 \end{array}$$

- 00ZR** 3. A functor $[\alpha]: C \times \mathbb{I} \rightarrow \mathcal{D}$ filling the diagram

$$\begin{array}{ccc}
 C & & \\
 \uparrow \text{ev}_0 & \searrow F & \\
 C \times \mathbb{I} & \xrightarrow{[\alpha]} & \mathcal{D}. \\
 \downarrow \text{ev}_1 & \nearrow G & \\
 C & &
 \end{array}$$

³⁵Taken from [MO 64365].

Proof. From Item 1 to Item 2 and Back: We may identify \mathcal{D}^* with $\text{Arr}(\mathcal{D})$. Given a natural transformation $\alpha: F \Rightarrow G$, we have a functor

$$\begin{aligned} [\alpha]: C &\longrightarrow \mathcal{D}^* \\ A &\longmapsto \alpha_A \\ (f: A \rightarrow B) &\longmapsto \left(\begin{array}{ccc} F_A & \xrightarrow{F_f} & F_B \\ \downarrow \alpha_A & & \downarrow \alpha_B \\ G_A & \xrightarrow{G_f} & G_B \end{array} \right) \end{aligned}$$

making the diagram in **Item 2** commute. Conversely, every such functor gives rise to a natural transformation from F to G , and these constructions are inverse to each other.

From Item 2 to Item 3 and Back: This follows from **Item 3** of [Proposition 11.6.1.1.2](#). \square

11.5.5 Natural Isomorphisms

00ZT Definition 11.5.5.1.1. A natural transformation $\alpha: F \Rightarrow G$ between functors $F, G: C \rightarrow \mathcal{D}$ between categories C and \mathcal{D} is a **natural isomorphism** if there exists a natural transformation $\alpha^{-1}: G \Rightarrow F$ such that

$$\begin{aligned} \alpha^{-1} \circ \alpha &= \text{id}_F, \\ \alpha \circ \alpha^{-1} &= \text{id}_G. \end{aligned}$$

00ZU Proposition 11.5.5.1.2. Let $\alpha: F \Rightarrow G$ be a natural transformation.

00ZV 1. *Characterisations.* The following conditions are equivalent:

00ZW (a) The natural transformation α is a natural isomorphism.

00ZX (b) For each $A \in \text{Obj}(C)$, the morphism $\alpha_A: F_A \rightarrow G_A$ is an isomorphism.

00ZY 2. *Componentwise Inverses of Natural Transformations Assemble Into Natural Transformations.* Let $\alpha^{-1}: G \Rightarrow F$ be a transformation such that, for each $A \in \text{Obj}(C)$, we have

$$\begin{aligned} \alpha_A^{-1} \circ \alpha_A &= \text{id}_{F(A)}, \\ \alpha_A \circ \alpha_A^{-1} &= \text{id}_{G(A)}. \end{aligned}$$

Then α^{-1} is a natural transformation.

Proof. Item 1, Characterisations: The implication **Item 1a** \Rightarrow **Item 1b** is clear, whereas the implication **Item 1b** \Rightarrow **Item 1a** follows from **Item 2**.

Item 2, Componentwise Inverses of Natural Transformations Assemble Into Natural Transformations: The naturality condition for α^{-1} corresponds to the commutativity of the

diagram

$$\begin{array}{ccc} G(A) & \xrightarrow{G(f)} & G(B) \\ \alpha_A^{-1} \downarrow & & \downarrow \alpha_B^{-1} \\ F(A) & \xrightarrow{F(f)} & F(B) \end{array}$$

for each $A, B \in \text{Obj}(C)$ and each $f \in \text{Hom}_C(A, B)$. Considering the diagram

$$\begin{array}{ccc} G(A) & \xrightarrow{G(f)} & G(B) \\ \alpha_A^{-1} \downarrow & (1) & \downarrow \alpha_B^{-1} \\ F(A) & \xrightarrow{F(f)} & F(B) \\ \alpha_A \downarrow & (2) & \downarrow \alpha_B \\ G(A) & \xrightarrow{G(f)} & G(B), \end{array}$$

where the boundary diagram as well as Subdiagram (2) commute, we have

$$\begin{aligned} G(f) &= G(f) \circ \text{id}_{G(A)} \\ &= G(f) \circ \alpha_A \circ \alpha_A^{-1} \\ &= \alpha_B \circ F(f) \circ \alpha_A^{-1}. \end{aligned}$$

Postcomposing both sides with α_B^{-1} , we get

$$\begin{aligned} \alpha_B^{-1} \circ G(f) &= \alpha_B^{-1} \circ \alpha_B \circ F(f) \circ \alpha_A^{-1} \\ &= \text{id}_{F(B)} \circ F(f) \circ \alpha_A^{-1} \\ &= F(f) \circ \alpha_A^{-1}, \end{aligned}$$

which is the naturality condition we wanted to show. Thus α^{-1} is a natural transformation.

□

11.6 Categories of Categories

11.6.1 Functor Categories

Let C be a category and \mathcal{D} be a small category.

0101 Definition 11.6.1.1.1. The **category of functors from C to \mathcal{D}** ³⁶ is the category $\text{Fun}(C, \mathcal{D})$ ³⁷ where

³⁶Further Terminology: Also called the **functor category** $\text{Fun}(C, \mathcal{D})$.

³⁷Further Notation: Also written \mathcal{D}^C and $[C, \mathcal{D}]$.

- *Objects.* The objects of $\text{Fun}(C, \mathcal{D})$ are functors from C to \mathcal{D} ;
- *Morphisms.* For each $F, G \in \text{Obj}(\text{Fun}(C, \mathcal{D}))$, we have

$$\text{Hom}_{\text{Fun}(C, \mathcal{D})}(F, G) \stackrel{\text{def}}{=} \text{Nat}(F, G);$$

- *Identities.* For each $F \in \text{Obj}(\text{Fun}(C, \mathcal{D}))$, the unit map

$$\text{id}_F^{\text{Fun}(C, \mathcal{D})} : \text{pt} \rightarrow \text{Nat}(F, F)$$

of $\text{Fun}(C, \mathcal{D})$ at F is given by

$$\text{id}_F^{\text{Fun}(C, \mathcal{D})} \stackrel{\text{def}}{=} \text{id}_F,$$

where $\text{id}_F : F \Rightarrow F$ is the identity natural transformation of F of [Example 11.5.1.1.3](#);

- *Composition.* For each $F, G, H \in \text{Obj}(\text{Fun}(C, \mathcal{D}))$, the composition map

$$\circ_{F, G, H}^{\text{Fun}(C, \mathcal{D})} : \text{Nat}(G, H) \times \text{Nat}(F, G) \rightarrow \text{Nat}(F, H)$$

of $\text{Fun}(C, \mathcal{D})$ at (F, G, H) is given by

$$\beta \circ_{F, G, H}^{\text{Fun}(C, \mathcal{D})} \alpha \stackrel{\text{def}}{=} \beta \circ \alpha,$$

where $\beta \circ \alpha$ is the vertical composition of α and β of [Item 1 of Proposition 11.5.2.1.2](#).

0102 Proposition 11.6.1.1.2. Let C and \mathcal{D} be categories and let $F : C \rightarrow \mathcal{D}$ be a functor.

0103 1. Functoriality. The assignments $C, \mathcal{D}, (C, \mathcal{D}) \mapsto \text{Fun}(C, \mathcal{D})$ define functors

$$\begin{aligned} \text{Fun}(C, -_2) &: \text{Cats} \rightarrow \text{Cats}, \\ \text{Fun}(-_1, \mathcal{D}) &: \text{Cats}^{\text{op}} \rightarrow \text{Cats}, \\ \text{Fun}(-_1, -_2) &: \text{Cats}^{\text{op}} \times \text{Cats} \rightarrow \text{Cats}. \end{aligned}$$

0104 2. 2-Functoriality. The assignments $C, \mathcal{D}, (C, \mathcal{D}) \mapsto \text{Fun}(C, \mathcal{D})$ define 2-functors

$$\begin{aligned} \text{Fun}(C, -_2) &: \text{Cats}_2 \rightarrow \text{Cats}_2, \\ \text{Fun}(-_1, \mathcal{D}) &: \text{Cats}_2^{\text{op}} \rightarrow \text{Cats}_2, \\ \text{Fun}(-_1, -_2) &: \text{Cats}_2^{\text{op}} \times \text{Cats}_2 \rightarrow \text{Cats}_2. \end{aligned}$$

0105 3. Adjointness. We have adjunctions

$$\begin{aligned} (C \times - \dashv \text{Fun}(C, -)) : \text{Cats} &\begin{array}{c} \xrightarrow{C \times -} \\ \perp \\ \xleftarrow{\text{Fun}(C, -)} \end{array} \text{Cats}, \\ (- \times \mathcal{D} \dashv \text{Fun}(\mathcal{D}, -)) : \text{Cats} &\begin{array}{c} \xrightarrow{- \times \mathcal{D}} \\ \perp \\ \xleftarrow{\text{Fun}(\mathcal{D}, -)} \end{array} \text{Cats}, \end{aligned}$$

witnessed by bijections of sets

$$\begin{aligned}\text{Hom}_{\text{Cats}}(C \times \mathcal{D}, \mathcal{E}) &\cong \text{Hom}_{\text{Cats}}(\mathcal{D}, \text{Fun}(C, \mathcal{E})), \\ \text{Hom}_{\text{Cats}}(C \times \mathcal{D}, \mathcal{E}) &\cong \text{Hom}_{\text{Cats}}(C, \text{Fun}(\mathcal{D}, \mathcal{E})),\end{aligned}$$

natural in $C, \mathcal{D}, \mathcal{E} \in \text{Obj}(\text{Cats})$.

0106 4. *2-Adjointness.* We have 2-adjunctions

$$\begin{aligned}(C \times - \dashv \text{Fun}(C, -)): \quad \text{Cats}_2 &\begin{array}{c} \xrightarrow{\quad C \times - \quad} \\ \begin{array}{c} \xleftarrow{\perp_2} \\ \text{Fun}(C, -) \end{array} \end{array} \text{Cats}_2, \\ (- \times \mathcal{D} \dashv \text{Fun}(\mathcal{D}, -)): \quad \text{Cats}_2 &\begin{array}{c} \xrightarrow{\quad - \times \mathcal{D} \quad} \\ \begin{array}{c} \xleftarrow{\perp_2} \\ \text{Fun}(\mathcal{D}, -) \end{array} \end{array} \text{Cats}_2,\end{aligned}$$

witnessed by isomorphisms of categories

$$\begin{aligned}\text{Fun}(C \times \mathcal{D}, \mathcal{E}) &\cong \text{Fun}(\mathcal{D}, \text{Fun}(C, \mathcal{E})), \\ \text{Fun}(C \times \mathcal{D}, \mathcal{E}) &\cong \text{Fun}(C, \text{Fun}(\mathcal{D}, \mathcal{E})),\end{aligned}$$

natural in $C, \mathcal{D}, \mathcal{E} \in \text{Obj}(\text{Cats}_2)$.

0107 5. *Trivial Functor Categories.* We have a canonical isomorphism of categories

$$\text{Fun}(\text{pt}, C) \cong C,$$

natural in $C \in \text{Obj}(\text{Cats})$.

0108 6. *Objectwise Computation of Co/Limits.* Let

$$D: \mathcal{I} \rightarrow \text{Fun}(C, \mathcal{D})$$

be a diagram in $\text{Fun}(C, \mathcal{D})$. We have isomorphisms

$$\begin{aligned}\lim(D)_A &\cong \lim_{i \in \mathcal{I}}(D_i(A)), \\ \text{colim}(D)_A &\cong \text{colim}_{i \in \mathcal{I}}(D_i(A)),\end{aligned}$$

naturally in $A \in \text{Obj}(C)$.

0109 7. *Bicompleteness.* If \mathcal{E} is co/complete, then so is $\text{Fun}(C, \mathcal{E})$.

010A 8. *Abelianness.* If \mathcal{E} is abelian, then so is $\text{Fun}(C, \mathcal{E})$.

010B 9. *Monomorphisms and Epimorphisms.* Let $\alpha: F \rightarrow G$ be a morphism of $\text{Fun}(C, \mathcal{D})$. The following conditions are equivalent:

(a) The natural transformation

$$\alpha: F \Rightarrow G$$

is a monomorphism (resp. epimorphism) in $\text{Fun}(C, \mathcal{D})$.

(b) For each $A \in \text{Obj}(C)$, the morphism

$$\alpha_A: F_A \rightarrow G_A$$

is a monomorphism (resp. epimorphism) in \mathcal{D} .

Proof. **Item 1, Functoriality:** Omitted.

Item 2, 2-Functoriality: Omitted.

Item 3, Adjointness: Omitted.

Item 4, 2-Adjointness: Omitted.

Item 5, Trivial Functor Categories: Omitted.

Item 6, Objectwise Computation of Co/Limits: Omitted.

Item 7, Bicompleteness: This follows from ??.

Item 8, Abelianness: Omitted.

Item 9, Monomorphisms and Epimorphisms: Omitted. □

11.6.2 The Category of Categories and Functors

010D Definition 11.6.2.1.1. The **category of (small) categories and functors** is the category Cats where

- **Objects.** The objects of Cats are small categories;
- **Morphisms.** For each $C, \mathcal{D} \in \text{Obj}(\text{Cats})$, we have

$$\text{Hom}_{\text{Cats}}(C, \mathcal{D}) \stackrel{\text{def}}{=} \text{Obj}(\text{Fun}(C, \mathcal{D}));$$

- **Identities.** For each $C \in \text{Obj}(\text{Cats})$, the unit map

$$\text{id}_C^{\text{Cats}}: \text{pt} \rightarrow \text{Hom}_{\text{Cats}}(C, C)$$

of Cats at C is defined by

$$\text{id}_C^{\text{Cats}} \stackrel{\text{def}}{=} \text{id}_C,$$

where $\text{id}_C: C \rightarrow C$ is the identity functor of C of [Example 11.4.1.1.2](#);

- **Composition.** For each $C, \mathcal{D}, \mathcal{E} \in \text{Obj}(\text{Cats})$, the composition map

$$\circ_{C, \mathcal{D}, \mathcal{E}}^{\text{Cats}}: \text{Hom}_{\text{Cats}}(\mathcal{D}, \mathcal{E}) \times \text{Hom}_{\text{Cats}}(C, \mathcal{D}) \rightarrow \text{Hom}_{\text{Cats}}(C, \mathcal{E})$$

of Cats at $(C, \mathcal{D}, \mathcal{E})$ is given by

$$G \circ_{C, \mathcal{D}, \mathcal{E}}^{\text{Cats}} F \stackrel{\text{def}}{=} G \circ F,$$

where $G \circ F: C \rightarrow \mathcal{E}$ is the composition of F and G of [Definition 11.4.1.1.3](#).

010E Proposition 11.6.2.1.2. Let C be a category.

010F 1. *Co/Completeness.* The category Cats is complete and cocomplete.

010G 2. *Cartesian Monoidal Structure.* The quadruple $(\text{Cats}, \times, \text{pt}, \text{Fun})$ is a Cartesian closed monoidal category.

Proof. *Item 1, Co/Completeness:* This follows from

Item 2, Cartesian Monoidal Structure: Omitted. \square

11.6.3 The 2-Category of Categories, Functors, and Natural Transformations

010J Definition 11.6.3.1.1. The **2-category of (small) categories, functors, and natural transformations** is the 2-category Cats_2 where

- *Objects.* The objects of Cats_2 are small categories;
- *Hom-Categories.* For each $C, D \in \text{Obj}(\text{Cats}_2)$, we have

$$\text{Hom}_{\text{Cats}_2}(C, D) \stackrel{\text{def}}{=} \text{Fun}(C, D);$$

- *Identities.* For each $C \in \text{Obj}(\text{Cats}_2)$, the unit functor

$$\text{id}_C^{\text{Cats}_2} : \text{pt} \rightarrow \text{Fun}(C, C)$$

of Cats_2 at C is the functor picking the identity functor $\text{id}_C : C \rightarrow C$ of C ;

- *Composition.* For each $C, D, E \in \text{Obj}(\text{Cats}_2)$, the composition bifunctor

$$\circ_{C, D, E}^{\text{Cats}_2} : \text{Hom}_{\text{Cats}_2}(D, E) \times \text{Hom}_{\text{Cats}_2}(C, D) \rightarrow \text{Hom}_{\text{Cats}_2}(C, E)$$

of Cats_2 at (C, D, E) is the functor where

- *Action on Objects.* For each object $(G, F) \in \text{Obj}(\text{Hom}_{\text{Cats}_2}(D, E) \times \text{Hom}_{\text{Cats}_2}(C, D))$, we have

$$\circ_{C, D, E}^{\text{Cats}_2}(G, F) \stackrel{\text{def}}{=} G \circ F;$$

- *Action on Morphisms.* For each morphism $(\beta, \alpha) : (K, H) \rightarrow (G, F)$ of $\text{Hom}_{\text{Cats}_2}(D, E) \times \text{Hom}_{\text{Cats}_2}(C, D)$, we have

$$\circ_{C, D, E}^{\text{Cats}_2}(\beta, \alpha) \stackrel{\text{def}}{=} \beta \star \alpha,$$

where $\beta \star \alpha$ is the horizontal composition of α and β of [Definition 11.5.3.1.1](#).

010K Proposition 11.6.3.1.2. Let C be a category.

010L 1. *2-Categorical Co/Completeness.* The 2-category Cats_2 is complete and cocomplete as a 2-category, having all 2-categorical and bicategorical co/limits.

Proof. *Item 1, Co/Completeness:* This follows from \square

11.6.4 The Category of Groupoids

010N Definition 11.6.4.1.1. The **category of (small) groupoids** is the full subcategory Grpd of Cats spanned by the groupoids.

11.6.5 The 2-Category of Groupoids

010Q Definition 11.6.5.1.1. The **2-category of (small) groupoids** is the full sub-2-category Grpd_2 of Cats_2 spanned by the groupoids.

11.7 Miscellany

11.7.1 Concrete Categories

010T Definition 11.7.1.1.1. A category C is **concrete** if there exists a faithful functor $F: C \rightarrow \text{Sets}$.

11.7.2 Balanced Categories

010V Definition 11.7.2.1.1. A category is **balanced** if every morphism which is both a monomorphism and an epimorphism is an isomorphism.

11.7.3 Monoid Actions on Objects of Categories

Let A be a monoid, let C be a category, and let $X \in \text{Obj}(C)$.

010X Definition 11.7.3.1.1. An **A -action on X** is a functor $\lambda: BA \rightarrow C$ with $\lambda(\star) = X$.

010Y Remark 11.7.3.1.2. In detail, an **A -action on X** is an A -action on $\text{End}_C(X)$, consisting of a morphism

$$\lambda: A \rightarrow \underbrace{\text{End}_C(X)}_{\stackrel{\text{def}}{=} \text{Hom}_C(X, X)}$$

satisfying the following conditions:

1. *Preservation of Identities.* We have

$$\lambda_{1_A} = \text{id}_X.$$

2. *Preservation of Composition.* For each $a, b \in A$, we have

$$\begin{array}{ccc} X & \xrightarrow{\lambda_a} & X \\ & \searrow \lambda_{ab} & \downarrow \lambda_b \\ & & X. \end{array}$$

$$\lambda_b \circ \lambda_a = \lambda_{ab},$$

11.7.4 Group Actions on Objects of Categories

Let G be a group, let C be a category, and let $X \in \text{Obj}(C)$.

0110 Definition 11.7.4.1.1. A G -action on X is a functor $\lambda: BG \rightarrow C$ with $\lambda(\star) = X$.

0111 Remark 11.7.4.1.2. In detail, a G -action on X is a G -action on $\text{Aut}_C(X)$, consisting of a morphism

$$\lambda: G \rightarrow \underbrace{\text{End}_C(X)}_{\stackrel{\text{def}}{=} \text{Hom}_C(X, X)}$$

satisfying the following conditions:

1. *Preservation of Identities.* We have

$$\lambda_{1_A} = \text{id}_X.$$

2. *Preservation of Composition.* For each $a, b \in A$, we have

$$\begin{array}{ccc} X & \xrightarrow{\lambda_a} & X \\ \lambda_b \circ \lambda_a = \lambda_{ab}, & \searrow \lambda_{ab} & \downarrow \lambda_b \\ & & X. \end{array}$$

Appendices

11.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets
5. Relations
6. Spans
7. Posets

Indexed and Fibred Sets

7. Indexed Sets

8. Fibred Sets

9. Un/Straightening for Indexed and Fibred Sets

Category Theory

11. Categories
12. Types of Morphisms in Categories
13. Adjunctions and the Yoneda Lemma
14. Constructions With Categories
15. Profunctors
16. Cartesian Closed Categories
17. Kan Extensions

Bicategories

- 18. Bicategories
- 19. Internal Adjunctions

Internal Category Theory

- 20. Internal Categories

Cyclic Stuff

- 21. The Cycle Category

Cubical Stuff

- 22. The Cube Category

Globular Stuff

- 23. The Globe Category

Cellular Stuff

- 24. The Cell Category

Monoids

- 25. Monoids
- 26. Constructions With Monoids

Monoids With Zero

- 27. Monoids With Zero
- 28. Constructions With Monoids With Zero

Groups

- 29. Groups
- 30. Constructions With Groups

Hyper Algebra

- 31. Hypermonoids

Hypergroups

- 32. Hypersemirings and Hyperrings
- 33. Quantales

Near-Rings

- 34. Near-Semirings
- 35. Near-Rings

Real Analysis

- 36. Real Analysis in One Variable
- 37. Real Analysis in Several Variables

Measure Theory

- 38. Measures and Integration

Probability Theory

- 39. Probability Theory

Stochastic Analysis

- 40. Stochastic Processes, Martingales, and Brownian Motion
- 41. Itô Calculus
- 42. Stochastic Differential Equations

Differential Geometry

- 43. Topological and Smooth Manifolds

Schemes

- 44. Schemes

11.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 12

Types of Morphisms in Categories

0112 Create tags (see [MSE 350788] for some of these):

1. ??
2. ??
3. ??
4. ??
5. write material on sections and retractions

Contents

12.1 Monomorphisms	359
12.1.1 Foundations	359
12.1.2 Monomorphism-Reflecting Functors	363
12.1.3 Split Monomorphisms.....	363
12.2 Epimorphisms	364
12.2.1 Foundations	364
12.2.2 Regular Epimorphisms	366
12.2.3 Effective Epimorphisms.....	367
12.2.4 Split Epimorphisms.....	367
12.A Other Chapters	367
12.2 Other Chapters	369

12.1 Monomorphisms

12.1.1 Foundations

Let C be a category.

0115 Definition 12.1.1.1. A morphism $m: A \rightarrow B$ of C is a **monomorphism** if, for each diagram of the form

$$m \circ f = m \circ g, \quad X \xrightarrow{\begin{smallmatrix} f \\ g \end{smallmatrix}} A \xrightarrow{m} B$$

in Cats , we have $f = g$.

0116 Example 12.1.1.2. Here are some examples of monomorphisms.

0117 1. Monomorphisms in Sets. The monomorphisms in Sets are precisely the injections.

Proof. **Item 1, Monomorphisms in Sets:** Let $f: A \rightarrow B$ be a morphism in Sets . Suppose that f is a monomorphism and consider the diagram

$$\{*\} \xrightarrow{\begin{smallmatrix} [x] \\ [y] \end{smallmatrix}} A \xrightarrow{f} B,$$

where $[x]$ and $[y]$ are the morphisms picking the elements x and y of A . If $f(x) = f(y)$, then $f \circ [x] = f \circ [y]$, and thus $[x] = [y]$ since f is a monomorphism. Hence $x = y$ and we see that f is injective.

Conversely, suppose that f is injective. Proceeding by contrapositive, we claim that given a pair of maps $g, h: X \rightrightarrows A$ such that $g \neq h$, then $f \circ g \neq f \circ h$. Indeed, as g and h are different maps, there must exist at least one element $x \in X$ such that $g(x) \neq h(x)$. But then we have $f(g(x)) \neq f(h(x))$, since f is injective. Thus $f \circ g \neq f \circ h$, and we are done, having showed that f is a monomorphism. \square

0118 Proposition 12.1.1.3. Let $f: A \rightarrow B$ be a morphism of C .

0119 1. Characterisations. The following conditions are equivalent:

- 011A** (a) The morphism f is a monomorphism.
- 011B** (b) For each $X \in \text{Obj}(C)$, the map of sets

$$f_*: \text{Hom}_{\text{Sets}}(X, A) \rightarrow \text{Hom}_{\text{Sets}}(X, B)$$

is injective.

- 011C** (c) The kernel pair of f is trivial, i.e. we have

$$\begin{array}{ccc} A & \xrightarrow{\text{id}_A} & A \\ A \times_B A \cong A, & \downarrow \text{id}_A & \downarrow f \\ A & \xrightarrow{f} & B. \end{array}$$

011D 2. Duality. The following conditions are equivalent:

- (a) The morphism $f: A \rightarrow B$ is a monomorphism in C .
- (b) The morphism $f^\dagger: B \rightarrow A$ is an epimorphism in C^{op} .

011E 3. *Monomorphisms vs. Injective Maps.* Let

- C be a concrete category as in ??;
- $\mathfrak{F}_C: C \rightarrow \text{Sets}$ be the forgetful functor from C to Sets;
- $f: A \rightarrow B$ be a morphism of C .

If \mathfrak{F}_C preserves pullbacks, then the following conditions are equivalent:

- (a) The morphism f is a monomorphism.
- (b) The morphism $\mathfrak{F}(f)_C$ is injective.

011F 4. *Stability Properties.* The class of all monomorphisms of C is stable under the following operations:

- (a) *Composition.* If f and g are monomorphisms, then so is $g \circ f$.¹
- (b) *Pullbacks.* Let

$$\begin{array}{ccc} A \times_C B & \longrightarrow & B \\ m' \downarrow & \lrcorner & \downarrow m \\ A & \longrightarrow & C \end{array}$$

be a diagram in C . If m is a monomorphism in C , then so is m' .

011G 5. *Morphisms From the Terminal Object Are Monomorphisms.* If C has a terminal object $\mathbb{1}_C$, then every morphism of C from $\mathbb{1}_C$ is a monomorphism.

Proof. **Item 1, Characterisations:** The equivalence between **Items 1a** and **1b** is clear. We claim that **Items 1a** and **1c** are equivalent:

1. **Item 1a \implies Item 1c:** Suppose that f is a monomorphism. Then A satisfies the universal property of the pullback:

$$\begin{array}{ccccc} P & \xrightarrow{\phi} & A & \xrightarrow{\text{id}_A} & A \\ \exists! \quad \phi \quad \dashv & \searrow & \downarrow & & \downarrow f \\ & & A & \xrightarrow{\text{id}_A} & B \\ & \phi \quad \swarrow & \downarrow & & \downarrow f \\ & & A & \xrightarrow{f} & B. \end{array}$$

¹Conversely, if $g \circ f$ is a monomorphism, then so is f .

2. *Item 1c \implies Item 1a:* Suppose that $A \cong A \times_B A$ and let $g, h: C \rightrightarrows A$ be a pair of morphisms. Consider the diagram

$$\begin{array}{ccccc} & & C & & \\ & \swarrow g & & \searrow h & \\ & & A & \xrightarrow{\text{id}_A} & A \\ & & \downarrow \lrcorner & & \downarrow f \\ & & A & \xrightarrow{f} & B. \end{array}$$

The universal property of the pullback says that there exists a unique morphism $C \rightarrow A$ making the diagram

$$\begin{array}{ccccc} & & C & & \\ & \searrow g & & \nearrow h & \\ & & A & \xrightarrow{\text{id}_A} & A \\ & \nearrow \exists! & & \downarrow \lrcorner & \downarrow f \\ & & A & \xrightarrow{f} & B \end{array}$$

commute, which implies $g = h$. Therefore, f is a monomorphism.

Item 3, Monomorphisms vs. Injective Maps: Assume that f is injective. As the forgetful functor from C to Sets is faithful, we see that [Proposition 12.1.2.1.2](#) together with ?? imply that f is a monomorphism.

Conversely, assume that f is a monomorphism. As F preserves pullbacks, it also preserves kernel pairs. By ??, we see that F preserves monomorphisms. Thus F_f is a monomorphism, and hence is injective by ??.

Item 4, Stability Properties: Let $f, g: X \rightrightarrows A \times_C B$ be two morphisms such that the diagram

$$X \xrightleftharpoons[g]{f} A \times_C B \xrightarrow{m'} A$$

commutes. It follows that the diagram

$$\begin{array}{ccccc}
 & X & & & \\
 & \swarrow f & \searrow \text{pr}_2 \circ f & & \\
 & A \times_C B & \xrightarrow{\quad \text{pr}_2 \quad} & B & \\
 & \downarrow m' & \lrcorner & \downarrow m & \\
 A & \xrightarrow{\psi} & C & &
 \end{array}$$

$m' \circ g$ is the diagonal arrow from X to $A \times_C B$.

also commutes. From the universal property of the pullback, it follows that there must be precisely one morphism from X to $A \times_C B$ making the above diagram commute. Thus $f = g$ and m' is a monomorphism.

Item 5, Morphisms From the Terminal Object Are Monomorphisms: Clear. \square

12.1.2 Monomorphism-Reflecting Functors

011J Definition 12.1.2.1.1. A functor $F: C \rightarrow \mathcal{D}$ **reflects monomorphisms** if, for each morphism f of C , whenever F_f is a monomorphism, so is f .

011K Proposition 12.1.2.1.2. Let $F: C \rightarrow \mathcal{D}$ be a functor. If F is faithful, then it reflects monomorphisms.

Proof. Let $f: A \rightarrow B$ be a morphism of C and suppose that $F_f: F_A \rightarrow F_B$ is a monomorphism. Let $g, h: B \rightrightarrows C$ be two morphisms of C such that $g \circ f = h \circ f$. As F is faithful, we must have

$$F_g \circ F_f = F_{g \circ f} = F_{h \circ f} = F_h \circ F_f,$$

but as F_f is a monomorphism, it must be that $F_g = F_h$. Using the faithfulness of F again, we see that $g = h$. Therefore f is a monomorphism. \square

12.1.3 Split Monomorphisms

Let C be a category.

011M Definition 12.1.3.1.1. A morphism $f: A \rightarrow B$ of C is a **split monomorphism**² if there exists a morphism $g: B \rightarrow A$ of C such that³

$$g \circ f = \text{id}_A.$$

²Further Terminology: Also called a **section**, or a **split monic** morphism.

³ **Warning:** There exist monomorphisms which are not split monomorphisms, e.g. $\mathbb{Z}/2 \hookrightarrow \mathbb{Z}/4$ in Ring.

011N Proposition 12.1.3.1.2. Let C be a category.

011P 1. *Split Monomorphisms are Monomorphisms.* If m is a split monomorphism, then m is a monomorphism.

Proof. Item 1, Split Monomorphisms are Monomorphisms: Let $m: A \rightarrow B$ be a split monomorphism of C , let $e: B \rightarrow A$ be a morphism of C with

$$e \circ m = \text{id}_A,$$

and let $f, g: C \rightrightarrows A$ be two morphisms of C such that the diagram

$$C \xrightarrow{\begin{smallmatrix} f \\ g \end{smallmatrix}} A \xrightarrow{m} B$$

commutes. Then we have

$$\begin{aligned} f &= \text{id}_A \circ f \\ &= (e \circ m) \circ f \\ &= e \circ (m \circ f) \\ &= e \circ (m \circ g) \\ &= (e \circ m) \circ g \\ &= \text{id}_A \circ g \\ &= g, \end{aligned}$$

showing m to be a monomorphism. □

12.2 Epimorphisms

12.2.1 Foundations

Let C be a category.

011S Definition 12.2.1.1.1. A morphism $f: A \rightarrow B$ of C is an **epimorphism** if for every commutative⁴ diagram of the form

$$A \xrightarrow{f} B \xrightarrow{\begin{smallmatrix} g \\ h \end{smallmatrix}} C,$$

we have $g = h$.

011T Example 12.2.1.1.2. Let $f: A \rightarrow B$ be a function. The following conditions are equivalent:

1. The function f is injective.

⁴That is, with $g \circ f = h \circ f$.

2. The function f is an epimorphism in Sets.

Proof. Suppose that f is surjective and let $g, h: B \rightrightarrows C$ be morphisms such that $g \circ f = h \circ f$. Then for each $a \in A$, we have

$$g(f(a)) = h(f(a)),$$

but this implies that

$$g(b) = h(b)$$

for each $b \in B$, as f is surjective. Thus $g = h$ and f is an epimorphism.

To prove the converse, we proceed by contrapositive. So suppose that f is not surjective and consider the diagram

$$A \xrightarrow{f} B \xrightleftharpoons[h]{g} C,$$

where h is the map defined by $h(b) = 0$ for each $b \in B$ and g is the map defined by

$$g(b) = \begin{cases} 1 & \text{if } b \in \text{Im}(f), \\ 0 & \text{otherwise.} \end{cases}$$

Then $h \circ f = g \circ f$, as $h(f(a)) = 1 = g(f(a))$ for each $a \in A$. However, for any $b \in B \setminus \text{Im}(f)$, we have

$$g(b) = 0 \neq 1 = h(b).$$

Therefore $g \neq h$ and f is not an epimorphism. \square

011U Proposition 12.2.1.1.3. Let C be a category.

011V 1. *Characterisations.* Let C be a category with pullbacks and $f: A \rightarrow B$ be a morphism of C . The following conditions are equivalent:

011W (a) The morphism f is an epimorphism.

011X (b) For each $X \in \text{Obj}(C)$, the map of sets

$$f^*: \text{Hom}_{\text{Sets}}(B, X) \rightarrow \text{Hom}_{\text{Sets}}(A, X)$$

is injective.

011Y (c) The cokernel pair of f is trivial, i.e. we have

$$\begin{array}{ccc} B & \xleftarrow{\quad} & B \\ \uparrow \lrcorner & & \uparrow f \\ B \coprod_A B & \cong & B \\ \uparrow & & \uparrow f \\ B & \xleftarrow{f} & A. \end{array}$$

011Z 2. *Epimorphisms vs. Surjective Maps.* Let

- C be a concrete category;
- $\mathfrak{F} : C \rightarrow \text{Sets}$ be the forgetful functor from C to Sets;
- $f : A \rightarrow B$ be a morphism of C .

If \mathfrak{F} preserves pushouts, then the following conditions are equivalent:

- The morphism f is a epimorphism.
- The morphism f is surjective.

0120 3. *Stability Properties.* The class of all epimorphisms of C is stable under the following operations:

- Composition.* If f and g are epimorphisms, then so is $g \circ f$.⁵
- Pushouts.* Let

$$\begin{array}{ccc} A \coprod_C B & \xleftarrow{\quad} & B \\ \uparrow e' \quad \lrcorner & & \uparrow e \\ A & \longrightarrow & C \end{array}$$

be a diagram in C . If m is an epimorphism in C , then so is e' .

0121 4. *Morphisms to the Initial Object Are Monomorphisms.* If C has an initial object \emptyset_C , then every morphism of C to \emptyset_C is a epimorphism.

Proof. This is dual to [Proposition 12.1.1.3](#). □

12.2.2 Regular Epimorphisms

0123 **Proposition 12.2.2.1.1.** Let C be a category.

1. *Stability Under Pullbacks.* Consider the diagram

$$\begin{array}{ccc} A \times_C B & \longrightarrow & B \\ \downarrow e' \quad \lrcorner & & \downarrow e \\ A & \longrightarrow & C \end{array}$$

in C . If e is a regular epimorphism, then so is e' .

Proof. Epimorphisms Need Not Be Stable Under Pullback. : Regular Epimorphisms Are Stable Under Pullback. : □

⁵Conversely, if $g \circ f$ is a epimorphism, then so is g .

12.2.3 Effective Epimorphisms

Let C be a category.

0125 Definition 12.2.3.1.1. An epimorphism $f: A \rightarrow B$ of C is **effective** if we have an isomorphism

$$B \cong \text{CoEq}(A \times_B A \rightrightarrows A).$$

12.2.4 Split Epimorphisms

Let C be a category.

0127 Definition 12.2.4.1.1. A morphism $f: A \rightarrow B$ of C is a **retraction**⁶ if there is an arrow $g: B \rightarrow A$ such that $f \circ g = \text{id}_B$.

0128 Proposition 12.2.4.1.2. Let $f: A \rightarrow B$ be a morphism of C .

1. Every split epimorphism is an epimorphism.⁷

Proof. This is dual to ??.

□

Appendices

12.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets
5. Relations
6. Spans
7. Posets

Indexed and Fibred Sets

7. Indexed Sets
8. Fibred Sets

Un/Straightening for Indexed and Fibred Sets

Category Theory

11. Categories
12. Types of Morphisms in Categories
13. Adjunctions and the Yoneda Lemma
14. Constructions With Categories
15. Profunctors
16. Cartesian Closed Categories
17. Kan Extensions

Bicategories

18. Bicategories

⁶Further Terminology: Also called a **split epimorphism**.

⁷⚠️ Warning: There are epimorphisms which are not split epimorphisms, however, e.g. $\mathbb{Z} \hookrightarrow \mathbb{Z}/2$.

- 19. Internal Adjunctions
- 32. Hypergroups
- Internal Category Theory**
- 33. Hypersemirings and Hyperrings
- 20. Internal Categories
- 34. Quantales
- Cyclic Stuff**
- Near-Rings**
- 21. The Cycle Category
- 35. Near-Semirings
- Cubical Stuff**
- 36. Near-Rings
- 22. The Cube Category
- Real Analysis**
- Globular Stuff**
- 37. Real Analysis in One Variable
- 23. The Globe Category
- 38. Real Analysis in Several Variables
- Cellular Stuff**
- Measure Theory**
- 24. The Cell Category
- 39. Measurable Spaces
- Monoids**
- 40. Measures and Integration
- 25. Monoids
- Probability Theory**
- 26. Constructions With Monoids
- 40. Probability Theory
- Monoids With Zero**
- Stochastic Analysis**
- 27. Monoids With Zero
- 41. Stochastic Processes, Martingales,
and Brownian Motion
- 28. Constructions With Monoids With
Zero
- 42. Itô Calculus
- Groups**
- 43. Stochastic Differential Equations
- 29. Groups
- Differential Geometry**
- 30. Constructions With Groups
- 44. Topological and Smooth Manifolds
- Hyper Algebra**
- Schemes**
- 31. Hypermonoids
- 45. Schemes

12.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 13

Adjunctions and the Yoneda Lemma

0129

Contents

13.1 Other Chapters	371
---------------------------	-----

13.1 Other Chapters

Sets	12. Types of Morphisms in Categories
1. Sets	13. Adjunctions and the Yoneda Lemma
2. Constructions With Sets	14. Constructions With Categories
3. Pointed Sets	15. Profunctors
4. Tensor Products of Pointed Sets	16. Cartesian Closed Categories
5. Relations	17. Kan Extensions
6. Spans	Bicategories
7. Posets	18. Bicategories
Indexed and Fibred Sets	19. Internal Adjunctions
7. Indexed Sets	Internal Category Theory
8. Fibred Sets	20. Internal Categories
9. Un/Straightening for Indexed and Fibred Sets	Cyclic Stuff
Category Theory	21. The Cycle Category
11. Categories	Cubical Stuff
	22. The Cube Category
	Globular Stuff

23. The Globe Category	35. Near-Semirings
Cellular Stuff	36. Near-Rings
24. The Cell Category	Real Analysis
Monoids	37. Real Analysis in One Variable
25. Monoids	38. Real Analysis in Several Variables
26. Constructions With Monoids	Measure Theory
Monoids With Zero	39. Measurable Spaces
27. Monoids With Zero	40. Measures and Integration
28. Constructions With Monoids With Zero	Probability Theory
Groups	40. Probability Theory
29. Groups	Stochastic Analysis
30. Constructions With Groups	41. Stochastic Processes, Martingales, and Brownian Motion
Hyper Algebra	42. Itô Calculus
31. Hypermonoids	43. Stochastic Differential Equations
32. Hypergroups	Differential Geometry
33. Hypersemirings and Hyperrings	44. Topological and Smooth Manifolds
34. Quantales	Schemes
Near-Rings	45. Schemes

Chapter 14

Constructions With Categories

012A

Contents

14.A Other Chapters	373
14.2 Other Chapters	375

Appendices

14.A Other Chapters

Sets

- 1. Sets
- 2. Constructions With Sets
- 3. Pointed Sets
- 4. Tensor Products of Pointed Sets
- 5. Relations
- 6. Spans
- 7. Posets

Indexed and Fibred Sets

- 7. Indexed Sets
- 8. Fibred Sets
- 9. Un/Straightening for Indexed and Fibred Sets

Category Theory

- 11. Categories
- 12. Types of Morphisms in Categories
- 13. Adjunctions and the Yoneda Lemma
- 14. Constructions With Categories
- 15. Profunctors
- 16. Cartesian Closed Categories
- 17. Kan Extensions

Bicategories

- 18. Bicategories
- 19. Internal Adjunctions

Internal Category Theory

- 20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With
Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales,
and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

14.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
Category Theory	27. Monoids With Zero
11. Categories	28. Constructions With Monoids With Zero
12. Types of Morphisms in Categories	Groups
13. Adjunctions and the Yoneda Lemma	29. Groups
14. Constructions With Categories	30. Constructions With Groups
15. Profunctors	Hyper Algebra
16. Cartesian Closed Categories	31. Hypermonoids
17. Kan Extensions	32. Hypergroups
Bicategories	33. Hypersemirings and Hyperrings
18. Bicategories	34. Quantales
19. Internal Adjunctions	Near-Rings
Internal Category Theory	35. Near-Semirings
20. Internal Categories	36. Near-Rings
	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 15

Profunctors

012B

Contents

15.1	Profunctors.....	377
15.1.1	Foundations	377
15.2	Operations With Profunctors	378
15.2.1	The Domain and Range of a Profunctor.....	378
15.2.2	Composition of Profunctors	379
15.2.3	Representable Profunctors.....	379
15.2.4	Collages.....	380
15.3	Categories of Profunctors.....	384
15.3.1	The Bicategory of Profunctors.....	384
15.3.2	Properties of Prof.....	384
15.4	Other Chapters	386

15.1 Profunctors

15.1.1 Foundations

Let C and \mathcal{D} be categories.

012E **Definition 15.1.1.1.** A **profunctor**¹ $p: C \nrightarrow \mathcal{D}$ from C to \mathcal{D} is a functor $p: \mathcal{D}^{\text{op}} \times C \rightarrow \text{Sets}$.

012F **Remark 15.1.1.2.** Equivalently, we may define a profunctor from C to \mathcal{D} as:

012G 1. A functor $p: \mathcal{D}^{\text{op}} \times C \rightarrow \text{Sets}$;

012H 2. A functor $p: C \rightarrow \text{PSh}(\mathcal{D})$;

¹Further Terminology: Also called a **distributor**, a **bimodule**, a **correspondence**, or a **relator**.

- 012J** 3. A functor $\mathbf{p}: \mathcal{D}^{\text{op}} \rightarrow \text{Fun}(C, \text{Sets})$;
- 012K** 4. A cocontinuous functor $\mathbf{p}: \text{PSh}(C) \rightarrow \text{PSh}(\mathcal{D})$;

That is, we have isomorphisms of categories

$$\begin{aligned}\text{Prof}(C, \mathcal{D}) &\cong \text{Fun}(C, \text{PSh}(\mathcal{D})), \\ &\cong \text{Fun}(\mathcal{D}^{\text{op}}, \text{CoPSh}(C)), \\ &\cong \text{CoContFun}(\text{PSh}(C), \text{PSh}(\mathcal{D})),\end{aligned}$$

natural in $C, \mathcal{D} \in \text{Obj}(\text{Cats})$.

Proof. We claim that **Items 1 to 4** are indeed equivalent:

- The equivalence between **Items 1** and **2** is an instance of currying, following from the isomorphisms of categories

$$\text{Fun}(\mathcal{D}^{\text{op}} \times C, \text{Sets}) \cong \text{Fun}(C, \text{Fun}(\mathcal{D}^{\text{op}}, \text{Sets})) \stackrel{\text{def}}{=} \text{Fun}(C, \text{PSh}(\mathcal{D})).$$

- The equivalence between **Items 1** and **3** is also an instance of currying, following from the isomorphisms of categories

$$\text{Fun}(\mathcal{D}^{\text{op}} \times C, \text{Sets}) \cong \text{Fun}(\mathcal{D}^{\text{op}}, \text{Fun}(C, \text{Sets})) \stackrel{\text{def}}{=} \text{Fun}(\mathcal{D}^{\text{op}}, \text{Fun}(C, \text{Sets})).$$

- The equivalence between **Items 1** and **4** follows from the universal property of the category $\text{PSh}(C)$ of presheaves on C as the free cocompletion of C via the Yoneda embedding

$$\mathfrak{J}: C^{\text{op}} \hookrightarrow \text{PSh}(C)$$

of C into $\text{PSh}(C)$ (**??** of **??**).

This finishes the proof. □

15.2 Operations With Profunctors

15.2.1 The Domain and Range of a Profunctor

- 012N** **Definition 15.2.1.1.1.** Let $\mathbf{p}: C \dashv \mathcal{D}$ be a profunctor.²

1. The **domain** of \mathbf{p} is the presheaf $\text{dom}(\mathbf{p}): \mathcal{D}^{\text{op}} \rightarrow \text{Sets}$ on \mathcal{D} defined by

$$\text{dom}(\mathbf{p})^- \stackrel{\text{def}}{=} \underset{B \in \mathcal{D}}{\text{colim}} (\mathbf{p}_B^-).$$

²In other words, the domain and range of \mathbf{p} are the functors

$$\begin{aligned}\text{dom}(\mathbf{p}) &: \mathcal{D}^{\text{op}} \rightarrow \text{Sets}, \\ \text{range}(\mathbf{p}) &: C \rightarrow \text{Sets}\end{aligned}$$

2. The **range** of \mathbf{p} is the copresheaf $\text{range}(\mathbf{p}): C \rightarrow \text{Sets}$ on C defined by

$$\text{range}(\mathbf{p})_- \stackrel{\text{def}}{=} \underset{A \in \mathcal{D}}{\text{colim}} (\mathbf{p}_-^A).$$

15.2.2 Composition of Profunctors

Let C , \mathcal{D} , and \mathcal{E} be categories and let $\mathbf{p}: C \nrightarrow \mathcal{D}$ and $\mathbf{q}: \mathcal{D} \nrightarrow \mathcal{E}$ be profunctors.

012Q Definition 15.2.2.1.1. The **composition** of \mathbf{p} and \mathbf{q} is the profunctor $\mathbf{q} \diamond \mathbf{p}: C \nrightarrow \mathcal{E}$ defined by³

$$(\mathbf{q} \diamond \mathbf{p})_{-2}^{-1} \stackrel{\text{def}}{=} \int^{B \in \mathcal{D}} \mathbf{q}_B^{-1} \times \mathbf{p}_{-2}^B.$$

15.2.3 Representable Profunctors

012S Definition 15.2.3.1.1. The **representable profunctor associated to a functor** $F: C \rightarrow \mathcal{D}$ is the profunctor $\widehat{F}^*: C \nrightarrow \mathcal{D}$ defined as the adjunct of the composition

$$C \xrightarrow{F} \mathcal{D} \xrightarrow{\mathfrak{X}} \text{PSh}(\mathcal{D})$$

under the adjunction

$$\text{Fun}(\mathcal{D}^{\text{op}} \times C, \text{Sets}) \cong \text{Fun}(C, \text{PSh}(\mathcal{D}))$$

of ?? of ??.⁴

012T Definition 15.2.3.1.2. A profunctor is **representable** if it is isomorphic to a representable profunctor.

defined by

$$\begin{array}{ccc} \mathcal{D}^{\text{op}} & \xrightarrow{\mathbf{p}^\dagger} & \text{PSh}(\mathcal{D}) \\ \text{dom}(\mathbf{p}) \searrow & \downarrow \text{colim} & \\ & \text{Sets}, & \end{array} \quad \begin{array}{c} \text{dom}(\mathbf{p}) \stackrel{\text{def}}{=} \text{colim} \circ \mathbf{p}^\dagger, \\ \text{range}(\mathbf{p}) \stackrel{\text{def}}{=} \text{colim} \circ \mathbf{p}^\ddagger, \end{array} \quad \begin{array}{ccc} C & \xrightarrow{\mathbf{p}^\ddagger} & \text{Fun}(C, \text{Sets}) \\ \text{range}(\mathbf{p}) \searrow & \downarrow \text{colim} & \\ & \text{Sets}. & \end{array}$$

³Alternatively, we may define $\mathbf{q} \diamond \mathbf{p}$ (using the equivalent definition of Item 2 of Remark 15.1.1.2) by

$$\begin{array}{ccc} & \mathcal{D} & \xrightarrow{\mathbf{p}^\dagger} \text{PSh}(C). \\ & \text{Lan}_{\mathfrak{X}}(\mathbf{p}^\dagger) \circ \mathbf{q}^\dagger, & \downarrow \mathfrak{X} \\ & \mathcal{E} & \xrightarrow{\mathbf{q}^\dagger} \text{PSh}(\mathcal{D}) \end{array}$$

⁴That is, we have

$$\widehat{F}^* \stackrel{\text{def}}{=} \text{Hom}_{\mathcal{D}}(-_1, F_{-2}).$$

012U Definition 15.2.3.1.3. The **corepresentable⁵ profunctor associated to a functor** $F: C \rightarrow \mathcal{D}$ is the profunctor $\widehat{F}_*: \mathcal{D} \nrightarrow C$ defined as the adjunct of the composition

$$C^{\text{op}} \xrightarrow{F^{\text{op}}} \mathcal{D}^{\text{op}} \xrightarrow{\text{?}} \text{CoPSh}(\mathcal{D})$$

under the adjunction

$$\text{Fun}(C^{\text{op}} \times \mathcal{D}, \text{Sets}) \cong \text{Fun}(C^{\text{op}}, \text{CoPSh}(\mathcal{D}))$$

of ?? of ??⁶

012V Definition 15.2.3.1.4. A profunctor is **corepresentable** if it is isomorphic to a corepresentable profunctor.

15.2.4 Collages

Let C and \mathcal{D} be categories.

012X Definition 15.2.4.1.1. The **collage** of a profunctor $\mathbf{p}: C \nrightarrow \mathcal{D}$ is the category $\text{Coll}(\mathbf{p})$ ⁷ where⁸

- *Objects.* We have

$$\text{Obj}(\text{Coll}(\mathbf{p})) \stackrel{\text{def}}{=} \text{Obj}(C) \sqcup \text{Obj}(\mathcal{D});$$

⁵Some authors call both \widehat{F}^* and \widehat{F}_* the **representable profunctors associated to F** .

⁶That is:

$$\widehat{F}_* \stackrel{\text{def}}{=} \text{Hom}_{\mathcal{D}}(F_{-1}, -_2).$$

⁷Further Notation: Also written $C \star^{\mathbf{p}} \mathcal{D}$, notably in [HigherToposTheory].

⁸We also have a functor $\phi: \text{Coll}(\mathbf{p}) \rightarrow \mathbb{M}$ where

- *Actions on Objects.* For each $A \in \text{Obj}(\text{Coll}(\mathbf{p}))$, we have

$$\phi_A \stackrel{\text{def}}{=} \begin{cases} [0] & \text{if } A \in \text{Obj}(C), \\ [1] & \text{if } A \in \text{Obj}(\mathcal{D}). \end{cases}$$

- *Actions on Morphisms.* For each $A, B \in \text{Obj}(\text{Coll}(\mathbf{p}))$, the action on morphisms

$$\phi_{A,B}: \text{Hom}_{\text{Coll}(\mathbf{p})}(A, B) \rightarrow \text{Hom}_{\text{Coll}(\mathbf{p})}(\phi_A, \phi_B)$$

of ϕ at (A, B) is given by

$$\phi_{A,B}(f) \stackrel{\text{def}}{=} \begin{cases} \text{id}_{[0]} & \text{if } A, B \in \text{Obj}(C), \\ \text{id}_{[1]} & \text{if } A, B \in \text{Obj}(\mathcal{D}), \\ [0] \rightarrow [1] & \text{if } A \in \text{Obj}(C) \text{ and } B \in \text{Obj}(\mathcal{D}). \end{cases}$$

If $A \in \text{Obj}(\mathcal{D})$ and $B \in \text{Obj}(C)$, we have $\phi_{A,B} \stackrel{\text{def}}{=} \text{id}_{\emptyset}$.

- *Morphisms.* For each $A, B \in \text{Obj}(\text{Coll}(\mathbf{p}))$, we have

$$\text{Hom}_{\text{Coll}(\mathbf{p})}(A, B) \stackrel{\text{def}}{=} \begin{cases} \text{Hom}_C(A, B) & \text{if } A, B \in \text{Obj}(C), \\ \text{Hom}_{\mathcal{D}}(A, B) & \text{if } A, B \in \text{Obj}(\mathcal{D}), \\ \mathbf{p}(A, B) & \text{if } A \in \text{Obj}(C) \text{ and } B \in \text{Obj}(\mathcal{D}), \\ \emptyset & \text{if } A \in \text{Obj}(\mathcal{D}) \text{ and } B \in \text{Obj}(C); \end{cases}$$

- *Identities.* For each $A \in \text{Obj}(\text{Coll}(\mathbf{p}))$, the unit map

$$\mathbb{1}_A^{\text{Coll}(\mathbf{p})} : \text{pt} \rightarrow \text{Hom}_{\text{Coll}(\mathbf{p})}(A, A)$$

of $\text{Coll}(\mathbf{p})$ at A is defined by

$$\text{id}_A \stackrel{\text{def}}{=} \begin{cases} \text{id}_A^C & \text{if } A \in \text{Obj}(C), \\ \text{id}_A^{\mathcal{D}} & \text{if } A \in \text{Obj}(\mathcal{D}); \end{cases}$$

- *Composition.* For each $A, B, C \in \text{Obj}(\text{Coll}(\mathbf{p}))$, the composition map

$$\circ_{A,B,C}^{\text{Coll}(\mathbf{p})} : \text{Hom}_{\text{Coll}(\mathbf{p})}(B, C) \times \text{Hom}_{\text{Coll}(\mathbf{p})}(A, B) \rightarrow \text{Hom}_{\text{Coll}(\mathbf{p})}(A, C)$$

of $\text{Coll}(\mathbf{p})$ at (A, B, C) is defined by⁹

$$\circ_{A,B,C}^{\text{Coll}(\mathbf{p})} \stackrel{\text{def}}{=} \begin{cases} \circ_{A,B,C}^C & \text{if } A, B, C \in \text{Obj}(C), \\ \mathbf{p}_C^{A,B} & \text{if } A, B \in \text{Obj}(C) \text{ and } C \in \text{Obj}(\mathcal{D}), \\ \iota & \text{if } A, C \in \text{Obj}(C) \text{ and } B \in \text{Obj}(\mathcal{D}), \\ \iota & \text{if } B, C \in \text{Obj}(C) \text{ and } A \in \text{Obj}(\mathcal{D}), \\ \mathbf{p}_{B,C}^A & \text{if } A \in \text{Obj}(C) \text{ and } B, C \in \text{Obj}(\mathcal{D}), \\ \iota & \text{if } B \in \text{Obj}(C) \text{ and } A, C \in \text{Obj}(\mathcal{D}), \\ \iota & \text{if } C \in \text{Obj}(C) \text{ and } A, B \in \text{Obj}(\mathcal{D}), \\ \circ_{A,B,C}^{\mathcal{D}} & \text{if } A, B, C \in \text{Obj}(\mathcal{D}). \end{cases}$$

012Y Example 15.2.4.1.2. If \mathbf{p} is the constant functor $\Delta_{\text{pt}} : \mathcal{D}^{\text{op}} \times \mathcal{C} \rightarrow \text{Sets}$ with value pt , then $\text{Coll}(\mathbf{p})$ is the join $\mathcal{C} \star \mathcal{D}$ of \mathcal{C} and \mathcal{D} of ??.

012Z Proposition 15.2.4.1.3. Let $\mathbf{p} : \mathcal{C} \rightarrow \mathcal{D}$ be a profunctor.

⁹Here the maps $\mathbf{p}_C^{A,B}$ and $\mathbf{p}_{B,C}^A$ are the maps

$$\begin{aligned} \mathbf{p}_C^{A,B} : \mathbf{p}_C^B \times \text{Hom}_C(A, B) &\rightarrow \mathbf{p}_C^A, \\ \mathbf{p}_{B,C}^A : \text{Hom}_{\mathcal{D}}(B, C) \times \mathbf{p}_B^A &\rightarrow \mathbf{p}_C^A \end{aligned}$$

coming from the profunctor structure of \mathbf{p} and the ι 's are inclusions of the empty set into the appropriate Hom sets.

0130 1. *Functionality.* The assignment $\mathbf{p} \mapsto \text{Coll}(\mathbf{p})$ defines a functor¹⁰

$$\text{Coll}_{\mathcal{C}, \mathcal{D}} : \text{Prof}(\mathcal{C}, \mathcal{D}) \rightarrow \text{Cats}_{/\mathbb{K}}(\mathcal{C}, \mathcal{D}),$$

where

- *Action on Objects.* For each $\mathbf{p} \in \text{Obj}(\text{Prof}(\mathcal{C}, \mathcal{D}))$, we have

$$[\text{Coll}](\mathbf{p}) \stackrel{\text{def}}{=} \text{Coll}(\mathbf{p});$$

- *Action on Morphisms.* For each $\mathbf{p}, \mathbf{q} \in \text{Obj}(\text{Prof}(\mathcal{C}, \mathcal{D}))$, the action on Hom-sets

$$\text{Coll}_{\mathbf{p}, \mathbf{q}} : \text{Nat}(\mathbf{p}, \mathbf{q}) \rightarrow \text{Fun}_{/\mathbb{K}}(\text{Coll}(\mathbf{p}), \text{Coll}(\mathbf{q}))$$

of Coll at (\mathbf{p}, \mathbf{q}) is the function sending a natural transformation $\alpha : \mathbf{p} \Rightarrow \mathbf{q}$ to the functor

$$\text{Coll}(\alpha) : \text{Coll}(\mathbf{p}) \rightarrow \text{Coll}(\mathbf{q})$$

over \mathbb{K} where

- *Action on Objects.* For each $X \in \text{Obj}(\text{Coll}(\mathbf{p}))$, we have

$$[\text{Coll}(\alpha)](X) \stackrel{\text{def}}{=} X;$$

- *Action on Morphisms.* For each $X, Y \in \text{Obj}(\text{Coll}(\mathbf{p}))$, the action on Hom-sets


$$\text{Coll}(\alpha)_{X, Y} : \text{Hom}_{\text{Coll}(\mathbf{p})}(X, Y) \rightarrow \underbrace{\text{Hom}_{\text{Coll}(\mathbf{q})}([\text{Coll}(\alpha)](X), [\text{Coll}(\alpha)](Y))}_{\stackrel{\text{def}}{=} \text{Hom}_{\text{Coll}(\mathbf{q})}(X, Y)}$$

of $\text{Coll}(\alpha)$ at (X, Y) is defined as follows:

¹⁰Here $\text{Cats}_{/\mathbb{K}}(\mathcal{C}, \mathcal{D})$ is the category defined as the pullback

$$\text{Cats}_{/\mathbb{K}}(\mathcal{C}, \mathcal{D}) \stackrel{\text{def}}{=} \text{pt}_{[C], \text{Cats}, \text{fib}_0} \times_{\text{fib}_1, \text{Cats}, [\mathcal{D}]} \text{pt},$$

as in the diagram



* If $X, Y \in \text{Obj}(\mathcal{C})$ or $X, Y \in \text{Obj}(\mathcal{D})$, then we have

$$\text{Coll}(\alpha)_{X,Y}(f) \stackrel{\text{def}}{=} f$$

for each $f \in \text{Hom}_{\text{Coll}(\mathfrak{p})}(X, Y)$.

* If $X \in \text{Obj}(\mathcal{C})$ and $Y \in \text{Obj}(\mathcal{D})$, then

$$\text{Coll}(\alpha)_{X,Y}: \underbrace{\text{Hom}_{\text{Coll}(\mathfrak{p})}(X, Y)}_{\stackrel{\text{def}}{=} \mathfrak{p}_Y^X} \rightarrow \underbrace{\text{Hom}_{\text{Coll}(\mathfrak{q})}(X, Y)}_{\stackrel{\text{def}}{=} \mathfrak{q}_Y^X}$$

is defined by

$$\text{Coll}(\alpha)_{X,Y}(f) \stackrel{\text{def}}{=} \alpha_Y^X;$$

* If $Y \in \text{Obj}(\mathcal{C})$ and $X \in \text{Obj}(\mathcal{D})$, then we have

$$\text{Coll}(\alpha)_{X,Y}(f) \stackrel{\text{def}}{=} \text{id}_\emptyset.$$

0131 2. *Collages as Lax Colimits.* We have an isomorphism of categories

$$\text{Coll}(\mathfrak{p}) \cong \text{colim}^{\text{lax}}(\mathfrak{p}),$$

functorial in \mathfrak{p} , where the above lax colimit is taken in the bicategory Prof .

0132 3. *Profunctors vs.Collages.* We have an equivalence of categories

$$(\text{Coll} \dashv \Gamma): \quad \text{Prof}(\mathcal{C}, \mathcal{D}) \begin{array}{c} \xrightarrow{\text{Coll}} \\[-1ex] \xleftarrow[\Gamma]{\perp_{\text{eq}}} \end{array} \text{Cats}_{/\mathbb{M}},$$

where $\Gamma: \text{Cats}_{/\mathbb{M}} \rightarrow \text{Prof}(\mathcal{C}, \mathcal{D})$ is the functor sending a functor $\mathcal{E} \rightarrow \mathbb{M}$ to the profunctor

$$\Gamma(\mathfrak{p}): \mathcal{C} \dashrightarrow \mathcal{D}$$

given on objects by

$$\Gamma(\mathfrak{p})_B^A \stackrel{\text{def}}{=} \text{Hom}_{\mathcal{E}}(A, B)$$

for each $A, B \in \text{Obj}(\mathcal{E})$.

Proof. **Item 1, Functoriality:** Omitted.

Item 2, Collages as Lax Colimits: See [[collages-as-lax-colimits](#)].

Item 3, Profunctors vs.Collages: See [[joyal:distributors-and-barrels](#)].

□

15.3 Categories of Profunctors

15.3.1 The Bicategory of Profunctors

0135 Definition 15.3.1.1.1. The **bicategory of profunctors** is the bicategory Prof where¹¹

1. *Objects.* The objects of Prof are categories;
2. *1-Morphisms.* The 1-morphisms of Prof are profunctors;
3. *2-Morphisms.* The 2-morphisms of Prof are natural transformations between profunctors;
4. *Identities.* For each $C \in \text{Obj}(\text{Prof})$, we have

$$\text{id}_C^{\text{Prof}} \stackrel{\text{def}}{=} \text{Hom}_C(-, -);$$

5. *Composition.* For each $C, D, E \in \text{Obj}(\text{Prof})$, the composition bifunctor

$$\diamond: \text{Prof}(D, E) \times \text{Prof}(C, D) \rightarrow \text{Prof}(C, E)$$

is defined on objects by sending profunctors $p: C \nrightarrow D$ and $q: D \nrightarrow E$ to the profunctor $q \diamond p$ of [Definition 15.2.2.1.1](#).

Proof. See [Definition 15.3.1.1.1](#). □

15.3.2 Properties of Prof

0137 Proposition 15.3.2.1.1. Let C and D be categories.

0138 1. Self-Duality. The bicategory Prof is self-dual: we have a biequivalence of bicategories

$$(-)^{\text{op}}: \text{Prof} \xrightarrow{\cong} \text{Prof}^{\text{op}}$$

where

- *Action on Objects.* The functor $(-)^{\text{op}}$ sends categories to their opposites;
- *Action on 1-Morphisms.* The functor $(-)^{\text{op}}$ sends profunctors to itself under the identification

$$\begin{aligned} \text{Prof}(C, D) &\stackrel{\text{def}}{=} \text{Fun}(D^{\text{op}} \times C, \text{Sets}), \\ &\cong \text{Fun}(C \times D^{\text{op}}, \text{Sets}), \\ &\stackrel{\text{def}}{=} \text{Prof}(D^{\text{op}}, C^{\text{op}}); \end{aligned}$$

¹¹The bicategory Prof admits a nice strictification to a 2-category: it is biequivalent to the sub-bicategory of Cats spanned by the presheaf categories, cocontinuous functors between them, and natural transformation between these.

- *Action on 2-Morphisms.* The functor $(-)^{\text{op}}$ sends natural transformations between profunctors to themselves.

0139 2. *Relation to Cats.* The co/representable profunctor constructions of Definitions 15.2.3.1.1 and 15.2.3.1.3 define embeddings of bicategories

$$\begin{aligned}\text{Cats}^{\text{op}} &\hookrightarrow \text{Prof}, \\ \text{Cats}^{\text{co}} &\hookrightarrow \text{Prof}.\end{aligned}$$

013A 3. *Equivalences in Prof and Cauchy Completions.* Every category is equivalent to its Cauchy completion in Prof.

013B 4. *Equivalences in Prof.* The following conditions are equivalent:

- (a) The categories C and \mathcal{D} are equivalent in Prof.
- (b) The categories $\text{PSh}(C)$ and $\text{PSh}(\mathcal{D})$ are equivalent in Cats_2 .
- (c) The Cauchy completions of C and \mathcal{D} are equivalent in Cats_2 .

013C 5. *Adjunctions in Prof.* Let C and \mathcal{D} be categories. The following data are equivalent:

- (a) An adjunction in Prof from C to \mathcal{D} .
- (b) A functor from C to the Cauchy completion $\overline{\mathcal{D}}$ of \mathcal{D} .
- (c) A **semifunctor** from C to \mathcal{D} .

013D 6. *As a Kleisli Bicategory.* We have a biequivalence of bicategories

$$\text{Prof} \cong \text{FreePsAlg}_{\text{PSh}},$$

where PSh is the presheaf category relative pseudomonad of [relative-pseudomonads-kleisli-bicategories-and-adjunctions]

013E 7. *Closedness.* The bicategory Prof is a closed bicategory, where given a profunctor $p: C \nrightarrow \mathcal{D}$ and a category X :

- *Right Kan Extensions.* The right adjoint

$$\text{Ran}_p: \text{Rel}(C, X) \rightarrow \text{Rel}(\mathcal{D}, X)$$

to the precomposition functor $p^*: \text{Rel}(\mathcal{D}, X) \rightarrow \text{Rel}(C, X)$ is given by

$$\text{Ran}_p(q) \stackrel{\text{def}}{=} \int_{A \in C} \text{Sets}(p_A^{-2}, q_A^{-1})$$

for each $q \in \text{Rel}(C, X)$.

- *Right Kan Lifts.* The right adjoint to the postcomposition functor

$$\text{Rift}_p: \text{Rel}(X, \mathcal{D}) \rightarrow \text{Rel}(X, C)$$

to the postcomposition functor $p_*: \text{Rel}(X, C) \rightarrow \text{Rel}(X, \mathcal{D})$ is given by

$$\text{Rift}_p(q) \stackrel{\text{def}}{=} \int_{B \in \mathcal{D}} \text{Sets}(p_{-1}^B, q_{-2}^B)$$

for each $q \in \text{Rel}(X, \mathcal{D})$.

- 013F 8. *Un/Straightening for Profunctors: Two-Sided Discrete Fibrations.* We have an equivalence of categories

$$\text{Prof}(C, \mathcal{D}) \cong \text{DFib}(C, \mathcal{D}).$$

Proof. [Item 1, Self-Duality](#): See [[lorelian2020coend](#)].

[Item 2, Relation to Cats](#): See [[lorelian2020coend](#)].

[Item 3, Equivalences in Prof and Cauchy Completions](#): See [[borceux-2](#)].

[Item 4, Equivalences in Prof](#): See [[borceux-2](#)].

[Item 5, Adjunctions in Prof](#): Omitted.

[Item 6, As a Kleisli Bicategory](#): See [[relative-pseudomonads-kleisli-bicategories-and-substitution-monoidal-structure](#)]

[Item 7, Closedness](#): Omitted.

[Item 8, Un/Straightening for Profunctors: Two-Sided Discrete Fibrations](#): See [[riehl:two-sided-discrete-fibrations](#)]

□

15.4 Other Chapters

Sets	14. Constructions With Categories
1. Sets	15. Profunctors
2. Constructions With Sets	16. Cartesian Closed Categories
3. Pointed Sets	17. Kan Extensions
4. Tensor Products of Pointed Sets	Bicategories
5. Relations	18. Bicategories
6. Spans	19. Internal Adjunctions
7. Posets	Internal Category Theory
Indexed and Fibred Sets	20. Internal Categories
7. Indexed Sets	Cyclic Stuff
8. Fibred Sets	21. The Cycle Category
9. Un/Straightening for Indexed and Fibred Sets	Cubical Stuff
Category Theory	22. The Cube Category
11. Categories	Globular Stuff
12. Types of Morphisms in Categories	23. The Globe Category
13. Adjunctions and the Yoneda Lemma	Cellular Stuff
	24. The Cell Category
	Monoids

25. Monoids	Real Analysis
26. Constructions With Monoids	37. Real Analysis in One Variable
Monoids With Zero	38. Real Analysis in Several Variables
27. Monoids With Zero	Measure Theory
28. Constructions With Monoids With Zero	39. Measurable Spaces
Groups	40. Measures and Integration
29. Groups	Probability Theory
30. Constructions With Groups	40. Probability Theory
Hyper Algebra	Stochastic Analysis
31. Hypermonoids	41. Stochastic Processes, Martingales, and Brownian Motion
32. Hypergroups	42. Itô Calculus
33. Hypersemirings and Hyperrings	43. Stochastic Differential Equations
34. Quantales	Differential Geometry
Near-Rings	44. Topological and Smooth Manifolds
35. Near-Semirings	Schemes
36. Near-Rings	45. Schemes

Chapter 16

Cartesian Closed Categories

013G Create tags (see [MSE 350788] for some of these):

1. define bicategory $\text{Adj}(C)$
2. [https://www.google.com/search?q=category+of+categories+is+no
t+locally+cartesian+closed](https://www.google.com/search?q=category+of+categories+is+not+locally+cartesian+closed)
3. [https://math.stackexchange.com/questions/2864916/are-there-i
mportant-locally-cartesian-closed-categories-that-actually-a
re-not-ca](https://math.stackexchange.com/questions/2864916/are-there-important-locally-cartesian-closed-categories-that-actually-are-not-ca)
4. Cats is not locally Cartesian closed: f^* does have a left adjoint (the proof for fibred sets seems to apply for any category with pullbacks), but does not have a right adjoint, as proved in example 1.7 of [https://sinhp.github.io/files/
CT/notes_on_lcccs.pdf](https://sinhp.github.io/files/CT/notes_on_lcccs.pdf)
5. internal **Hom** in categories of co/Cartesian fibrations
6. [https://mathoverflow.net/questions/460146/universal-propert
y-of-isbell-duality](https://mathoverflow.net/questions/460146/universal-property-of-isbell-duality)
7. <http://www.tac.mta.ca/tac/volumes/36/12/36-12abs.html>
8. Cartesian closed categories and locally Cartesian closed categories
 - (a) [https://ncatlab.org/nlab/show/locally+cartesian+closed+f
unctor](https://ncatlab.org/nlab/show/locally+cartesian+closed+functor)
 - (b) <https://ncatlab.org/nlab/show/cartesian+closed+functor>
 - (c) [https://ncatlab.org/nlab/show/locally+cartesian+closed+c
ategory](https://ncatlab.org/nlab/show/locally+cartesian+closed+category)
 - (d) <https://ncatlab.org/nlab/show/Frobenius+reciprocity>
9. [https://math.stackexchange.com/questions/3657046/the-inver](https://math.stackexchange.com/questions/3657046/the-inverse)

- [se-of-a-natural-isomorphism-is-a-natural-isomorphism](#) to justify adjunctions via homs
10. <https://ncatlab.org/nlab/show/enrichment+versus+internalisation>
 11. <https://mathoverflow.net/questions/382239/proof-that-a-cartesian-category-is-monoidal>
 12. <https://arxiv.org/pdf/2004.08964.pdf>

Create tags:

1. <https://www.google.com/search?q=category+of+categories+is+not+locally+cartesian+closed>
2. <https://math.stackexchange.com/questions/2864916/are-there-important-locally-cartesian-closed-categories-that-actually-are-not-ca>
3. Cats is not locally Cartesian closed: f^* does have a left adjoint (the proof for fibred sets seems to apply for any category with pullbacks), but does not have a right adjoint, as proved in example 1.7 of https://sinhp.github.io/files/CT/notes_on_lcccs.pdf
4. Cartesian closed categories and locally Cartesian closed categories
 - (a) <https://ncatlab.org/nlab/show/locally+cartesian+closed+functor>
 - (b) <https://ncatlab.org/nlab/show/cartesian+closed+functor>
 - (c) <https://ncatlab.org/nlab/show/locally+cartesian+closed+category>
 - (d) <https://ncatlab.org/nlab/show/Frobenius+reciprocity>
5. <https://mathoverflow.net/questions/382239/proof-that-a-cartesian-category-is-monoidal>

Contents

16.1	Cartesian Closed Categories.....	389
16.A	Other Chapters	390
16.2	Other Chapters	391

16.1 Cartesian Closed Categories

Appendices

16.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets
5. Relations
6. Spans
7. Posets

Indexed and Fibred Sets

7. Indexed Sets
8. Fibred Sets
9. Un/Straightening for Indexed and Fibred Sets

Category Theory

11. Categories
12. Types of Morphisms in Categories
13. Adjunctions and the Yoneda Lemma
14. Constructions With Categories
15. Profunctors
16. Cartesian Closed Categories
17. Kan Extensions

Bicategories

18. Bicategories
19. Internal Adjunctions

Internal Category Theory

20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

41. Stochastic Processes, Martingales, and Brownian Motion

Measure Theory

39. Measurable Spaces

42. Itô Calculus

40. Measures and Integration

43. Stochastic Differential Equations

Probability Theory

40. Probability Theory

Differential Geometry

44. Topological and Smooth Manifolds

Stochastic Analysis

Schemes

45. Schemes

16.2 Other Chapters

Sets

15. Profunctors

1. Sets

16. Cartesian Closed Categories

2. Constructions With Sets

17. Kan Extensions

3. Pointed Sets

Bicategories

4. Tensor Products of Pointed Sets

18. Bicategories

5. Relations

19. Internal Adjunctions

6. Spans

Internal Category Theory

7. Posets

20. Internal Categories

Indexed and Fibred Sets

Cyclic Stuff

7. Indexed Sets

21. The Cycle Category

8. Fibred Sets

Cubical Stuff

9. Un/Straightening for Indexed and Fibred Sets

22. The Cube Category

Category Theory

Globular Stuff

11. Categories

23. The Globe Category

12. Types of Morphisms in Categories

Cellular Stuff

13. Adjunctions and the Yoneda Lemma

24. The Cell Category

14. Constructions With Categories

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero	37. Real Analysis in One Variable
27. Monoids With Zero	38. Real Analysis in Several Variables
28. Constructions With Monoids With Zero	Measure Theory
Groups	39. Measurable Spaces
29. Groups	40. Measures and Integration
30. Constructions With Groups	Probability Theory
Hyper Algebra	40. Probability Theory
31. Hypermonoids	Stochastic Analysis
32. Hypergroups	41. Stochastic Processes, Martingales, and Brownian Motion
33. Hypersemirings and Hyperrings	42. Itô Calculus
34. Quantales	43. Stochastic Differential Equations
Near-Rings	Differential Geometry
35. Near-Semirings	44. Topological and Smooth Manifolds
36. Near-Rings	Schemes
Real Analysis	45. Schemes

Chapter 17

Kan Extensions

013J

Contents

17.A Other Chapters	393
17.2 Other Chapters	395

Appendices

17.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff[21. The Cycle Category](#)**Cubical Stuff**[22. The Cube Category](#)**Globular Stuff**[23. The Globe Category](#)**Cellular Stuff**[24. The Cell Category](#)**Monoids**[25. Monoids](#)[26. Constructions With Monoids](#)**Monoids With Zero**[27. Monoids With Zero](#)[28. Constructions With Monoids With Zero](#)**Groups**[29. Groups](#)[30. Constructions With Groups](#)**Hyper Algebra**[31. Hypermonoids](#)[32. Hypergroups](#)[33. Hypersemirings and Hyperrings](#)[34. Quantales](#)**Near-Rings**[35. Near-Semirings](#)[36. Near-Rings](#)**Real Analysis**[37. Real Analysis in One Variable](#)[38. Real Analysis in Several Variables](#)**Measure Theory**[39. Measurable Spaces](#)[40. Measures and Integration](#)**Probability Theory**[40. Probability Theory](#)**Stochastic Analysis**[41. Stochastic Processes, Martingales, and Brownian Motion](#)[42. Itô Calculus](#)[43. Stochastic Differential Equations](#)**Differential Geometry**[44. Topological and Smooth Manifolds](#)**Schemes**[45. Schemes](#)

17.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part IV

Bicategories

Chapter 18

Bicategories

013K

Contents

18.1	Monomorphisms in Bicategories	398
18.1.1	Faithful Monomorphisms.....	398
18.1.2	Full Monomorphisms	399
18.1.3	Fully Faithful Monomorphisms	400
18.1.4	Strict Monomorphisms.....	400
18.2	Epimorphisms in Bicategories	401
18.2.1	Faithful Epimorphisms.....	401
18.2.2	Full Epimorphisms	401
18.2.3	Fully Faithful Epimorphisms	402
18.2.4	Strict Epimorphisms.....	403
18.3	bicategories of spans	403
18.A	Other Chapters	405
18.2	Other Chapters	406

Create tags and TODO:

1. spans in bicategories: add Proposition 7 here: <https://arxiv.org/abs/1903.03890>
2. add fact: internal adjunctions in $\text{PseudoFun}(C, \mathcal{D})$ are precisely the invertible strong transformations as in [JY21, Example 6.2.7]. What are the internal adjunctions?

18.1 Monomorphisms in Bicategories

18.1.1 Faithful Monomorphisms

Let C be a bicategory.

013N Definition 18.1.1.1. A 1-morphism $f: A \rightarrow B$ is a **faithful monomorphism in C** if the following equivalent conditions are satisfied:

013P 1. For each $X \in \text{Obj}(C)$, the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by f is faithful.

013Q 2. Given a diagram in C of the form

$$\begin{array}{ccc} X & \xrightarrow{\phi} & A \xrightarrow{f} B \\ \alpha \Downarrow \beta & \Downarrow & \\ & \psi & \end{array}$$

if we have $\text{id}_f \star \alpha = \text{id}_f \star \beta$, then $\alpha = \beta$.

013R Example 18.1.1.2. Here are some examples of faithful monomorphisms.

013S 1. *Full Monomorphisms in Cats_2 .*

013T 2. *Full Monomorphisms in Rel .*

013U 3. *Full Monomorphisms in Span .*

18.1.2 Full Monomorphisms

Let C be a bicategory.

013W Definition 18.1.2.1.1. A 1-morphism $f: A \rightarrow B$ is a **full monomorphism in C** if the following equivalent conditions are satisfied:

013X 1. For each $X \in \text{Obj}(C)$, the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by f is full.

014F 2. For each $X \in \text{Obj}(C)$ and each 2-morphism

$$\gamma: f \circ \phi \Rightarrow f \circ \psi, \quad X \xrightarrow{\begin{array}{c} f \circ \phi \\ \gamma \\ f \circ \psi \end{array}} B$$

of C , there exists a 2-morphism $\alpha: \phi \Rightarrow \psi$ of C such that we have an equality

$$X \xrightarrow{\begin{array}{c} \phi \\ \alpha \\ \psi \end{array}} A \xrightarrow{f} B = X \xrightarrow{\begin{array}{c} f \circ \phi \\ \gamma \\ f \circ \psi \end{array}} B$$

of pasting diagrams in C , i.e. such that we have

$$\gamma = \text{id}_f \star \alpha.$$

013Z Example 18.1.2.1.2. Here are some examples of full monomorphisms.

- 0140** 1. *Full Monomorphisms in Cats₂.*
- 0141** 2. *Full Monomorphisms in Rel.*
- 0142** 3. *Full Monomorphisms in Span.*

18.1.3 Fully Faithful Monomorphisms

Let C be a bicategory.

0144 Definition 18.1.3.1.1. A 1-morphism $f: A \rightarrow B$ is a **fully faithful monomorphism** in C if the following equivalent conditions are satisfied:

- 0145** 1. The 1-morphism f is fully and faithful.
- 0146** 2. For each $X \in \text{Obj}(C)$, the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by f is fully faithful.

- 0147** 3. The conditions in Item 1 of Definition 18.1.1.1 and Item 1 of Definition 18.1.2.1.1 hold.

0148 Example 18.1.3.1.2. Here are some examples of fully faithful monomorphisms.

- 0149** 1. *Fully Faithful Monomorphisms in Cats₂.*
- 014A** 2. *Fully Faithful Monomorphisms in Rel.*
- 014B** 3. *Fully Faithful Monomorphisms in Span.*

18.1.4 Strict Monomorphisms

Let C be a bicategory.

014D Definition 18.1.4.1.1. A 1-morphism $f: A \rightarrow B$ is a **strict monomorphism** in C if the following equivalent conditions are satisfied:

- 014E** 1. For each $X \in \text{Obj}(C)$, the action on objects

$$f_*: \text{Obj}(\text{Hom}_C(X, A)) \rightarrow \text{Obj}(\text{Hom}_C(X, B))$$

of the functor

$$f_*: \text{Hom}_C(X, A) \rightarrow \text{Hom}_C(X, B)$$

given by postcomposition by f is injective.

014F 2. For each diagram in C of the form

$$X \xrightarrow[\psi]{\phi} A \xrightarrow{f} B,$$

if $f \circ \phi = f \circ \psi$, then $\phi = \psi$.

014G **Example 18.1.4.1.2.** Here are some examples of strict monomorphisms.

- 014H 1. *Strict Monomorphisms in Cats₂*.
- 014J 2. *Strict Monomorphisms in Rel*.
- 014K 3. *Strict Monomorphisms in Span*.

18.2 Epimorphisms in Bicategories

18.2.1 Faithful Epimorphisms

Let C be a bicategory.

014N **Definition 18.2.1.1.1.** A 1-morphism $f: A \rightarrow B$ is a **faithful epimorphism in C** if the following equivalent conditions are satisfied:

014P 1. For each $X \in \text{Obj}(C)$, the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by f is faithful.

014Q 2. Given a diagram in C of the form

$$A \xrightarrow{f} B \begin{array}{c} \nearrow \phi \\ \alpha \Downarrow \beta \\ \psi \end{array} X,$$

if we have $\alpha \star \text{id}_f = \beta \star \text{id}_f$, then $\alpha = \beta$.

014R **Example 18.2.1.1.2.** Here are some examples of faithful epimorphisms.

- 014S 1. *Full Epimorphisms in Cats₂*.
- 014T 2. *Full Epimorphisms in Rel*.
- 014U 3. *Full Epimorphisms in Span*.

18.2.2 Full Epimorphisms

Let C be a bicategory.

014W Definition 18.2.2.1.1. A 1-morphism $f: A \rightarrow B$ is a **full epimorphism in C** if the following equivalent conditions are satisfied:

014X 1. For each $X \in \text{Obj}(C)$, the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by f is full.

015F 2. For each $X \in \text{Obj}(C)$ and each 2-morphism

$$\gamma: \phi \circ f \Rightarrow \psi \circ f, \quad X \xrightarrow[\psi \circ f]{\gamma \Downarrow} B$$

of C , there exists a 2-morphism $\alpha: \phi \Rightarrow \psi$ of C such that we have an equality

$$A \xrightarrow{f} B \xrightarrow[\psi]{\alpha \Downarrow} X = A \xrightarrow[\psi \circ f]{\phi \Downarrow} X$$

of pasting diagrams in C , i.e. such that we have

$$\gamma = \alpha \star \text{id}_f.$$

014Z Example 18.2.2.1.2. Here are some examples of full epimorphisms.

0150 1. *Full Epimorphisms in Cats_2 .*

0151 2. *Full Epimorphisms in Rel .*

0152 3. *Full Epimorphisms in Span .*

18.2.3 Fully Faithful Epimorphisms

Let C be a bicategory.

0154 Definition 18.2.3.1.1. A 1-morphism $f: A \rightarrow B$ is a **fully faithful epimorphism in C** if the following equivalent conditions are satisfied:

0155 1. The 1-morphism f is fully and faithful.

0156 2. For each $X \in \text{Obj}(C)$, the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by f is fully faithful.

0157 3. The conditions in Item 1 of **Definition 18.2.1.1.1** and Item 1 of **Definition 18.2.2.1.1** hold.

0158 Example 18.2.3.1.2. Here are some examples of fully faithful epimorphisms.

0159 1. *Fully Faithful Epimorphisms in Cats_2 .*

015A 2. *Fully Faithful Epimorphisms in Rel .*

015B 3. *Fully Faithful Epimorphisms in Span .*

18.2.4 Strict Epimorphisms

Let C be a bicategory.

015D Definition 18.2.4.1.1. A 1-morphism $f: A \rightarrow B$ is a **strict epimorphism in C** if the following equivalent conditions are satisfied:

015E 1. For each $X \in \text{Obj}(C)$, the action on objects

$$f^*: \text{Obj}(\text{Hom}_C(B, X)) \rightarrow \text{Obj}(\text{Hom}_C(A, X))$$

of the functor

$$f^*: \text{Hom}_C(B, X) \rightarrow \text{Hom}_C(A, X)$$

given by precomposition by f is injective.

015F 2. For each diagram in C of the form

$$A \xrightarrow{f} B \xrightarrow[\psi]{\phi} X,$$

if $\phi \circ f = \psi \circ f$, then $\phi = \psi$.

015G Example 18.2.4.1.2. Here are some examples of strict epimorphisms.

015H 1. *Strict Epimorphisms in Cats_2 .*

015J 2. *Strict Epimorphisms in Rel .*

015K 3. *Strict Epimorphisms in Span .*

18.3 bicategories of spans

015L Proposition 18.3.0.1.1. Let A and B be objects of C .

015M 1. *As a Pullback.* We have an isomorphism of categories

$$\begin{array}{ccc} \text{Span}(A, B) & \longrightarrow & C_{/B} \\ \downarrow \lrcorner & & \downarrow \text{忘} \\ \text{Span}_C(A, B) & \cong & C_{/A} \times_C C_{/B}, \\ \downarrow & & \downarrow \\ C_{/A} & \xrightarrow{\text{忘}} & C. \end{array}$$

Proof. Item 1, As a Pullback: In detail, the pullback $C_{/A} \times_C C_{/B}$ is the category where

- *Objects.* The objects of $C_{/A} \times_C C_{/B}$ consist of pairs $((S, f), (S', g))$ of objects of C consisting of

- A pair (S, f) in $\text{Obj}(C/A)$ consisting of an object S of C and a morphism $f: S \rightarrow A$ of C ;
- A pair (S', g) in $\text{Obj}(C/B)$ consisting of an object S' of C and a morphism $g: S' \rightarrow B$ of C ;

such that

$$\underbrace{\mathfrak{f}(S, f)}_{\stackrel{\text{def}}{=} S} = \underbrace{\mathfrak{f}(S', g)}_{\stackrel{\text{def}}{=} S'}.$$

Thus the objects of $C/A \times_C C/B$ are the same as spans in C from A to B .

- *Morphisms.* A morphism of $C/A \times_C C/B$ from (S, f, g) to (S', f', g') consists of a pair of morphisms

$$\begin{aligned}\phi: S &\rightarrow S' \\ \psi: S &\rightarrow S'\end{aligned}$$

such that the diagrams

$$\begin{array}{ccc} S & \xrightarrow{\phi} & S' \\ f \searrow & & \swarrow f' \\ & A & \end{array} \quad \begin{array}{ccc} S & \xrightarrow{\psi} & S' \\ g \searrow & & \swarrow g' \\ & B & \end{array}$$

such that

$$\underbrace{\mathfrak{f}(\phi)}_{\stackrel{\text{def}}{=} \phi} = \underbrace{\mathfrak{f}(\psi)}_{\stackrel{\text{def}}{=} \psi}.$$

Thus the morphisms of $C/A \times_C C/B$ are also the same as morphisms of spans in C from (S, f, g) to (S', f', g') .

- *Identities and Composition.* The identities and composition of $C/A \times_C C/B$ are also the same as those in $\text{Span}_C(A, B)$.

This finishes the proof. □

Appendices

18.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets
5. Relations
6. Spans
7. Posets

Indexed and Fibred Sets

7. Indexed Sets
8. Fibred Sets
9. Un/Straightening for Indexed and Fibred Sets

Category Theory

11. Categories
12. Types of Morphisms in Categories
13. Adjunctions and the Yoneda Lemma
14. Constructions With Categories
15. Profunctors
16. Cartesian Closed Categories
17. Kan Extensions

Bicategories

18. Bicategories
19. Internal Adjunctions

Internal Category Theory

20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

41. Stochastic Processes, Martingales, and Brownian Motion

Measure Theory

39. Measurable Spaces

42. Itô Calculus

40. Measures and Integration

43. Stochastic Differential Equations

Probability Theory

40. Probability Theory

Differential Geometry

44. Topological and Smooth Manifolds

Stochastic Analysis

Schemes

45. Schemes

18.2 Other Chapters

Sets

15. Profunctors

1. Sets

16. Cartesian Closed Categories

2. Constructions With Sets

17. Kan Extensions

3. Pointed Sets

Bicategories

4. Tensor Products of Pointed Sets

18. Bicategories

5. Relations

19. Internal Adjunctions

6. Spans

Internal Category Theory

7. Posets

20. Internal Categories

Indexed and Fibred Sets

Cyclic Stuff

7. Indexed Sets

21. The Cycle Category

8. Fibred Sets

Cubical Stuff

9. Un/Straightening for Indexed and Fibred Sets

22. The Cube Category

Category Theory

Globular Stuff

11. Categories

23. The Globe Category

12. Types of Morphisms in Categories

Cellular Stuff

13. Adjunctions and the Yoneda Lemma

24. The Cell Category

14. Constructions With Categories

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero	37. Real Analysis in One Variable
27. Monoids With Zero	38. Real Analysis in Several Variables
28. Constructions With Monoids With Zero	Measure Theory
Groups	39. Measurable Spaces
29. Groups	40. Measures and Integration
30. Constructions With Groups	Probability Theory
Hyper Algebra	40. Probability Theory
31. Hypermonoids	Stochastic Analysis
32. Hypergroups	41. Stochastic Processes, Martingales, and Brownian Motion
33. Hypersemirings and Hyperrings	42. Itô Calculus
34. Quantales	43. Stochastic Differential Equations
Near-Rings	Differential Geometry
35. Near-Semirings	44. Topological and Smooth Manifolds
36. Near-Rings	Schemes
Real Analysis	45. Schemes

Chapter 19

Internal Adjunctions

015N Create tags:

1. <https://www.google.com/search?q=mate+of+an+adjunction>
2. Moreover, by uniqueness of adjoints (Item 2 of Proposition 19.1.2.1.4), this implies also that $S = f^{-1}$.
3. define bicategory $\text{Adj}(C)$
4. walking monad
5. proposition: 2-functors preserve unitors and associators
6. <https://ncatlab.org/nlab/show/2-category+of+adjunctions>. Is there a 3-category too?
7. <https://ncatlab.org/nlab/show/free+monad>
8. <https://ncatlab.org/nlab/show/CatAdj>
9. <https://ncatlab.org/nlab/show/Adj>
10. $\text{Adj}(\text{Adj}(C))$

Contents

19.1 Internal Adjunctions	409
19.1.1 The Walking Adjunction	409
19.1.2 Internal Adjunctions	410
19.1.3 Internal Adjoint Equivalences	416
19.1.4 Mates	418
19.2 Morphisms of Internal Adjunctions	420
19.2.1 Lax Morphisms of Internal Adjunctions	420
19.2.2 Oplax Morphisms of Internal Adjunctions	422

19.2.3	Strong Morphisms of Internal Adjunctions	423
19.2.4	Strict Morphisms of Internal Adjunctions.....	423
19.3	2-Morphisms Between Morphisms of Internal Adjunctions ...	424
19.3.1	2-Morphisms Between Lax Morphisms of Internal Adjunctions.....	424
19.3.2	2-Morphisms Between Oplax Morphisms of Internal Adjunctions.....	424
19.3.3	2-Morphisms Between Strong Morphisms of Internal Adjunctions.....	425
19.3.4	2-Morphisms Between Strict Morphisms of Internal Adjunctions.....	425
19.4	Bicategories of Internal Adjunctions in a Bicategory.....	426
19.A	Other Chapters	426
19.2	Other Chapters	427

19.1 Internal Adjunctions

19.1.1 The Walking Adjunction

015R **Definition 19.1.1.1.** The **walking adjunction** is the bicategory Adj freely generated by¹

- *Objects.* A pair of objects A and B ;
- *Morphisms.* A pair of morphisms

$$\begin{aligned} L: A &\rightarrow B, \\ R: B &\rightarrow A; \end{aligned}$$

- *2-Morphisms.* A pair of 2-morphisms

$$\begin{aligned} \eta: \text{id}_A &\rightarrow R \circ L, \\ \epsilon: L \circ R &\rightarrow \text{id}_B; \end{aligned}$$

¹See [SS86] for an explicit description of the 2-category (as opposed to a bicategory) version of Adj in terms of finite ordinals, similar to the description of the 2-category version of the walking monad (??) as a subcategory of Δ .

subject to the equalities

$$\begin{array}{ccc}
 \begin{array}{c} B \\ \swarrow L \quad \nearrow R \\ \parallel \quad \parallel \\ A \xrightarrow{\text{id}_A} A \end{array} & = & \begin{array}{c} B \xrightarrow{\text{id}_B} B \\ \swarrow L \quad \nearrow R \\ \parallel \quad \parallel \\ A \xrightarrow{\text{id}_A} A \end{array} \\
 \begin{array}{c} A \\ \swarrow R \quad \nearrow L \\ \parallel \quad \parallel \\ B \xrightarrow{\text{id}_B} B \end{array} & = & \begin{array}{c} A \xrightarrow{\text{id}_A} A \\ \swarrow R \quad \nearrow L \\ \parallel \quad \parallel \\ B \xrightarrow{\text{id}_B} B \end{array}
 \end{array}$$

of pasting diagrams, which are equivalent to the following conditions:

1. *The Left Triangle Identity.* The diagram

$$\begin{array}{ccccc}
 L \circ \text{id}_A & \xrightarrow{\text{id}_L * \eta} & L \circ (R \circ L) & \xrightarrow{(\alpha_{L,R,L}^{\text{Adj}})^{-1}} & (L \circ R) \circ L \\
 & \searrow & & & \downarrow \epsilon * \text{id}_L \\
 & & & & \text{id}_B \circ L \\
 & & \rho_L^{\text{Adj}} & & \downarrow \lambda_L^{\text{Adj}} \\
 & & & & L
 \end{array}$$

commutes.

2. *The Right Triangle Identity.* The diagram

$$\begin{array}{ccccc}
 \text{id}_A \circ R & \xrightarrow{\eta * \text{id}_R} & (R \circ L) \circ R & \xrightarrow{\alpha_{R,L,R}^{\text{Adj}}} & R \circ (L \circ R) \\
 & \searrow & & & \downarrow \text{id}_R * \epsilon \\
 & & \lambda_R^{\text{Adj}} & & \downarrow \rho_R^{\text{Adj}} \\
 & & & & R
 \end{array}$$

19.1.2 Internal Adjunctions

Let C be a bicategory.

015T Definition 19.1.2.1.1. An **internal adjunction** in $C^{2,3}$ is a 2-functor $\text{Adj} \rightarrow C$.

015U Remark 19.1.2.1.2. In detail, an **internal adjunction** in C consists of

- **Objects.** A pair of objects A and B of C ;
- **Morphisms.** A pair of morphisms

$$L: A \rightarrow B,$$

$$R: B \rightarrow A$$

of C ;

- **2-Morphisms.** A pair of 2-morphisms

$$\eta: \text{id}_A \rightarrow R \circ L,$$

$$\epsilon: L \circ R \rightarrow \text{id}_B$$

of C ;

subject to the equalities

$$\begin{array}{ccc}
 \begin{array}{c}
 \begin{array}{ccccc}
 & B & \xrightarrow{\text{id}_B} & B & \\
 & \swarrow L & \uparrow \eta & \searrow R & \\
 A & \xrightarrow{\text{id}_A} & & \xrightarrow{\epsilon} & A \\
 & \parallel & & \parallel & \\
 & \uparrow & & \downarrow & \\
 & B & \xrightarrow{\text{id}_B} & B &
 \end{array}
 \end{array}
 & = &
 \begin{array}{c}
 \begin{array}{ccccc}
 & B & \xrightarrow{\text{id}_B} & B & \\
 & \swarrow L & \Downarrow \text{id}_L & \searrow R & \\
 A & \xrightarrow{\text{id}_A} & & & A \\
 & \parallel & & & \\
 & \uparrow & & \downarrow & \\
 & B & \xrightarrow{\text{id}_B} & B &
 \end{array}
 \end{array}
 \\
 \begin{array}{ccc}
 \begin{array}{c}
 \begin{array}{ccccc}
 & A & \xrightarrow{\text{id}_A} & A & \\
 & \swarrow R & \uparrow \epsilon & \searrow L & \\
 B & \xrightarrow{\text{id}_B} & & \xrightarrow{\eta} & A \\
 & \parallel & & \parallel & \\
 & \uparrow & & \downarrow & \\
 & A & \xrightarrow{\text{id}_A} & A &
 \end{array}
 \end{array}
 & = &
 \begin{array}{c}
 \begin{array}{ccccc}
 & A & \xrightarrow{\text{id}_A} & A & \\
 & \swarrow R & \Downarrow \text{id}_R & \searrow L & \\
 B & \xrightarrow{\text{id}_B} & & & A \\
 & \parallel & & & \\
 & \uparrow & & \downarrow & \\
 & A & \xrightarrow{\text{id}_A} & A &
 \end{array}
 \end{array}
 \end{array}$$

of pasting diagrams in C , which are equivalent to the following conditions:⁴

²Further Terminology: Also called an **adjunction internal to C** .

³Further Terminology: In this situation, we also call (g, f) an **adjoint pair**, f the **left adjoint** of the pair, g the **right adjoint** of the pair, η the **unit** of the adjunction, and ϵ the **counit** of the adjunction.

⁴When C is a 2-category, these diagrams take the following form:

$$\begin{array}{ccc}
 \begin{array}{c}
 \begin{array}{ccc}
 L & \xrightarrow{\text{id}_L * \eta} & L \circ R \circ L \\
 \swarrow \text{id}_L & & \downarrow \epsilon * \text{id}_L \\
 & & L
 \end{array}
 \end{array}
 & &
 \begin{array}{c}
 \begin{array}{ccc}
 R & \xrightarrow{\text{id}_R * \eta} & R \circ L \circ R \\
 \swarrow \text{id}_L & & \downarrow \epsilon * \text{id}_R \\
 & & R
 \end{array}
 \end{array}
 \end{array}$$

1. *The Left Triangle Identity.* The diagram

$$\begin{array}{ccccc}
 L \circ \text{id}_A & \xrightarrow{\text{id}_L \star \eta} & L \circ (R \circ L) & \xrightarrow{(\alpha_{L,R,L}^C)^{-1}} & (L \circ R) \circ L \\
 & \searrow \rho_L^C & & & \downarrow \epsilon \star \text{id}_L \\
 & & & & \text{id}_B \circ L \\
 & & & & \downarrow \lambda_L^C \\
 & & & & L
 \end{array}$$

commutes.

2. *The Right Triangle Identity.* The diagram

$$\begin{array}{ccccc}
 \text{id}_A \circ R & \xrightarrow{\eta \star \text{id}_R} & (R \circ L) \circ R & \xrightarrow{\alpha_{R,L,R}^C} & R \circ (L \circ R) \\
 & \searrow \lambda_R^C & & & \downarrow \text{id}_R \star \epsilon \\
 & & & & R \circ \text{id}_B \\
 & & & & \downarrow \rho_R^C \\
 & & & & R.
 \end{array}$$

015V Example 19.1.2.1.3. Here are some examples of internal adjunctions.

- 015W** 1. *Internal Adjunctions in Cats_2 .* The internal adjunctions in the 2-category Cats_2 of categories, functors, and natural transformations are precisely the adjunctions of ??.
- 015X** 2. *Internal Adjunctions in Rel .* The internal adjunctions in Rel are precisely the relations of the form $\text{Gr}(f) \dashv f^{-1}$ with f a function; see ?? of ??.
- 015Y** 3. *Internal Adjunctions in Span .* The internal adjunctions in Span are precisely the spans of the form

$$\begin{array}{ccc}
 & S & \\
 \phi \swarrow & & \searrow g \\
 A & & B
 \end{array}$$

with ϕ an isomorphism; see ?? of ??.

015Z Proposition 19.1.2.1.4. Let C be a bicategory.

- 0160 1. *Duality.* Let (f, g, η, ϵ) be an internal adjunction in C .
- The quadruple (g, f, η, ϵ) is an internal adjunction in C^{op} .
 - The quadruple (g, f, ϵ, η) is an internal adjunction in C^{co} .
 - The quadruple (f, g, η, ϵ) is an internal adjunction in C^{coop} .
- 0161 2. *Uniqueness of Adjoints.* Let (f, g, η, ϵ) and $(f', g', \eta', \epsilon')$ be internal adjunctions in C . We have a canonical isomorphism⁵

$$g \xrightarrow{(\lambda_g^C)^{-1}} \text{id}_A \circ g \xrightarrow{\eta' \star \text{id}_g} (g' \circ f) \circ g \xrightarrow{\alpha_{g', f, g}^C} g' \circ (f \circ g) \xrightarrow{\text{id}_{g'} \star \epsilon} g' \circ \text{id}_B \xrightarrow{(\rho_{g'}^C)^{-1}} g'$$

with inverse

$$g' \xrightarrow{(\lambda_{g'}^C)^{-1}} \text{id}_B \circ g' \xrightarrow{\eta \star \text{id}_{g'}} (g \circ f) \circ g' \xrightarrow{\alpha_{g', f, g}^C} g \circ (f \circ g') \xrightarrow{\text{id}_{g'} \star \epsilon'} g \circ \text{id}_B \xrightarrow{(\lambda_g^C)^{-1}} g.$$

- 0162 3. *Carrying Internal Adjunctions Through Pseudofunctors.* Let $F: C \rightarrow \mathcal{D}$ be a pseudofunctor and (f, g, η, ϵ) be an internal adjunction in C . There is an induced internal adjunction⁶

$$(F(f), F(g), \bar{\eta}, \bar{\epsilon})$$

in \mathcal{D} , where:

- (a) The unit

$$\bar{\eta}: \text{id}_{F(A)} \Rightarrow F(g) \circ F(f)$$

is the composition

$$\text{id}_{F(A)} \xrightarrow{F_A} F(\text{id}_A) \xrightarrow{F(\eta)} F(g \circ f) \xrightarrow{F_{g,f}^{-1}} F(g) \circ F(f).$$

- (b) The counit

$$\bar{\epsilon}: F(f) \circ F(g) \Rightarrow \text{id}_{F(B)}$$

is the composition

$$F(f) \circ F(g) \xrightarrow{F_{f,g}} F(f \circ g) \xrightarrow{F(\epsilon)} F(\text{id}_B) \xrightarrow{F_B} \text{id}_{F(B)}.$$

Proof. Item 1, Duality: Omitted.⁷

Item 2, Uniqueness of Adjoints:⁸ Consider the diagram (if you *really* want to consider it I


⁵ *Slogan:* Left adjoints are unique up to canonical isomorphism. Dually, so are right adjoints.

⁶  *Warning:* Lax or oplax functors which are not pseudofunctors need not preserve internal adjunctions.

⁷ Reference: [JY21, Exercise 6.6.2].

⁸ Reference: [JY21, Lemma 6.1.6].

fear you will need to zoom in)



In this diagram:


1. The morphisms in green are the composition $g \xrightarrow{\cong} g' \xrightarrow{\cong} g$;
2. The morphisms in red are equal to λ_g^C by the right triangle identity for (f, g, η, ϵ) . Hence the composition of the morphism in blue with the morphisms in red is the identity;
3. Subdiagrams (1), (2), (10), (11), (29), (31), and (43) commute by the naturality of the left unit of C and its inverse;
4. Subdiagrams (8), (19), and (21) commute by the naturality of the right unit of C and its inverse;
5. Subdiagrams (6), (13), (17), (18), (20), (22), (32), (33), (36), (38), (40), (41), and (45) commute by the naturality of the associator of C and its inverse;
6. Subdiagrams (37), (39), and (42) commute by the pentagon identity for C ;
7. Subdiagrams (3), (4), (7), (12), (25), (30), and (48) commute by ?? of ??;
8. Subdiagrams (5), (14), (23), (24), (34), and (35) commute by middle-four exchange;
9. Subdiagrams (9), (15), (16), (27), (28), (44), (46), (49), and (50) commute trivially;
10. Subdiagram (26) commutes by ??? of ??;

11. Subdiagram (47) commutes by ?? of ?? and the naturality of the left unitor of right unitor of C .

Hence $g \cong g'$.

Item 3, Carrying Internal Adjunctions Through Pseudofunctors: ⁹We claim that the left and right triangle identities for $(F(f), F(g), \bar{\eta}, \bar{\epsilon})$ hold:

1. The left triangle identity for $(F(f), F(g), \bar{\eta}, \bar{\epsilon})$ is the condition that the boundary diagram of the diagram (you may need to zoom in)



commutes. Since


- (a) Subdiagrams (1) and (7) commute by applying middle-four exchange twice,
- (b) Subdiagrams (2) and (8) commute by the left and right lax unity conditions for F ,
- (c) Subdiagrams (3) and (6) commute by the naturality of the lax functoriality constraints of F ,
- (d) Subdiagram (4) commutes by the lax associativity condition for F , and
- (e) Subdiagram (5) commutes by the left triangle identity for (f, g, η, ϵ) ,

so does the boundary diagram.

2. The right triangle identity for $(F(f), F(g), \bar{\eta}, \bar{\epsilon})$ is the condition that the boundary

⁹Reference: [JY21, Proposition 6.1.7].

diagram of the diagram (you may need to zoom in)



commutes. Since

- (a) Subdiagrams (1) and (7) commute by applying middle-four exchange twice,
- (b) Subdiagrams (2) and (8) commute by the left and right lax unity conditions for F ,
- (c) Subdiagrams (3) and (6) commute by the naturality of the lax functoriality constraints of F ,
- (d) Subdiagram (4) commutes by the lax associativity condition for F , and
- (e) Subdiagram (5) commutes by the right triangle identity for (f, g, η, ϵ) ,

so does the boundary diagram.

This finishes the proof. \square

19.1.3 Internal Adjoint Equivalences

Let C be a bicategory.

0164 Definition 19.1.3.1.1. An internal adjunction (f, g, η, ϵ) in C is an **internal adjoint equivalence** if η and ϵ are isomorphisms in C .

0165 Example 19.1.3.1.2. Here are some examples of internal adjoint equivalences.

0166 1. *Internal Adjoint Equivalences in Cats₂*. The internal adjoint equivalences in the

2-category Cats_2 of categories, functors, and natural transformations are precisely the adjoint equivalences of ?? .¹⁰

- 0167 2. *Internal Adjoint Equivalences in Mod.* The internal adjoint equivalences in Mod are precisely the invertible R -modules; see ?? .¹¹
- 0168 3. *Internal Adjoint Equivalences in $\text{PseudoFun}(C, \mathcal{D})$.* The internal adjoint equivalences in $\text{PseudoFun}(C, \mathcal{D})$ are precisely the invertible strong transformations; see ?? .¹²
- 0169 4. *Internal Adjoint Equivalences in Rel .* The internal adjoint equivalences in Rel are precisely the relations of the form $\text{Gr}(f) \dashv f^{-1}$ with f an isomorphism; see ?? .
- 016A 5. *Internal Adjoint Equivalences in Span .* The internal adjoint equivalences in Span are precisely the spans of the form $A \xleftarrow{\phi} S \xrightarrow{\psi} B$ with ϕ and ψ isomorphisms; see ?? .

016B **Proposition 19.1.3.1.3.** Let C be a bicategory.

- 016C 1. *Carrying Internal Adjoint Equivalences Through Pseudofunctors.* Let $F: C \rightarrow \mathcal{D}$ be a pseudofunctor and (f, g, η, ϵ) be an internal adjunction in C . If (f, g, η, ϵ) is an internal adjoint equivalence in C , then the induced internal adjunction

$$(F(f), F(g), \bar{\eta}, \bar{\epsilon})$$

in \mathcal{D} of Item 3 of Proposition 19.1.2.1.4 is an internal adjoint equivalence as well.

- 016D 2. *Internal Adjunctions Always Refine to Internal Adjoint Equivalences.* Let (f, g, η, ϵ) be an internal adjunction in C . If f is an equivalence, then there exist 2-morphisms

$$\begin{aligned}\bar{\eta}: \text{id}_A &\Longrightarrow g \circ f \\ \bar{\epsilon}: f \circ g &\Longrightarrow \text{id}_B\end{aligned}$$

of C such that $(f, g, \bar{\eta}, \bar{\epsilon})$ is an internal adjoint equivalence.

Proof. Item 1, Carrying Internal Adjoint Equivalences Through Pseudofunctors: See [JY21, Proposition 6.2.3].

Item 2, Internal Adjunctions Always Refine to Internal Adjoint Equivalences: See [JY21, Proposition 6.2.4]. \square

¹⁰Reference: [JY21, Examples 6.2.5].

¹¹Reference: [JY21, Examples 6.2.6].

¹²Reference: [JY21, Examples 6.2.7].

19.1.4 Mates

Let C be a bicategory, let (f, g, η, ϵ) and $(f', g', \eta', \epsilon')$ be adjunctions, and let h and k be morphisms of C as in the diagram

$$\begin{array}{ccc} & f & \\ A & \begin{array}{c} \swarrow \perp \searrow \\ g \end{array} & B \\ h \downarrow & & \downarrow k \\ C & \begin{array}{c} \swarrow \perp \searrow \\ g' \end{array} & D. \end{array}$$


016F Definition 19.1.4.1.1. The **mates** of a pair of 2-morphisms

$$\begin{array}{ccc} A & \xrightarrow{f} & B \\ h \downarrow & \nearrow \omega & \downarrow k \\ C & \xrightarrow{f'} & D \end{array} \quad \begin{array}{l} \omega: f' \circ h \Rightarrow k \circ f, \\ v: h \circ g \Rightarrow g' \circ k \end{array} \quad \begin{array}{ccc} A & \xleftarrow{g} & B \\ h \downarrow & \nearrow v & \downarrow k \\ C & \xleftarrow{g'} & D \end{array}$$

are the 2-morphisms

$$\begin{array}{ccc} A & \xleftarrow{g} & B \\ h \downarrow & \nearrow \omega^\dagger & \downarrow k \\ C & \xleftarrow{g'} & D \end{array} \quad \begin{array}{l} \omega^\dagger: h \circ g \Rightarrow g' \circ k, \\ v^\dagger: f' \circ h \Rightarrow k \circ f \end{array} \quad \begin{array}{ccc} A & \xrightarrow{f} & B \\ h \downarrow & \nearrow v^\dagger & \downarrow k \\ C & \xrightarrow{f'} & D \end{array}$$

defined as the pastings of the diagrams¹³



016G Proposition 19.1.4.1.2. Let $\omega: f' \circ h \Rightarrow k \circ f$ and $v: h \circ g \Rightarrow g' \circ k$ be 2-morphisms.

016H 1. *The Mate Correspondence.* The map

$$(-)^\dagger: \text{Hom}_{\text{Hom}_C(A,C)}(f' \circ h, k \circ f) \longrightarrow \text{Hom}_{\text{Hom}_C(B,D)}(h \circ g, g' \circ k)$$

$$\omega \longmapsto \omega^\dagger$$

is a bijection.

Proof. Item 1, The Mate Correspondence: Here we give a proof for 2-categories (which indirectly proves also the general case by ??). A proof for general bicategories can be found in [JY21, Lemma 6.1.13].

¹³If C is a 2-category, these pasting diagrams become the following:

$$\begin{array}{ccc}
 \begin{array}{c}
 \begin{array}{ccc}
 B & \xrightarrow{g} & A \\
 k \downarrow & \swarrow \omega^\dagger & \downarrow h \\
 D & \xrightarrow{g'} & C
 \end{array}
 \end{array}
 & = &
 \begin{array}{c}
 \begin{array}{ccccc}
 B & \xrightarrow{g} & A & \xrightarrow{h} & C \\
 \downarrow \text{id}_B & \swarrow \epsilon & \downarrow f & \swarrow \omega & \downarrow \text{id}_C \\
 B & \xrightarrow{k} & D & \xrightarrow{f'} & C
 \end{array}
 \end{array}
 \\
 \begin{array}{c}
 \begin{array}{ccc}
 C & \xrightarrow{f'} & D \\
 h \uparrow & \searrow v^\dagger & \uparrow k \\
 A & \xrightarrow{f} & B
 \end{array}
 \end{array}
 & = &
 \begin{array}{c}
 \begin{array}{ccccc}
 A & \xrightarrow{\text{id}_A} & B & \xrightarrow{g} & C & \xrightarrow{f'} & D \\
 \downarrow \eta & \swarrow \text{id}_B & \downarrow \text{id}_B & \swarrow \nu & \downarrow \text{id}_C & \swarrow \epsilon' & \downarrow \text{id}_D \\
 A & \xrightarrow{f} & B & \xrightarrow{k} & C & \xrightarrow{f'} & D
 \end{array}
 \end{array}
 \end{array}$$

Let

$$v: h \circ g \Rightarrow g' \circ k$$

be a 2-morphism of C . The mate v^\dagger of v is then given by

and the mate of v^\dagger is the 2-morphism $(v^\dagger)^\dagger: f' \circ h \Rightarrow k \circ f$ given by

Similarly, $(\omega)^\dagger = \omega$. □

19.2 Morphisms of Internal Adjunctions

19.2.1 Lax Morphisms of Internal Adjunctions

Let C be a bicategory and let $(A, B, F, G, \eta, \epsilon)$ and $(A', B', F', G', \eta', \epsilon')$ be internal adjunctions in C .

016L Definition 19.2.1.1.1. A **lax morphism of internal adjunctions** from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$ is a lax transformation between these viewed as 2-functors from the walking adjunction.

016M Remark 19.2.1.1.2. In detail, a **lax morphism of internal adjunctions** from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$ consists of

- **1-Morphisms.** A pair of 1-morphisms

$$\phi: A \rightarrow A',$$

$$\psi: B \rightarrow B'$$

of \mathcal{C} ;

- **2-Morphisms.** A pair of 2-morphisms

$$\begin{array}{ccc} A & \xrightarrow{F} & B \\ \phi \downarrow & \nearrow \alpha & \downarrow \psi \\ A' & \xrightarrow{F'} & B' \end{array}$$

$$\begin{array}{l} \alpha: F' \circ \phi \Rightarrow \psi \circ F, \\ \beta: G' \circ \phi \Rightarrow \psi \circ G \end{array}$$

$$\begin{array}{ccc} A & \xleftarrow{G} & B \\ \phi \downarrow & \nearrow \beta & \downarrow \psi \\ A' & \xleftarrow{G'} & B' \end{array}$$

of \mathcal{C} ;

satisfying the following conditions:

1. **Compatibility With Units.** We have an equality

$$\begin{array}{ccc} \text{Diagram showing compatibility with units: } & = & \text{Diagram showing compatibility with counits: } \\ \begin{array}{c} \text{A complex commutative diagram involving } A, B, A', B', F, G, \eta, \epsilon, \phi, \psi, \lambda_\phi^C, \rho_\phi^{C,-1}, \text{ and id}_A, \text{id}_{A'}, \text{id}_B, \text{id}_{B'} \text{.} \\ \text{It shows the equality of two pasting diagrams in } \mathcal{C}. \end{array} & = & \begin{array}{c} \text{A complex commutative diagram involving } A, B, A', B', F, G, \eta, \epsilon, \phi, \psi, \alpha, \beta, \lambda_\phi^C, \rho_\phi^{C,-1}, \text{ and id}_A, \text{id}_{A'}, \text{id}_B, \text{id}_{B'} \text{.} \\ \text{It shows the equality of two pasting diagrams in } \mathcal{C}. \end{array} \end{array}$$

of pasting diagrams in \mathcal{C} ;

2. **Compatibility With Counits.** We have an equality

$$\begin{array}{ccc} \text{Diagram showing compatibility with counits: } & = & \text{Diagram showing compatibility with units: } \\ \begin{array}{c} \text{A complex commutative diagram involving } B, A, B', A', \psi, \phi, \epsilon, \epsilon', \alpha, \beta, \lambda_\psi^C, \rho_\psi^{C,-1}, \text{ and id}_B, \text{id}_{B'}, \text{id}_A, \text{id}_{A'} \text{.} \\ \text{It shows the equality of two pasting diagrams in } \mathcal{C}. \end{array} & = & \begin{array}{c} \text{A complex commutative diagram involving } B, A, B', A', \psi, \phi, \epsilon, \epsilon', \alpha, \beta, \lambda_\psi^C, \rho_\psi^{C,-1}, \text{ and id}_B, \text{id}_{B'}, \text{id}_A, \text{id}_{A'} \text{.} \\ \text{It shows the equality of two pasting diagrams in } \mathcal{C}. \end{array} \end{array}$$

of pasting diagrams in \mathcal{C} .

19.2.2 Oplax Morphisms of Internal Adjunctions

Let C be a bicategory and let $(A, B, F, G, \eta, \epsilon)$ and $(A', B', F', G', \eta', \epsilon')$ be internal adjunctions in C .

016P Definition 19.2.2.1.1. An **oplax morphism of internal adjunctions** from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$ is an oplax transformation between these viewed as 2-functors from the walking adjunction.

016Q Remark 19.2.2.1.2. In detail, an **oplax morphism of internal adjunctions** from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$ consists of

- **1-Morphisms.** A pair of 1-morphisms

$$\begin{aligned}\phi: A &\rightarrow A', \\ \psi: B &\rightarrow B'\end{aligned}$$

of C ;

- **2-Morphisms.** A pair of 2-morphisms

$$\begin{array}{ccc} \begin{array}{ccc} A & \xrightarrow{F} & B \\ \phi \downarrow & \swarrow \alpha & \downarrow \psi \\ A' & \xrightarrow{F'} & B' \end{array} & \alpha: \psi \circ F \Rightarrow F' \circ \phi, & \begin{array}{ccc} A & \xleftarrow{G} & B \\ \phi \downarrow & \swarrow \beta & \downarrow \psi \\ A' & \xleftarrow{G'} & B' \end{array} \\ \beta: \psi \circ G \Rightarrow G' \circ \phi & & \end{array}$$

of C ;

satisfying the following conditions:

1. **Compatibility With Units.** We have an equality

$$\begin{array}{ccc} \begin{array}{c} \text{Diagram showing compatibility with units: } \psi \circ F = G \circ \phi \text{ and } \psi \circ G = F \circ \phi. \\ \text{It includes 1-morphisms } \phi, \psi, F, G, \text{ and 2-morphisms } \alpha, \beta, \lambda_\phi^{C,-1}, \rho_\phi^C, \text{ and } \epsilon, \epsilon'. \end{array} & = & \begin{array}{c} \text{Diagram showing compatibility with units: } \psi \circ F = G \circ \phi \text{ and } \psi \circ G = F \circ \phi. \\ \text{It includes 1-morphisms } \phi, \psi, F, G, \text{ and 2-morphisms } \alpha, \beta, \lambda_\phi^{C,-1}, \rho_\phi^C, \text{ and } \epsilon, \epsilon'. \end{array} \end{array}$$

of pasting diagrams in C ;

2. *Compatibility With Counits.* We have an equality

of pasting diagrams in \mathcal{C} .

19.2.3 Strong Morphisms of Internal Adjunctions

Let \mathcal{C} be a bicategory and let $(A, B, F, G, \eta, \epsilon)$ and $(A', B', F', G', \eta', \epsilon')$ be internal adjunctions in \mathcal{C} .

016S Definition 19.2.3.1.1. A **strong morphism of internal adjunctions** from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$ is a strong transformation between these viewed as 2-functors from the walking adjunction.

016T Remark 19.2.3.1.2. In detail, a **strong morphism of internal adjunctions** from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$ is equivalently:

1. A lax morphism of internal adjunctions as in [Remark 19.2.1.1.2](#) whose 2-morphisms are invertible.
2. An oplax morphism of internal adjunctions as in [Remark 19.2.2.1.2](#) whose 2-morphisms are invertible.

19.2.4 Strict Morphisms of Internal Adjunctions

Let \mathcal{C} be a bicategory and let $(A, B, F, G, \eta, \epsilon)$ and $(A', B', F', G', \eta', \epsilon')$ be internal adjunctions in \mathcal{C} .

016V Definition 19.2.4.1.1. A **strict morphism of internal adjunctions** from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$ is a strict transformation between these viewed as 2-functors from the walking adjunction.

016W Remark 19.2.4.1.2. In detail, a **strict morphism of internal adjunctions** from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$ is equivalently:

1. A lax morphism of internal adjunctions as in [Remark 19.2.1.1.2](#) whose 2-morphisms are identities.
2. An oplax morphism of internal adjunctions as in [Remark 19.2.2.1.2](#) whose 2-morphisms are identities.

19.3 2-Morphisms Between Morphisms of Internal Adjunctions

19.3.1 2-Morphisms Between Lax Morphisms of Internal Adjunctions

Let C be a bicategory, let $(A, B, F, G, \eta, \epsilon)$ and $(A', B', F', G', \eta', \epsilon')$ be internal adjunctions in C , and let $(\phi_1, \psi_1, \alpha_1, \beta_1)$ and $(\phi_2, \psi_2, \alpha_2, \beta_2)$ be lax morphisms of internal adjunctions from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$.

016Z Definition 19.3.1.1.1. A **2-morphism from** $(\phi_1, \psi_1, \alpha_1, \beta_1)$ **to** $(\phi_2, \psi_2, \alpha_2, \beta_2)$ is a modification between these viewed as lax transformations.

0170 Remark 19.3.1.1.2. In detail, a **2-morphism from** $(\phi_1, \psi_1, \alpha_1, \beta_1)$ **to** $(\phi_2, \psi_2, \alpha_2, \beta_2)$ consist of 2-morphisms

$$\begin{aligned}\Gamma: \phi_1 &\Rightarrow \phi_2 \\ \Sigma: \psi_1 &\Rightarrow \psi_2\end{aligned}$$

of C such that we have equalities

$$\begin{array}{ccc} \begin{array}{c} A \xrightarrow{F} B \\ \phi_1 \left(\begin{array}{c} \xrightarrow{\Gamma} \\ \Downarrow \end{array} \right) \phi_2 \quad \Downarrow \alpha_2 \quad \Downarrow \psi_2 \\ A' \xrightarrow{F'} B' \end{array} & = & \begin{array}{c} A \xrightarrow{F} B \\ \phi_1 \left(\begin{array}{c} \Downarrow \alpha_1 \\ \xrightarrow{\Sigma} \end{array} \right) \psi_1 \quad \Downarrow \psi_2 \\ A' \xrightarrow{F'} B' \end{array} \\ \begin{array}{c} B \xrightarrow{G} A \\ \psi_1 \left(\begin{array}{c} \xrightarrow{\Sigma} \\ \Downarrow \end{array} \right) \psi_2 \quad \Downarrow \beta_2 \quad \Downarrow \phi_2 \\ B' \xrightarrow{G'} A' \end{array} & = & \begin{array}{c} B \xrightarrow{G} A \\ \psi_1 \left(\begin{array}{c} \Downarrow \beta_1 \\ \xrightarrow{\Gamma} \end{array} \right) \phi_1 \quad \Downarrow \phi_2 \\ B' \xrightarrow{G'} A' \end{array} \end{array}$$

of pasting diagrams in C .

19.3.2 2-Morphisms Between Oplax Morphisms of Internal Adjunctions

Let C be a bicategory, let $(A, B, F, G, \eta, \epsilon)$ and $(A', B', F', G', \eta', \epsilon')$ be internal adjunctions in C , and let $(\phi_1, \psi_1, \alpha_1, \beta_1)$ and $(\phi_2, \psi_2, \alpha_2, \beta_2)$ be oplax morphisms of internal adjunctions from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$.

0172 Definition 19.3.2.1.1. A **2-morphism from** $(\phi_1, \psi_1, \alpha_1, \beta_1)$ **to** $(\phi_2, \psi_2, \alpha_2, \beta_2)$ is a modification between these viewed as oplax transformations.

0173 Remark 19.3.2.1.2. In detail, a **2-morphism from** $(\phi_1, \psi_1, \alpha_1, \beta_1)$ **to** $(\phi_2, \psi_2, \alpha_2, \beta_2)$ consist of 2-morphisms

$$\begin{aligned}\Gamma: \phi_1 &\Rightarrow \phi_2 \\ \Sigma: \psi_1 &\Rightarrow \psi_2\end{aligned}$$

of C such that we have equalities

$$\begin{array}{ccc}
 \begin{array}{c} A \xrightarrow{F} B \\ \phi_2 \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} \right) \phi_1 \quad \alpha_1 \swarrow \searrow \\ A' \xrightarrow{F'} B' \end{array} & = & \begin{array}{c} A \xrightarrow{F} B \\ \phi_2 \left(\begin{array}{c} \text{---} \\ \alpha_2 \swarrow \psi_2 \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} \right) \psi_1 \end{array} \right) \end{array} \\
 \begin{array}{c} B \xrightarrow{G} A \\ \psi_2 \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} \right) \psi_1 \quad \beta_1 \swarrow \searrow \\ B' \xrightarrow{G'} A' \end{array} & = & \begin{array}{c} B \xrightarrow{G} A \\ \psi_2 \left(\begin{array}{c} \text{---} \\ \beta_2 \swarrow \phi_2 \left(\begin{array}{c} \text{---} \\ \text{---} \end{array} \right) \phi_1 \end{array} \right) \end{array}
 \end{array}$$

of pasting diagrams in C .

19.3.3 2-Morphisms Between Strong Morphisms of Internal Adjunctions

Let C be a bicategory, let $(A, B, F, G, \eta, \epsilon)$ and $(A', B', F', G', \eta', \epsilon')$ be internal adjunctions in C , and let $(\phi_1, \psi_1, \alpha_1, \beta_1)$ and $(\phi_2, \psi_2, \alpha_2, \beta_2)$ be strong morphisms of internal adjunctions from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$.

0175 Definition 19.3.3.1.1. A **2-morphism from** $(\phi_1, \psi_1, \alpha_1, \beta_1)$ **to** $(\phi_2, \psi_2, \alpha_2, \beta_2)$ is a modification between these viewed as strong transformations.

0176 Remark 19.3.3.1.2. In detail, a **2-morphism from** $(\phi_1, \psi_1, \alpha_1, \beta_1)$ **to** $(\phi_2, \psi_2, \alpha_2, \beta_2)$ is equivalently:

- A 2-morphism (Γ, Σ) from $(\phi_1, \psi_1, \alpha_1, \beta_1)$ to $(\phi_2, \psi_2, \alpha_2, \beta_2)$ viewed as lax transformations as in [Remark 19.3.1.1.2](#).
- A 2-morphism (Γ, Σ) from $(\phi_1, \psi_1, \alpha_1, \beta_1)$ to $(\phi_2, \psi_2, \alpha_2, \beta_2)$ viewed as oplax transformations as in [Remark 19.3.2.1.2](#).

19.3.4 2-Morphisms Between Strict Morphisms of Internal Adjunctions

Let C be a bicategory, let $(A, B, F, G, \eta, \epsilon)$ and $(A', B', F', G', \eta', \epsilon')$ be internal adjunctions in C , and let $(\phi_1, \psi_1, \alpha_1, \beta_1)$ and $(\phi_2, \psi_2, \alpha_2, \beta_2)$ be lax morphisms of internal adjunctions from $(A, B, F, G, \eta, \epsilon)$ to $(A', B', F', G', \eta', \epsilon')$.

0178 Definition 19.3.4.1.1. A **2-morphism from** $(\phi_1, \psi_1, \alpha_1, \beta_1)$ **to** $(\phi_2, \psi_2, \alpha_2, \beta_2)$ is a modification between these viewed as strict transformations.

0179 Remark 19.3.4.1.2. In detail, a **2-morphism from** $(\phi_1, \psi_1, \alpha_1, \beta_1)$ **to** $(\phi_2, \psi_2, \alpha_2, \beta_2)$ is equivalently:

- A 2-morphism (Γ, Σ) from $(\phi_1, \psi_1, \alpha_1, \beta_1)$ to $(\phi_2, \psi_2, \alpha_2, \beta_2)$ viewed as lax transformations as in [Remark 19.3.1.1.2](#).

- A 2-morphism (Γ, Σ) from $(\phi_1, \psi_1, \alpha_1, \beta_1)$ to $(\phi_2, \psi_2, \alpha_2, \beta_2)$ viewed as oplax transformations as in Remark 19.3.2.1.2.
- A 2-morphism (Γ, Σ) from $(\phi_1, \psi_1, \alpha_1, \beta_1)$ to $(\phi_2, \psi_2, \alpha_2, \beta_2)$ viewed as strong transformations as in Remark 19.3.3.1.2.

19.4 Bicategories of Internal Adjunctions in a Bicategory

Appendices

19.A Other Chapters

Sets	16. Cartesian Closed Categories
1. Sets	17. Kan Extensions
2. Constructions With Sets	Bicategories
3. Pointed Sets	18. Bicategories
4. Tensor Products of Pointed Sets	19. Internal Adjunctions
5. Relations	Internal Category Theory
6. Spans	20. Internal Categories
7. Posets	Cyclic Stuff
Indexed and Fibred Sets	21. The Cycle Category
7. Indexed Sets	Cubical Stuff
8. Fibred Sets	22. The Cube Category
9. Un/Straightening for Indexed and Fibred Sets	Globular Stuff
Category Theory	23. The Globe Category
11. Categories	Cellular Stuff
12. Types of Morphisms in Categories	24. The Cell Category
13. Adjunctions and the Yoneda Lemma	Monoids
14. Constructions With Categories	25. Monoids
15. Profunctors	26. Constructions With Monoids
	Monoids With Zero
	27. Monoids With Zero

-
28. Constructions With Monoids With Zero 38. Real Analysis in Several Variables

Groups

29. Groups
30. Constructions With Groups

Hyper Algebra

31. Hypermonoids
32. Hypergroups
33. Hypersemirings and Hyperrings
34. Quantales

Near-Rings

35. Near-Semirings
36. Near-Rings
37. Real Analysis in One Variable

19.2 Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets
5. Relations
6. Spans
7. Posets

Indexed and Fibred Sets

7. Indexed Sets
8. Fibred Sets

Measure Theory

39. Measurable Spaces
40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales, and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

9. Un/Straightening for Indexed and Fibred Sets

Category Theory

11. Categories
12. Types of Morphisms in Categories
13. Adjunctions and the Yoneda Lemma
14. Constructions With Categories
15. Profunctors
16. Cartesian Closed Categories
17. Kan Extensions

Bicategories

18. Bicategories

19. Internal Adjunctions	32. Hypergroups
Internal Category Theory	33. Hypersemirings and Hyperrings
20. Internal Categories	34. Quantales
Cyclic Stuff	Near-Rings
21. The Cycle Category	35. Near-Semirings
Cubical Stuff	36. Near-Rings
22. The Cube Category	Real Analysis
Globular Stuff	37. Real Analysis in One Variable
23. The Globe Category	38. Real Analysis in Several Variables
Cellular Stuff	Measure Theory
24. The Cell Category	39. Measurable Spaces
Monoids	40. Measures and Integration
25. Monoids	Probability Theory
26. Constructions With Monoids	40. Probability Theory
Monoids With Zero	Stochastic Analysis
27. Monoids With Zero	41. Stochastic Processes, Martingales, and Brownian Motion
28. Constructions With Monoids With Zero	42. Itô Calculus
Groups	43. Stochastic Differential Equations
29. Groups	Differential Geometry
30. Constructions With Groups	44. Topological and Smooth Manifolds
Hyper Algebra	Schemes
31. Hypermonoids	45. Schemes

Part V

Internal Category Theory

Chapter 20

Internal Categories

017B

Contents

20.A Other Chapters	430
20.2 Other Chapters	432

Appendices

20.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff	34. Quantales
21. The Cycle Category	
Cubical Stuff	Near-Rings
22. The Cube Category	35. Near-Semirings
Globular Stuff	36. Near-Rings
23. The Globe Category	Real Analysis
Cellular Stuff	37. Real Analysis in One Variable
24. The Cell Category	38. Real Analysis in Several Variables
Monoids	Measure Theory
25. Monoids	39. Measurable Spaces
26. Constructions With Monoids	40. Measures and Integration
Monoids With Zero	Probability Theory
27. Monoids With Zero	40. Probability Theory
28. Constructions With Monoids With Zero	Stochastic Analysis
Groups	41. Stochastic Processes, Martingales, and Brownian Motion
29. Groups	42. Itô Calculus
30. Constructions With Groups	43. Stochastic Differential Equations
Hyper Algebra	Differential Geometry
31. Hypermonoids	44. Topological and Smooth Manifolds
32. Hypergroups	Schemes
33. Hypersemirings and Hyperrings	45. Schemes

20.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
Category Theory	27. Monoids With Zero
11. Categories	28. Constructions With Monoids With Zero
12. Types of Morphisms in Categories	Groups
13. Adjunctions and the Yoneda Lemma	29. Groups
14. Constructions With Categories	30. Constructions With Groups
15. Profunctors	Hyper Algebra
16. Cartesian Closed Categories	31. Hypermonoids
17. Kan Extensions	32. Hypergroups
Bicategories	33. Hypersemirings and Hyperrings
18. Bicategories	34. Quantales
19. Internal Adjunctions	Near-Rings
Internal Category Theory	35. Near-Semirings
20. Internal Categories	36. Near-Rings
	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part VI

Cyclic Stuff

Chapter 21

The Cycle Category

017C

Contents

21.A Other Chapters	435
21.2 Other Chapters	437

Appendices

21.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff	34. Quantales
21. The Cycle Category	
Cubical Stuff	Near-Rings
22. The Cube Category	35. Near-Semirings
Globular Stuff	36. Near-Rings
23. The Globe Category	Real Analysis
Cellular Stuff	37. Real Analysis in One Variable
24. The Cell Category	38. Real Analysis in Several Variables
Monoids	Measure Theory
25. Monoids	39. Measurable Spaces
26. Constructions With Monoids	40. Measures and Integration
Monoids With Zero	Probability Theory
27. Monoids With Zero	40. Probability Theory
28. Constructions With Monoids With Zero	Stochastic Analysis
Groups	41. Stochastic Processes, Martingales, and Brownian Motion
29. Groups	42. Itô Calculus
30. Constructions With Groups	43. Stochastic Differential Equations
Hyper Algebra	Differential Geometry
31. Hypermonoids	44. Topological and Smooth Manifolds
32. Hypergroups	Schemes
33. Hypersemirings and Hyperrings	45. Schemes

21.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part VII

Cubical Stuff

Chapter 22

The Cube Category

017D

Contents

22.A Other Chapters	440
22.2 Other Chapters	442

Appendices

22.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff[21. The Cycle Category](#)**Cubical Stuff**[22. The Cube Category](#)**Globular Stuff**[23. The Globe Category](#)**Cellular Stuff**[24. The Cell Category](#)**Monoids**[25. Monoids](#)[26. Constructions With Monoids](#)**Monoids With Zero**[27. Monoids With Zero](#)[28. Constructions With Monoids With Zero](#)**Groups**[29. Groups](#)[30. Constructions With Groups](#)**Hyper Algebra**[31. Hypermonoids](#)[32. Hypergroups](#)[33. Hypersemirings and Hyperrings](#)[34. Quantales](#)**Near-Rings**[35. Near-Semirings](#)[36. Near-Rings](#)**Real Analysis**[37. Real Analysis in One Variable](#)[38. Real Analysis in Several Variables](#)**Measure Theory**[39. Measurable Spaces](#)[40. Measures and Integration](#)**Probability Theory**[40. Probability Theory](#)**Stochastic Analysis**[41. Stochastic Processes, Martingales, and Brownian Motion](#)[42. Itô Calculus](#)[43. Stochastic Differential Equations](#)**Differential Geometry**[44. Topological and Smooth Manifolds](#)**Schemes**[45. Schemes](#)

22.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part VIII

Globular Stuff

Chapter 23

The Globe Category

017E

Contents

23.A Other Chapters	445
23.2 Other Chapters	447

Appendices

23.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With
Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales,
and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

23.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
Category Theory	27. Monoids With Zero
11. Categories	28. Constructions With Monoids With Zero
12. Types of Morphisms in Categories	Groups
13. Adjunctions and the Yoneda Lemma	29. Groups
14. Constructions With Categories	30. Constructions With Groups
15. Profunctors	Hyper Algebra
16. Cartesian Closed Categories	31. Hypermonoids
17. Kan Extensions	32. Hypergroups
Bicategories	33. Hypersemirings and Hyperrings
18. Bicategories	34. Quantales
19. Internal Adjunctions	Near-Rings
Internal Category Theory	35. Near-Semirings
20. Internal Categories	36. Near-Rings
	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part IX

Cellular Stuff

Chapter 24

The Cell Category

017F

Contents

24.A Other Chapters	450
24.2 Other Chapters	452

Appendices

24.A Other Chapters

Sets

- 1. Sets
- 2. Constructions With Sets
- 3. Pointed Sets
- 4. Tensor Products of Pointed Sets
- 5. Relations
- 6. Spans
- 7. Posets

Indexed and Fibred Sets

- 7. Indexed Sets
- 8. Fibred Sets
- 9. Un/Straightening for Indexed and Fibred Sets

Category Theory

- 11. Categories
- 12. Types of Morphisms in Categories
- 13. Adjunctions and the Yoneda Lemma
- 14. Constructions With Categories
- 15. Profunctors
- 16. Cartesian Closed Categories
- 17. Kan Extensions

Bicategories

- 18. Bicategories
- 19. Internal Adjunctions

Internal Category Theory

- 20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With
Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales,
and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

24.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part X

Monoids

Chapter 25

Monoids

017G

Contents

25.A Other Chapters	455
25.2 Other Chapters	457

Appendices

25.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff[21. The Cycle Category](#)**Cubical Stuff**[22. The Cube Category](#)**Globular Stuff**[23. The Globe Category](#)**Cellular Stuff**[24. The Cell Category](#)**Monoids**[25. Monoids](#)[26. Constructions With Monoids](#)**Monoids With Zero**[27. Monoids With Zero](#)[28. Constructions With Monoids With Zero](#)**Groups**[29. Groups](#)[30. Constructions With Groups](#)**Hyper Algebra**[31. Hypermonoids](#)[32. Hypergroups](#)[33. Hypersemirings and Hyperrings](#)[34. Quantales](#)**Near-Rings**[35. Near-Semirings](#)[36. Near-Rings](#)**Real Analysis**[37. Real Analysis in One Variable](#)[38. Real Analysis in Several Variables](#)**Measure Theory**[39. Measurable Spaces](#)[40. Measures and Integration](#)**Probability Theory**[40. Probability Theory](#)**Stochastic Analysis**[41. Stochastic Processes, Martingales, and Brownian Motion](#)[42. Itô Calculus](#)[43. Stochastic Differential Equations](#)**Differential Geometry**[44. Topological and Smooth Manifolds](#)**Schemes**[45. Schemes](#)

25.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
Category Theory	27. Monoids With Zero
11. Categories	28. Constructions With Monoids With Zero
12. Types of Morphisms in Categories	Groups
13. Adjunctions and the Yoneda Lemma	29. Groups
14. Constructions With Categories	30. Constructions With Groups
15. Profunctors	Hyper Algebra
16. Cartesian Closed Categories	31. Hypermonoids
17. Kan Extensions	32. Hypergroups
Bicategories	33. Hypersemirings and Hyperrings
18. Bicategories	34. Quantales
19. Internal Adjunctions	Near-Rings
Internal Category Theory	35. Near-Semirings
20. Internal Categories	36. Near-Rings
	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 26

Constructions With Monoids

017H

Contents

26.A Other Chapters	459
26.2 Other Chapters	461

Appendices

26.A Other Chapters

Sets

- 1. Sets
- 2. Constructions With Sets
- 3. Pointed Sets
- 4. Tensor Products of Pointed Sets
- 5. Relations
- 6. Spans
- 7. Posets

Indexed and Fibred Sets

- 7. Indexed Sets
- 8. Fibred Sets
- 9. Un/Straightening for Indexed and Fibred Sets

Category Theory

- 11. Categories
- 12. Types of Morphisms in Categories
- 13. Adjunctions and the Yoneda Lemma
- 14. Constructions With Categories
- 15. Profunctors
- 16. Cartesian Closed Categories
- 17. Kan Extensions

Bicategories

- 18. Bicategories
- 19. Internal Adjunctions

Internal Category Theory

- 20. Internal Categories

Cyclic Stuff[21. The Cycle Category](#)**Cubical Stuff**[22. The Cube Category](#)**Globular Stuff**[23. The Globe Category](#)**Cellular Stuff**[24. The Cell Category](#)**Monoids**[25. Monoids](#)[26. Constructions With Monoids](#)**Monoids With Zero**[27. Monoids With Zero](#)[28. Constructions With Monoids With Zero](#)**Groups**[29. Groups](#)[30. Constructions With Groups](#)**Hyper Algebra**[31. Hypermonoids](#)[32. Hypergroups](#)[33. Hypersemirings and Hyperrings](#)[34. Quantales](#)**Near-Rings**[35. Near-Semirings](#)[36. Near-Rings](#)**Real Analysis**[37. Real Analysis in One Variable](#)[38. Real Analysis in Several Variables](#)**Measure Theory**[39. Measurable Spaces](#)[40. Measures and Integration](#)**Probability Theory**[40. Probability Theory](#)**Stochastic Analysis**[41. Stochastic Processes, Martingales, and Brownian Motion](#)[42. Itô Calculus](#)[43. Stochastic Differential Equations](#)**Differential Geometry**[44. Topological and Smooth Manifolds](#)**Schemes**[45. Schemes](#)

26.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
Category Theory	27. Monoids With Zero
11. Categories	28. Constructions With Monoids With Zero
12. Types of Morphisms in Categories	Groups
13. Adjunctions and the Yoneda Lemma	29. Groups
14. Constructions With Categories	30. Constructions With Groups
15. Profunctors	Hyper Algebra
16. Cartesian Closed Categories	31. Hypermonoids
17. Kan Extensions	32. Hypergroups
Bicategories	33. Hypersemirings and Hyperrings
18. Bicategories	34. Quantales
19. Internal Adjunctions	Near-Rings
Internal Category Theory	35. Near-Semirings
20. Internal Categories	36. Near-Rings
	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part XI

Monoids With Zero

Chapter 27

Monoids With Zero

017J

Contents

27.A Other Chapters	464
27.2 Other Chapters	466

Appendices

27.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With
Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales,
and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

27.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 28

Constructions With Monoids With Zero

017K

Contents

28.A Other Chapters	468
28.2 Other Chapters	470

Appendices

28.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets
5. Relations
6. Spans
7. Posets

Indexed and Fibred Sets

7. Indexed Sets
8. Fibred Sets

9. Un/Straightening for Indexed and Fibred Sets

Category Theory

11. Categories
12. Types of Morphisms in Categories
13. Adjunctions and the Yoneda Lemma
14. Constructions With Categories
15. Profunctors
16. Cartesian Closed Categories

17. Kan Extensions

Bicategories

18. Bicategories

19. Internal Adjunctions	32. Hypergroups
Internal Category Theory	33. Hypersemirings and Hyperrings
20. Internal Categories	34. Quantales
Cyclic Stuff	Near-Rings
21. The Cycle Category	35. Near-Semirings
Cubical Stuff	36. Near-Rings
22. The Cube Category	Real Analysis
Globular Stuff	37. Real Analysis in One Variable
23. The Globe Category	38. Real Analysis in Several Variables
Cellular Stuff	Measure Theory
24. The Cell Category	39. Measurable Spaces
Monoids	40. Measures and Integration
25. Monoids	Probability Theory
26. Constructions With Monoids	40. Probability Theory
Monoids With Zero	Stochastic Analysis
27. Monoids With Zero	41. Stochastic Processes, Martingales, and Brownian Motion
28. Constructions With Monoids With Zero	42. Itô Calculus
Groups	43. Stochastic Differential Equations
29. Groups	Differential Geometry
30. Constructions With Groups	44. Topological and Smooth Manifolds
Hyper Algebra	Schemes
31. Hypermonoids	45. Schemes

28.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	27. Monoids With Zero
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part XII

Groups

Chapter 29

Groups

017L

Contents

29.A Other Chapters	473
29.2 Other Chapters	475

Appendices

29.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff	34. Quantales
21. The Cycle Category	
Cubical Stuff	Near-Rings
22. The Cube Category	35. Near-Semirings
Globular Stuff	36. Near-Rings
23. The Globe Category	Real Analysis
Cellular Stuff	37. Real Analysis in One Variable
24. The Cell Category	38. Real Analysis in Several Variables
Monoids	Measure Theory
25. Monoids	39. Measurable Spaces
26. Constructions With Monoids	40. Measures and Integration
Monoids With Zero	Probability Theory
27. Monoids With Zero	40. Probability Theory
28. Constructions With Monoids With Zero	Stochastic Analysis
Groups	41. Stochastic Processes, Martingales, and Brownian Motion
29. Groups	42. Itô Calculus
30. Constructions With Groups	43. Stochastic Differential Equations
Hyper Algebra	Differential Geometry
31. Hypermonoids	44. Topological and Smooth Manifolds
32. Hypergroups	Schemes
33. Hypersemirings and Hyperrings	45. Schemes

29.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
Category Theory	27. Monoids With Zero
11. Categories	28. Constructions With Monoids With Zero
12. Types of Morphisms in Categories	Groups
13. Adjunctions and the Yoneda Lemma	29. Groups
14. Constructions With Categories	30. Constructions With Groups
15. Profunctors	Hyper Algebra
16. Cartesian Closed Categories	31. Hypermonoids
17. Kan Extensions	32. Hypergroups
Bicategories	33. Hypersemirings and Hyperrings
18. Bicategories	34. Quantales
19. Internal Adjunctions	Near-Rings
Internal Category Theory	35. Near-Semirings
20. Internal Categories	36. Near-Rings
	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 30

Constructions With Groups

017M

Contents

30.A Other Chapters	477
30.2 Other Chapters	479

Appendices

30.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With
Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales,
and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

30.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
Category Theory	27. Monoids With Zero
11. Categories	28. Constructions With Monoids With Zero
12. Types of Morphisms in Categories	Groups
13. Adjunctions and the Yoneda Lemma	29. Groups
14. Constructions With Categories	30. Constructions With Groups
15. Profunctors	Hyper Algebra
16. Cartesian Closed Categories	31. Hypermonoids
17. Kan Extensions	32. Hypergroups
Bicategories	33. Hypersemirings and Hyperrings
18. Bicategories	34. Quantales
19. Internal Adjunctions	Near-Rings
Internal Category Theory	35. Near-Semirings
20. Internal Categories	36. Near-Rings
	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part XIII

Hyper Algebra

Chapter 31

Hypermonoids

017N

Contents

31.A Other Chapters	482
31.2 Other Chapters	484

Appendices

31.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With
Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales,
and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

31.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 32

Hypergroups

017P

Contents

32.A Other Chapters	486
32.2 Other Chapters	488

Appendices

32.A Other Chapters

Sets

- 1. Sets
- 2. Constructions With Sets
- 3. Pointed Sets
- 4. Tensor Products of Pointed Sets
- 5. Relations
- 6. Spans
- 7. Posets

Indexed and Fibred Sets

- 7. Indexed Sets
- 8. Fibred Sets
- 9. Un/Straightening for Indexed and Fibred Sets

Category Theory

- 11. Categories
- 12. Types of Morphisms in Categories
- 13. Adjunctions and the Yoneda Lemma
- 14. Constructions With Categories
- 15. Profunctors
- 16. Cartesian Closed Categories
- 17. Kan Extensions

Bicategories

- 18. Bicategories
- 19. Internal Adjunctions

Internal Category Theory

- 20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With
Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales,
and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

32.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 33

Hypersemirings and Hyperrings

017Q

Contents

33.A Other Chapters	490
33.2 Other Chapters	492

Appendices

33.A Other Chapters

Sets

- 1. Sets
- 2. Constructions With Sets
- 3. Pointed Sets
- 4. Tensor Products of Pointed Sets
- 5. Relations
- 6. Spans
- 7. Posets

Indexed and Fibred Sets

- 7. Indexed Sets
- 8. Fibred Sets
- 9. Un/Straightening for Indexed and Fibred Sets

Category Theory

- 11. Categories
- 12. Types of Morphisms in Categories
- 13. Adjunctions and the Yoneda Lemma
- 14. Constructions With Categories
- 15. Profunctors
- 16. Cartesian Closed Categories
- 17. Kan Extensions

Bicategories

- 18. Bicategories
- 19. Internal Adjunctions

Internal Category Theory

- 20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With
Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales,
and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

33.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 34

Quantales

017R

Contents

34.A Other Chapters	494
34.2 Other Chapters	496

Appendices

34.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With
Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales**Near-Rings**

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales,
and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

34.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
Category Theory	27. Monoids With Zero
11. Categories	28. Constructions With Monoids With Zero
12. Types of Morphisms in Categories	Groups
13. Adjunctions and the Yoneda Lemma	29. Groups
14. Constructions With Categories	30. Constructions With Groups
15. Profunctors	Hyper Algebra
16. Cartesian Closed Categories	31. Hypermonoids
17. Kan Extensions	32. Hypergroups
Bicategories	33. Hypersemirings and Hyperrings
18. Bicategories	34. Quantales
19. Internal Adjunctions	Near-Rings
Internal Category Theory	35. Near-Semirings
20. Internal Categories	36. Near-Rings
	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part XIV

Near-Rings

Chapter 35

Near-Semirings

017S

Contents

35.A Other Chapters	499
35.2 Other Chapters	501

Appendices

35.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With
Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales,
and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

35.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 36

Near-Rings

017T

Contents

36.A Other Chapters	503
36.2 Other Chapters	505

Appendices

36.A Other Chapters

Sets

- 1. Sets
- 2. Constructions With Sets
- 3. Pointed Sets
- 4. Tensor Products of Pointed Sets
- 5. Relations
- 6. Spans
- 7. Posets

Indexed and Fibred Sets

- 7. Indexed Sets
- 8. Fibred Sets
- 9. Un/Straightening for Indexed and Fibred Sets

Category Theory

- 11. Categories
- 12. Types of Morphisms in Categories
- 13. Adjunctions and the Yoneda Lemma
- 14. Constructions With Categories
- 15. Profunctors
- 16. Cartesian Closed Categories
- 17. Kan Extensions

Bicategories

- 18. Bicategories
- 19. Internal Adjunctions

Internal Category Theory

- 20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With
Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales,
and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

36.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part XV

Real Analysis

Chapter 37

Real Analysis in One Variable

017U

Contents

37.A Other Chapters	508
37.2 Other Chapters	510

Appendices

37.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff

21. [The Cycle Category](#)

Cubical Stuff

22. [The Cube Category](#)

Globular Stuff

23. [The Globe Category](#)

Cellular Stuff

24. [The Cell Category](#)

Monoids

25. [Monoids](#)

26. [Constructions With Monoids](#)

Monoids With Zero

27. [Monoids With Zero](#)

28. [Constructions With Monoids With Zero](#)

Groups

29. [Groups](#)

30. [Constructions With Groups](#)

Hyper Algebra

31. [Hypermonoids](#)

32. [Hypergroups](#)

33. [Hypersemirings and Hyperrings](#)

34. Quantales**Near-Rings**

35. [Near-Semirings](#)

36. [Near-Rings](#)

Real Analysis

37. [Real Analysis in One Variable](#)

38. [Real Analysis in Several Variables](#)

Measure Theory

39. [Measurable Spaces](#)

40. [Measures and Integration](#)

Probability Theory

40. [Probability Theory](#)

Stochastic Analysis

41. [Stochastic Processes, Martingales, and Brownian Motion](#)

42. [Itô Calculus](#)

43. [Stochastic Differential Equations](#)

Differential Geometry

44. [Topological and Smooth Manifolds](#)

Schemes

45. [Schemes](#)

37.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	27. Monoids With Zero
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 38

Real Analysis in Several Variables

017V

Contents

38.A Other Chapters	512
38.2 Other Chapters	514

Appendices

38.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With
Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales,
and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

38.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part XVI

Measure Theory

Chapter 39

Measurable Spaces

017W

Contents

39.A Other Chapters	517
39.2 Other Chapters	519

Appendices

39.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With
Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales,
and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

39.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
Category Theory	27. Monoids With Zero
11. Categories	28. Constructions With Monoids With Zero
12. Types of Morphisms in Categories	Groups
13. Adjunctions and the Yoneda Lemma	29. Groups
14. Constructions With Categories	30. Constructions With Groups
15. Profunctors	Hyper Algebra
16. Cartesian Closed Categories	31. Hypermonoids
17. Kan Extensions	32. Hypergroups
Bicategories	33. Hypersemirings and Hyperrings
18. Bicategories	34. Quantales
19. Internal Adjunctions	Near-Rings
Internal Category Theory	35. Near-Semirings
20. Internal Categories	36. Near-Rings
	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 40

Measures and Integration

017X

Contents

40.A Other Chapters	521
40.2 Other Chapters	523

Appendices

40.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With
Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales,
and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

40.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
Category Theory	27. Monoids With Zero
11. Categories	28. Constructions With Monoids With Zero
12. Types of Morphisms in Categories	Groups
13. Adjunctions and the Yoneda Lemma	29. Groups
14. Constructions With Categories	30. Constructions With Groups
15. Profunctors	Hyper Algebra
16. Cartesian Closed Categories	31. Hypermonoids
17. Kan Extensions	32. Hypergroups
Bicategories	33. Hypersemirings and Hyperrings
18. Bicategories	34. Quantales
19. Internal Adjunctions	Near-Rings
Internal Category Theory	35. Near-Semirings
20. Internal Categories	36. Near-Rings
	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part XVII

Probability Theory

Chapter 41

Probability Theory

017Y

Contents

41.A Other Chapters	526
41.2 Other Chapters	528

Appendices

41.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With
Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales,
and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

41.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part XVIII

Stochastic Analysis

Chapter 42

Stochastic Processes, Martingales, and Brownian Motion

017Z

Contents

42.A Other Chapters	531
42.2 Other Chapters	533

Appendices

42.A Other Chapters

Sets

- 1. Sets
- 2. Constructions With Sets
- 3. Pointed Sets
- 4. Tensor Products of Pointed Sets
- 5. Relations
- 6. Spans
- 7. Posets

Indexed and Fibred Sets

- 7. Indexed Sets
- 8. Fibred Sets

Un/Straightening for Indexed and Fibred Sets

Category Theory

- 11. Categories
- 12. Types of Morphisms in Categories
- 13. Adjunctions and the Yoneda Lemma
- 14. Constructions With Categories
- 15. Profunctors
- 16. Cartesian Closed Categories

- 17. Kan Extensions

Bicategories

- 18. Bicategories

- 19. Internal Adjunctions
- 32. Hypergroups
- Internal Category Theory**
- 33. Hypersemirings and Hyperrings
- 20. Internal Categories
- 34. Quantales
- Cyclic Stuff**
- Near-Rings**
- 21. The Cycle Category
- 35. Near-Semirings
- Cubical Stuff**
- 36. Near-Rings
- 22. The Cube Category
- Real Analysis**
- Globular Stuff**
- 37. Real Analysis in One Variable
- 23. The Globe Category
- 38. Real Analysis in Several Variables
- Cellular Stuff**
- Measure Theory**
- 24. The Cell Category
- 39. Measurable Spaces
- Monoids**
- 40. Measures and Integration
- 25. Monoids
- Probability Theory**
- 26. Constructions With Monoids
- 40. Probability Theory
- Monoids With Zero**
- Stochastic Analysis**
- 27. Monoids With Zero
- 41. Stochastic Processes, Martingales,
and Brownian Motion
- 28. Constructions With Monoids With
Zero
- 42. Itô Calculus
- Groups**
- 43. Stochastic Differential Equations
- 29. Groups
- Differential Geometry**
- 30. Constructions With Groups
- 44. Topological and Smooth Manifolds
- Hyper Algebra**
- Schemes**
- 31. Hypermonoids
- 45. Schemes

42.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 43

Itô Calculus

0180

Contents

43.A Other Chapters	535
43.2 Other Chapters	537

Appendices

43.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff	34. Quantales
21. The Cycle Category	
Cubical Stuff	Near-Rings
22. The Cube Category	35. Near-Semirings
Globular Stuff	36. Near-Rings
23. The Globe Category	Real Analysis
Cellular Stuff	37. Real Analysis in One Variable
24. The Cell Category	38. Real Analysis in Several Variables
Monoids	Measure Theory
25. Monoids	39. Measurable Spaces
26. Constructions With Monoids	40. Measures and Integration
Monoids With Zero	Probability Theory
27. Monoids With Zero	40. Probability Theory
28. Constructions With Monoids With Zero	Stochastic Analysis
Groups	41. Stochastic Processes, Martingales, and Brownian Motion
29. Groups	42. Itô Calculus
30. Constructions With Groups	43. Stochastic Differential Equations
Hyper Algebra	Differential Geometry
31. Hypermonoids	44. Topological and Smooth Manifolds
32. Hypergroups	Schemes
33. Hypersemirings and Hyperrings	45. Schemes

43.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Chapter 44

Stochastic Differential Equations

0181

Contents

44.A Other Chapters	539
44.2 Other Chapters	541

Appendices

44.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff[21. The Cycle Category](#)**Cubical Stuff**[22. The Cube Category](#)**Globular Stuff**[23. The Globe Category](#)**Cellular Stuff**[24. The Cell Category](#)**Monoids**[25. Monoids](#)[26. Constructions With Monoids](#)**Monoids With Zero**[27. Monoids With Zero](#)[28. Constructions With Monoids With Zero](#)**Groups**[29. Groups](#)[30. Constructions With Groups](#)**Hyper Algebra**[31. Hypermonoids](#)[32. Hypergroups](#)[33. Hypersemirings and Hyperrings](#)[34. Quantales](#)**Near-Rings**[35. Near-Semirings](#)[36. Near-Rings](#)**Real Analysis**[37. Real Analysis in One Variable](#)[38. Real Analysis in Several Variables](#)**Measure Theory**[39. Measurable Spaces](#)[40. Measures and Integration](#)**Probability Theory**[40. Probability Theory](#)**Stochastic Analysis**[41. Stochastic Processes, Martingales, and Brownian Motion](#)[42. Itô Calculus](#)[43. Stochastic Differential Equations](#)**Differential Geometry**[44. Topological and Smooth Manifolds](#)**Schemes**[45. Schemes](#)

44.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
Category Theory	27. Monoids With Zero
11. Categories	28. Constructions With Monoids With Zero
12. Types of Morphisms in Categories	Groups
13. Adjunctions and the Yoneda Lemma	29. Groups
14. Constructions With Categories	30. Constructions With Groups
15. Profunctors	Hyper Algebra
16. Cartesian Closed Categories	31. Hypermonoids
17. Kan Extensions	32. Hypergroups
Bicategories	33. Hypersemirings and Hyperrings
18. Bicategories	34. Quantales
19. Internal Adjunctions	Near-Rings
Internal Category Theory	35. Near-Semirings
20. Internal Categories	36. Near-Rings
	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part XIX

Differential Geometry

Chapter 45

Topological and Smooth Manifolds

0182

Contents

45.A Other Chapters	544
45.2 Other Chapters	546

Appendices

45.A Other Chapters

Sets	Category Theory
1. Sets	11. Categories
2. Constructions With Sets	12. Types of Morphisms in Categories
3. Pointed Sets	13. Adjunctions and the Yoneda Lemma
4. Tensor Products of Pointed Sets	14. Constructions With Categories
5. Relations	15. Profunctors
6. Spans	16. Cartesian Closed Categories
7. Posets	17. Kan Extensions
Indexed and Fibred Sets	Bicategories
7. Indexed Sets	18. Bicategories
8. Fibred Sets	19. Internal Adjunctions
9. Un/Straightening for Indexed and Fibred Sets	Internal Category Theory
	20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With
Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

Measure Theory

39. Measurable Spaces

40. Measures and Integration

Probability Theory

40. Probability Theory

Stochastic Analysis

41. Stochastic Processes, Martingales,
and Brownian Motion

42. Itô Calculus

43. Stochastic Differential Equations

Differential Geometry

44. Topological and Smooth Manifolds

Schemes

45. Schemes

45.2 Other Chapters

Sets	Cyclic Stuff
1. Sets	21. The Cycle Category
2. Constructions With Sets	Cubical Stuff
3. Pointed Sets	22. The Cube Category
4. Tensor Products of Pointed Sets	Globular Stuff
5. Relations	23. The Globe Category
6. Spans	Cellular Stuff
7. Posets	24. The Cell Category
Indexed and Fibred Sets	Monoids
7. Indexed Sets	25. Monoids
8. Fibred Sets	26. Constructions With Monoids
9. Un/Straightening for Indexed and Fibred Sets	Monoids With Zero
27. Monoids With Zero	
Category Theory	28. Constructions With Monoids With Zero
11. Categories	Groups
12. Types of Morphisms in Categories	29. Groups
13. Adjunctions and the Yoneda Lemma	30. Constructions With Groups
14. Constructions With Categories	Hyper Algebra
15. Profunctors	31. Hypermonoids
16. Cartesian Closed Categories	32. Hypergroups
17. Kan Extensions	33. Hypersemirings and Hyperrings
Bicategories	34. Quantales
18. Bicategories	Near-Rings
19. Internal Adjunctions	35. Near-Semirings
Internal Category Theory	36. Near-Rings
20. Internal Categories	Real Analysis
	37. Real Analysis in One Variable

38. Real Analysis in Several Variables
Measure Theory
39. Measurable Spaces
40. Measures and Integration
Probability Theory
40. Probability Theory
- Stochastic Analysis
41. Stochastic Processes, Martingales,
and Brownian Motion
42. Itô Calculus
43. Stochastic Differential Equations
Differential Geometry
44. Topological and Smooth Manifolds
Schemes
45. Schemes

Part XX

Schemes

Chapter 46

Schemes

0183 46.1 Introduction

In this document we define schemes. A basic reference is [**EGA**].

46.2 Other Chapters

Sets	12. Types of Morphisms in Categories
1. Sets	13. Adjunctions and the Yoneda Lemma
2. Constructions With Sets	14. Constructions With Categories
3. Pointed Sets	15. Profunctors
4. Tensor Products of Pointed Sets	16. Cartesian Closed Categories
5. Relations	17. Kan Extensions
6. Spans	Bicategories
7. Posets	18. Bicategories
Indexed and Fibred Sets	19. Internal Adjunctions
7. Indexed Sets	Internal Category Theory
8. Fibred Sets	20. Internal Categories
9. Un/Straightening for Indexed and Fibred Sets	Cyclic Stuff
Category Theory	21. The Cycle Category
11. Categories	Cubical Stuff
	22. The Cube Category
	Globular Stuff
	23. The Globe Category

Cellular Stuff	36. Near-Rings
24. The Cell Category	
Monoids	Real Analysis
25. Monoids	37. Real Analysis in One Variable
26. Constructions With Monoids	38. Real Analysis in Several Variables
Monoids With Zero	Measure Theory
27. Monoids With Zero	39. Measurable Spaces
28. Constructions With Monoids With Zero	40. Measures and Integration
Groups	Probability Theory
29. Groups	40. Probability Theory
30. Constructions With Groups	Stochastic Analysis
Hyper Algebra	41. Stochastic Processes, Martingales, and Brownian Motion
31. Hypermonoids	42. Itô Calculus
32. Hypergroups	43. Stochastic Differential Equations
33. Hypersemirings and Hyperrings	Differential Geometry
34. Quantales	44. Topological and Smooth Manifolds
Near-Rings	Schemes
35. Near-Semirings	45. Schemes

Part XXI

Secret Part

Chapter 47

To Do List

0185 This chapter contains some material about relations and constructions with them. Notably, we discuss and explore:

Contents

47.1 Notes to Self.....	552
47.1.1 Things To Ask On MO/Zulip	552
47.1.2 Things To Explore/Add.....	553
47.1.3 Random Cool Papers.....	554
47.1.4 Omitted Proofs To Add	554
47.A Other Chapters	555
47.2 Other Chapters	556

47.1 Notes to Self

47.1.1 Things To Ask On MO/Zulip

0188 Remark 47.1.1.1. Here is a list of things to be asked on MO/Zulip.

1. What are
 - (a) Cartesian bicategories
 - (b) Double categories of relations (<https://arxiv.org/abs/2107.07621>)
 - (c) Categories of relations
 - (d) Allegories
 - (e) 1-Category equipped with relations (<https://ncatlab.org/nlab/show/1-category+equipped+with+relations>)

good for? What have these notions been developed for, why are they important, and what have they lead to?

47.1.2 Things To Explore/Add

018A **Remark 47.1.2.1.1.** Here is a list of things to be explored.

1. <https://mathoverflow.net/a/461814>
2. there's some cool stuff in <https://arxiv.org/abs/2312.00990>, e.g. on cofunctors.
3. internal adjunctions in Mod as in [JY21, Section 6.3]; see [JY21, Example 6.2.6].
4. write the “profunctors” equivalent of the relations chapter
5. change χ_B notation throughout the notes
6. maybe note that skew monoidal structures on $\mathbf{Rel}(A, B)$ satisfy coherence trivially since the 2-morphisms are inclusions
7. reconsider notation $\text{FreeAlg}_{\mathcal{P}}$ in [Relations](#)
8. Constructions With Sets: Isbell duality for powersets
9. Categories: comma category notation as in <https://mathoverflow.net/questions/455630>
10. Universal property of the bicategory of spans, <https://ncatlab.org/nlab/show/span>
11. Codensity monad $\text{Ran}_J(J)$ of a relation (What about $\text{Rift}_J(J)$?)
12. Relative comonads in [Rel](#).
13. Write proper sections on straightening for lax functors from sets to Rel or Span (displayed sets) when I study the corresponding notions for categories
14. Write about cospans.
15. CoCartesian fibration classifying $\text{Fun}(F, G)$, <https://mathoverflow.net/questions/457533/cocartesian-fibration-classifying-\mathrm{fun}\ f-g>
16. Constructions With Sets: functoriality of limits/colimits, like functoriality of pullbacks
17. <https://ncatlab.org/nlab/show/adjoint+lifting+theorem>
18. <https://ncatlab.org/nlab/show/Gabriel%20%93Ulmer+duality>

47.1.3 Random Cool Papers

018C **Remark 47.1.3.1.1.** Here are some random cool papers that appeared on arXiv and that I want to check eventually.

1. A Derived Geometric Approach to Propagation of Solution Singularities for Non-linear PDEs I: Foundations
2. The Fundamental Theorem of Calculus point-free, with applications to exponentials and logarithms

47.1.4 Omitted Proofs To Add

Не так благоврона истина, как зловредна ее видимость.

Даниил Данковский

Truth does not do as much good in the world as the appearance of truth does evil.

Daniil Dankovsky

There's a very large number of omitted proofs throughout these notes; here I list some of the ones that I really want to add to the notes at some point.

018E **Remark 47.1.4.1.1.** Here is a list of omitted proofs that I really want to eventually write up or add a reference to.

- ?? of ??

Appendices

47.A Other Chapters

Sets

1. Sets
2. Constructions With Sets
3. Pointed Sets
4. Tensor Products of Pointed Sets
5. Relations
6. Spans
7. Posets

Indexed and Fibred Sets

7. Indexed Sets
8. Fibred Sets
9. Un/Straightening for Indexed and Fibred Sets

Category Theory

11. Categories
12. Types of Morphisms in Categories
13. Adjunctions and the Yoneda Lemma
14. Constructions With Categories
15. Profunctors
16. Cartesian Closed Categories
17. Kan Extensions

Bicategories

18. Bicategories
19. Internal Adjunctions

Internal Category Theory

20. Internal Categories

Cyclic Stuff

21. The Cycle Category

Cubical Stuff

22. The Cube Category

Globular Stuff

23. The Globe Category

Cellular Stuff

24. The Cell Category

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero

27. Monoids With Zero

28. Constructions With Monoids With Zero

Groups

29. Groups

30. Constructions With Groups

Hyper Algebra

31. Hypermonoids

32. Hypergroups

33. Hypersemirings and Hyperrings

34. Quantales

Near-Rings

35. Near-Semirings

36. Near-Rings

Real Analysis

37. Real Analysis in One Variable

38. Real Analysis in Several Variables

41. Stochastic Processes, Martingales, and Brownian Motion

Measure Theory

39. Measurable Spaces

42. Itô Calculus

40. Measures and Integration

43. Stochastic Differential Equations

Probability Theory

40. Probability Theory

Differential Geometry

44. Topological and Smooth Manifolds

Stochastic Analysis

Schemes

45. Schemes

47.2 Other Chapters

Sets

15. Profunctors

1. Sets

16. Cartesian Closed Categories

2. Constructions With Sets

17. Kan Extensions

3. Pointed Sets

Bicategories

4. Tensor Products of Pointed Sets

18. Bicategories

5. Relations

19. Internal Adjunctions

6. Spans

Internal Category Theory

7. Posets

20. Internal Categories

Indexed and Fibred Sets

Cyclic Stuff

7. Indexed Sets

21. The Cycle Category

8. Fibred Sets

Cubical Stuff

9. Un/Straightening for Indexed and Fibred Sets

22. The Cube Category

Category Theory

Globular Stuff

11. Categories

23. The Globe Category

12. Types of Morphisms in Categories

Cellular Stuff

13. Adjunctions and the Yoneda Lemma

24. The Cell Category

14. Constructions With Categories

Monoids

25. Monoids

26. Constructions With Monoids

Monoids With Zero	37. Real Analysis in One Variable
27. Monoids With Zero	38. Real Analysis in Several Variables
28. Constructions With Monoids With Zero	Measure Theory
Groups	39. Measurable Spaces
29. Groups	40. Measures and Integration
30. Constructions With Groups	Probability Theory
Hyper Algebra	40. Probability Theory
31. Hypermonoids	Stochastic Analysis
32. Hypergroups	41. Stochastic Processes, Martingales, and Brownian Motion
33. Hypersemirings and Hyperrings	42. Itô Calculus
34. Quantales	43. Stochastic Differential Equations
Near-Rings	Differential Geometry
35. Near-Semirings	44. Topological and Smooth Manifolds
36. Near-Rings	Schemes
Real Analysis	45. Schemes

Bibliography

- [MO 119454] user30818. *Category and the Axiom of Choice*. MathOverflow. URL: <https://mathoverflow.net/q/119454> (cit. on p. 336).
- [MO 350788] Peter LeFanu Lumsdaine. *Epimorphisms of relations*. MathOverflow. URL: <https://mathoverflow.net/q/455260> (cit. on p. 148).
- [MO 382264] Neil Strickland. *Proof that a Cartesian category is monoidal*. MathOverflow. URL: <https://mathoverflow.net/q/382264> (cit. on p. 19).
- [MO 64365] Giorgio Mossa. *Natural Transformations as Categorical Homotopies*. MathOverflow. URL: <https://mathoverflow.net/q/64365> (cit. on p. 347).
- [MSE 267469] Zhen Lin. *Show that the powerset partial order is a cartesian closed category*. Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/267469> (cit. on p. 47).
- [MSE 350788] Qiaochu Yuan. *Mono's and Epi's in the category Rel?* Mathematics Stack Exchange. URL: <https://math.stackexchange.com/q/350788> (cit. on pp. 145, 147, 309, 359, 388).
- [Bor94] Francis Borceux. *Handbook of Categorical Algebra I*. Vol. 50. Encyclopedia of Mathematics and its Applications. Basic Category Theory. Cambridge University Press, Cambridge, 1994, pp. xvi+345. ISBN: 0-521-44178-1 (cit. on p. 345).
- [BS10] John C. Baez and Michael Shulman. “Lectures on n -Categories and Cohomology”. In: *Towards higher categories*. Vol. 152. IMA Vol. Math. Appl. Springer, New York, 2010, pp. 1–68. DOI: [10.1007/978-1-4419-1524-5_1](https://doi.org/10.1007/978-1-4419-1524-5_1). URL: https://doi.org/10.1007/978-1-4419-1524-5_1 (cit. on pp. 2, 3).
- [Cie97] Krzysztof Ciesielski. *Set Theory for the Working Mathematician*. Vol. 39. London Mathematical Society Student Texts. Cambridge University Press, Cambridge, 1997, pp. xii+236. ISBN: 0-521-59441-3; 0-521-59465-0. DOI: [10.1017/CBO9781139173131](https://doi.org/10.1017/CBO9781139173131). URL: <https://doi.org/10.1017/CBO9781139173131> (cit. on p. 42).
- [GGN15] David Gepner, Moritz Groth, and Thomas Nikolaus. “Universality of Multiplicative Infinite Loop Space Machines”. In: *Algebr. Geom. Topol.*

- 15.6 (2015), pp. 3107–3153. ISSN: 1472-2747. DOI: [10.2140/agt.2015.15.3107](https://doi.org/10.2140/agt.2015.15.3107). URL: <https://doi.org/10.2140/agt.2015.15.3107> (cit. on p. 112).
- [JY21] Niles Johnson and Donald Yau. *2-Dimensional Categories*. Oxford University Press, Oxford, 2021, pp. xix+615. ISBN: 978-0-19-887138-5; 978-0-19-887137-8. DOI: [10.1093/oso/9780198871378.001.0001](https://doi.org/10.1093/oso/9780198871378.001.0001). URL: <https://doi.org/10.1093/oso/9780198871378.001.0001> (cit. on pp. 398, 413, 415, 417, 419, 553).
- [Nie04] Susan Niefield. “Change of Base for Relational Variable Sets”. In: *Theory Appl. Categ.* 12 (2004), No. 7, 248–261. ISSN: 1201-561X (cit. on p. 304).
- [nLa24a] nLab Authors. *Displayed Category*. <https://ncatlab.org/nlab/show/displayed+category>. Oct. 2024 (cit. on p. 305).
- [nLa24b] nLab Authors. *Groupoid*. <https://ncatlab.org/nlab/show/groupoid>. Oct. 2024 (cit. on p. 337).
- [nLab23] The nLab Authors. *Skeleton*. 2024. URL: <https://ncatlab.org/nlab/show/skeleton> (cit. on p. 316).
- [Pro24a] Proof Wiki Contributors. *Bijection Between $R \times (S \times T)$ and $(R \times S) \times T$ — Proof Wiki*. 2024. URL: [https://proofwiki.org/wiki/Bijection_between_R_x_\(S_x_T\)_and_\(R_x_S\)_x_T](https://proofwiki.org/wiki/Bijection_between_R_x_(S_x_T)_and_(R_x_S)_x_T) (cit. on p. 19).
- [Pro24b] Proof Wiki Contributors. *Bijection Between $S \times T$ and $T \times S$ — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Bijection_between_S_x_T_and_T_x_S (cit. on p. 19).
- [Pro24c] Proof Wiki Contributors. *Cartesian Product Distributes Over Set Difference — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Cartesian_Product_Distributes_over_Set_Difference (cit. on p. 19).
- [Pro24d] Proof Wiki Contributors. *Cartesian Product Distributes Over Symmetric Difference — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Cartesian_Product_Distributes_over_Symmetric_Difference (cit. on p. 19).
- [Pro24e] Proof Wiki Contributors. *Cartesian Product Distributes Over Union — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Cartesian_Product_Distributes_over_Union (cit. on p. 19).
- [Pro24f] Proof Wiki Contributors. *Cartesian Product Is Empty Iff Factor Is Empty — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Cartesian_Product_is_Empty_iff_Factor_is_Empty (cit. on p. 19).

- [Pro24g] Proof Wiki Contributors. *Cartesian Product of Intersections — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Cartesian_Product_of_Intersections (cit. on p. 19).
- [Pro24h] Proof Wiki Contributors. *Characteristic Function of Intersection — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Characteristic_Function_of_Intersection (cit. on p. 47).
- [Pro24i] Proof Wiki Contributors. *Characteristic Function of Set Difference — ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Characteristic_Function_of_Set_Difference (cit. on p. 50).
- [Pro24j] Proof Wiki Contributors. *Characteristic Function of Symmetric Difference — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Characteristic_Function_of_Symmetric_Difference (cit. on p. 55).
- [Pro24k] Proof Wiki Contributors. *Characteristic Function of Union — ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Characteristic_Function_of_Union (cit. on p. 44).
- [Pro24l] Proof Wiki Contributors. *Complement of Complement — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Complement_of_Complement (cit. on p. 51).
- [Pro24m] Proof Wiki Contributors. *Condition For Mapping from Quotient Set To Be A Surjection — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Condition_for_Mapping_from_Quotient_Set_to_be_Surjection (cit. on p. 181).
- [Pro24n] Proof Wiki Contributors. *Condition For Mapping From Quotient Set To Be An Injection — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Condition_for_Mapping_from_Quotient_Set_to_be_Injection (cit. on p. 181).
- [Pro24o] Proof Wiki Contributors. *Condition For Mapping From Quotient Set To Be Well-Defined — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Condition_for_Mapping_from_Quotient_Set_to_be_Well-Defined (cit. on p. 181).
- [Pro24p] Proof Wiki Contributors. *De Morgan's Laws (Set Theory) — Proof Wiki*. 2024. URL: [https://proofwiki.org/wiki/De_Morgan%5C27s_Laws_\(Set_Theory\)](https://proofwiki.org/wiki/De_Morgan%5C27s_Laws_(Set_Theory)) (cit. on pp. 50, 51).
- [Pro24q] Proof Wiki Contributors. *De Morgan's Laws (Set Theory)/Set Difference/Difference with Union — ProofWiki*. 2024. URL: [https://proofwiki.org/wiki/De_Morgan%5C27s_Laws_\(Set_Theory\)/Set_Difference/Difference_with_Union](https://proofwiki.org/wiki/De_Morgan%5C27s_Laws_(Set_Theory)/Set_Difference/Difference_with_Union) (cit. on p. 50).
- [Pro24r] Proof Wiki Contributors. *Equivalence of Definitions of Symmetric Difference — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Equivalence_of_Definitions_of_Symmetric_Difference

- [Pro24s] Proof Wiki Contributors. *Equivalence_of_Definitions_of_Symmetric_Difference* — Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Equivalence_of_Definitions_of_Symmetric_Difference (cit. on p. 55).
- [Pro24t] Proof Wiki Contributors. *Intersection_Distributes_Over_Symmetric_Difference* — Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Intersection_Distributes_Over_Symmetric_Difference (cit. on p. 55).
- [Pro24u] Proof Wiki Contributors. *Intersection_Is_Associative* — Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Intersection_Is_Associative (cit. on p. 47).
- [Pro24v] Proof Wiki Contributors. *Intersection_Is_Commutative* — Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Intersection_Is_Commutative (cit. on p. 47).
- [Pro24w] Proof Wiki Contributors. *Intersection_With_Empty_Set* — Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Intersection_With_Empty_Set (cit. on p. 47).
- [Pro24x] Proof Wiki Contributors. *Intersection_With_Subset_Is_Subset* — Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Intersection_With_Subset_Is_Subset (cit. on p. 47).
- [Pro24y] Proof Wiki Contributors. *Mapping_From_Quotient_Set_When_Defined_Is_Unique* — Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Mapping_From_Quotient_Set_When_Defined_Is_Unique (cit. on p. 181).
- [Pro24z] Proof Wiki Contributors. *Quotient_Mapping_Is_Coequalizer* — Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Quotient_Mapping_Is_Coequalizer (cit. on p. 39).
- [Pro24aa] Proof Wiki Contributors. *Set_Difference_as_Intersection_With_Complement* — Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Set_Difference_as_Intersection_With_Complement (cit. on p. 50).
- [Pro24ab] Proof Wiki Contributors. *Set_Difference_as_Symmetric_Difference_With_Intersection* — Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Set_Difference_as_Symmetric_Difference_With_Intersection (cit. on p. 50).
- [Pro24ac] Proof Wiki Contributors. *Set_Difference_Is_Right_Distributive_Over_Union* — Proof Wiki. 2024. URL: https://proofwiki.org/wiki/Set_Difference_Is_Right_Distributive_Over_Union (cit. on p. 50).

- [Pro24ad] ProofWiki Contributors. *Set Difference Over Subset — ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Set_Difference_over_Subset (cit. on p. 50).
- [Pro24ae] ProofWiki Contributors. *Set Difference With Empty Set Is Self — ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Set_Difference_with_Empty_Set_is_Self (cit. on p. 50).
- [Pro24af] ProofWiki Contributors. *Set Difference With Self Is Empty Set — ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Set_Difference_with_Self_is_Empty_Set (cit. on p. 50).
- [Pro24ag] ProofWiki Contributors. *Set Difference With Set Difference Is Union of Set Difference With Intersection — ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Set_Difference_with_Set_Difference_is_Union_of_Set_Difference_with_Intersection (cit. on p. 50).
- [Pro24ah] ProofWiki Contributors. *Set Difference With Subset Is Superset of Set Difference — ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Set_Difference_with_Subset_is_Superset_of_Set_Difference (cit. on p. 50).
- [Pro24ai] ProofWiki Contributors. *Set Difference With Union — ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Set_Difference_with_Union (cit. on p. 50).
- [Pro24aj] ProofWiki Contributors. *Set Intersection Distributes Over Union — ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Intersection_Distributes_over_Union (cit. on p. 47).
- [Pro24ak] ProofWiki Contributors. *Set Intersection Is Idempotent — ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Set_Intersection_is_Idempotent (cit. on p. 47).
- [Pro24al] ProofWiki Contributors. *Set Intersection Preserves Subsets — ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Set_Intersection_Preserves_Subsets (cit. on p. 47).
- [Pro24am] ProofWiki Contributors. *Set Union Is Idempotent — ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Set_Union_is_Idempotent (cit. on p. 44).
- [Pro24an] ProofWiki Contributors. *Set Union Preserves Subsets — ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Set_Union_Preserves_Subsets (cit. on p. 44).
- [Pro24ao] ProofWiki Contributors. *Symmetric Difference Is Associative — ProofWiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_is_Associative (cit. on p. 55).

- [Pro24ap] ProofWiki Contributors. *Symmetric Difference Is Commutative — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_is_Commutative (cit. on p. 55).
- [Pro24aq] ProofWiki Contributors. *Symmetric Difference of Complements — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_of_Complements (cit. on p. 55).
- [Pro24ar] ProofWiki Contributors. *Symmetric Difference on Power Set Forms Abelian Group — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_on_Power_Set_forms_Abelian_Group (cit. on p. 55).
- [Pro24as] ProofWiki Contributors. *Symmetric Difference With Complement — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Complement (cit. on p. 55).
- [Pro24at] ProofWiki Contributors. *Symmetric Difference With Empty Set — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Empty_Set (cit. on p. 55).
- [Pro24au] ProofWiki Contributors. *Symmetric Difference With Intersection Forms Ring — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Intersection_forms_Ring (cit. on p. 56).
- [Pro24av] ProofWiki Contributors. *Symmetric Difference With Self Is Empty Set — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Self_is_Empty_Set (cit. on p. 55).
- [Pro24aw] ProofWiki Contributors. *Symmetric Difference With Union Does Not Form Ring — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Union_does_not_form_Ring (cit. on p. 55).
- [Pro24ax] ProofWiki Contributors. *Symmetric Difference With Universe — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Symmetric_Difference_with_Universe (cit. on p. 55).
- [Pro24ay] ProofWiki Contributors. *Union as Symmetric Difference With Intersection — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Union_as_Symmetric_Difference_with_Intersection (cit. on p. 44).
- [Pro24az] ProofWiki Contributors. *Union Distributes Over Intersection — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Union_Distributes_over_Intersection (cit. on p. 44).
- [Pro24ba] ProofWiki Contributors. *Union Is Associative — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Union_is_Associative (cit. on p. 44).

- [Pro24bb] Proof Wiki Contributors. *Union Is Commutative — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Union_is_Commutative (cit. on p. 44).
- [Pro24bc] Proof Wiki Contributors. *Union of Symmetric Differences — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Union_of_Symmetric_Differences (cit. on p. 55).
- [Pro24bd] Proof Wiki Contributors. *Union With Empty Set — Proof Wiki*. 2024. URL: https://proofwiki.org/wiki/Union_with_Empty_Set (cit. on p. 44).
- [PS19] Maximilien Péroux and Brooke Shipley. “Coalgebras in Symmetric Monoidal Categories of Spectra”. In: *Homology Homotopy Appl.* 21.1 (2019), pp. 1–18. ISSN: 1532-0073. DOI: [10.4310/HHA.2019.v21.n1.a1](https://doi.org/10.4310/HHA.2019.v21.n1.a1). URL: <https://doi.org/10.4310/HHA.2019.v21.n1.a1> (cit. on p. 112).
- [Rie17] Emily Riehl. *Category Theory in Context*. Vol. 10. Aurora: Dover Modern Math Originals. Courier Dover Publications, 2017, pp. xviii+240. ISBN: 978-0486809038. URL: <http://www.math.jhu.edu/~eriehl/context.pdf> (cit. on pp. 330, 337).
- [SS86] Stephen Schanuel and Ross Street. “The Free Adjunction”. In: *Cahiers Topologie Géom. Différentielle Catég.* 27.1 (1986), pp. 81–83. ISSN: 0008-0004 (cit. on p. 409).

Index of Notation

A

- $[a]$, 178
- (A, B) , 42
- Adj , 409
- $A \coprod B$, 30
- $A \coprod_C B$, 32
- $A \triangle B$, 52
- $A \times B$, 14
- $A \times_C B$, 20
- $A \times_{f,C,g} B$, 20
- $A \cup B$, 42

- $\text{Coim}(f)$, 180
- $\text{Coll}(\mathfrak{p})$, 380
- $\text{Coll}(R)$, 168
- $\text{Coll}(R)$, 168
- $\coprod_{i \in I} A_i$, 28
- $\text{Core}(C)$, 327
- C^\simeq , 327

D

- \mathcal{D}^C , 349
- $\text{dom}(\mathfrak{p})$, 378
- $\text{dom}(R)$, 156

B

- $\beta \circ \alpha$, 341
- $\beta \star \alpha$, 344

C

- $C(A, B)$, 311
- Cats , 352
- Cats_2 , 353
- $[C, \mathcal{D}]$, 349
- $\chi_{(-)}$, 57
- χ_{-2}^{-1} , 56
- χ_U , 56
- χ_x , 56
- $\chi_X(-_1, -_2)$, 56
- $\chi_X(-, U)$, 56
- $\chi_X(-, x)$, 56
- $\chi_X(U, -)$, 56
- $\chi_X(x, -)$, 56
- $\text{CoEq}(f, g)$, 38

E

- \emptyset_{cat} , 313
- \emptyset , 41
- $\text{Eq}(f, g)$, 24
- $\text{eq}(f, g)$, 24

F

- \widehat{F}_* , 380
- \widehat{F}^* , 379
- FibSets , 274
- $\text{FibSets}(K)$, 273
- f^{-1} , 70
- $f^{-1}(V)$, 70
- \mathbb{F}_1 , 84
- $f_!$, 73
- $f_{!,\text{cp}}(U)$, 75
- $f_{!,\text{im}}(U)$, 74
- $f_!(U)$, 73
- f^* , 65, 317

f_x^* , 272	
$f(U)$, 66	
$\text{Fun}(C, \mathcal{D})$, 349	
G	P
$\text{Gr}(A)$, 151	$\mathfrak{p}: C \dashrightarrow \mathcal{D}$, 377
Grpd , 354	$\phi^{-1}(x)$, 272
Grpd_2 , 354	ϕ_x , 272
H	$\Pi_f(X)$, 278
$\text{Hom}_{\text{Sets}_*}^{\otimes, \text{L}}(X \times Y, Z)$, 96	$\Pi_\phi(X)$, 266
$\text{Hom}_{\text{Sets}_*}^{\otimes, \text{R}}(X \times Y, Z)$, 97	$\pi_0(C)$, 320
$\text{Hom}_{\text{Sets}_*}^\otimes(X \times Y, Z)$, 98	\mathcal{P}^{-1} , 61, 202
I	$\prod_{i \in I} A_i$, 12
id_C , 331	Prof , 384
id_F , 340	$\mathcal{P}_!$, 61, 202
$\bigcup_{i \in I} R_i$, 159	\mathcal{P}_* , 61, 202
$\bigcap_{X \in \mathcal{F}} X$, 44	pt , 84
ISets , 260	pt , 313
$\text{ISets}(K)$, 260	$\mathcal{P}(X)$, 60
K	Q
$\text{K}_0(C)$, 324	$\mathfrak{q} \diamond \mathfrak{p}$, 384
L	$\mathfrak{q} \diamond \mathfrak{p}$, 379
Λ , 222	
M	R
$\text{Mor}(C)$, 311	$\text{range}(\mathfrak{p})$, 379
N	$\text{range}(R)$, 156
\ltimes , 314	R^\dagger , 162
$\text{Nat}(F, G)$, 340	Rel , 126
O	Rel , 131
ω^\dagger , 418	$\text{Rel}(A, B)$, 120
	$\text{Rel}(A, B)$, 120
	Rel^{dbl} , 132
	$\text{Rel}^{\text{eq}}(A, B)$, 177
	$\text{Rel}^{\text{eq}}(A, B)$, 177
	$\text{Rel}^{\text{refl}}(A, A)$, 170
	$\text{Rel}^{\text{refl}}(A, A)$, 170
	$\text{Rel}^{\text{symm}}(A, A)$, 172
	$\text{Rel}^{\text{symm}}(A, A)$, 172
	$\text{Rel}^{\text{trans}}(A)$, 174
	$\text{Rel}^{\text{trans}}(A)$, 174
	R^{eq} , 177
	$R \cap S$, 158
	R^{-1} , 192
	R_{-1} , 187

$R^{-1}(V)$, 192	$\text{St}_{A,B}$, 241
$R_{-1}(V)$, 187	St_K , 299
R^{refl} , 170	S^0 , 84
$R_!$, 196	
$R_!(U)$, 196	T
R^* , 182	$\{\text{t}, \text{f}\}$, 3
R^{symm} , 172	$\prod_{i \in I} R_i$, 162
$R \times S$, 160	$\{\text{true}, \text{false}\}$, 3
R^{trans} , 175	
$R(U)$, 182	
$R \cup S$, 157	
S	
S_{ab} , 241	U^c , 51
$S \diamond R$, 163	$\bigcup_{i \in I} A_i$, 42
$\text{Sets}_{/K}$, 273	$\bigcup_{i \in I} R_i$, 158
Sets_* , 85	Un_K , 301
$\text{Hom}_{\text{Sets}_*}(X, Y)$, 88	$\text{UnNat}(F, G)$, 339
$\Sigma_f(X)$, 277	
$\Sigma_\phi(X)$, 265	X
\sim_{cotriv} , 124	$\{X\}$, 41
\sim_{id} , 56	X_{disc} , 321
\sim_R^{refl} , 170	X_{indisc} , 323
\sim_R^{symm} , 172	$X \cap Y$, 45
\sim_R^{eq} , 177	$X \setminus Y$, 48
\sim_R^{trans} , 175	X / \sim_R , 178
\sim_{triv} , 124	$X \otimes_{\mathbb{F}_1} Y$, 108
Span , 227	X^+ , 88
$\text{Span}(A, B)$, 225	$X \vee Y$, 89
Span^{dbl} , 229	$X \wedge Y$, 108
	$\{X, Y\}$, 41

Index of Set Theory

B

bilinear morphism
of pointed sets, 97
bilinear morphism of pointed sets
left, 96
right, 96
binary intersection, 45

C

Cartesian product, 12, 14
category of relations, 126
 associator of, 128
 internal Hom of, 129
 left unit of, 128
 monoidal product of, 127
 monoidal unit of, 128
 right unit of, 129
 symmetry of, 129
characteristic embedding, 57
characteristic function
 of a set, 56
 of an element, 56
characteristic relation, 56
coequaliser of sets, 38
coimage, 180
complement of a set, 51

D

difference of sets, 48
disjoint union, 30
 of a family of sets, 28

E

empty set, 41
equaliser of sets, 24
equivalence class, 178
equivalence relation
 kernel, 177

F

fibre coproduct of sets, 32
fibre product of sets, 20
fibred set, 272
 category of, 273, 274
 change of base, 275
 dependent product, 278
 dependent sum, 277
 fibre over an element, 272
 internal **Hom** of, 282
 morphism of, 272
 transport map on fibres, 272
field with one element
module over, 84
module over, morphism of, 84
function
 associated direct image function,
 65
 associated direct image with
 compact support function, 73
 associated direct image with
 compact support function,
 complement part, 75
 associated direct image with
 compact support function,

image part, 74
 associated inverse image function, 70
 coimage of, 180
 graph of, 151
 inverse of, 154
 kernel of, 177

I

indexed set, 259
 category of, 260
 category of K -indexed sets, 260
 change of indexing, 264
 coequaliser, 264
 coproduct, 263
 dependent product, 266
 dependent sum, 265
 equaliser, 262, 263
 internal **Hom** of, 267
 morphism of, 259
 product, 261, 262
 pullback, 262, 263
 pushout, 263
 indicator function, *see* characteristic function
 intersection of a family of sets, 44

L

left tensor product of pointed sets
 skew associator, 101
 skew left unit, 102
 skew right unit, 102

M

(-1) -category, 2
 (-2) -category, 2
 module
 underlying pointed set of, 84

O

ordered pairing, 42

P

pairing of two sets, 41
 pointed function, *see* pointed set, morphism of
 pointed set, 84
 category of, 85
 coequaliser of, 87
 coproduct of, 87
 cotensor by a set, 99
 equaliser of, 87
 \mathbb{F}_1 , 84
 free, 88
 left tensor product, 100
 morphism of, 84
 of morphisms of pointed sets, 88
 product of, 86
 pullback of, 87
 pushout of, 87
 right tensor product, 104
 tensor by a set, 98
 trivial, 84
 underlying a module, 84
 underlying a semimodule, 84
 wedge sum of, 89
 0-sphere, 84
 pointed sets
 left-skew monoidal category
 structure on, 103
 right-skew monoidal category
 structure on, 107
 poset
 of (-1) -categories, 3
 of truth values, 3
 powerset, 60
 product of a family of sets, 12
 product of sets, 14
 pullback of sets, 20
 pushout of sets, 32

Q

quotient
 by an equivalence relation, 178

R

relation, 120
associated direct image function, 182
associated direct image with compact support function, 196
associated strong inverse image function, 187
associated to a span, 245
associated weak inverse image function, 192
category of, 126
closed symmetric monoidal category of, 130
collage of, 168
composition of, 163
corepresentable, 156
cotrivial, 124
domain of, 156
double category of, 132
equivalence closure of, 177
equivalence relation, 176
equivalence, set of, 177
functional, 125
intersection of, 158
intersection of a family of, 159
inverse of, 162
on powersets associated to a relation, 203
partial equivalence relation, 176
poset of, 120, 172, 174, 177
product of, 160
product of a family of, 162
range of, 156
reflexive, 170
reflexive closure of, 170
reflexive, poset of, 170
reflexive, set of, 170
representable, 155
set of, 120
symmetric, 172
symmetric closure of, 172

symmetric, set of, 172
total, 126
transitive, 174
transitive closure of, 175
transitive, set of, 174
trivial, 124
2-category of, 131
union of, 157
union of a family of, 158
right tensor product of pointed sets
skew associator, 105
skew left unit, 106
skew right unit, 106

S

semimodule
underlying pointed set of, 84
set of bilinear morphisms of pointed sets, 98
left, 96
right, 97
singleton set, 41
smash product
of pointed sets, 108
span, 222
associated relation, 245
bicategory of, 227
category of, 225
composition of, 239
corepresentable, 239
functional, 223
horizontal composition of morphisms of, 240
morphism of, 223
representable, 239
total, 224
straightening
of a fibred set, 299
of a span, 241
symmetric difference of sets, 52

U

union, 42
of a family of sets, 42
unstraightening
for spans, 242
of an indexed set, 301

Z
0-category, 6
0-groupoid, 6