

# Constructions With Sets

December 3, 2023

000D This chapter contains some material relating to constructions with sets. Notably, it contains:

1. Explicit descriptions of the major types of co/limits in Sets, including in particular pushouts and coequalisers (see [Definitions 2.3.1](#) and [2.4.1](#) and [Remarks 2.3.2](#) and [2.4.2](#));
2. A discussion of powersets as decategorifications of categories of presheaves ([Remarks 4.1.2](#) and [4.2.2](#));
3. A lengthy discussion of the adjoint triple

$$f_* \dashv f^{-1} \dashv f_! : \mathcal{P}(A) \xrightarrow{\cong} \mathcal{P}(B)$$

of functors (morphisms of posets) between  $\mathcal{P}(A)$  and  $\mathcal{P}(B)$  induced by a map of sets  $f : A \rightarrow B$ , along with a discussion of the properties of  $f_*$ ,  $f^{-1}$ , and  $f_!$ .

## Contents

<b>1</b>	<b>Limits of Sets .....</b>	<b>2</b>
1.1	Products of Families of Sets .....	2
1.2	Binary Products of Sets .....	2
1.3	Pullbacks .....	6
1.4	Equalisers .....	8
<b>2</b>	<b>Colimits of Sets .....</b>	<b>10</b>
2.1	Coproducts of Families of Sets .....	11
2.2	Binary Coproducts .....	11
2.3	Pushouts .....	13
2.4	Coequalisers .....	16

<b>3</b>	<b>Operations With Sets</b>	<b>20</b>
3.1	The Empty Set	20
3.2	Singleton Sets	20
3.3	Pairings of Sets	20
3.4	Unions of Families	21
3.5	Binary Unions	21
3.6	Intersections of Families	24
3.7	Binary Intersections	24
3.8	Differences	28
3.9	Complements	31
3.10	Symmetric Differences	33
3.11	Ordered Pairs	38
<b>4</b>	<b>Powersets</b>	<b>38</b>
4.1	Characteristic Functions	38
4.2	Powersets	43
4.3	Direct Images	49
4.4	Inverse Images	55
4.5	Direct Images With Compact Support	60
<b>A</b>	<b>Other Chapters</b>	<b>70</b>

## 000E 1 Limits of Sets

### 000F 1.1 Products of Families of Sets

Let  $\{A_i\}_{i \in I}$  be a family of sets.

#### 000G DEFINITION 1.1.1 ► THE PRODUCT OF A FAMILY OF SETS

The **product**<sup>1</sup> of  $\{A_i\}_{i \in I}$  is the set  $\prod_{i \in I} A_i$  defined by

$$\prod_{i \in I} A_i \stackrel{\text{def}}{=} \left\{ f \in \text{Sets} \left( I, \bigcup_{i \in I} A_i \right) \mid \begin{array}{l} \text{for each } i \in I, \text{ we} \\ \text{have } f(i) \in A_i \end{array} \right\}.$$

<sup>1</sup>Further Terminology: Also called the **Cartesian product** of  $\{A_i\}_{i \in I}$ .

### 000H 1.2 Binary Products of Sets

Let  $A$  and  $B$  be sets.

000J

**DEFINITION 1.2.1 ► PRODUCTS OF SETS**

The **product**<sup>1</sup> of  $A$  and  $B$  is the set  $A \times B$  defined by

$$\begin{aligned} A \times B &\stackrel{\text{def}}{=} \prod_{z \in \{A, B\}} z \\ &\stackrel{\text{def}}{=} \{f \in \text{Sets}(\{0, 1\}, A \cup B) \mid \text{we have } f(0) \in A \text{ and } f(1) \in B\} \\ &\cong \{\{\{a\}, \{a, b\}\} \in \mathcal{P}(\mathcal{P}(A \cup B)) \mid \text{we have } a \in A \text{ and } b \in B\}. \end{aligned}$$

<sup>1</sup>*Further Terminology:* Also called the **Cartesian product of  $A$  and  $B$**  or the **binary Cartesian product of  $A$  and  $B$** , for emphasis.

This can also be thought of as the  $(\mathbb{E}_{-1}, \mathbb{E}_{-1})$ -**tensor product of  $A$  and  $B$** .

000K

**PROPOSITION 1.2.2 ► PROPERTIES OF PRODUCTS OF SETS**

Let  $A, B, C$ , and  $X$  be sets.

000L

1. *Functoriality.* The assignments  $A, B, (A, B) \mapsto A \times B$  define functors

$$\begin{aligned} A \times -_2 &: \text{Sets} \rightarrow \text{Sets}, \\ -_1 \times B &: \text{Sets} \rightarrow \text{Sets}, \\ -_1 \times -_2 &: \text{Sets} \times \text{Sets} \rightarrow \text{Sets}, \end{aligned}$$

where  $-_1 \times -_2$  is the functor where

- *Action on Objects.* For each  $(A, B) \in \text{Obj}(\text{Sets} \times \text{Sets})$ , we have

$$[-_1 \times -_2](A, B) \stackrel{\text{def}}{=} A \times B;$$

- *Action on Morphisms.* For each  $(A, B), (X, Y) \in \text{Obj}(\text{Sets})$ , the action on Hom-sets

$$\times_{(A, B), (X, Y)} : \text{Sets}(A, X) \times \text{Sets}(B, Y) \rightarrow \text{Sets}(A \times B, X \times Y)$$

of  $\times$  at  $((A, B), (X, Y))$  is defined by sending  $(f, g)$  to the function

$$f \times g : A \times B \rightarrow X \times Y$$

defined by

$$[f \times g](a, b) \stackrel{\text{def}}{=} (f(a), g(b))$$

for each  $(a, b) \in A \times B$ ;

and where  $A \times -$  and  $- \times B$  are the partial functors of  $-_1 \times -_2$  at  $A, B \in \text{Obj}(\text{Sets})$ .

000M

2. *Adjointness.* We have adjunctions

$$(A \times - \dashv \text{Sets}(A, -)): \text{Sets} \begin{array}{c} \xrightarrow{A \times -} \\ \perp \\ \xleftarrow{\text{Sets}(A, -)} \end{array} \text{Sets},$$

$$(- \times B \dashv \text{Sets}(B, -)): \text{Sets} \begin{array}{c} \xrightarrow{- \times B} \\ \perp \\ \xleftarrow{\text{Sets}(B, -)} \end{array} \text{Sets},$$

witnessed by bijections

$$\begin{aligned} \text{Sets}(A \times B, C) &\cong \text{Sets}(A, \text{Sets}(B, C)), \\ \text{Sets}(A \times B, C) &\cong \text{Sets}(B, \text{Sets}(A, C)), \end{aligned}$$

natural in  $A, B, C \in \text{Obj}(\text{Sets})$ .

000N

3. *Associativity.* We have an isomorphism of sets

$$(A \times B) \times C \cong A \times (B \times C),$$

natural in  $A, B, C \in \text{Obj}(\text{Sets})$ .

000P

4. *Unitality.* We have isomorphisms of sets

$$\begin{aligned} \text{pt} \times A &\cong A, \\ A \times \text{pt} &\cong A, \end{aligned}$$

natural in  $A \in \text{Obj}(\text{Sets})$ .

000Q

5. *Commutativity.* We have an isomorphism of sets

$$A \times B \cong B \times A,$$

natural in  $A, B \in \text{Obj}(\text{Sets})$ .

000R

6. *Annihilation With the Empty Set.* We have isomorphisms of sets

$$A \times \emptyset \cong \emptyset,$$

$$\emptyset \times A \cong \emptyset,$$

natural in  $A \in \text{Obj}(\text{Sets})$ .

000S

7. *Distributivity Over Unions.* We have isomorphisms of sets

$$A \times (B \cup C) = (A \times B) \cup (A \times C),$$

$$(A \cup B) \times C = (A \times C) \cup (B \times C),$$

natural in  $A, B, C \in \text{Obj}(\text{Sets})$ .

000T

8. *Distributivity Over Intersections.* We have isomorphisms of sets

$$A \times (B \cap C) = (A \times B) \cap (A \times C),$$

$$(A \cap B) \times C = (A \times C) \cap (B \times C),$$

natural in  $A, B, C \in \text{Obj}(\text{Sets})$ .

000U

9. *Distributivity Over Differences.* We have isomorphisms of sets

$$A \times (B \setminus C) = (A \times B) \setminus (A \times C),$$

$$(A \setminus B) \times C = (A \times C) \setminus (B \times C),$$

natural in  $A, B, C \in \text{Obj}(\text{Sets})$ .

000V

10. *Distributivity Over Symmetric Differences.* We have isomorphisms of sets

$$A \times (B \triangle C) = (A \times B) \triangle (A \times C),$$

$$(A \triangle B) \times C = (A \times C) \triangle (B \times C),$$

natural in  $A, B, C \in \text{Obj}(\text{Sets})$ .

000W

11. *Symmetric Monoidality.* The triple  $(\text{Sets}, \times, \text{pt})$  is a symmetric monoidal category.

000X

12. *Symmetric Bimonoidality.* The quintuple  $(\text{Sets}, \coprod, \emptyset, \times, \text{pt})$  is a symmetric bimonoidal category.

**PROOF 1.2.3 ► PROOF OF PROPOSITION 1.2.2**

Item 1: Functoriality

Omitted.

Item 2: Adjointness

Omitted.

Item 3: Associativity

Clear.

Item 4: Unitality

Clear.

Item 5: Commutativity

Clear.

Item 6: Annihilation With the Empty Set

Clear.

Item 7: Distributivity Over Unions

Omitted.

Item 8: Distributivity Over Intersections

Omitted.

Item 9: Distributivity Over Differences

Omitted.

Item 10: Distributivity Over Symmetric Differences

Omitted.

Item 11: Symmetric Monoidality

Omitted.

Item 12: Symmetric Bimonoidality

Omitted.

**000Y 1.3 Pullbacks**

Let  $A$ ,  $B$ , and  $C$  be sets and let  $f: A \rightarrow C$  and  $g: B \rightarrow C$  be functions.

000Z

**DEFINITION 1.3.1 ► PULLBACKS OF SETS**

The **pullback of  $A$  and  $B$  over  $C$  along  $f$  and  $g$** <sup>1</sup> is the set  $A \times_C B$  defined by

$$A \times_C B \stackrel{\text{def}}{=} \{(a, b) \in A \times B \mid f(a) = g(b)\}.$$

<sup>1</sup>*Further Terminology:* Also called the **fibre product of  $A$  and  $B$  over  $C$  along  $f$  and  $g$** .

0010

**EXAMPLE 1.3.2 ► EXAMPLES OF PULLBACKS OF SETS**

Here are some examples of pullbacks of sets.

0011

1. *Unions via Intersections.* Let  $A, B \subset X$ . We have a bijection of sets

$$A \cap B \cong A \times_{A \cup B} B.$$

0012

**PROPOSITION 1.3.3 ► PROPERTIES OF PULLBACKS OF SETS**

Let  $A, B, C$ , and  $X$  be sets.

0013

1. *Associativity.* We have an isomorphism of sets

$$(A \times_X B) \times_X C \cong A \times_X (B \times_X C),$$

natural in  $A, B, C, X \in \text{Obj}(\text{Sets})$ .

0014

2. *Unitality.* We have isomorphisms of sets

$$X \times_X A \cong A,$$

$$A \times_X X \cong A,$$

natural in  $A, X \in \text{Obj}(\text{Sets})$ .

0015

3. *Commutativity.* We have an isomorphism of sets

$$A \times_X B \cong B \times_X A,$$

natural in  $A, B, X \in \text{Obj}(\text{Sets})$ .

0016

4. *Annihilation With the Empty Set.* We have isomorphisms of sets

$$A \times_X \emptyset \cong \emptyset,$$

$$\emptyset \times_X A \cong \emptyset,$$

natural in  $A, X \in \text{Obj}(\text{Sets})$ .

0017

5. *Symmetric Monoidality.* The triple  $(\text{Sets}, \times_X, X)$  is a symmetric monoidal category.

**PROOF 1.3.4 ► PROOF OF PROPOSITION 1.3.3**

Item 1: Associativity

Clear.

Item 2: Unitality

Clear.


Item 3: Commutativity

Clear.

Item 4: Annihilation With the Empty Set

Clear.

Item 5: Symmetric Monoidality

Omitted. 

**0018 1.4 Equalisers**

Let  $A$  and  $B$  be sets and let  $f, g: A \rightrightarrows B$  be functions.

0019

**DEFINITION 1.4.1 ► EQUALISERS OF SETS**

The **equaliser of  $f$  and  $g$**  is the set  $\text{Eq}(f, g)$  defined by

$$\text{Eq}(f, g) \stackrel{\text{def}}{=} \{a \in A \mid f(a) = g(a)\}.$$

001A

**PROPOSITION 1.4.2 ► PROPERTIES OF EQUALISERS OF SETS**

Let  $A$ ,  $B$ , and  $C$  be sets.

001B

1. *Associativity.* We have an isomorphism of sets<sup>1</sup>

$$\underbrace{\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h))}_{=\text{Eq}(f \circ \text{eq}(g, h), h \circ \text{eq}(g, h))} \cong \text{Eq}(f, g, h) \cong \underbrace{\text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g))}_{=\text{Eq}(g \circ \text{eq}(f, g), h \circ \text{eq}(f, g))},$$



where  $\text{Eq}(f, g, h)$  is the limit of the diagram

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{-g} \\ \xrightarrow{h} \end{array} B$$

in Sets.

001C

2. *Unitality.* We have an isomorphism of sets

$$\text{Eq}(f, f) \cong A.$$

001D

3. *Commutativity.* We have an isomorphism of sets

$$\text{Eq}(f, g) \cong \text{Eq}(g, f).$$

001E

4. *Interaction With Composition.* Let

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} B \begin{array}{c} \xrightarrow{h} \\ \xrightarrow{k} \end{array} C$$

be functions. We have an inclusion of sets

$$\text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g)) \subset \text{Eq}(h \circ f, k \circ g),$$

where  $\text{Eq}(h \circ f \circ \text{eq}(f, g), k \circ g \circ \text{eq}(f, g))$  is the equaliser of the composition

$$\text{Eq}(f, g) \xrightarrow{\text{eq}(f, g)} A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} B \begin{array}{c} \xrightarrow{h} \\ \xrightarrow{k} \end{array} C.$$

---

<sup>1</sup>That is: the following constructions give the same result:

- (a) Take the equaliser of  $(f, g, h)$ , i.e. the limit of the diagram

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{-g} \\ \xrightarrow{h} \end{array} B$$

in Sets.

- (b) First take the equaliser of  $f$  and  $g$ , forming a diagram

$$\text{Eq}(f, g) \xrightarrow{\text{eq}(f, g)} A \begin{matrix} \xrightarrow{f} \\ \xrightarrow{g} \end{matrix} B$$

and then take the equaliser of the composition

$$\text{Eq}(f, g) \xrightarrow{\text{eq}(f, g)} A \begin{matrix} \xrightarrow{f} \\ \xrightarrow{h} \end{matrix} B,$$

obtaining a subset

$$\text{Eq}(f \circ \text{eq}(f, g), h \circ \text{eq}(f, g)) = \text{Eq}(g \circ \text{eq}(f, g), h \circ \text{eq}(f, g))$$

of  $\text{Eq}(f, g)$ .

- (c) First take the equaliser of  $g$  and  $h$ , forming a diagram

$$\text{Eq}(g, h) \xrightarrow{\text{eq}(g, h)} A \begin{matrix} \xrightarrow{g} \\ \xrightarrow{h} \end{matrix} B$$

and then take the equaliser of the composition

$$\text{Eq}(g, h) \xrightarrow{\text{eq}(g, h)} A \begin{matrix} \xrightarrow{f} \\ \xrightarrow{g} \end{matrix} B,$$

obtaining a subset

$$\text{Eq}(f \circ \text{eq}(g, h), g \circ \text{eq}(g, h)) = \text{Eq}(f \circ \text{eq}(g, h), h \circ \text{eq}(g, h))$$

of  $\text{Eq}(g, h)$ .

#### PROOF 1.4.3 ► PROOF OF PROPOSITION 1.4.2

Item 1: Associativity

Clear.

Item 2: Unitality

Clear.

Item 3: Commutativity

Clear.

Item 4: Interaction With Composition

Omitted.



## 001G 2.1 Coproducts of Families of Sets

Let  $\{A_i\}_{i \in I}$  be a family of sets.

## 001H DEFINITION 2.1.1 ► DISJOINT UNIONS OF FAMILIES

The **disjoint union of the family**  $\{A_i\}_{i \in I}$  is the set  $\coprod_{i \in I} A_i$  defined by

$$\coprod_{i \in I} A_i \stackrel{\text{def}}{=} \bigcup_{i \in I} \left\{ (x, i) \in \left( \bigcup_{i \in I} A_i \right) \times I \mid x \in A_i \right\}.$$

## 001J 2.2 Binary Coproducts

Let  $A$  and  $B$  be sets.

## 001K DEFINITION 2.2.1 ► COPRODUCTS OF SETS

The **coproduct**<sup>1</sup> of  $A$  and  $B$  is the set  $A \coprod B$  defined by

$$\begin{aligned} A \coprod B &\stackrel{\text{def}}{=} \coprod_{z \in \{A, B\}} z \\ &\stackrel{\text{def}}{=} \{(a, 0) \mid a \in A\} \cup \{(b, 1) \mid b \in B\}. \end{aligned}$$

<sup>1</sup>Further Terminology: Also called the **disjoint union of  $A$  and  $B$** , or the **binary disjoint union of  $A$  and  $B$** , for emphasis.

## 001L PROPOSITION 2.2.2 ► PROPERTIES OF COPRODUCTS OF SETS

Let  $A, B, C$ , and  $X$  be sets.

- 001M 1. *Functoriality.* The assignment  $A, B, (A, B) \mapsto A \coprod B$  defines functors

$$\begin{aligned} A \coprod -_2 &: \text{Sets} \rightarrow \text{Sets}, \\ -_1 \coprod B &: \text{Sets} \rightarrow \text{Sets}, \\ -_1 \coprod -_2 &: \text{Sets} \times \text{Sets} \rightarrow \text{Sets}, \end{aligned}$$

where  $-_1 \coprod -_2$  is the functor where

- *Action on Objects.* For each  $(A, B) \in \text{Obj}(\text{Sets} \times \text{Sets})$ , we have

$$[-_1 \coprod -_2](A, B) \stackrel{\text{def}}{=} A \coprod B;$$

- *Action on Morphisms.* For each  $(A, B), (X, Y) \in \text{Obj}(\text{Sets})$ , the action on Hom-sets

$$\coprod_{(A,B),(X,Y)} : \text{Sets}(A, X) \times \text{Sets}(B, Y) \rightarrow \text{Sets}(A \coprod B, X \coprod Y)$$

of  $\coprod$  at  $((A, B), (X, Y))$  is defined by sending  $(f, g)$  to the function

$$f \coprod g : A \coprod B \rightarrow X \coprod Y$$

defined by

$$[f \coprod g](x) \stackrel{\text{def}}{=} \begin{cases} f(x) & \text{if } x \in A, \\ g(x) & \text{if } x \in B \end{cases}$$

for each  $x \in A \coprod B$ ;

and where  $A \coprod -$  and  $- \coprod B$  are the partial functors of  $-_1 \coprod -_2$  at  $A, B \in \text{Obj}(\text{Sets})$ .

001N

2. *Associativity.* We have an isomorphism of sets

$$(A \coprod B) \coprod C \cong A \coprod (B \coprod C),$$

natural in  $A, B, C \in \text{Obj}(\text{Sets})$ .

001P

3. *Unitality.* We have isomorphisms of sets

$$A \coprod \emptyset \cong A,$$

$$\emptyset \coprod A \cong A,$$

natural in  $A \in \text{Obj}(\text{Sets})$ .

001Q

4. *Commutativity.* We have an isomorphism of sets

$$A \coprod B \cong B \coprod A,$$

natural in  $A, B \in \text{Obj}(\text{Sets})$ .

001R

5. *Symmetric Monoidality.* The triple  $(\text{Sets}, \coprod, \emptyset)$  is a symmetric monoidal category.

## PROOF 2.2.3 ► PROOF OF PROPOSITION 2.2.2

Item 1: Functoriality

Omitted.

Item 2: Associativity

Clear.

Item 3: Unitality

Clear.

Item 4: Commutativity

Clear.

Item 5: Symmetric Monoidality

Omitted. 

## 001S 2.3 Pushouts

Let  $A, B$ , and  $C$  be sets and let  $f: C \rightarrow A$  and  $g: C \rightarrow B$  be functions.

## 001T DEFINITION 2.3.1 ► PUSHOUTS OF SETS

The **pushout of  $A$  and  $B$  over  $C$  along  $f$  and  $g$** <sup>1</sup> is the set  $A \amalg_C B$  defined by

$$A \amalg_C B \stackrel{\text{def}}{=} A \amalg B / \sim_C,$$

where  $\sim_C$  is the equivalence relation on  $A \amalg B$  generated by  $f(c) \sim_C g(c)$ .

<sup>1</sup>Further Terminology: Also called the **fibre coproduct of  $A$  and  $B$  over  $C$  along  $f$  and  $g$** .

## 001U REMARK 2.3.2 ► UNWINDING DEFINITION 2.3.1

In detail, the relation  $\sim$  of Definition 2.3.1 is given by declaring  $a \sim b$  iff one of the following conditions is satisfied:

- We have  $a, b \in A$  and  $a = b$ ;
- We have  $a, b \in B$  and  $a = b$ ;
- There exist  $x_1, \dots, x_n \in A \amalg B$  such that  $a \sim' x_1 \sim' \dots \sim' x_n \sim' b$ , where we declare  $x \sim' y$  if one of the following conditions is satisfied:

1. There exists  $c \in C$  such that  $x = f(c)$  and  $y = g(c)$ .
2. There exists  $c \in C$  such that  $x = g(c)$  and  $y = f(c)$ .

That is: we require the following condition to be satisfied:

- (★) There exist  $x_1, \dots, x_n \in A \coprod B$  satisfying the following conditions:
1. There exists  $c_0 \in C$  satisfying one of the following conditions:
    - (a) We have  $a = f(c_0)$  and  $x_1 = g(c_0)$ .
    - (b) We have  $a = g(c_0)$  and  $x_1 = f(c_0)$ .
  2. For each  $1 \leq i \leq n - 1$ , there exists  $c_i \in C$  satisfying one of the following conditions:
    - (a) We have  $x_i = f(c_i)$  and  $x_{i+1} = g(c_i)$ .
    - (b) We have  $x_i = g(c_i)$  and  $x_{i+1} = f(c_i)$ .
  3. There exists  $c_n \in C$  satisfying one of the following conditions:
    - (a) We have  $x_n = f(c_n)$  and  $b = g(c_n)$ .
    - (b) We have  $x_n = g(c_n)$  and  $b = f(c_n)$ .

001V

#### EXAMPLE 2.3.3 ► EXAMPLES OF PUSHOUTS OF SETS

Here are some examples of pushouts of sets.

001W

1. *Wedge Sums of Pointed Sets.* The wedge sum of two pointed sets of ?? is an example of a pushout of sets.

001X

2. *Intersections via Unions.* Let  $A, B \subset X$ . We have a bijection of sets

$$A \cup B \cong A \coprod_{A \cap B} B.$$

001Y

#### PROPOSITION 2.3.4 ► PROPERTIES OF PUSHOUTS OF SETS

Let  $A, B, C$ , and  $X$  be sets.

001Z

1. *Associativity.* We have an isomorphism of sets

$$(A \coprod_X B) \coprod_X C \cong A \coprod_X (B \coprod_X C),$$

natural in  $A, B, C, X \in \text{Obj}(\text{Sets})$ .

0020

2. *Unitality.* We have isomorphisms of sets

$$\begin{aligned}\emptyset \coprod_X A &\cong A, \\ A \coprod_X \emptyset &\cong A,\end{aligned}$$

natural in  $A, X \in \text{Obj}(\text{Sets})$ .

0021

3. *Commutativity.* We have an isomorphism of sets

$$A \coprod_X B \cong B \coprod_X A,$$

natural in  $A, B, X \in \text{Obj}(\text{Sets})$ .

0022

4. *Annihilation With the Empty Set.* We have isomorphisms of sets

$$\begin{aligned}A \coprod_X \emptyset &\cong \emptyset, \\ \emptyset \coprod_X A &\cong \emptyset,\end{aligned}$$

natural in  $A, X \in \text{Obj}(\text{Sets})$ .

0023

5. *Symmetric Monoidality.* The triple  $(\text{Sets}, \coprod_X, \emptyset)$  is a symmetric monoidal category.

#### PROOF 2.3.5 ► PROOF OF PROPOSITION 2.3.4

Item 1: Associativity

Clear.

Item 2: Unitality

Clear.

Item 3: Commutativity

Clear.

Item 4: Annihilation With the Empty Set

Clear.

Item 5: Symmetric Monoidality

Omitted.



**0024 2.4 Coequalisers**

Let  $A$  and  $B$  be sets and let  $f, g: A \rightrightarrows B$  be functions.



0025

## DEFINITION 2.4.1 ► COEQUALISERS OF SETS

The **coequaliser of  $f$  and  $g$**  is the set  $\text{CoEq}(f, g)$  defined by

$$\text{CoEq}(f, g) \stackrel{\text{def}}{=} B/\sim,$$

where  $\sim$  is the equivalence relation on  $B$  generated by  $f(a) \sim g(a)$ .

0026

## REMARK 2.4.2 ► UNWINDING DEFINITION 2.4.1

In detail, the relation  $\sim$  of **Definition 2.4.1** is given by declaring  $a \sim b$  iff one of the following conditions is satisfied:

- We have  $a = b$ ;
- There exist  $x_1, \dots, x_n \in B$  such that  $a \sim' x_1 \sim' \dots \sim' x_n \sim' b$ , where we declare  $x \sim' y$  if one of the following conditions is satisfied:
  1. There exists  $z \in A$  such that  $x = f(z)$  and  $y = g(z)$ .
  2. There exists  $z \in A$  such that  $x = g(z)$  and  $y = f(z)$ .

That is: we require the following condition to be satisfied:

- (★) There exist  $x_1, \dots, x_n \in B$  satisfying the following conditions:
1. There exists  $z_0 \in A$  satisfying one of the following conditions:
    - (a) We have  $a = f(z_0)$  and  $x_1 = g(z_0)$ .
    - (b) We have  $a = g(z_0)$  and  $x_1 = f(z_0)$ .
  2. For each  $1 \leq i \leq n - 1$ , there exists  $z_i \in A$  satisfying one of the following conditions:
    - (a) We have  $x_i = f(z_i)$  and  $x_{i+1} = g(z_i)$ .
    - (b) We have  $x_i = g(z_i)$  and  $x_{i+1} = f(z_i)$ .
  3. There exists  $z_n \in A$  satisfying one of the following conditions:
    - (a) We have  $x_n = f(z_n)$  and  $b = g(z_n)$ .
    - (b) We have  $x_n = g(z_n)$  and  $b = f(z_n)$ .

0027 **EXAMPLE 2.4.3 ► EXAMPLES OF COEQUALISERS OF SETS**

Here are some examples of coequalisers of sets.

- 0028 1. *Quotients by Equivalence Relations.* Let  $R$  be an equivalence relation on a set  $X$ . We have a bijection of sets

$$X/\sim_R \cong \text{CoEq}\left(R \hookrightarrow X \times X \begin{array}{c} \xrightarrow{\text{pr}_1} \\ \xrightarrow{\text{pr}_2} \end{array} X\right).$$

0029 **PROPOSITION 2.4.4 ► PROPERTIES OF COEQUALISERS OF SETS**

Let  $A$ ,  $B$ , and  $C$  be sets.

- 002A 1. *Associativity.* We have an isomorphism of sets<sup>1</sup>

$$\underbrace{\text{CoEq}(\text{coeq}(f, g) \circ f, \text{coeq}(f, g) \circ h)}_{=\text{CoEq}(\text{coeq}(f, g) \circ g, \text{coeq}(f, g) \circ h)} \cong \text{CoEq}(f, g, h) \cong \underbrace{\text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ g)}_{=\text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ h)},$$

where  $\text{CoEq}(f, g, h)$  is the colimit of the diagram

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{-g-} \\ \xrightarrow{h} \end{array} B$$

in Sets.

- 002B 2. *Unitality.* We have an isomorphism of sets

$$\text{CoEq}(f, f) \cong B.$$

- 002C 3. *Commutativity.* We have an isomorphism of sets

$$\text{CoEq}(f, g) \cong \text{CoEq}(g, f).$$

- 002D 4. *Interaction With Composition.* Let

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} B \begin{array}{c} \xrightarrow{h} \\ \xrightarrow{k} \end{array} C$$

be functions. We have a surjection

$$\text{CoEq}(h \circ f, k \circ g) \twoheadrightarrow \text{CoEq}(\text{coeq}(h, k) \circ h \circ f, \text{coeq}(h, k) \circ k \circ g)$$

exhibiting  $\text{CoEq}(\text{coeq}(h, k) \circ h \circ f, \text{coeq}(h, k) \circ k \circ g)$  as a quotient of  $\text{CoEq}(h \circ f, k \circ g)$  by the relation generated by declaring  $h(y) \sim k(y)$  for each  $y \in B$ .

<sup>1</sup>That is: the following constructions give the same result:

- (a) Take the coequaliser of  $(f, g, h)$ , i.e. the colimit of the diagram

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \\ \xrightarrow{h} \end{array} B$$

in Sets.

- (b) First take the coequaliser of  $f$  and  $g$ , forming a diagram

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} B \xrightarrow{\text{coeq}(f, g)} \text{CoEq}(f, g)$$

and then take the coequaliser of the composition

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \\ \xrightarrow{h} \end{array} B \xrightarrow{\text{coeq}(f, g)} \text{CoEq}(f, g),$$

obtaining a quotient

$$\text{CoEq}(\text{coeq}(f, g) \circ f, \text{coeq}(f, g) \circ h) = \text{CoEq}(\text{coeq}(f, g) \circ g, \text{coeq}(f, g) \circ h)$$

of  $\text{CoEq}(f, g)$

- (c) First take the coequaliser of  $g$  and  $h$ , forming a diagram

$$A \begin{array}{c} \xrightarrow{g} \\ \xrightarrow{h} \end{array} B \xrightarrow{\text{coeq}(g, h)} \text{CoEq}(g, h)$$

and then take the coequaliser of the composition

$$A \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \\ \xrightarrow{h} \end{array} B \xrightarrow{\text{coeq}(g, h)} \text{CoEq}(g, h),$$

obtaining a quotient

$$\text{CoEq}(\text{coeq}(g, h) \circ f, \text{coeq}(g, h) \circ g) = \text{CoEq}(\text{coeq}(g, h) \circ h, \text{coeq}(g, h) \circ g)$$

of  $\text{CoEq}(g, h)$ .

#### PROOF 2.4.5 ► PROOF OF PROPOSITION 2.4.4

Item 1: Associativity

Omitted.

Item 2: Unitality

Clear.

Item 3: Commutativity

Clear.

Item 4: Interaction With Composition

Omitted.



## 002E 3 Operations With Sets

### 002F 3.1 The Empty Set

#### 002G DEFINITION 3.1.1 ► THE EMPTY SET

The **empty set** is the set  $\emptyset$  defined by

$$\emptyset \stackrel{\text{def}}{=} \{x \in X \mid x \neq x\},$$

where  $A$  is the set in the set existence axiom, ?? of ??.

### 002H 3.2 Singleton Sets

Let  $X$  be a set.

#### 002J DEFINITION 3.2.1 ► SINGLETON SETS

The **singleton set containing**  $X$  is the set  $\{X\}$  defined by

$$\{X\} \stackrel{\text{def}}{=} \{X, X\},$$

where  $\{X, X\}$  is the pairing of  $X$  with itself (Definition 3.3.1).

### 002K 3.3 Pairings of Sets

Let  $X$  and  $Y$  be sets.

002L

**DEFINITION 3.3.1 ► PAIRINGS OF SETS**

The **pairing of  $X$  and  $Y$**  is the set  $\{X, Y\}$  defined by

$$\{X, Y\} \stackrel{\text{def}}{=} \{x \in A \mid x = X \text{ or } x = Y\},$$

where  $A$  is the set in the axiom of pairing, ?? of ??.

002M

**3.4 Unions of Families**

Let  $\{A_i\}_{i \in I}$  be a family of sets.

002N

**DEFINITION 3.4.1 ► UNIONS OF FAMILIES**

The **union of the family  $\{A_i\}_{i \in I}$**  is the set  $\bigcup_{i \in I} A_i$  defined by

$$\bigcup_{i \in I} A_i \stackrel{\text{def}}{=} \{x \in F \mid \text{there exists some } i \in I \text{ such that } x \in A_i\},$$

where  $F$  is the set in the axiom of union, ?? of ??.

002P

**3.5 Binary Unions**

Let  $A$  and  $B$  be sets.

002Q

**DEFINITION 3.5.1 ► BINARY UNIONS**

The **union<sup>1</sup> of  $A$  and  $B$**  is the set  $A \cup B$  defined by

$$A \cup B \stackrel{\text{def}}{=} \bigcup_{z \in \{A, B\}} z.$$

<sup>1</sup>Further Terminology: Also called the **binary union of  $A$  and  $B$** , for emphasis.

002R

**PROPOSITION 3.5.2 ► PROPERTIES OF BINARY UNIONS**

Let  $X$  be a set.

002S

1. *Functoriality.* The assignments  $U, V, (U, V) \mapsto U \cup V$  define functors

$$\begin{aligned} U \cup -: (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ - \cup V: (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ -_1 \cup -_2: (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \subset) &\rightarrow (\mathcal{P}(X), \subset), \end{aligned}$$

where  $-_1 \cup -_2$  is the functor where

· *Action on Objects.* For each  $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(X)$ , we have

$$[-_1 \cup -_2](U, V) \stackrel{\text{def}}{=} U \cup V;$$

· *Action on Morphisms.* For each pair of morphisms

$$\begin{aligned} \iota_U: U &\hookrightarrow U', \\ \iota_V: V &\hookrightarrow V' \end{aligned}$$

of  $\mathcal{P}(X) \times \mathcal{P}(X)$ , the image

$$\iota_U \cup \iota_V: U \cup V \hookrightarrow U' \cup V'$$

of  $(\iota_U, \iota_V)$  by  $\cup$  is the inclusion

$$U \cup V \subset U' \cup V'$$

i.e. where we have

(★) If  $U \subset U'$  and  $V \subset V'$ , then  $U \cup V \subset U' \cup V'$ ;

and where  $U \cup -$  and  $- \cup V$  are the partial functors of  $-_1 \cup -_2$  at  $U, V \in \mathcal{P}(X)$ .

002T

2. *Via Intersections and Symmetric Differences.* We have an equality of sets

$$U \cup V = (U \Delta V) \Delta (U \cap V)$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V \in \mathcal{P}(X)$ .

002U

3. *Associativity.* We have an equality of sets

$$(U \cup V) \cup W = U \cup (V \cup W)$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V, W \in \mathcal{P}(X)$ .

002V

4. *Unitality.* We have equalities of sets

$$U \cup \emptyset = U,$$

$$\emptyset \cup U = U$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U \in \mathcal{P}(X)$ .

002W

5. *Commutativity.* We have an equality of sets

$$U \cup V = V \cup U$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V \in \mathcal{P}(X)$ .

002X

6. *Idempotency.* We have an equality of sets

$$U \cup U = U$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U \in \mathcal{P}(X)$ .

002Y

7. *Distributivity Over Intersections.* We have equalities of sets

$$U \cup (V \cap W) = (U \cup V) \cap (U \cup W),$$

$$(U \cap V) \cup W = (U \cup W) \cap (V \cup W)$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V, W \in \mathcal{P}(X)$ .

002Z

8. *Interaction With Powersets and Semirings.* The quintuple  $(\mathcal{P}(X), \cup, \cap, \emptyset, X)$  is an idempotent commutative semiring.

#### PROOF 3.5.3 ► PROOF OF PROPOSITION 3.5.2

Item 1: Functoriality

Omitted.

Item 2: Via Intersections and Symmetric Differences

Omitted.

Item 3: Associativity

Clear.

Item 4: Unitality

Clear.

Item 5: Commutativity

Clear.

Item 6: Idempotency

Clear.

Item 7: Distributivity Over Intersections

Omitted.

Item 8: Interaction With Powersets and Semirings

This follows from **Items 3 to 6** and **Items 3 to 5, 7 and 8** of **Proposition 3.7.2**. 

### 0030 3.6 Intersections of Families

Let  $\mathcal{F}$  be a family of sets.

#### 0031 DEFINITION 3.6.1 ► INTERSECTIONS OF FAMILIES

The **intersection of a family  $\mathcal{F}$  of sets** is the set  $\bigcap_{X \in \mathcal{F}} X$  defined by

$$\bigcap_{X \in \mathcal{F}} X \stackrel{\text{def}}{=} \left\{ z \in \bigcup_{X \in \mathcal{F}} X \mid \text{for each } X \in \mathcal{F}, \text{ we have } z \in X \right\}.$$

### 0032 3.7 Binary Intersections

Let  $X$  and  $Y$  be sets.

#### 0033 DEFINITION 3.7.1 ► BINARY INTERSECTIONS

The **intersection<sup>1</sup> of  $X$  and  $Y$**  is the set  $X \cap Y$  defined by

$$X \cap Y \stackrel{\text{def}}{=} \bigcap_{z \in \{X, Y\}} z.$$

<sup>1</sup>*Further Terminology:* Also called the **binary intersection of  $X$  and  $Y$** , for emphasis.



0034

**PROPOSITION 3.7.2 ► PROPERTIES OF BINARY INTERSECTIONS**

Let  $X$  be a set.

0035

1. *Functoriality.* The assignments  $U, V, (U, V) \mapsto U \cap V$  define functors

$$\begin{aligned} U \cap -: (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ - \cap V: (\mathcal{P}(X), \subset) &\rightarrow (\mathcal{P}(X), \subset), \\ -_1 \cap -_2: (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \subset) &\rightarrow (\mathcal{P}(X), \subset), \end{aligned}$$

where  $-_1 \cap -_2$  is the functor where

- *Action on Objects.* For each  $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(X)$ , we have

$$[-_1 \cap -_2](U, V) \stackrel{\text{def}}{=} U \cap V;$$

- *Action on Morphisms.* For each pair of morphisms

$$\begin{aligned} \iota_U: U &\hookrightarrow U', \\ \iota_V: V &\hookrightarrow V' \end{aligned}$$

of  $\mathcal{P}(X) \times \mathcal{P}(X)$ , the image

$$\iota_U \cap \iota_V: U \cap V \hookrightarrow U' \cap V'$$

of  $(\iota_U, \iota_V)$  by  $\cap$  is the inclusion

$$U \cap V \subset U' \cap V'$$

i.e. where we have

$$(\star) \text{ If } U \subset U' \text{ and } V \subset V', \text{ then } U \cap V \subset U' \cap V';$$

and where  $U \cap -$  and  $- \cap V$  are the partial functors of  $-_1 \cap -_2$  at  $U, V \in \mathcal{P}(X)$ .

0036

2. *Adjointness.* We have adjunctions

$$(U \cap - \dashv \mathbf{Hom}_{\mathcal{P}(X)}(U, -)): \mathcal{P}(X) \begin{array}{c} \xrightarrow{U \cap -} \\ \perp \\ \xleftarrow{\mathbf{Hom}_{\mathcal{P}(X)}(U, -)} \end{array} \mathcal{P}(X),$$

$$(- \cap V \dashv \mathbf{Hom}_{\mathcal{P}(X)}(V, -)): \mathcal{P}(X) \begin{array}{c} \xrightarrow{- \cap V} \\ \perp \\ \xleftarrow{\mathbf{Hom}_{\mathcal{P}(X)}(V, -)} \end{array} \mathcal{P}(X),$$

where

$$\mathbf{Hom}_{\mathcal{P}(X)}(-_1, -_2): \mathcal{P}(X)^{\text{op}} \times \mathcal{P}(X) \rightarrow \mathcal{P}(X)$$

is the bifunctor defined by<sup>1</sup>

$$\mathbf{Hom}_{\mathcal{P}(X)}(U, V) \stackrel{\text{def}}{=} (X \setminus U) \cup V$$

witnessed by bijections

$$\text{Hom}_{\mathcal{P}(X)}(U \cap V, W) \cong \text{Hom}_{\mathcal{P}(X)}(U, \mathbf{Hom}_{\mathcal{P}(X)}(V, W)),$$

$$\text{Hom}_{\mathcal{P}(X)}(U \cap V, W) \cong \text{Hom}_{\mathcal{P}(X)}(V, \mathbf{Hom}_{\mathcal{P}(X)}(U, W)),$$

natural in  $U, V, W \in \mathcal{P}(X)$ , i.e. where:

(a) The following conditions are equivalent:

- i. We have  $U \cap V \subset W$ .
- ii. We have  $U \subset \mathbf{Hom}_{\mathcal{P}(X)}(V, W)$ .
- iii. We have  $U \subset (X \setminus V) \cup W$ .

(b) The following conditions are equivalent:

- i. We have  $V \cap U \subset W$ .
- ii. We have  $V \subset \mathbf{Hom}_{\mathcal{P}(X)}(U, W)$ .
- iii. We have  $V \subset (X \setminus U) \cup W$ .

0037

3. *Associativity.* We have an equality of sets

$$(U \cap V) \cap W = U \cap (V \cap W)$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V, W \in \mathcal{P}(X)$ .

0038

4. *Unitality.* Let  $X$  be a set and let  $U \in \mathcal{P}(X)$ . We have equalities of sets

$$X \cap U = U,$$

$$U \cap X = U$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U \in \mathcal{P}(X)$ .

0039 5. *Commutativity.* We have an equality of sets

$$U \cap V = V \cap U$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V \in \mathcal{P}(X)$ .

003A 6. *Idempotency.* We have an equality of sets

$$U \cap U = U$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U \in \mathcal{P}(X)$ .

003B 7. *Distributivity Over Unions.* We have equalities of sets

$$\begin{aligned} U \cap (V \cup W) &= (U \cap V) \cup (U \cap W), \\ (U \cup V) \cap W &= (U \cap W) \cup (V \cap W) \end{aligned}$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V, W \in \mathcal{P}(X)$ .

003C 8. *Annihilation With the Empty Set.* We have an equality of sets

$$\begin{aligned} \emptyset \cap X &= \emptyset, \\ X \cap \emptyset &= \emptyset \end{aligned}$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U \in \mathcal{P}(X)$ .

003D 9. *Interaction With Powersets and Monoids With Zero.* The quadruple  $((\mathcal{P}(X), \emptyset), \cap, X)$  is a commutative monoid with zero.

003E 10. *Interaction With Powersets and Semirings.* The quintuple  $(\mathcal{P}(X), \cup, \cap, \emptyset, X)$  is an idempotent commutative semiring.

---

<sup>1</sup>*Intuition:* Since intersections are the products in  $\mathcal{P}(X)$ , the left adjoint  $\mathbf{Hom}_{\mathcal{P}(X)}(U, V)$  works as a function type  $U \rightarrow V$ .

Now, under the Curry–Howard correspondence, the function type  $U \rightarrow V$  corresponds to implication  $U \implies V$ , which is logically equivalent to the statement  $\neg U \vee V$ , which in turn corresponds to the set  $U^c \vee V \stackrel{\text{def}}{=} (X \setminus U) \cup V$ .

## PROOF 3.7.3 ► PROOF OF PROPOSITION 3.7.2

Item 1: Functoriality

Omitted.

Item 2: Adjointness

See [MSE 267469].

Item 3: Associativity

Clear.

Item 4: Unitality

Clear.

Item 5: Commutativity

Clear.

Item 6: Idempotency

Clear.

Item 7: Distributivity Over Unions

Omitted.

Item 8: Annihilation With the Empty Set

Clear.

Item 9: Interaction With Powersets and Monoids With Zero

This follows from Items 3 to 5 and 8.

Item 10: Interaction With Powersets and Semirings

This follows from Items 3 to 6 and Items 3 to 5, 7 and 8 of Proposition 3.7.2. 

## 003F 3.8 Differences

Let  $X$  and  $Y$  be sets.

## 003G DEFINITION 3.8.1 ► DIFFERENCES

The **difference of  $X$  and  $Y$**  is the set  $X \setminus Y$  defined by

$$X \setminus Y \stackrel{\text{def}}{=} \{a \in X \mid a \notin Y\}.$$

003H

**PROPOSITION 3.8.2 ► PROPERTIES OF DIFFERENCES**

Let  $X$  be a set.

003J

1. *Functoriality.* The assignments  $U, V, (U, V) \mapsto U \cap V$  define functors

$$\begin{aligned} U \setminus - &: (\mathcal{P}(X), \supset) \rightarrow (\mathcal{P}(X), \subset), \\ - \setminus V &: (\mathcal{P}(X), \subset) \rightarrow (\mathcal{P}(X), \subset), \\ -_1 \setminus -_2 &: (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \supset) \rightarrow (\mathcal{P}(X), \subset), \end{aligned}$$

where  $-_1 \setminus -_2$  is the functor where

- *Action on Objects.* For each  $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(X)$ , we have

$$[-_1 \setminus -_2](U, V) \stackrel{\text{def}}{=} U \setminus V;$$

- *Action on Morphisms.* For each pair of morphisms

$$\begin{aligned} \iota_A &: A \hookrightarrow B, \\ \iota_U &: U \hookrightarrow V \end{aligned}$$

of  $\mathcal{P}(X) \times \mathcal{P}(X)$ , the image

$$\iota_U \setminus \iota_V : A \setminus V \hookrightarrow B \setminus U$$

of  $(\iota_U, \iota_V)$  by  $\setminus$  is the inclusion

$$A \setminus V \subset B \setminus U$$

i.e. where we have

$$(\star) \text{ If } A \subset B \text{ and } U \subset V, \text{ then } A \setminus V \subset B \setminus U;$$

and where  $U \setminus -$  and  $- \setminus V$  are the partial functors of  $-_1 \setminus -_2$  at  $U, V \in \mathcal{P}(X)$ .

003K

2. *De Morgan's Laws.* We have equalities of sets

$$\begin{aligned} X \setminus (U \cup V) &= (X \setminus U) \cap (X \setminus V), \\ X \setminus (U \cap V) &= (X \setminus U) \cup (X \setminus V) \end{aligned}$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V \in \mathcal{P}(X)$ .

003L

3. *Interaction With Unions I.* We have equalities of sets

$$(U \setminus V) \cup W = (U \cup W) \setminus (V \setminus W)$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V, W \in \mathcal{P}(X)$ .

003M

4. *Interaction With Unions II.* We have equalities of sets

$$(U \cup V) \setminus W = (U \setminus W) \cup (V \setminus W)$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V, W \in \mathcal{P}(X)$ .

003N

5. *Interaction With Intersections.* We have equalities of sets

$$\begin{aligned} (U \setminus V) \cap W &= (U \cap W) \setminus V \\ &= U \cap (W \setminus V) \end{aligned}$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V, W \in \mathcal{P}(X)$ .

003P

6. *Triple Differences.* We have

$$U \setminus (V \setminus W) = (U \cap W) \cup (U \setminus V)$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V, W \in \mathcal{P}(X)$ .

003Q

7. *Left Annihilation.* We have

$$\emptyset \setminus U = \emptyset$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U \in \mathcal{P}(X)$ .

003R

8. *Right Unitality.* We have

$$U \setminus \emptyset = U$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U \in \mathcal{P}(X)$ .

003S

9. *Invertibility.* We have

$$U \setminus U = \emptyset$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U \in \mathcal{P}(X)$ .

003T

10. *Interaction With Containment.* The following conditions are equivalent:(a) We have  $V \setminus U \subset W$ .(b) We have  $V \setminus W \subset U$ .

**PROOF 3.8.3 ► PROOF OF PROPOSITION 3.8.2**

Item 1: Functoriality

Omitted.

Item 2: De Morgan's Laws

Omitted.

Item 3: Interaction With Unions I

Omitted.

Item 4: Interaction With Unions II

Omitted.

Item 5: Interaction With Intersections

Omitted.

Item 6: Triple Differences

Omitted.

Item 7: Left Annihilation

Clear.

Item 8: Right Unitality

Clear.

Item 9: Invertibility

Clear.

Item 10: Interaction With Containment

Omitted. **003U 3.9 Complements**Let  $X$  be a set and let  $U \in \mathcal{P}(X)$ .**003V DEFINITION 3.9.1 ► COMPLEMENTS**The **complement of  $U$**  is the set  $U^c$  defined by

$$\begin{aligned}
 U^c &\stackrel{\text{def}}{=} X \setminus U \\
 &\stackrel{\text{def}}{=} \{a \in X \mid a \notin U\}.
 \end{aligned}$$

003W

**PROPOSITION 3.9.2 ► PROPERTIES OF COMPLEMENTS**

Let  $X$  be a set.

003X

1. *Functoriality.* The assignment  $U \mapsto U^c$  defines a functor

$$(-)^c: \mathcal{P}(X)^{\text{op}} \rightarrow \mathcal{P}(X),$$

where

- *Action on Objects.* For each  $U \in \mathcal{P}(X)$ , we have

$$[(-)^c](U) \stackrel{\text{def}}{=} U^c;$$

- *Action on Morphisms.* For each morphism  $\iota_U: U \hookrightarrow V$  of  $\mathcal{P}(X)$ , the image

$$\iota_U^c: V^c \hookrightarrow U^c$$

of  $\iota_U$  by  $(-)^c$  is the inclusion

$$V^c \subset U^c$$

i.e. where we have

(★) If  $U \subset V$ , then  $V^c \subset U^c$ .

003Y

2. *De Morgan's Laws.* We have equalities of sets

$$(U \cup V)^c = U^c \cap V^c,$$

$$(U \cap V)^c = U^c \cup V^c$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V \in \mathcal{P}(X)$ .

003Z

3. *Involutority.* We have

$$(U^c)^c = U$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U \in \mathcal{P}(X)$ .



## PROOF 3.9.3 ► PROOF OF PROPOSITION 3.9.2

Item 1: Functoriality

Clear.

Item 2: De Morgan's Laws

Omitted.

Item 3: Involutority

Clear.



## 0040 3.10 Symmetric Differences

Let  $A$  and  $B$  be sets.

## 0041 DEFINITION 3.10.1 ► SYMMETRIC DIFFERENCES

The **symmetric difference of  $A$  and  $B$**  is the set  $A \triangle B$  defined by

$$A \triangle B \stackrel{\text{def}}{=} (A \setminus B) \cup (B \setminus A).$$

## 0042 PROPOSITION 3.10.2 ► PROPERTIES OF SYMMETRIC DIFFERENCES

Let  $X$  be a set.

- 0043 1. *Lack of Functoriality.* The assignment  $(U, V) \mapsto U \triangle V$  **does not** define a functor

$$-_1 \triangle -_2: (\mathcal{P}(X) \times \mathcal{P}(X), \subset \times \subset) \rightarrow (\mathcal{P}(X), \subset).$$

- 0044 2. *Via Unions and Intersections.* We have<sup>1</sup>

$$U \triangle V = (U \cup V) \setminus (U \cap V)$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V \in \mathcal{P}(X)$ .

- 0045 3. *Associativity.* We have<sup>2</sup>

$$(U \triangle V) \triangle W = U \triangle (V \triangle W)$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V, W \in \mathcal{P}(X)$ .

0046 4. *Unitality*. We have

$$\begin{aligned} U \Delta \emptyset &= U, \\ \emptyset \Delta U &= U \end{aligned}$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U \in \mathcal{P}(X)$ .

0047 5. *Invertibility*. We have

$$U \Delta U = \emptyset$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U \in \mathcal{P}(X)$ .

0048 6. *Commutativity*. We have

$$U \Delta V = V \Delta U$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V \in \mathcal{P}(X)$ .

0049 7. *“Transitivity”*. We have

$$(U \Delta V) \Delta (V \Delta W) = U \Delta W$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V, W \in \mathcal{P}(X)$ .

004A 8. *The Triangle Inequality for Symmetric Differences*. We have

$$U \Delta W \subset U \Delta V \cup V \Delta W$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V, W \in \mathcal{P}(X)$ .

004B 9. *Distributivity Over Intersections*. We have

$$\begin{aligned} U \cap (V \Delta W) &= (U \cap V) \Delta (U \cap W), \\ (U \Delta V) \cap W &= (U \cap W) \Delta (V \cap W) \end{aligned}$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V, W \in \mathcal{P}(X)$ .

004C 10. *Interaction With Indicator Functions*. We have

$$\chi_{U \Delta V} \equiv \chi_U + \chi_V \pmod{2}$$

for each  $X \in \text{Obj}(\text{Sets})$  and each  $U, V \in \mathcal{P}(X)$ .

004D

11. *Bijectivity.* Given  $A, B \subset \mathcal{P}(X)$ , the maps

$$\begin{aligned} A \triangle -: \mathcal{P}(X) &\rightarrow \mathcal{P}(X), \\ - \triangle B: \mathcal{P}(X) &\rightarrow \mathcal{P}(X) \end{aligned}$$

are bijections with inverses given by

$$\begin{aligned} (A \triangle -)^{-1} &= - \cup (A \cap -), \\ (- \triangle B)^{-1} &= - \cup (B \cap -). \end{aligned}$$

Moreover, the map

$$C \mapsto C \triangle (A \triangle B)$$

is a bijection of  $\mathcal{P}(X)$  onto itself sending  $A$  to  $B$  and  $B$  to  $A$ .

004E

12. *Interaction With Powersets and Groups I.* The quadruple  $(\mathcal{P}(X), \triangle, \emptyset, \text{id}_{\mathcal{P}(X)})$  is an abelian group.<sup>3,4,5</sup>

004F

13. *Interaction With Powersets and Groups II.* Every element of  $\mathcal{P}(X)$  has order 2 with respect to  $\triangle$ , and thus  $\mathcal{P}(X)$  is a *Boolean group* (i.e. an abelian 2-group).

004G

14. *Interaction With Powersets and Vector Spaces I.* The pair  $(\mathcal{P}(X), \alpha_{\mathcal{P}(X)})$  consisting of

- The group  $\mathcal{P}(X)$  of **Item 12**;
- The map  $\alpha_{\mathcal{P}(X)}: \mathbb{F}_2 \times \mathcal{P}(X) \rightarrow \mathcal{P}(X)$  defined by

$$\begin{aligned} 0 \cdot U &\stackrel{\text{def}}{=} \emptyset, \\ 1 \cdot U &\stackrel{\text{def}}{=} U; \end{aligned}$$

is an  $\mathbb{F}_2$ -vector space.

004H

15. *Interaction With Powersets and Vector Spaces II.* If  $X$  is finite, then:

(a) The set of singletons sets on the elements of  $X$  forms a basis for the  $\mathbb{F}_2$ -vector space  $(\mathcal{P}(X), \alpha_{\mathcal{P}(X)})$  of **Item 14**.

(b) We have

$$\dim(\mathcal{P}(X)) = \# \mathcal{P}(X).$$


004J

16. *Interaction With Powersets and Rings.* The quintuple  $(\mathcal{P}(X), \Delta, \cap, \emptyset, X)$  is a commutative ring.<sup>6</sup>

<sup>1</sup>Illustration:



<sup>2</sup>Illustration:



<sup>3</sup>Example: When  $X = \emptyset$ , we have an isomorphism of groups between  $\mathcal{P}(\emptyset)$  and the trivial group:

$$(\mathcal{P}(\emptyset), \Delta, \emptyset, \text{id}_{\mathcal{P}(\emptyset)}) \cong \text{pt.}$$

<sup>4</sup>Example: When  $X = \text{pt}$ , we have an isomorphism of groups between  $\mathcal{P}(\text{pt})$  and  $\mathbb{Z}_2$ :

$$(\mathcal{P}(\text{pt}), \Delta, \emptyset, \text{id}_{\mathcal{P}(\text{pt})}) \cong \mathbb{Z}_2.$$

<sup>5</sup>Example: When  $X = \{0, 1\}$ , we have an isomorphism of groups between  $\mathcal{P}(\{0, 1\})$  and  $\mathbb{Z}_2 \times \mathbb{Z}_2$ :

$$(\mathcal{P}(\{0, 1\}), \Delta, \emptyset, \text{id}_{\mathcal{P}(\{0, 1\})}) \cong \mathbb{Z}_2 \times \mathbb{Z}_2.$$



<sup>6</sup>Warning: The analogous statement replacing intersections by unions (i.e. that the quintuple  $(\mathcal{P}(X), \Delta, \cup, \emptyset, X)$  is a ring) is false, however. See [Pro23b] for a proof.

#### PROOF 3.10.3 ► PROOF OF PROPOSITION 3.10.2

Item 1: Lack of Functoriality

Omitted.

Item 2: Via Unions and Intersections

Omitted.

Item 3: Associativity

Omitted.

Item 4: Unitality

Clear.

Item 5: Invertibility

Clear.

Item 6: Commutativity

Clear.

Item 7: “Transitivity”

We have

$$\begin{aligned}
 (U \triangle V) \triangle (V \triangle W) &= U \triangle (V \triangle (V \triangle W)) && \text{(by Item 3)} \\
 &= U \triangle ((V \triangle V) \triangle W) && \text{(by Item 3)} \\
 &= U \triangle (\emptyset \triangle W) && \text{(by Item 5)} \\
 &= U \triangle W && \text{(by Item 4)}
 \end{aligned}$$

Item 8: The Triangle Inequality for Symmetric Differences

This follows from Items 2 and 7.

Item 9: Distributivity Over Intersections

Omitted.

Item 10: Interaction With Indicator Functions

Clear.

Item 11: Bijectivity

Clear.

Item 12: Interaction With Powersets and Groups I

This follows from Items 3 to 6.

Item 13: Interaction With Powersets and Groups II

This follows from Item 5.

Item 14: Interaction With Powersets and Vector Spaces I

Clear.

Item 15: Interaction With Powersets and Vector Spaces II

Omitted.

Item 16: Interaction With Powersets and Rings

This follows from Items 9 and 12 and Items 8 and 9 of Proposition 3.7.2.<sup>1</sup>



<sup>1</sup>Reference: [Pro23a].

**004K 3.11 Ordered Pairs**

Let  $A$  and  $B$  be sets.

**004L DEFINITION 3.11.1 ► ORDERED PAIRS**

The **ordered pair associated to  $A$  and  $B$**  is the set  $(A, B)$  defined by

$$(A, B) \stackrel{\text{def}}{=} \{\{A\}, \{A, B\}\}.$$

**004M PROPOSITION 3.11.2 ► PROPERTIES OF ORDERED PAIRS**

Let  $A$  and  $B$  be sets.

- 004N**
1. *Uniqueness.* Let  $A, B, C$ , and  $D$  be sets. The following conditions are equivalent:
    - (a) We have  $(A, B) = (C, D)$ .
    - (b) We have  $A = C$  and  $B = D$ .

**PROOF 3.11.3 ► PROOF OF PROPOSITION 3.11.2**

Item 1: Uniqueness

See [Cie97, Theorem 1.2.3].

**004P 4 Powersets****004Q 4.1 Characteristic Functions**

Let  $X$  be a set.

**004R DEFINITION 4.1.1 ► CHARACTERISTIC FUNCTIONS**

Let  $U \subset X$  and let  $x \in X$ .

- 004S**
1. The **characteristic function of  $U$** <sup>1</sup> is the function<sup>2</sup>

$$\chi_U: X \rightarrow \{\text{t}, \text{f}\}$$

defined by

$$\chi_U(x) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x \in U, \\ \text{false} & \text{if } x \notin U \end{cases}$$

for each  $x \in X$ .

004T

2. The **characteristic function of  $x$**  is the function<sup>3</sup>

$$\chi_x: X \rightarrow \{\text{t}, \text{f}\}$$

defined by

$$\chi_x \stackrel{\text{def}}{=} \chi_{\{x\}},$$

i.e. by

$$\chi_x(y) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x = y, \\ \text{false} & \text{if } x \neq y \end{cases}$$

for each  $y \in X$ .

004U

3. The **characteristic relation on  $X$** <sup>4</sup> is the relation<sup>5</sup>

$$\chi_X(-, -): X \times X \rightarrow \{\text{t}, \text{f}\}$$

on  $X$  defined by<sup>6</sup>

$$\chi_X(x, y) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x = y, \\ \text{false} & \text{if } x \neq y \end{cases}$$

for each  $x, y \in X$ .

004V

4. The **characteristic embedding**<sup>7</sup> of  $X$  into  $\mathcal{P}(X)$  is the function

$$\chi(-): X \hookrightarrow \mathcal{P}(X)$$

defined by

$$\chi(-)(x) \stackrel{\text{def}}{=} \chi_x$$

for each  $x \in X$ .

<sup>1</sup>Further Terminology: Also called the **indicator function of  $U$** .

<sup>2</sup>Further Notation: Also written  $\chi_X(U, -)$  or  $\chi_X(-, U)$ .

<sup>3</sup>Further Notation: Also written  $\chi_x$ ,  $\chi_X(x, -)$ , or  $\chi_X(-, x)$ .

<sup>4</sup>Further Terminology: Also called the **identity relation on  $X$** .

<sup>5</sup>Further Notation: Also written  $\chi_X^{-1}$ , or  $\sim_{\text{id}}$  in the context of relations.

<sup>6</sup>As a subset of  $X \times X$ , the relation  $\chi_X$  corresponds to the diagonal  $\Delta_X \subset X \times X$  of  $X$ .

<sup>7</sup>The name “characteristic *embedding*” comes from the fact that there is an analogue of fully faithfulness for  $\chi(-)$ : given a set  $X$ , we have

$$\text{Hom}_{\mathcal{P}(X)}(\chi_x, \chi_y) = \chi_X(x, y),$$

for each  $x, y \in X$ .

004W

#### REMARK 4.1.2 ► CHARACTERISTIC FUNCTIONS AS DECATEGORIFICATIONS OF PRESHEAVES

The definitions in [Definition 4.1.1](#) are decategorifications of co/presheaves, representable co/presheaves, Hom profunctors, and the Yoneda embedding:<sup>1</sup>

1. A function

$$f: X \rightarrow \{\mathbf{t}, \mathbf{f}\}$$

is a decategorification of a presheaf

$$\mathcal{F}: C^{\text{op}} \rightarrow \text{Sets},$$

with the characteristic functions  $\chi_U$  of the subsets of  $X$  being the primordial examples (and, in fact, all examples) of these.

2. The characteristic function

$$\chi_x: X \rightarrow \{\mathbf{t}, \mathbf{f}\}$$

of an *element*  $x$  of  $X$  is a decategorification of the representable presheaf

$$h_X: C^{\text{op}} \rightarrow \text{Sets}$$

of an *object*  $x$  of a category  $C$ .

3. The characteristic relation

$$\chi_X(-_1, -_2): X \times X \rightarrow \{\mathbf{t}, \mathbf{f}\}$$

of  $X$  is a decategorification of the Hom profunctor

$$\text{Hom}_C(-_1, -_2): C^{\text{op}} \times C \rightarrow \text{Sets}$$

of a category  $C$ .



## 4. The characteristic embedding

$$\chi_{(-)} : X \hookrightarrow \mathcal{P}(X)$$

of  $X$  into  $\mathcal{P}(X)$  is a decategorification of the Yoneda embedding

$$\mathcal{Y} : C^{\text{op}} \hookrightarrow \text{PSh}(C)$$

of a category  $C$  into  $\text{PSh}(C)$ .

## 5. There is also a direct parallel between unions and colimits:

- An element of  $\mathcal{P}(X)$  is a union of elements of  $X$ , viewed as one-point subsets  $\{x\} \in \mathcal{P}(A)$ ;
- An object of  $\text{PSh}(C)$  is a colimit of objects of  $C$ , viewed as representable presheaves  $h_X \in \text{Obj}(\text{PSh}(C))$ .

<sup>1</sup>These statements can be made precise by using the embeddings

$$\begin{aligned} (-)_{\text{disc}} : \text{Sets} &\hookrightarrow \text{Cats}, \\ (-)_{\text{disc}} : \{t, f\}_{\text{disc}} &\hookrightarrow \text{Sets} \end{aligned}$$

of sets into categories and of classical truth values into sets.

For instance, in this approach the characteristic function

$$\chi_x : X \rightarrow \{t, f\}$$

of an element  $x$  of  $X$ , defined by

$$\chi_x(y) \stackrel{\text{def}}{=} \begin{cases} \text{true} & \text{if } x = y, \\ \text{false} & \text{if } x \neq y \end{cases}$$

for each  $y \in X$ , is recovered as the representable presheaf

$$\text{Hom}_{X_{\text{disc}}}(-, x) : X_{\text{disc}} \rightarrow \text{Sets}$$

of the corresponding object  $x$  of  $X_{\text{disc}}$ , defined on objects by

$$\text{Hom}_{X_{\text{disc}}}(y, x) \stackrel{\text{def}}{=} \begin{cases} \text{pt} & \text{if } x = y, \\ \emptyset & \text{if } x \neq y \end{cases}$$

for each  $y \in \text{Obj}(X_{\text{disc}})$ .

004X


**PROPOSITION 4.1.3 ► THE INCLUSION OF CHARACTERISTIC RELATIONS ASSOCIATED TO A FUNCTION**

Let  $f: A \rightarrow B$  be a function. We have an inclusion

$$\begin{array}{ccc} A \times A & \xrightarrow{\chi_A(-1, -2)} & \{\text{true}, \text{false}\} \\ \downarrow f \times f & \subset & \downarrow \text{id}_{\{\text{true}, \text{false}\}} \\ B \times B & \xrightarrow{\chi_B(-1, -2)} & \{\text{true}, \text{false}\}. \end{array}$$

$\chi_B \circ (f \times f) \subset \chi_A$

**PROOF 4.1.4 ► PROOF OF PROPOSITION 4.1.3**

The inclusion  $\chi_B(f(a), f(b)) \subset \chi_A(a, b)$  is equivalent to the statement “if  $a = b$ , then  $f(a) = f(b)$ ”, which is true. 

004Y

**PROPOSITION 4.1.5 ► THE YONEDA LEMMA FOR SETS**

Let  $X$  be a set and let  $U \subset X$  be a subset of  $X$ . We have

$$\text{Hom}_{\mathcal{P}(X)}(\chi_x, \chi_U) = \chi_U(x)$$

for each  $x \in X$ , giving an equality of functions

$$\text{Hom}_{\mathcal{P}(X)}(\chi_{(-)}, \chi_U) = \chi_U.$$

**PROOF 4.1.6 ► PROOF OF PROPOSITION 4.1.5**

Clear. 

004Z

**COROLLARY 4.1.7 ► THE CHARACTERISTIC EMBEDDING IS FULLY FAITHFUL**

The characteristic embedding is fully faithful, i.e., we have

$$\text{Hom}_{\mathcal{P}(X)}(\chi_x, \chi_y) = \chi_X(x, y)$$

for each  $x, y \in X$ .

## PROOF 4.1.8 ► PROOF OF COROLLARY 4.1.7

This follows from [Proposition 4.1.5](#).



## 0050 4.2 Powersets

Let  $X$  be a set.

## 0051 DEFINITION 4.2.1 ► POWERSETS

The **powerset of  $X$**  is the set  $\mathcal{P}(X)$  defined by

$$\mathcal{P}(X) \stackrel{\text{def}}{=} \{U \in P \mid U \subset X\},$$

where  $P$  is the set in the axiom of powerset, ?? of ??.

## 0052 REMARK 4.2.2 ► POWERSETS AS DECATEGORIFICATIONS OF Co/PRESHEAF CATEGORIES

The powerset of a set is a decategorification of the category of presheaves of a category: while<sup>1</sup>

- The powerset of a set  $X$  is equivalently ([Item 6 of Proposition 4.2.3](#)) the set

$$\text{Sets}(X, \{t, f\})$$

of functions from  $X$  to the set  $\{t, f\}$  of classical truth values;

- The category of presheaves on a category  $C$  is the category

$$\text{Fun}(C^{\text{op}}, \text{Sets})$$

of functors from  $C^{\text{op}}$  to the category Sets of sets.

<sup>1</sup>This parallel is based on the following comparison:

- A category is enriched over the category

$$\text{Sets} \stackrel{\text{def}}{=} \text{Cats}_0$$

of sets (i.e. “0-categories”), with presheaves taking values on it;

- A set is enriched over the set

$$\{t, f\} \stackrel{\text{def}}{=} \text{Cats}_{-1}$$

of classical truth values (i.e. “(−1)-categories”), with characteristic functions taking values on it.

0053

**PROPOSITION 4.2.3 ► PROPERTIES OF POWERSETS**

Let  $X$  be a set.

0054

1. *Functoriality.* The assignment  $X \mapsto \mathcal{P}(X)$  defines functors

$$\begin{aligned}\mathcal{P}_* &: \mathbf{Sets} \rightarrow \mathbf{Sets}, \\ \mathcal{P}^{-1} &: \mathbf{Sets}^{\text{op}} \rightarrow \mathbf{Sets}, \\ \mathcal{P}_! &: \mathbf{Sets} \rightarrow \mathbf{Sets}\end{aligned}$$

where

- *Action on Objects.* For each  $A \in \text{Obj}(\mathbf{Sets})$ , we have

$$\begin{aligned}\mathcal{P}_*(A) &\stackrel{\text{def}}{=} \mathcal{P}(A), \\ \mathcal{P}^{-1}(A) &\stackrel{\text{def}}{=} \mathcal{P}(A), \\ \mathcal{P}_!(A) &\stackrel{\text{def}}{=} \mathcal{P}(A);\end{aligned}$$

- *Action on Morphisms.* For each morphism  $f: A \rightarrow B$  of  $\mathbf{Sets}$ , the images

$$\begin{aligned}\mathcal{P}_*(f) &: \mathcal{P}(A) \rightarrow \mathcal{P}(B), \\ \mathcal{P}^{-1}(f) &: \mathcal{P}(B) \rightarrow \mathcal{P}(A), \\ \mathcal{P}_!(f) &: \mathcal{P}(A) \rightarrow \mathcal{P}(B)\end{aligned}$$

of  $f$  by  $\mathcal{P}_*$ ,  $\mathcal{P}^{-1}$ , and  $\mathcal{P}_!$  are defined by

$$\begin{aligned}\mathcal{P}_*(f) &\stackrel{\text{def}}{=} f_*, \\ \mathcal{P}^{-1}(f) &\stackrel{\text{def}}{=} f^{-1}, \\ \mathcal{P}_!(f) &\stackrel{\text{def}}{=} f_!,\end{aligned}$$

as in [Definitions 4.3.1](#), [4.4.1](#) and [4.5.1](#).

0055

2. *Adjointness I.* We have an adjunction

$$(\mathcal{P}^{-1} \dashv \mathcal{P}^{-1, \text{op}}): \quad \mathbf{Sets}^{\text{op}} \begin{array}{c} \xrightarrow{\mathcal{P}^{-1}} \\ \perp \\ \xleftarrow{\mathcal{P}^{-1, \text{op}}} \end{array} \mathbf{Sets},$$

witnessed by a bijection

$$\underbrace{\text{Sets}^{\text{op}}(\mathcal{P}(X), Y)}_{\stackrel{\text{def}}{=} \text{Sets}(Y, \mathcal{P}(X))} \cong \text{Sets}(X, \mathcal{P}(Y)),$$

natural in  $X \in \text{Obj}(\text{Sets})$  and  $Y \in \text{Obj}(\text{Sets}^{\text{op}})$ .

0056

3. *Adjointness II.* We have an adjunction

$$(\text{Gr} \dashv \mathcal{P}_*) : \text{Sets} \begin{array}{c} \xrightarrow{\text{Gr}} \\ \perp \\ \xleftarrow{\mathcal{P}_*} \end{array} \text{Rel},$$

witnessed by a bijection of sets

$$\text{Rel}(\text{Gr}(A), B) \cong \text{Sets}(A, \mathcal{P}(B))$$

natural in  $A \in \text{Obj}(\text{Sets})$  and  $B \in \text{Obj}(\text{Rel})$ , where  $\text{Gr}$  is the graph functor of [Relations, Item 1](#) of [Proposition 3.1.2](#).

0057

4. *Symmetric Strong Monoidality With Respect to Coproducts.* The powerset functor  $\mathcal{P}_*$  of [Item 1](#) has a symmetric strong monoidal structure

$$(\mathcal{P}_*, \mathcal{P}_*^{\amalg}, \mathcal{P}_*^{\amalg}) : (\text{Sets}, \amalg, \emptyset) \rightarrow (\text{Sets}, \times, \text{pt})$$

being equipped with isomorphisms

$$\begin{aligned} \mathcal{P}_*^{\amalg} : \mathcal{P}(X) \times \mathcal{P}(Y) &\xrightarrow{\cong} \mathcal{P}(X \amalg Y), \\ \mathcal{P}_*^{\amalg} : \text{pt} &\xrightarrow{=} \mathcal{P}(\emptyset), \end{aligned}$$

natural in  $X, Y \in \text{Obj}(\text{Sets})$ .

0058

5. *Symmetric Lax Monoidality With Respect to Products.* The powerset functor  $\mathcal{P}_*$  of [Item 1](#) has a symmetric lax monoidal structure

$$(\mathcal{P}_*, \mathcal{P}_*^{\otimes}, \mathcal{P}_*^{\otimes}) : (\text{Sets}, \times, \text{pt}) \rightarrow (\text{Sets}, \times, \text{pt})$$

being equipped with isomorphisms

$$\begin{aligned}\mathcal{P}_{*|X,Y}^{\otimes} &: \mathcal{P}(X) \times \mathcal{P}(Y) \rightarrow \mathcal{P}(X \times Y), \\ \mathcal{P}_{*|\mathbb{P}}^{\otimes} &: \text{pt} \xrightarrow{=} \mathcal{P}(\emptyset),\end{aligned}$$

natural in  $X, Y \in \text{Obj}(\text{Sets})$ , where  $\mathcal{P}_{*|X,Y}^{\otimes}$  is given by

$$\mathcal{P}_{*|X,Y}^{\otimes}(U, V) \stackrel{\text{def}}{=} U \times V$$

for each  $(U, V) \in \mathcal{P}(X) \times \mathcal{P}(Y)$ .

0059 6. *Powersets as Sets of Functions.* The assignment  $U \mapsto \chi_U$  defines a bijection<sup>1</sup>

$$\chi_{(-)}: \mathcal{P}(X) \xrightarrow{\cong} \text{Sets}(X, \{\text{t}, \text{f}\}),$$

natural in  $X \in \text{Obj}(\text{Sets})$ .

005A 7. *Powersets as Sets of Relations.* We have bijections

$$\begin{aligned}\mathcal{P}(X) &\cong \text{Rel}(\text{pt}, X), \\ \mathcal{P}(X) &\cong \text{Rel}(X, \text{pt}),\end{aligned}$$

natural in  $X \in \text{Obj}(\text{Sets})$ .

005B 8. *As a Free Cocompletion: Universal Property.* The pair  $(\mathcal{P}(X), \chi_{(-)})$  consisting of

- The powerset  $\mathcal{P}(X)$  of  $X$ ;
- The characteristic embedding  $\chi_{(-)}: X \hookrightarrow \mathcal{P}(X)$  of  $X$  into  $\mathcal{P}(X)$ ;

satisfies the following universal property:

- (★) Given another pair  $(Y, f)$  consisting of
  - A cocomplete poset  $(Y, \leq)$ ;
  - A function  $f: X \rightarrow Y$ ;

there exists a unique cocontinuous morphism of posets  $(\mathcal{P}(X), \subset) \xrightarrow{\exists!} (Y, \leq)$  making the diagram

$$\begin{array}{ccc} & \mathcal{P}(X) & \\ \chi_X \nearrow & \downarrow \exists! & \\ X & \xrightarrow{f} & Y \end{array}$$

commute.

005C

9. *As a Free Cocompletion: Adjointness.* We have an adjunction<sup>2</sup>

$$(\chi_{(-)} \dashv \text{忘}) : \text{Sets} \begin{array}{c} \xrightarrow{\chi_{(-)}} \\ \perp \\ \xleftarrow{\text{忘}} \end{array} \text{Pos}^{\text{cocomp.}},$$

witnessed by a bijection

$$\text{Pos}^{\text{cocomp.}}((\mathcal{P}(X), \subset), (Y, \leq)) \cong \text{Sets}(X, Y),$$

natural in  $X \in \text{Obj}(\text{Sets})$  and  $(Y, \leq) \in \text{Obj}(\text{Pos})$ , where

· We have a natural map

$$\chi_X^* : \text{Pos}^{\text{cocomp.}}((\mathcal{P}(X), \subset), (Y, \leq)) \rightarrow \text{Sets}(X, Y)$$

defined by

$$\chi_X^*(f) \stackrel{\text{def}}{=} f \circ \chi_X,$$

i.e. by sending a cocontinuous morphism of posets  $f : \mathcal{P}(X) \rightarrow Y$  to the composition

$$X \xrightarrow{\chi_X} \mathcal{P}(X) \xrightarrow{f} Y;$$

· We have a natural map

$$\text{Lan}_{\chi_X} : \text{Sets}(X, Y) \rightarrow \text{Pos}^{\text{cocomp.}}((\mathcal{P}(X), \subset), (Y, \leq))$$

computed by

$$\begin{aligned}
 [\text{Lan}_{\chi_X}(f)](U) &\cong \int^{x \in X} \chi_{\mathcal{P}(X)}(\chi_x, U) \odot f(x) \\
 &\cong \int^{x \in X} \chi_U(x) \odot f(x) && \text{(by Proposition 4.1.5)} \\
 &\cong \bigvee_{x \in X} (\chi_U(x) \odot f(x))
 \end{aligned}$$

for each  $U \in \mathcal{P}(X)$ , where:

- $\bigvee$  is the join in  $(Y, \leq)$ ;
- We have

$$\begin{aligned}
 \text{true} \odot f(x) &\stackrel{\text{def}}{=} f(x), \\
 \text{false} \odot f(x) &\stackrel{\text{def}}{=} \emptyset_Y,
 \end{aligned}$$

where  $\emptyset_Y$  is the minimal element of  $(Y, \leq)$ .

<sup>1</sup>This bijection is a decategorified form of the equivalence

$$\text{PSh}(C) \stackrel{\text{eq.}}{\cong} \text{DFib}(C)$$

of Fibred Categories, ?? of ??, with  $\chi_{(-)}$  being a decategorified version of the category of elements construction of Fibred Categories, ??.

See also ?? of ??.

<sup>2</sup>In this sense,  $\mathcal{P}(A)$  is the free cocompletion of  $A$ . (Note that, despite its name, however, this is not an idempotent operation, as we have  $\mathcal{P}(\mathcal{P}(A)) \neq \mathcal{P}(A)$ .)

#### PROOF 4.2.4 ► PROOF OF PROPOSITION 4.2.3

Item 1: Functoriality

This follows from Items 3 and 4 of Proposition 4.3.5, Items 3 and 4 of Proposition 4.4.5, and Items 3 and 4 of Proposition 4.5.7.

Item 2: Adjointness I

Omitted.

Item 3: Adjointness II

Omitted.



Item 4: Symmetric Strong Monoidality With Respect to Coproducts

Omitted.

Item 5: Symmetric Lax Monoidality With Respect to Products

Omitted.

Item 6: Powersets as Sets of Functions

Omitted.

Item 7: Powersets as Sets of Relations

Omitted.

Item 8: As a Free Cocompletion: Universal Property

This is a rephrasing of ??.

Item 9: As a Free Cocompletion: Adjointness

Omitted.



### 005D 4.3 Direct Images

Let  $A$  and  $B$  be sets and let  $f: A \rightarrow B$  be a function.

#### 005E DEFINITION 4.3.1 ► DIRECT IMAGES

The **direct image function associated to  $f$**  is the function<sup>1</sup>

$$f_*: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by<sup>2,3</sup>

$$\begin{aligned} f_*(U) &\stackrel{\text{def}}{=} f(U) \\ &\stackrel{\text{def}}{=} \left\{ b \in B \mid \text{there exists some } a \in U \text{ such that } b = f(a) \right\} \\ &= \{ f(a) \in B \mid a \in U \} \end{aligned}$$

for each  $U \in \mathcal{P}(A)$ .

<sup>1</sup>*Further Notation:* Also written  $\exists_f: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$ . This notation comes from the fact that the following statements are equivalent, where  $b \in B$  and  $U \in \mathcal{P}(A)$ :

- We have  $b \in \exists_f(U)$ .

- There exists some  $a \in U$  such that  $f(a) = b$ .

<sup>2</sup>*Further Terminology:* The set  $f(U)$  is called the **direct image of  $U$  by  $f$** .

<sup>3</sup>We also have

$$f_*(U) = B \setminus f_1(A \setminus U);$$

see [Item 7](#) of [Proposition 4.3.3](#).

005F

**REMARK 4.3.2 ► UNWINDING DEFINITION 4.3.1**

Identifying subsets of  $A$  with functions from  $A$  to  $\{\text{true}, \text{false}\}$  via [Item 6](#) of [Proposition 4.2.3](#), we see that the direct image function associated to  $f$  is equivalently the function

$$f_*: \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by

$$\begin{aligned} f_*(\chi_U) &\stackrel{\text{def}}{=} \text{Lan}_f(\chi_U) \\ &= \text{colim} \left( \left( f \times \begin{smallmatrix} \rightarrow \\ (-1) \end{smallmatrix} \right) \xrightarrow{\text{pr}} A \xrightarrow{\chi_U} \{\text{t}, \text{f}\} \right) \\ &= \text{colim}_{\substack{a \in A \\ f(a) = -1}} (\chi_U(a)) \\ &= \bigvee_{\substack{a \in A \\ f(a) = -1}} (\chi_U(a)). \end{aligned}$$

So, in other words, we have

$$\begin{aligned} [f_*(\chi_U)](b) &= \bigvee_{\substack{a \in A \\ f(a) = b}} (\chi_U(a)) \\ &= \begin{cases} \text{true} & \text{if there exists some } a \in A \text{ such} \\ & \text{that } f(a) = b \text{ and } a \in U, \\ \text{false} & \text{otherwise} \end{cases} \\ &= \begin{cases} \text{true} & \text{if there exists some } a \in U \\ & \text{such that } f(a) = b, \\ \text{false} & \text{otherwise} \end{cases} \end{aligned}$$

for each  $b \in B$ .

005G

**PROPOSITION 4.3.3 ► PROPERTIES OF DIRECT IMAGES I**

Let  $f: A \rightarrow B$  be a function.

005H

1. *Functoriality.* The assignment  $U \mapsto f_*(U)$  defines a functor

$$f_*: (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

where

- *Action on Objects.* For each  $U \in \mathcal{P}(A)$ , we have

$$[f_*)(U) \stackrel{\text{def}}{=} f_*(U);$$

- *Action on Morphisms.* For each  $U, V \in \mathcal{P}(A)$ :  
 (★) If  $U \subset V$ , then  $f_*(U) \subset f_*(V)$ .

005J

2. *Triple Adjointness.* We have a triple adjunction

$$(f_* \dashv f^{-1} \dashv f_!): \quad \mathcal{P}(A) \begin{array}{c} \xrightarrow{f_*} \\ \perp \\ \xleftarrow{f^{-1}} \\ \perp \\ \xrightarrow{f_!} \end{array} \mathcal{P}(B),$$

witnessed by bijections of sets

$$\begin{aligned} \text{Hom}_{\mathcal{P}(B)}(f_*(U), V) &\cong \text{Hom}_{\mathcal{P}(A)}(U, f^{-1}(V)), \\ \text{Hom}_{\mathcal{P}(A)}(f^{-1}(U), V) &\cong \text{Hom}_{\mathcal{P}(A)}(U, f_!(V)), \end{aligned}$$

natural in  $U \in \mathcal{P}(A)$  and  $V \in \mathcal{P}(B)$  and (respectively)  $V \in \mathcal{P}(A)$  and  $U \in \mathcal{P}(B)$ , i.e. where:

- (a) The following conditions are equivalent:
- We have  $f_*(U) \subset V$ .
  - We have  $U \subset f^{-1}(V)$ .
- (b) The following conditions are equivalent:
- We have  $f^{-1}(U) \subset V$ .
  - We have  $U \subset f_!(V)$ .

005K

3. *Preservation of Colimits.* We have an equality of sets

$$f_*\left(\bigcup_{i \in I} U_i\right) = \bigcup_{i \in I} f_*(U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$ . In particular, we have equalities

$$\begin{aligned} f_*(U) \cup f_*(V) &= f_*(U \cup V), \\ f_*(\emptyset) &= \emptyset, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(A)$ .

005L

4. *Oplax Preservation of Limits.* We have an inclusion of sets

$$f_*\left(\bigcap_{i \in I} U_i\right) \subset \bigcap_{i \in I} f_*(U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$ . In particular, we have inclusions

$$\begin{aligned} f_*(U \cap V) &\subset f_*(U) \cap f_*(V), \\ f_*(A) &\subset B, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(A)$ .

005M

5. *Symmetric Strict Monoidality With Respect to Unions.* The direct image function of **Item 1** has a symmetric strict monoidal structure

$$(f_*, f_*^\otimes, f_{*|\mathbb{K}}^\otimes): (\mathcal{P}(A), \cup, \emptyset) \rightarrow (\mathcal{P}(B), \cup, \emptyset),$$

being equipped with equalities

$$\begin{aligned} f_{*|U,V}^\otimes: f_*(U) \cup f_*(V) &\xrightarrow{=} f_*(U \cup V), \\ f_{*|\mathbb{K}}^\otimes: \emptyset &\xrightarrow{=} \emptyset, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(A)$ .

005N

6. *Symmetric Oplax Monoidality With Respect to Intersections.* The direct image function of **Item 1** has a symmetric oplax monoidal structure

$$(f_*, f_*^\otimes, f_{*|\mathbb{P}}^\otimes): (\mathcal{P}(A), \cap, A) \rightarrow (\mathcal{P}(B), \cap, B),$$

being equipped with inclusions

$$\begin{aligned} f_{*|U,V}^\otimes: f_*(U \cap V) &\hookrightarrow f_*(U) \cap f_*(V), \\ f_{*|\mathbb{P}}^\otimes: f_*(A) &\hookrightarrow B, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(A)$ .

005P

7. *Relation to Direct Images With Compact Support.* We have

$$f_*(U) = B \setminus f_!(A \setminus U)$$

for each  $U \in \mathcal{P}(A)$ .

#### PROOF 4.3.4 ► PROOF OF PROPOSITION 4.3.3

Item 1: Functoriality

Clear.

Item 2: Triple Adjointness

This follows from **Kan Extensions**, ?? of ??.

Item 3: Preservation of Colimits

This follows from **Item 2** and **Categories**, ?? of ??.

Item 4: Oplax Preservation of Limits

Omitted.

Item 5: Symmetric Strict Monoidality With Respect to Unions

This follows from **Item 3**.

Item 6: Symmetric Oplax Monoidality With Respect to Intersections

This follows from ??.


Item 7: Relation to Direct Images With Compact Support

Applying  $f_!$  of  $f_!$  to  $A \setminus U$ , we have

$$\begin{aligned} f_!(A \setminus U) &= B \setminus f_*(A \setminus (A \setminus U)) \\ &= B \setminus f_*(U). \end{aligned}$$

Taking complements, we then obtain

$$\begin{aligned} f_*(U) &= B \setminus (B \setminus f_*(U)), \\ &= B \setminus f_!(A \setminus U), \end{aligned}$$

which finishes the proof. 

005Q

### PROPOSITION 4.3.5 ► PROPERTIES OF DIRECT IMAGES II

Let  $f: A \rightarrow B$  be a function.

005R

1. *Functionality I.* The assignment  $f \mapsto f_*$  defines a function

$$(-)_{*|A,B}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B)).$$

005S

2. *Functionality II.* The assignment  $f \mapsto f_*$  defines a function

$$(-)_{*|A,B}: \text{Sets}(A, B) \rightarrow \text{Pos}((\mathcal{P}(A), \subset), (\mathcal{P}(B), \subset)).$$

005T

3. *Interaction With Identities.* For each  $A \in \text{Obj}(\text{Sets})$ , we have

$$(\text{id}_A)_* = \text{id}_{\mathcal{P}(A)};$$

005U

4. *Interaction With Composition.* For each pair of composable functions  $f: A \rightarrow B$  and  $g: B \rightarrow C$ , we have

$$(g \circ f)_* = g_* \circ f_*,$$

$$\begin{array}{ccc} \mathcal{P}(A) & \xrightarrow{f_*} & \mathcal{P}(B) \\ & \searrow (g \circ f)_* & \downarrow g_* \\ & & \mathcal{P}(C). \end{array}$$

## PROOF 4.3.6 ► PROOF OF PROPOSITION 4.3.5

Item 1: Functionality I

Clear.

Item 2: Functionality II

Clear.

Item 3: Interaction With Identities

This follows from **Kan Extensions**, ?? of ??.

Item 4: Interaction With Composition

This follows from **Kan Extensions**, ?? of ??.

## 005V 4.4 Inverse Images

Let  $A$  and  $B$  be sets and let  $f: A \rightarrow B$  be a function.

## 005W DEFINITION 4.4.1 ► INVERSE IMAGES

The **inverse image function associated to  $f$**  is the function<sup>1</sup>

$$f^{-1}: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

defined by<sup>2</sup>

$$f^{-1}(V) \stackrel{\text{def}}{=} \{a \in A \mid \text{we have } f(a) \in V\}$$

for each  $V \in \mathcal{P}(B)$ .<sup>1</sup>Further Notation: Also written  $f^*: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$ .<sup>2</sup>Further Terminology: The set  $f^{-1}(V)$  is called the **inverse image of  $V$  by  $f$** .

## 005X REMARK 4.4.2 ► UNWINDING DEFINITION 4.4.1

Identifying subsets of  $B$  with functions from  $B$  to  $\{\text{true}, \text{false}\}$  via **Item 6 of Proposition 4.2.3**, we see that the inverse image function associated to  $f$  is equivalently the function

$$f^*: \mathcal{P}(B) \rightarrow \mathcal{P}(A)$$

defined by

$$f^*(\chi_V) \stackrel{\text{def}}{=} \chi_V \circ f$$

for each  $\chi_V \in \mathcal{P}(B)$ , where  $\chi_V \circ f$  is the composition

$$A \xrightarrow{f} B \xrightarrow{\chi_V} \{\text{true}, \text{false}\}$$

in Sets.

005Y

### PROPOSITION 4.4.3 ► PROPERTIES OF INVERSE IMAGES I

Let  $f: A \rightarrow B$  be a function.

005Z

1. *Functoriality.* The assignment  $V \mapsto f^{-1}(V)$  defines a functor

$$f^{-1}: (\mathcal{P}(B), \subset) \rightarrow (\mathcal{P}(A), \subset)$$

where

- *Action on Objects.* For each  $V \in \mathcal{P}(B)$ , we have

$$[f^{-1}](V) \stackrel{\text{def}}{=} f^{-1}(V);$$

- *Action on Morphisms.* For each  $U, V \in \mathcal{P}(B)$ :

$$(\star) \text{ If } U \subset V, \text{ then } f^{-1}(U) \subset f^{-1}(V).$$

0060

2. *Triple Adjointness.* We have a triple adjunction

$$(f_* \dashv f^{-1} \dashv f_!): \mathcal{P}(A) \begin{array}{c} \xrightarrow{f_*} \\ \perp \\ \xleftarrow{f^{-1}} \\ \perp \\ \xrightarrow{f_!} \end{array} \mathcal{P}(B),$$

witnessed by bijections of sets

$$\begin{aligned} \text{Hom}_{\mathcal{P}(B)}(f_*(U), V) &\cong \text{Hom}_{\mathcal{P}(A)}(U, f^{-1}(V)), \\ \text{Hom}_{\mathcal{P}(A)}(f^{-1}(U), V) &\cong \text{Hom}_{\mathcal{P}(A)}(U, f_!(V)), \end{aligned}$$

natural in  $U \in \mathcal{P}(A)$  and  $V \in \mathcal{P}(B)$  and (respectively)  $V \in \mathcal{P}(A)$  and  $U \in \mathcal{P}(B)$ , i.e. where:



(a) The following conditions are equivalent:

- i. We have  $f_*(U) \subset V$ ;
- ii. We have  $U \subset f^{-1}(V)$ ;

(b) The following conditions are equivalent:

- i. We have  $f^{-1}(U) \subset V$ .
- ii. We have  $U \subset f_!(V)$ .

0061

3. *Preservation of Colimits.* We have an equality of sets

$$f^{-1}\left(\bigcup_{i \in I} U_i\right) = \bigcup_{i \in I} f^{-1}(U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(B)^{\times I}$ . In particular, we have equalities

$$\begin{aligned} f^{-1}(U) \cup f^{-1}(V) &= f^{-1}(U \cup V), \\ f^{-1}(\emptyset) &= \emptyset, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(B)$ .

0062

4. *Preservation of Limits.* We have an equality of sets

$$f^{-1}\left(\bigcap_{i \in I} U_i\right) = \bigcap_{i \in I} f^{-1}(U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(B)^{\times I}$ . In particular, we have equalities

$$\begin{aligned} f^{-1}(U) \cap f^{-1}(V) &= f^{-1}(U \cap V), \\ f^{-1}(B) &= A, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(B)$ .

0063

5. *Symmetric Strict Monoidality With Respect to Unions.* The inverse image function of [Item 1](#) has a symmetric strict monoidal structure

$$(f^{-1}, f^{-1, \otimes}, f_{\#}^{-1, \otimes}) : (\mathcal{P}(B), \cup, \emptyset) \rightarrow (\mathcal{P}(A), \cup, \emptyset),$$

being equipped with equalities

$$f_{U,V}^{-1,\otimes} : f^{-1}(U) \cup f^{-1}(V) \xrightarrow{=} f^{-1}(U \cup V),$$

$$f_{\mathbb{K}}^{-1,\otimes} : \emptyset \xrightarrow{=} f^{-1}(\emptyset),$$

natural in  $U, V \in \mathcal{P}(B)$ .

0064

6. *Symmetric Strict Monoidality With Respect to Intersections.* The inverse image function of [Item 1](#) has a symmetric strict monoidal structure

$$(f^{-1}, f^{-1,\otimes}, f_{\mathbb{K}}^{-1,\otimes}) : (\mathcal{P}(B), \cap, B) \rightarrow (\mathcal{P}(A), \cap, A),$$

being equipped with equalities

$$f_{U,V}^{-1,\otimes} : f^{-1}(U) \cap f^{-1}(V) \xrightarrow{=} f^{-1}(U \cap V),$$

$$f_{\mathbb{K}}^{-1,\otimes} : A \xrightarrow{=} f^{-1}(B),$$

natural in  $U, V \in \mathcal{P}(B)$ .

#### PROOF 4.4.4 ► PROOF OF PROPOSITION 4.4.3

Item 1: Functoriality

Clear.

Item 2: Triple Adjointness

This follows from [Kan Extensions](#), ?? of ??.

Item 3: Preservation of Colimits

This follows from [Item 2](#) and [Categories](#), ?? of ??.

Item 4: Preservation of Limits

This follows from [Item 2](#) and [Categories](#), ?? of ??.

Item 5: Symmetric Strict Monoidality With Respect to Unions

This follows from [Item 3](#).

Item 6: Symmetric Strict Monoidality With Respect to Intersections

This follows from [Item 4](#).



## 0065 PROPOSITION 4.4.5 ► PROPERTIES OF INVERSE IMAGES II

Let  $f: A \rightarrow B$  be a function.

0066 1. *Functionality I.* The assignment  $f \mapsto f^{-1}$  defines a function

$$(-)_{A,B}^{-1}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(B), \mathcal{P}(A)).$$

0067 2. *Functionality II.* The assignment  $f \mapsto f^{-1}$  defines a function

$$(-)_{A,B}^{-1}: \text{Sets}(A, B) \rightarrow \text{Pos}((\mathcal{P}(B), \subset), (\mathcal{P}(A), \subset)).$$

0068 3. *Interaction With Identities.* For each  $A \in \text{Obj}(\text{Sets})$ , we have

$$\text{id}_A^{-1} = \text{id}_{\mathcal{P}(A)};$$

0069 4. *Interaction With Composition.* For each pair of composable functions  $f: A \rightarrow B$  and  $g: B \rightarrow C$ , we have

$$(g \circ f)^{-1} = f^{-1} \circ g^{-1},$$

$$\begin{array}{ccc} \mathcal{P}(C) & \xrightarrow{g^{-1}} & \mathcal{P}(B) \\ & \searrow (g \circ f)^{-1} & \downarrow f^{-1} \\ & & \mathcal{P}(A). \end{array}$$

## PROOF 4.4.6 ► PROOF OF PROPOSITION 4.4.5

Item 1: Functionality I

Clear.

Item 2: Functionality II

Clear.

Item 3: Interaction With Identities

This follows from **Categories**, ?? of ??.

Item 4: Interaction With Composition

This follows from **Categories**, ?? of ??.



## 006A 4.5 Direct Images With Compact Support

Let  $A$  and  $B$  be sets and let  $f: A \rightarrow B$  be a function.

## 006B DEFINITION 4.5.1 ► DIRECT IMAGES WITH COMPACT SUPPORT

The **direct image with compact support function associated to  $f$**  is the function<sup>1</sup>

$$f_! : \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by<sup>2,3</sup>

$$\begin{aligned} f_!(U) &\stackrel{\text{def}}{=} \left\{ b \in B \mid \begin{array}{l} \text{for each } a \in A, \text{ if we have} \\ f(a) = b, \text{ then } a \in U \end{array} \right\} \\ &= \{ b \in B \mid \text{we have } f^{-1}(b) \subset U \} \end{aligned}$$

for each  $U \in \mathcal{P}(A)$ .

<sup>1</sup>*Further Notation:* Also written  $\forall_f : \mathcal{P}(A) \rightarrow \mathcal{P}(B)$ . This notation comes from the fact that the following statements are equivalent, where  $b \in B$  and  $U \in \mathcal{P}(A)$ :

- We have  $b \in \forall_f(U)$ .
- For each  $a \in A$ , if  $b = f(a)$ , then  $a \in U$ .

<sup>2</sup>*Further Terminology:* The set  $f_!(U)$  is called the **direct image with compact support of  $U$  by  $f$** .

<sup>3</sup>We also have

$$f_!(U) = B \setminus f_*(A \setminus U);$$

see Item 7 of Proposition 4.5.5.

## 006C REMARK 4.5.2 ► UNWINDING DEFINITION 4.5.1

Identifying subsets of  $A$  with functions from  $A$  to  $\{\text{true}, \text{false}\}$  via Item 6 of Proposition 4.2.3, we see that the direct image with compact support function associated to  $f$  is equivalently the function

$$f_! : \mathcal{P}(A) \rightarrow \mathcal{P}(B)$$

defined by

$$\begin{aligned}
 f_!(\chi_U) &\stackrel{\text{def}}{=} \text{Ran}_f(\chi_U) \\
 &= \lim \left( \left( \underline{-1} \right) \vec{\times} f \right) \xrightarrow{\text{pr}} A \xrightarrow{\chi_U} \{\text{true}, \text{false}\} \\
 &= \lim_{\substack{a \in A \\ f(a) = -1}} (\chi_U(a)) \\
 &= \bigwedge_{\substack{a \in A \\ f(a) = -1}} (\chi_U(a)).
 \end{aligned}$$

So, in other words, we have

$$\begin{aligned}
 [f_!(\chi_U)](b) &= \bigwedge_{\substack{a \in A \\ f(a) = b}} (\chi_U(a)) \\
 &= \begin{cases} \text{true} & \text{if, for each } a \in A \text{ such that} \\ & f(a) = b, \text{ we have } a \in U, \\ \text{false} & \text{otherwise} \end{cases} \\
 &= \begin{cases} \text{true} & \text{if } f^{-1}(b) \subset U \\ \text{false} & \text{otherwise} \end{cases}
 \end{aligned}$$

for each  $b \in B$ .

006D

#### DEFINITION 4.5.3 ► THE IMAGE AND COMPLEMENT PARTS OF $f_!$

Let  $U$  be a subset of  $A$ .<sup>1,2</sup>

1. The **image part of the direct image with compact support**  $f_!(U)$  of  $U$  is the set  $f_{!,\text{im}}(U)$  defined by

$$\begin{aligned}
 f_{!,\text{im}}(U) &\stackrel{\text{def}}{=} f_!(U) \cap \text{Im}(f) \\
 &= \left\{ b \in B \mid \begin{array}{l} \text{we have } f^{-1}(b) \subset U \\ \text{and } f^{-1}(b) \neq \emptyset \end{array} \right\}.
 \end{aligned}$$

2. The **complement part of the direct image with compact support**  $f_!(U)$  of

$U$  is the set  $f_{!,\text{cp}}(U)$  defined by

$$\begin{aligned} f_{!,\text{cp}}(U) &\stackrel{\text{def}}{=} f_!(U) \cap (B \setminus \text{Im}(f)) \\ &= B \setminus \text{Im}(f) \\ &= \left\{ b \in B \mid \begin{array}{l} \text{we have } f^{-1}(b) \subset U \\ \text{and } f^{-1}(b) = \emptyset \end{array} \right\} \\ &= \{ b \in B \mid f^{-1}(b) = \emptyset \}. \end{aligned}$$

<sup>1</sup>Note that we have

$$f_!(U) = f_{!,\text{im}}(U) \cup f_{!,\text{cp}}(U),$$

as

$$\begin{aligned} f_!(U) &= f_!(U) \cap B \\ &= f_!(U) \cap (\text{Im}(f) \cup (B \setminus \text{Im}(f))) \\ &= (f_!(U) \cap \text{Im}(f)) \cup (f_!(U) \cap (B \setminus \text{Im}(f))) \\ &\stackrel{\text{def}}{=} f_{!,\text{im}}(U) \cup f_{!,\text{cp}}(U). \end{aligned}$$

<sup>2</sup>In terms of the meet computation of  $f_!(U)$  of [Remark 4.5.2](#), namely

$$f_!(\chi_U) = \bigwedge_{\substack{a \in A \\ f(a) = -1}} (\chi_U(a)),$$

we see that  $f_{!,\text{im}}$  corresponds to meets indexed over nonempty sets, while  $f_{!,\text{cp}}$  corresponds to meets indexed over the empty set.

006E

#### EXAMPLE 4.5.4 ► EXAMPLES OF DIRECT IMAGES WITH COMPACT SUPPORT

Here are some examples of direct images with compact support.

1. *The Multiplication by Two Map on the Natural Numbers.* Consider the function  $f: \mathbb{N} \rightarrow \mathbb{N}$  given by

$$f(n) \stackrel{\text{def}}{=} 2n$$

for each  $n \in \mathbb{N}$ . Since  $f$  is injective, we have

$$\begin{aligned} f_{!,\text{im}}(U) &= f_*(U) \\ f_{!,\text{cp}}(U) &= \{\text{odd natural numbers}\} \end{aligned}$$

for any  $U \subset \mathbb{N}$ .

2. *Parabolas.* Consider the function  $f: \mathbb{R} \rightarrow \mathbb{R}$  given by

$$f(x) \stackrel{\text{def}}{=} x^2$$

for each  $x \in \mathbb{R}$ . We have

$$f_{!,\text{cp}}(U) = \mathbb{R}_{<0}$$

for any  $U \subset \mathbb{R}$ . Moreover, since  $f^{-1}(x) = \{-\sqrt{x}, \sqrt{x}\}$ , we have e.g.:

$$\begin{aligned} f_{!,\text{im}}([0, 1]) &= \{0\}, \\ f_{!,\text{im}}([-1, 1]) &= [0, 1], \\ f_{!,\text{im}}([1, 2]) &= \emptyset, \\ f_{!,\text{im}}([-2, -1] \cup [1, 2]) &= [1, 4]. \end{aligned}$$

3. *Circles*. Consider the function  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  given by

$$f(x, y) \stackrel{\text{def}}{=} x^2 + y^2$$

for each  $(x, y) \in \mathbb{R}^2$ . We have

$$f_{!,\text{cp}}(U) = \mathbb{R}_{<0}$$

for any  $U \subset \mathbb{R}^2$ , and since

$$f^{-1}(r) = \begin{cases} \text{a circle of radius } r \text{ about the origin} & \text{if } r > 0, \\ \{(0, 0)\} & \text{if } r = 0, \\ \emptyset & \text{if } r < 0, \end{cases}$$

we have e.g.:

$$\begin{aligned} f_{!,\text{im}}([-1, 1] \times [-1, 1]) &= [0, 1], \\ f_{!,\text{im}}(([-1, 1] \times [-1, 1]) \setminus [-1, 1] \times \{0\}) &= \emptyset. \end{aligned}$$

006F

#### PROPOSITION 4.5.5 ► PROPERTIES OF DIRECT IMAGES WITH COMPACT SUPPORT I

Let  $f: A \rightarrow B$  be a function.

006G

1. *Functoriality*. The assignment  $U \mapsto f_!(U)$  defines a functor

$$f_!: (\mathcal{P}(A), \subset) \rightarrow (\mathcal{P}(B), \subset)$$

where

- *Action on Objects.* For each  $U \in \mathcal{P}(A)$ , we have

$$[f_!](U) \stackrel{\text{def}}{=} f_!(U);$$

- *Action on Morphisms.* For each  $U, V \in \mathcal{P}(A)$ :

(★) If  $U \subset V$ , then  $f_!(U) \subset f_!(V)$ .

006H

2. *Triple Adjointness.* We have a triple adjunction

$$(f_* \dashv f^{-1} \dashv f_!): \quad \mathcal{P}(A) \begin{array}{c} \xrightarrow{f_*} \\ \perp \\ \xleftarrow{f^{-1}} \\ \perp \\ \xrightarrow{f_!} \end{array} \mathcal{P}(B),$$

witnessed by bijections of sets

$$\begin{aligned} \text{Hom}_{\mathcal{P}(B)}(f_*(U), V) &\cong \text{Hom}_{\mathcal{P}(A)}(U, f^{-1}(V)), \\ \text{Hom}_{\mathcal{P}(A)}(f^{-1}(U), V) &\cong \text{Hom}_{\mathcal{P}(A)}(U, f_!(V)), \end{aligned}$$

natural in  $U \in \mathcal{P}(A)$  and  $V \in \mathcal{P}(B)$  and (respectively)  $V \in \mathcal{P}(A)$  and  $U \in \mathcal{P}(B)$ , i.e. where:

- (a) The following conditions are equivalent:

- We have  $f_*(U) \subset V$ ;
- We have  $U \subset f^{-1}(V)$ ;

- (b) The following conditions are equivalent:

- We have  $f^{-1}(U) \subset V$ .
- We have  $U \subset f_!(V)$ .

006J

3. *Lax Preservation of Colimits.* We have an inclusion of sets

$$\bigcup_{i \in I} f_!(U_i) \subset f_!\left(\bigcup_{i \in I} U_i\right),$$



natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$ . In particular, we have inclusions

$$\begin{aligned} f_!(U) \cup f_!(V) &\hookrightarrow f_!(U \cup V), \\ \emptyset &\hookrightarrow f_!(\emptyset), \end{aligned}$$

natural in  $U, V \in \mathcal{P}(A)$ .

006K

4. *Preservation of Limits.* We have an equality of sets

$$f_!\left(\bigcap_{i \in I} U_i\right) = \bigcap_{i \in I} f_!(U_i),$$

natural in  $\{U_i\}_{i \in I} \in \mathcal{P}(A)^{\times I}$ . In particular, we have equalities

$$\begin{aligned} f^{-1}(U \cap V) &= f_!(U) \cap f^{-1}(V), \\ f_!(A) &= B, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(A)$ .

006L

5. *Symmetric Lax Monoidality With Respect to Unions.* The direct image with compact support function of **Item 1** has a symmetric lax monoidal structure

$$(f_!, f_!^\otimes, f_{!|\mathbb{K}}^\otimes): (\mathcal{P}(A), \cup, \emptyset) \rightarrow (\mathcal{P}(B), \cup, \emptyset),$$

being equipped with inclusions

$$\begin{aligned} f_{!|U,V}^\otimes: f_!(U) \cup f_!(V) &\hookrightarrow f_!(U \cup V), \\ f_{!|\mathbb{K}}^\otimes: \emptyset &\hookrightarrow f_!(\emptyset), \end{aligned}$$

natural in  $U, V \in \mathcal{P}(A)$ .

006M

6. *Symmetric Strict Monoidality With Respect to Intersections.* The direct image function of **Item 1** has a symmetric strict monoidal structure

$$(f_!, f_!^\otimes, f_{!|\mathbb{K}}^\otimes): (\mathcal{P}(A), \cap, A) \rightarrow (\mathcal{P}(B), \cap, B),$$

being equipped with equalities

$$\begin{aligned} f_{!|U,V}^\otimes: f_!(U \cap V) &\xrightarrow{=} f_!(U) \cap f_!(V), \\ f_{!|\mathbb{K}}^\otimes: f_!(A) &\xrightarrow{=} B, \end{aligned}$$

natural in  $U, V \in \mathcal{P}(A)$ .

006N

7. *Relation to Direct Images.* We have

$$f_!(U) = B \setminus f_*(A \setminus U)$$

for each  $U \in \mathcal{P}(A)$ .

006P

8. *Interaction With Injections.* If  $f$  is injective, then we have

$$f_{!,\text{im}}(U) = f_*(U),$$

$$f_{!,\text{cp}}(U) = B \setminus \text{Im}(f),$$

$$\begin{aligned} f_!(U) &= f_{!,\text{im}}(U) \cup f_{!,\text{cp}}(U) \\ &= f_*(U) \cup (B \setminus \text{Im}(f)) \end{aligned}$$

for each  $U \in \mathcal{P}(A)$ .

006Q

9. *Interaction With Surjections.* If  $f$  is surjective, then we have

$$f_{!,\text{im}}(U) \subset f_*(U),$$

$$f_{!,\text{cp}}(U) = \emptyset,$$

$$f_!(U) \subset f_*(U)$$

for each  $U \in \mathcal{P}(A)$ .

## PROOF 4.5.6 ► PROOF OF PROPOSITION 4.5.5

Item 1: Functoriality

Clear.

Item 2: Triple Adjointness

This follows from [Kan Extensions](#), ?? of ??.

Item 3: Lax Preservation of Colimits

Omitted.

Item 4: Preservation of Limits

Omitted. This follows from [Item 2](#) and [Categories](#), ?? of ??.

Item 5: Symmetric Lax Monoidality With Respect to Unions

This follows from ??.

Item 6: Symmetric Strict Monoidality With Respect to Intersections

This follows from Item 4.

Item 7: Relation to Direct Images

We claim that  $f_!(U) = B \setminus f_*(A \setminus U)$ .

- *The First Implication.* We claim that

$$f_!(U) \subset B \setminus f_*(A \setminus U).$$

Let  $b \in f_!(U)$ . We need to show that  $b \notin f_*(A \setminus U)$ , i.e. that there is no  $a \in A \setminus U$  such that  $f(a) = b$ .

This is indeed the case, as otherwise we would have  $a \in f^{-1}(b)$  and  $a \notin U$ , contradicting  $f^{-1}(b) \subset U$  (which holds since  $b \in f_!(U)$ ).

Thus  $b \in B \setminus f_*(A \setminus U)$ .

- *The Second Implication.* We claim that

$$B \setminus f_*(A \setminus U) \subset f_!(U).$$

Let  $b \in B \setminus f_*(A \setminus U)$ . We need to show that  $b \in f_!(U)$ , i.e. that  $f^{-1}(b) \subset U$ .

Since  $b \notin f_*(A \setminus U)$ , there exists no  $a \in A \setminus U$  such that  $b = f(a)$ , and hence  $f^{-1}(b) \subset U$ .

Thus  $b \in f_!(U)$ .

This finishes the proof of Item 7.

Item 8: Interaction With Injections

Clear.

Item 9: Interaction With Surjections

Clear.



006R

**PROPOSITION 4.5.7 ► PROPERTIES OF DIRECT IMAGES WITH COMPACT SUPPORT II**

Let  $f: A \rightarrow B$  be a function.

006S

1. *Functionality I.* The assignment  $f \mapsto f_!$  defines a function

$$(-)_{!|A,B}: \text{Sets}(A, B) \rightarrow \text{Sets}(\mathcal{P}(A), \mathcal{P}(B)).$$

006T

2. *Functionality II.* The assignment  $f \mapsto f_!$  defines a function

$$(-)_{!|A,B}: \text{Sets}(A, B) \rightarrow \text{Pos}((\mathcal{P}(A), \subset), (\mathcal{P}(B), \subset)).$$

006U

3. *Interaction With Identities.* For each  $A \in \text{Obj}(\text{Sets})$ , we have

$$(\text{id}_A)_! = \text{id}_{\mathcal{P}(A)};$$

006V

4. *Interaction With Composition.* For each pair of composable functions  $f: A \rightarrow B$  and  $g: B \rightarrow C$ , we have

$$(g \circ f)_! = g_! \circ f_!,$$

$$\begin{array}{ccc} \mathcal{P}(A) & \xrightarrow{f_!} & \mathcal{P}(B) \\ & \searrow (g \circ f)_! & \downarrow g_! \\ & & \mathcal{P}(C). \end{array}$$

**PROOF 4.5.8 ► PROOF OF PROPOSITION 4.5.7**

Item 1: Functionality I

Clear.

Item 2: Functionality II

Clear.

Item 3: Interaction With Identities

This follows from **Kan Extensions**, ?? of ??.

Item 4: Interaction With Composition

This follows from **Kan Extensions**, ?? of ??.



## Appendices

## A Other Chapters

### Set Theory

1. [Sets](#)
2. [Constructions With Sets](#)
3. [Pointed Sets](#)
4. [Tensor Products of Pointed Sets](#)
5. [Indexed and Fibred Sets](#)
6. [Relations](#)
7. [Spans](#)
8. [Posets](#)

### Category Theory

9. [Categories](#)
10. [Constructions With Categories](#)
11. [Kan Extensions](#)

### Bicategories

12. [Bicategories](#)
13. [Internal Adjunctions](#)

### Internal Category Theory

14. [Internal Categories](#)

### Cyclic Stuff

15. [The Cycle Category](#)

### Cubical Stuff

16. [The Cube Category](#)

### Globular Stuff

17. [The Globe Category](#)

### Cellular Stuff

18. [The Cell Category](#)

### Monoids

19. [Monoids](#)
20. [Constructions With Monoids](#)

### Monoids With Zero

21. [Monoids With Zero](#)
22. [Constructions With Monoids With Zero](#)

### Groups

23. [Groups](#)
24. [Constructions With Groups](#)

### Hyper Algebra

25. [Hypermonoids](#)
26. [Hypergroups](#)
27. [Hypersemirings and Hyperrings](#)
28. [Quantaes](#)

### Near-Rings

29. [Near-Semirings](#)
30. [Near-Rings](#)

### Real Analysis

31. [Real Analysis in One Variable](#)
32. [Real Analysis in Several Variables](#)

### Measure Theory

33. [Measurable Spaces](#)

34. Measures and Integration

36. Itô Calculus

**Probability Theory**

37. Stochastic Differential Equations

34. Probability Theory

**Differential Geometry**

**Stochastic Analysis**

38. Topological and Smooth Manifolds

35. Stochastic Processes, Martingales,  
and Brownian Motion

**Schemes**

39. Schemes