

# JEE Main 2020 Paper

Date of Exam: 7<sup>th</sup> January (Shift II)

Time: 2:30 pm – 5:30 pm

Subject: Physics

---

1. A box weighs  $196\text{ N}$  on a spring balance at the North Pole. Its weight recorded on the same balance if it is shifted to the equator is close to (Take  $g = 10\text{ m/s}^2$  at the North Pole and radius of the Earth =  $6400\text{ km}$ )

- |                      |                      |
|----------------------|----------------------|
| a. $194.32\text{ N}$ | b. $194.66\text{ N}$ |
| c. $195.32\text{ N}$ | d. $195.66\text{ N}$ |

2. In a building, there are 15 bulbs of  $45\text{ W}$ , 15 bulbs of  $100\text{ W}$ , 15 small fans of  $10\text{ W}$  and 2 heaters of  $1\text{ kW}$ . The voltage of electric main is  $220\text{ V}$ . The minimum fuse capacity (rated value) of the building will be approximately

- |                  |                  |
|------------------|------------------|
| a. $10\text{ A}$ | b. $20\text{ A}$ |
| c. $25\text{ A}$ | d. $15\text{ A}$ |

# JEE Main 2020 Paper

3. Under an adiabatic process, the volume of an ideal gas gets doubled. Consequently, the mean collision time between the gas molecules changes from  $\tau_1$  to  $\tau_2$ . If  $\frac{C_p}{C_v} = \gamma$  for this gas, then a good estimate for  $\frac{\tau_2}{\tau_1}$  is given by
- a.  $\frac{1}{2}$
- b.  $\left(\frac{1}{2}\right)^{\frac{\gamma+1}{2}}$
- c.  $\left(\frac{1}{2}\right)^\gamma$
- d. 2

4. A mass of  $10\text{ kg}$  is suspended by a rope of length  $4\text{ m}$ , from the ceiling. A force  $F$  is applied horizontally at the mid-point of the rope such that the top half of the rope makes an angle of  $45^\circ$  with the vertical. Then  $F$  equals (Take  $g = 10\text{ m/s}^2$  and rope to be massless)
- a.  $100\text{ N}$                       b.  $90\text{ N}$
- c.  $75\text{ N}$                         d.  $70\text{ N}$

## JEE Main 2020 Paper

5. Mass per unit area of a circular disc of radius  $a$  depends on the distance  $r$  from its centre as  $\sigma(r) = A + Br$ . The moment of inertia of the disc about the axis, perpendicular to the plane and passing through its centre is

a.  $2\pi a^4 \left( \frac{A}{4} + \frac{aB}{5} \right)$

b.  $2\pi a^4 \left( \frac{aA}{4} + \frac{B}{5} \right)$

c.  $\pi a^4 \left( \frac{A}{4} + \frac{aB}{5} \right)$

d.  $2\pi a^4 \left( \frac{A}{4} + \frac{B}{5} \right)$

6. Two ideal Carnot engines operate in cascade (all heat given up by one engine is used by the other engine to produce work) between temperatures  $T_1$  and  $T_2$ . The temperature of the hot reservoir of the first engine is  $T_1$  and the temperature of the cold reservoir of the second engine is  $T_2$ .  $T$  is the temperature of the sink of first engine which is also the source for the second engine. How is  $T$  related to  $T_1$  and  $T_2$  if both the engines perform equal amount of work?

a.  $T = \frac{2T_1 T_2}{T_1 + T_2}$

b.  $T = \frac{T_1 + T_2}{2}$

c.  $T = 0$

d.  $T = \sqrt{T_1 T_2}$

# JEE Main 2020 Paper

7. The activity of a radioactive substance falls from  $700 \text{ s}^{-1}$  to  $500 \text{ s}^{-1}$  in 30 minutes. Its half-life is close to

- a. 66 min
- c. 52 min

- b. 62 min
- d. 72 min

8. In a Young's double slit experiment, the separation between the slits is  $0.15 \text{ mm}$ . In the experiment, a source of light of wavelength  $589 \text{ nm}$  is used and the interference pattern is observed on a screen kept  $1.5 \text{ m}$  away. The separation between the successive bright fringes on the screen is

- a.  $5.9 \text{ mm}$
- c.  $6.9 \text{ mm}$

- b.  $3.9 \text{ mm}$
- d.  $4.9 \text{ mm}$

## JEE Main 2020 Paper

9. An ideal fluid flows (laminar flow) through a pipe of non-uniform diameter. The maximum and minimum diameters of the pipes are  $6.4\text{ cm}$  and  $4.8\text{ cm}$ , respectively. The ratio of minimum and maximum velocities of fluid in this pipe is

a.  $\sqrt{\frac{3}{2}}$

b.  $\frac{9}{16}$

c.  $\frac{3}{4}$

d.  $\frac{3}{4}$


10. In the figure, potential difference between  $a$  and  $b$  is

a.  $0\text{ V}$

b.  $15\text{ V}$

c.  $10\text{ V}$

d.  $5\text{ V}$



## JEE Main 2020 Paper

11. A particle of mass  $m$  and charge  $q$  has an initial velocity  $\vec{v} = v_0 \hat{j}$ . If an electric field  $\vec{E} = E_0 \hat{i}$  and magnetic field  $\vec{B} = B_0 \hat{i}$  act on the particle, its speed will double after a time

a.  $\frac{\sqrt{3}mv_0}{qE_0}$   
c.  $\frac{3mv_0}{qE_0}$

b.  $\frac{\sqrt{2}mv_0}{qE_0}$   
d.  $\frac{2mv_0}{2qE_0}$

12. A stationary observer receives sound from two identical tuning forks, one of which approaches and the other one recedes with the same speed (much less than the speed of sound). The observer hears 2 beats/sec. The oscillation frequency of each tuning fork is  $\nu_0 = 1400 \text{ Hz}$  and the velocity of sound in air is  $350 \text{ m/s}$ . The speed of each tuning fork is close to

a.  $\frac{1}{4} \text{ m/s}$   
c.  $\frac{1}{2} \text{ m/s}$

b.  $1 \text{ m/s}$   
d.  $\frac{1}{8} \text{ m/s}$

# JEE Main 2020 Paper

13. An electron (of mass  $m$ ) and a photon have the same energy  $E$  in the range of few  $eV$ . The ratio of the de Broglie wavelength associated with the electron and the wavelength of the photon is. ( $c$  = speed of light in vacuum)

a.  $\left(\frac{E}{2m}\right)^{\frac{1}{2}}$

b.  $\frac{1}{c} \left( \frac{2E}{m} \right)^{\frac{1}{2}}$

$$\text{c. } c(2mE)^{\frac{1}{2}}$$

d.  $\frac{1}{c} \left( \frac{E}{2m} \right)^{\frac{1}{2}}$

14. A planar loop of wire rotates in a uniform magnetic field. Initially at  $t = 0$ , the plane of the loop is perpendicular to the magnetic field. If it rotates with a period of 10 s about an axis in its plane, then the magnitude of induced emf will be maximum and minimum, respectively at

a. 2.5 *sec* and 5 *sec*

b. 5 *sec* and 7.5 *sec*

c. 2.5 sec and 7.5 sec

d. 5 *sec* and 10 *sec*

15. The electric field of a plane electromagnetic wave is given by  $\vec{E}(t) = E_0 \frac{\hat{i}+j}{\sqrt{2}} \cos(kz + \omega t)$ . At  $t = 0$ , a positively charged particle is at the point  $(x, y, z) = (0, 0, \pi/k)$ . If its instantaneous velocity at  $t = 0$  is  $v_0 \hat{k}$ , the force acting on it due to the wave is

## JEE Main 2020 Paper

- a. zero
- b. antiparallel to  $\frac{i+j}{\sqrt{2}}$
- c. parallel to  $\frac{i+j}{\sqrt{2}}$
- d. parallel to  $\hat{k}$

16. A thin lens made of glass (refractive index = 1.5) of focal length  $f = 16 \text{ cm}$  is immersed in a liquid of refractive index 1.42. If its focal length in liquid is  $f_l$ , then the ratio  $f_l/f$  is closest to the integer

- a. 9
- b. 17
- c. 1
- d. 5

17. An elevator in a building can carry a maximum of 10 persons, with the average mass of each person being  $68 \text{ kg}$ . The mass of the elevator itself is  $920 \text{ kg}$  and it moves with a constant speed of  $3 \text{ m/s}$ . The frictional force opposing the motion is  $6000 \text{ N}$ . If the elevator is moving up with its full capacity, the power delivered by the motor to the elevator ( $g = 10 \text{ m/s}^2$ ) must be at least


- a.  $66000 \text{ W}$
- b.  $63360 \text{ W}$
- c.  $48000 \text{ W}$
- d.  $56300 \text{ W}$



## JEE Main 2020 Paper

18. The figure gives experimentally measured  $B$  vs  $H$  variation in a ferromagnetic material. The retentivity, coercivity and saturation, respectively, of the material are

- a. 1.5 T, 50 A/m, 1 T
- b. 1 T, 50 A/m, 1.5 T
- c. 1.5 T, 50 A/m, 1 T
- d. 150 A/m, 1 T, 1.5 T



## JEE Main 2020 Paper

19. An emf of  $20\text{ V}$  is applied at time  $t = 0$  to a circuit containing in series  $10\text{ mH}$  inductor and  $5\ \Omega$  resistor. The ratio of the currents at time  $t = \infty$  and  $t = 40\text{ s}$  is close to (take  $e^2 = 7.389$ )

- |         |         |
|---------|---------|
| a. 1.06 | b. 1.46 |
| c. 1.15 | d. 0.84 |

20. The dimension of  $\frac{B^2}{2\mu_0}$ , where  $B$  is magnetic field and  $\mu_0$  is the magnetic permeability of vacuum, is


- |                    |                 |
|--------------------|-----------------|
| a. $ML^{-1}T^{-2}$ | b. $ML^2T^{-2}$ |
| c. $MLT^{-2}$      | d. $ML^2T^{-1}$ |

21. A  $60\text{ pF}$  capacitor is fully charged by a  $20\text{ V}$  supply. It is then disconnected from the supply and is connected to another uncharged  $60\text{ pF}$  capacitor in parallel. The electrostatic energy that is lost in this process by the time the charge is redistributed between them is (in  $nJ$ ) \_\_\_\_\_.

## JEE Main 2020 Paper

22.  $M$  grams of steam at  $100^{\circ}\text{C}$  is mixed with  $200\text{ g}$  of ice at its melting point in a thermally insulated container. If it produces liquid water at  $40^{\circ}\text{C}$  [heat of vaporization of water is  $540\text{ cal/g}$  and heat of fusion of ice is  $80\text{ cal/g}$ ], the value of  $M$  is \_\_\_\_\_.
23. Consider a uniform cubical box of side  $a$  on a rough floor that is to be moved by applying minimum possible force  $F$  at a point  $b$  above its centre of mass (see figure). If the coefficient of friction is  $\mu = 0.4$ , the maximum value of  $100 \times \frac{b}{a}$  for the box not to topple before moving is \_\_\_\_\_.

## JEE Main 2020 Paper



24. The sum of two forces  $\vec{P}$  and  $\vec{Q}$  is  $\vec{R}$  such that  $|\vec{R}| = |\vec{P}|$ . The angle  $\theta$  (in degrees) that the resultant of  $2\vec{P}$  and  $\vec{Q}$  will make with  $\vec{Q}$  is \_\_\_\_\_

## JEE Main 2020 Paper

25. The balancing length for a cell is  $560\text{ cm}$  in a potentiometer experiment. When an external resistance of  $10\ \Omega$  is connected in parallel to the cell, the balancing length changes by  $60\text{ cm}$ . If the internal resistance of the cell is  $\frac{N}{10}\ \Omega$ , the value of  $N$  is \_\_\_\_\_

# JEE Main 2020 Paper


Date: 7<sup>th</sup> January 2020

Time: 02:30 PM – 05:30 PM


Subject: Chemistry

1. Consider the following reactions:


A.




B.



C.



D.



Which of these reactions are possible?

a. A and D


b. B and D

c. B, C and D

d. A and B


# JEE Main 2020 Paper

2. In the following reaction sequence,




The major product B is:


a.




c.



b.




d.




# JEE Main 2020 Paper

3. For the following reactions,



Where,



$k_s$  and  $k_e$  are, respectively, the rate constants for substitution and elimination, and  $\mu = \frac{k_s}{k_e}$ , the correct option is \_\_\_\_.

- $\mu_A > \mu_B$  and  $k_e(\text{A}) > k_e(\text{B})$
- $\mu_B > \mu_A$  and  $k_e(\text{A}) > k_e(\text{B})$
- $\mu_A > \mu_B$  and  $k_e(\text{B}) > k_e(\text{A})$
- $\mu_B > \mu_A$  and  $k_e(\text{B}) > k_e(\text{A})$


4. Which of the following statements is correct?

- Gluconic acid can form cyclic (acetal/hemiacetal) structure
- Gluconic acid is a dicarboxylic acid
- Gluconic acid is obtained by oxidation of glucose with  $\text{HNO}_3$
- Gluconic acid is a partial oxidation product of glucose




# JEE Main 2020 Paper


5. The correct order of stability for the following alkoxides is:



(A)



(B)



(C)

a. (C) > (A) > (B)


b. (C) > (B) > (A)

c. (B) > (A) > (C)


d. (B) > (C) > (A)

# JEE Main 2020 Paper


6. In the following reaction sequence, structures of A and B, respectively will be:




a.




b.



c.



d.



7. A chromatography column, packed with silica gel as stationary phase, was used to separate a mixture of compounds consisting of (A) benzanilide, (B) aniline and (C) acetophenone. When the column is eluted with a mixture of solvents, hexane : ethyl acetate (20:80), the sequence of obtained compounds is:

- (B), (A) and (C)
- (C), (A) and (B)
- (B), (C) and (A)
- (A), (B) and (C)

# JEE Main 2020 Paper

8. The number of possible optical isomers for the complexes  $[MA_2B_2]$  with  $sp^3$  or  $dsp^2$  hybridized metal atom, respectively, is:

Note: A and B are unidentate neutral and unidentate monoanionic ligands, respectively.

- |            |            |
|------------|------------|
| a. 0 and 1 | c. 2 and 2 |
| b. 0 and 0 | d. 0 and 2 |

9. The bond order and magnetic characteristics of  $CN^-$  are:


- |                     |                      |
|---------------------|----------------------|
| a. 3, paramagnetic  | c. 3, diamagnetic    |
| b. 2.5, diamagnetic | d. 2.5, paramagnetic |

# JEE Main 2020 Paper


10. The equation that is incorrect is:

- a.  $\Delta_m^0 \text{NaBr} - \Delta_m^0 \text{NaI} = \Delta_m^0 \text{KBr} - \Delta_m^0 \text{NaBr}$
- b.  $\Delta_m^0 \text{NaBr} - \Delta_m^0 \text{NaCl} = \Delta_m^0 \text{KBr} - \Delta_m^0 \text{KCl}$
- c.  $\Delta_m^0 \text{KCl} - \Delta_m^0 \text{NaCl} = \Delta_m^0 \text{KBr} - \Delta_m^0 \text{NaBr}$
- d.  $\Delta_m^0 \text{H}_2\text{O} = \Delta_m^0 \text{HCl} + \Delta_m^0 \text{NaOH} - \Delta_m^0 \text{NaBr}$

11. In the following reactions, product (A) and (B), respectively, are:



(hot & conc.)



(dry)

- a.  $\text{NaClO}_3$  and  $\text{Ca(ClO}_3)_2$
- b.  $\text{NaOCl}$  and  $\text{Ca(ClO}_3)_2$
- c.  $\text{NaOCl}$  and  $\text{Ca(OCl)}_2$
- d.  $\text{NaClO}_3$  and  $\text{Ca(OCl)}_2$

12. Two open beakers one containing a solvent and the other containing a mixture of that solvent with a non-volatile solute are together sealed in a container. Over time:

- a. the volume of the solution and the solvent does not change
- b. the volume of the solution increases and the volume of the solvent decreases
- c. the volume of the solution decreases and the volume of the solvent increases
- d. the volume of the solution does not change and the volume of the solvent decreases

# JEE Main 2020 Paper

13. The refining method used when the metal and the impurities have low and high melting temperatures, respectively, is:

- |                          |                  |
|--------------------------|------------------|
| a. vapour phase refining | b. distillation  |
| c. liquation             | d. zone refining |

14. Among statements I-IV, the correct ones are:

- |  |                       |
|--|-----------------------|
| I. Decomposition of hydrogen peroxide gives dioxygen   |                       |
| II. Like hydrogen peroxide, compounds, such as $\text{KClO}_3$ , $\text{Pb}(\text{NO}_3)_2$ and $\text{NaNO}_3$ when heated liberate dioxygen. |                       |
| III. 2-Ethylanthraquinone is useful for the industrial preparation of hydrogen peroxide.   |                       |
| IV. Hydrogen peroxide is used for the manufacture of sodium perborate  |                       |
| a. I, II, III and IV   | b. I, II and III only |
| c. I, III and IV only  | d. I and III only     |


# JEE Main 2020 Paper

15. The redox reaction among the following is:

- a. formation of ozone from atmospheric oxygen in the presence of sunlight
- b. reaction of  $\text{H}_2\text{SO}_4$  with  $\text{NaOH}$
- c. combination of dinitrogen with dioxygen at 2000 K
- d. Reaction of  $[\text{Co}(\text{H}_2\text{O})_6]\text{Cl}_3$  with  $\text{AgNO}_3$

# JEE Main 2020 Paper


16. Identify the correct labels of A, B and C in the following graph from the options given below:



Root mean square speed ( $V_{\text{rms}}$ ); most probable speed ( $V_{\text{mp}}$ ); average speed ( $V_{\text{av}}$ )

- |   |   |
|---|---|
| a. $A = V_{\text{mp}}, B = V_{\text{av}}, C = V_{\text{rms}}$ | b. $A = V_{\text{mp}}, B = V_{\text{rms}}, C = V_{\text{av}}$ |
| c. $A = V_{\text{av}}, B = V_{\text{rms}}, C = V_{\text{mp}}$ | d. $A = V_{\text{rms}}, B = V_{\text{mp}}, C = V_{\text{av}}$ |

17. For the reaction,



The observed rate expression is,  $\text{rate} = k_f[\text{NO}]^2[\text{H}_2]$ . The rate expression for the reverse reaction is:

- a.  $k_b[\text{N}_2][\text{H}_2\text{O}]^2$
- b.  $k_b[\text{N}_2][\text{H}_2\text{O}]$
- c.  $k_b[\text{N}_2][\text{H}_2\text{O}]^2/[\text{H}_2]$
- d.  $k_b[\text{N}_2][\text{H}_2\text{O}]^2/[\text{NO}]$

c

# JEE Main 2020 Paper

18. Within each pair of elements F & Cl, S & Se and Li & Na, respectively, the elements that release more energy upon an electron gain are:

- |                  |                 |
|------------------|-----------------|
| a. Cl, Se and Na | b. Cl, S and Li |
| c. F, S and Li   | d. F, Se and Na |

19. Among the following statements A-D, the incorrect ones are:

- |  |                 |
|--|-----------------|
| A. Octahedral Co(III) complexes with strong field ligands have high magnetic moments   |                 |
| B. When $\Delta_o < P$ , the d- electron configuration of Co(III) in an octahedral complex is $t_{2g}^4 e_g^2$ .   |                 |
| C. Wavelength of light absorbed by $[\text{Co}(\text{en})_3]^{3+}$ is lower than that of $[\text{CoF}_6]^{3-}$ .   |                 |
| D. If the $\Delta_o$ for an octahedral complex of Co(III) is $18000 \text{ cm}^{-1}$ , the $\Delta_t$ for its tetrahedral complex with the same ligand will be $16000 \text{ cm}^{-1}$ . |                 |
| a. B and C only  | c. A and D only |
| b. A and B only  | d. C and D only |



# JEE Main 2020 Paper

20. The ammonia ( $\text{NH}_3$ ) released on quantitative reaction of 0.6 g urea ( $\text{NH}_2\text{CONH}_2$ ) with sodium hydroxide ( $\text{NaOH}$ ) can be neutralized by:

- |                        |                        |
|------------------------|------------------------|
| a. 200 mL of 0.2 N HCl | c. 100 mL of 0.1 N HCl |
| b. 200 mL of 0.4 N HCl | d. 100 mL of 0.2 N HCl |

21. Number of  $\text{sp}^2$  hybrid carbon atoms present in aspartame is \_\_\_\_.




# JEE Main 2020 Paper

22. 3 grams of acetic acid is added to 250 mL of 0.1 M HCl and the solution is made up to 500 mL. To 20 mL of this solution  $\frac{1}{2}$  mL of 5 M NaOH is added. The pH of this solution is \_\_\_\_.

(Given:  $\log 3 = 0.4771$ ,  $pK_a$  of acetic acid = 4.74, molar mass of acetic acid = 60 g/mole).

23. The flocculation value of HCl for  $As_2S_3$  sol is  $30 \text{ mmolL}^{-1}$ . If  $H_2SO_4$  is used for the flocculation of arsenic sulphide, the amount, in grams, of  $H_2SO_4$  in 250 mL required for the above purpose is \_\_\_\_.

24. Consider the following reactions :



The sum of the total number of atoms in one molecule of (A), (B) & (C) is \_\_\_\_.

# JEE Main 2020 Paper

25. The standard heat of formation ( $\Delta_f H_{298}^\circ$ ) of ethane (in kJ/mol), if the heat of combustion of ethane, hydrogen and graphite are -1560, -393.5 and -286 kJ/mol, respectively, is\_\_\_\_\_.

# JEE Main 2020 Paper

Date of Exam: 7<sup>th</sup> January 2020 (Shift 2)

Time: 2:30 P.M. to 5:30 P.M.

Subject: Mathematics

---

1. If  $3x + 4y = 12\sqrt{2}$  is a tangent to the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{9} = 1$ , for some  $a \in \mathbb{R}$  then the distance between the foci of the ellipse is :

- |                |                |
|----------------|----------------|
| a. $2\sqrt{5}$ | b. $2\sqrt{7}$ |
| c. $2\sqrt{2}$ | d. 4           |

2. Let A, B, C and D be four non-empty sets. The Contrapositive statement of "If  $A \subseteq B$  and  $B \subseteq D$  then  $A \subseteq C$ " is :

- a. If  $A \subseteq C$ , then  $B \subset A$  or  $D \subset B$
- b. If  $A \not\subseteq C$ , then  $A \subseteq B$  and  $B \subseteq D$
- c. If  $A \not\subseteq C$ , then  $A \not\subseteq B$  and  $B \subseteq D$
- d. If  $A \not\subseteq C$ , then  $A \not\subseteq B$  or  $B \not\subseteq D$

# JEE Main 2020 Paper

3. The coefficient of  $x^7$  in the expression  $(1+x)^{10} + x(1+x)^9 + x^2(1+x)^8 + \dots + x^{10}$  is :
- a. 420  
b. 330  
c. 210  
d. 120
4. In a workshop, there are five machines and the probability of any one of them to be out of service on a day is  $\frac{1}{4}$ . If the probability that at most two machines will be out of service on the same day is  $\left(\frac{3}{4}\right)^3 k$ , then k is equal to :
- a.  $\frac{17}{2}$   
b. 4  
c.  $\frac{17}{4}$   
d.  $\frac{17}{8}$

# JEE Main 2020 Paper

5. The locus of mid points of the perpendiculars drawn from points on the line  $x = 2y$  to the line  $x = y$  is :

a.  $2x - 3y = 0$

b.  $3x - 2y = 0$

c.  $5x - 7y = 0$

d.  $7x - 5y = 0$

6. The value of  $\alpha$  for which  $4\alpha \int_{-1}^2 e^{-\alpha|x|} dx = 5$ , is :

a.  $\log_e 2$

b.  $\log_e \sqrt{2}$

c.  $\log_e \left(\frac{4}{3}\right)$

d.  $\log_e \left(\frac{3}{2}\right)$

# JEE Main 2020 Paper

7. If the sum of the first 40 terms of the series,  $3 + 4 + 8 + 9 + 13 + 14 + 18 + 19 + \dots$  is  $(102)m$ , then  $m$  is equal to :
- a. 10  
b. 25  
c. 5  
d. 20
8. If  $\frac{3+i \sin \theta}{4-i \cos \theta}$ ,  $\theta \in [0, 2\pi]$ , is a real number, then the argument of  $\sin \theta + i \cos \theta$  is :
- a.  $\pi - \tan^{-1}\left(\frac{4}{3}\right)$   
b.  $-\tan^{-1}\left(\frac{3}{4}\right)$   
c.  $\pi - \tan^{-1}\left(\frac{4}{3}\right)$   
d.  $\tan^{-1}\left(\frac{4}{3}\right)$

# JEE Main 2020 Paper

9. Let  $A = [a_{ij}]$  and  $B = [b_{ij}]$  be two  $3 \times 3$  real matrices such that  $b_{ij} = (3)^{(i+j-2)}a_{ji}$ , where  $i, j = 1, 2, 3$ . If the determinant of  $B$  is 81, then the determinant of  $A$  is :

a.  $\frac{1}{9}$

b.  $\frac{1}{81}$

c.  $\frac{1}{3}$

d. 3

10. Let  $f(x)$  be a polynomial of degree 5 such that  $x = \pm 1$  are its critical points. If  $\lim_{x \rightarrow 0} \left( 2 + \frac{f(x)}{x^3} \right) = 4$ , then which one of the following is not true?
- a.  $f(1) - 4f(-1) = 4$
  - b.  $x = 1$  is a point of maxima and  $x = -1$  is a point of minimum of  $f$ .
  - c.  $f$  is an odd function.
  - d.  $x = 1$  is a point of minima and  $x = -1$  is a point of maxima of  $f$ .



# JEE Main 2020 Paper

11. The number of ordered pairs  $(r, k)$  for which  $6 \cdot {}^{35}C_r = (k^2 - 3) \cdot {}^{36}C_{r+1}$ , where  $k$  is an integer, is :

- a. 4
- c. 2

- b. 6
- d. 3

12. Let  $a_1, a_2, a_3, \dots$  be a G.P. such that  $a_1 < 0$ ,  $a_1 + a_2 = 4$  and  $a_3 + a_4 = 16$ . If  $\sum_{i=1}^9 a_i = 4\lambda$ , then  $\lambda$  is equal to :

- a. 171

- b.  $\frac{511}{3}$

- c. -171

- d. -513

# JEE Main 2020 Paper

13. Let  $\vec{a}, \vec{b}$  and  $\vec{c}$  be three unit vectors such that  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ . If  $\lambda = \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{c} + \vec{c} \cdot \vec{a}$  and  $\vec{d} = \vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a}$ , then the ordered pair  $(\lambda, \vec{d})$  is equal to :

- |   |   |
|---|---|
| a. $\left(\frac{3}{2}, 3\vec{a} \times \vec{c}\right)$  | b. $\left(-\frac{3}{2}, 3\vec{c} \times \vec{b}\right)$ |
| c. $\left(-\frac{3}{2}, 3\vec{a} \times \vec{b}\right)$ | d. $\left(\frac{3}{2}, 3\vec{b} \times \vec{c}\right)$  |

14. Let  $y = y(x)$  be the solution curve of the differential equation,  $(y^2 - x) \frac{dy}{dx} = 1$ , satisfying  $y(0) = 1$ . This curve intersects the  $x$  - axis at a point whose abscissa is :

- |            |         |
|------------|---------|
| a. $2 + e$ | b. $2$  |
| c. $2 - e$ | d. $-e$ |

# JEE Main 2020 Paper

15. If  $\theta_1$  and  $\theta_2$  be respectively the smallest and the largest values of  $\theta$  in  $(0, 2\pi) - \{\pi\}$  which satisfy the equation,  $2 \cot^2 \theta - \frac{5}{\sin \theta} + 4 = 0$ , then  $\int_{\theta_1}^{\theta_2} \cos^2 3\theta \, d\theta$  is equal to :

a.  $\frac{2\pi}{3}$

b.  $\frac{\pi}{3}$

c.  $\frac{\pi}{3} + \frac{1}{6}$

d.  $\frac{\pi}{9}$

16. Let  $\alpha$  and  $\beta$  are the roots of the equation  $x^2 - x - 1 = 0$ . If  $p_k = (\alpha)^k + (\beta)^k$ ,  $k \geq 1$  then which one of the following statements is not true?

a.  $(p_1 + p_2 + p_3 + p_4 + p_5) = 26$

b.  $p_5 = 11$

c.  $p_5 = p_2 \cdot p_3$

d.  $p_3 = p_5 - p_4$

# JEE Main 2020 Paper

17. The area (in sq. units) of the region  $\{(x, y) \in \mathbb{R}^2 \mid 4x^2 \leq y \leq 8x + 12\}$  is :

a.  $\frac{125}{3}$

b.  $\frac{128}{3}$

c.  $\frac{124}{3}$

d.  $\frac{127}{3}$

# JEE Main 2020 Paper

18. The value of  $c$  in Lagrange's mean value theorem for the function  $f(x) = x^3 - 4x^2 + 8x + 11$ , where  $x \in [0,1]$  is :

a.  $\frac{4-\sqrt{7}}{3}$

b.  $\frac{2}{3}$

c.  $\frac{\sqrt{7}-2}{3}$

d.  $\frac{4-\sqrt{5}}{3}$

19. Let  $y = y(x)$  be a function of  $x$  satisfying  $y\sqrt{1-x^2} = k - x\sqrt{1-y^2}$  where  $k$  is a constant and  $y\left(\frac{1}{2}\right) = -\frac{1}{4}$ . Then  $\frac{dy}{dx}$  at  $x = \frac{1}{2}$ , is equal to :

a.  $-\frac{\sqrt{5}}{2}$

b.  $\frac{\sqrt{5}}{2}$

c.  $-\frac{\sqrt{5}}{4}$

d.  $\frac{2}{\sqrt{5}}$

# JEE Main 2020 Paper

20. Let the tangents drawn from the origin to the circle,  $x^2 + y^2 - 8x - 4y + 16 = 0$  touch it at the points  $A$  and  $B$ . The  $(AB)^2$  is equal to :

a.  $\frac{32}{5}$

b.  $\frac{64}{5}$

c.  $\frac{52}{5}$

d.  $\frac{56}{5}$

# JEE Main 2020 Paper

21. If system of linear equations

$$x + y + z = 6$$

$$x + 2y + 3z = 10$$

$$3x + 2y + \lambda z = \mu$$

has more than two solutions, then  $\mu - \lambda^2$  is equal to \_\_\_\_\_.

22. If the foot of perpendicular drawn from the point  $(1, 0, 3)$  on a line passing through  $(\alpha, 7, 1)$  is

$\left(\frac{5}{3}, \frac{7}{3}, \frac{17}{3}\right)$ , then  $\alpha$  is equal to \_\_\_\_\_.

23. If the function  $f$  defined on  $\left(-\frac{1}{3}, \frac{1}{3}\right)$  by

$$f(x) = \begin{cases} \left(\frac{1}{x}\right) \log_e \left(\frac{1+3x}{1-2x}\right) & \text{when } x \neq 0 \\ k & \text{when } x = 0 \end{cases}$$

is continuous, the  $k$  is equal to \_\_\_\_\_.

# JEE Main 2020 Paper

24. If the mean and variance of eight numbers  $3, 7, 9, 12, 13, 20, x$  and  $y$  be 10 and 25 respectively then  $xy$  is equal to \_\_\_\_\_.
25. Let  $X = \{n \in \mathbf{N} : 1 \leq n \leq 50\}$ . If  $A = \{n \in X : n \text{ is a multiple of } 2\}$  and  $B = \{n \in X : n \text{ is a multiple of } 7\}$ , then the number of elements in the smallest subset of  $X$  containing both  $A$  and  $B$  is \_\_\_\_\_.