

# FINAL JEE-MAIN EXAMINATION – MARCH, 2021

(Held On Wednesday 17<sup>th</sup> March, 2021) TIME : 3 : 00 PM to 6 : 00 PM

## PHYSICS

## TEST PAPER WITH ANSWER & SOLUTION


### SECTION-A

1. A rubber ball is released from a height of 5 m above the floor. It bounces back repeatedly, always rising to  $\frac{81}{100}$  of the height through which it falls. Find the average speed of the ball. (Take  $g = 10 \text{ ms}^{-2}$ )  
 (1)  $3.0 \text{ ms}^{-1}$  (2)  $3.50 \text{ ms}^{-1}$   
 (3)  $2.0 \text{ ms}^{-1}$  (4)  $2.50 \text{ ms}^{-1}$

2. If one mole of the polyatomic gas is having two vibrational modes and  $\beta$  is the ratio of molar specific heats for polyatomic gas  $\left(\beta = \frac{C_p}{C_v}\right)$  then the value of  $\beta$  is :  
 (1) 1.02 (2) 1.2  
 (3) 1.25 (4) 1.35

3. A block of mass 1 kg attached to a spring is made to oscillate with an initial amplitude of 12 cm. After 2 minutes the amplitude decreases to 6 cm. Determine the value of the damping constant for this motion. (take  $\ln 2 = 0.693$ )  
 (1)  $0.69 \times 10^2 \text{ kg s}^{-1}$  (2)  $3.3 \times 10^2 \text{ kg s}^{-1}$   
 (3)  $1.16 \times 10^2 \text{ kg s}^{-1}$  (4)  $5.7 \times 10^{-3} \text{ kg s}^{-1}$

4. Which one of the following will be the output of the given circuit ?



- (1) NOR Gate (2) NAND Gate  
 (3) AND Gate (4) XOR Gate
5. An object is located at 2 km beneath the surface of the water. If the fractional compression  $\frac{\Delta V}{V}$  is 1.36% , the ratio of hydraulic stress to the corresponding hydraulic strain will be \_\_\_\_\_.  
 [Given : density of water is  $1000 \text{ kg m}^{-3}$  and  $g = 9.8 \text{ ms}^{-2}$ .]  
 (1)  $1.96 \times 10^7 \text{ Nm}^{-2}$  (2)  $1.44 \times 10^7 \text{ Nm}^{-2}$   
 (3)  $2.26 \times 10^9 \text{ Nm}^{-2}$  (4)  $1.44 \times 10^9 \text{ Nm}^{-2}$





6. A geostationary satellite is orbiting around an arbitrary planet 'P' at a height of  $11R$  above the surface of 'P',  $R$  being the radius of 'P'. The time period of another satellite in hours at a height of  $2R$  from the surface of 'P' is \_\_\_\_\_. 'P' has the time period of 24 hours.

(1)  $6\sqrt{2}$     (2)  $\frac{6}{\sqrt{2}}$     (3) 3    (4) 5

7. A sound wave of frequency 245 Hz travels with the speed of  $300 \text{ ms}^{-1}$  along the positive x-axis. Each point of the wave moves to and fro through a total distance of 6 cm. What will be the mathematical expression of this travelling wave ?

(1)  $Y(x,t) = 0.03 [\sin 5.1 x - (0.2 \times 10^3)t]$   
 (2)  $Y(x,t) = 0.06 [\sin 5.1 x - (1.5 \times 10^3)t]$   
 (3)  $Y(x,t) = 0.06 [\sin 0.8 x - (0.5 \times 10^3)t]$   
 (4)  $Y(x,t) = 0.03 [\sin 5.1 x - (1.5 \times 10^3)t]$

8. Which one is the correct option for the two different thermodynamic processes ?



- (1) (c) and (a)                      (2) (c) and (d)  
 (3) (a) only                          (4) (b) and (c)


9. The velocity of a particle is  $v = v_0 + gt + Ft^2$ . Its position is  $x = 0$  at  $t = 0$  ; then its displacement after time ( $t = 1$ ) is :

- (1)  $v_0 + g + F$                       (2)  $v_0 + \frac{g}{2} + \frac{F}{3}$   
 (3)  $v_0 + \frac{g}{2} + F$                       (4)  $v_0 + 2g + 3F$

10. A carrier signal  $C(t) = 25 \sin (2.512 \times 10^{10} t)$  is amplitude modulated by a message signal  $m(t) = 5 \sin (1.57 \times 10^8 t)$  and transmitted through an antenna. What will be the bandwidth of the modulated signal ?


- (1) 8 GHz  
 (2) 2.01 GHz  
 (3) 1987.5 MHz  
 (4) 50 MHz

11. Two cells of emf  $2E$  and  $E$  with internal resistance  $r_1$  and  $r_2$  respectively are connected in series to an external resistor  $R$  (see figure). The value of  $R$ , at which the potential difference across the terminals of the first cell becomes zero is




- (1)  $r_1 + r_2$                       (2)  $\frac{r_1}{2} - r_2$   
 (3)  $\frac{r_1}{2} + r_2$                       (4)  $r_1 - r_2$

12. A hairpin like shape as shown in figure is made by bending a long current carrying wire. What is the magnitude of a magnetic field at point  $P$  which lies on the centre of the semicircle ?



- (1)  $\frac{\mu_0 I}{4\pi r} (2 - \pi)$                       (2)  $\frac{\mu_0 I}{4\pi r} (2 + \pi)$   
 (3)  $\frac{\mu_0 I}{2\pi r} (2 + \pi)$                       (4)  $\frac{\mu_0 I}{2\pi r} (2 - \pi)$

13. The four arms of a Wheatstone bridge have resistances as shown in the figure. A galvanometer of  $15\ \Omega$  resistance is connected across BD. Calculate the current through the galvanometer when a potential difference of 10V is maintained across AC.



- (1)  $2.44\ \mu\text{A}$  (2)  $2.44\ \text{mA}$   
(3)  $4.87\ \text{mA}$  (4)  $4.87\ \mu\text{A}$

14. Two particles A and B of equal masses are suspended from two massless springs of spring constants  $K_1$  and  $K_2$  respectively. If the maximum velocities during oscillations are equal, the ratio of the amplitude of A and B is

- (1)  $\frac{K_2}{K_1}$  (2)  $\frac{K_1}{K_2}$  (3)  $\sqrt{\frac{K_1}{K_2}}$  (4)  $\sqrt{\frac{K_2}{K_1}}$

15. Match List-I with List-II

**List-I**


**List-II**

- |   |  |
|---|--|
| (a) Phase difference between current and voltage in a purely resistive AC circuit | (i) $\frac{\pi}{2}$ ; current leads voltage      |
| (b) Phase difference between current and voltage in a pure inductive AC circuit   | (ii) zero  |
| (c) Phase difference between current and voltage in a pure capacitive AC circuit  | (iii) $\frac{\pi}{2}$ ; current lags voltage     |
| (d) Phase difference between current and voltage in an LCR series circuit         | (iv) $\tan^{-1}\left(\frac{X_C - X_L}{R}\right)$ |

Choose the most appropriate answer from the options given below :

- (1) (a)–(i), (b)–(iii), (c)–(iv), (d)–(ii)  
(2) (a)–(ii), (b)–(iv), (c)–(iii), (d)–(i)  
(3) (a)–(ii), (b)–(iii), (c)–(iv), (d)–(i)  
(4) (a)–(ii), (b)–(iii), (c)–(i), (d)–(iv)

16. Two identical blocks A and B each of mass  $m$  resting on the smooth horizontal floor are connected by a light spring of natural length  $L$  and spring constant  $K$ . A third block C of mass  $m$  moving with a speed  $v$  along the line joining A and B collides with A. The maximum compression in the spring is



- (1)  $v\sqrt{\frac{M}{2K}}$                       (2)  $\sqrt{\frac{mv}{2K}}$   
 (3)  $\sqrt{\frac{mv}{K}}$                       (4)  $\sqrt{\frac{m}{2K}}$

17. The atomic hydrogen emits a line spectrum consisting of various series. Which series of hydrogen atomic spectra is lying in the visible region ?
- (1) Brackett series                      (2) Paschen series  
 (3) Lyman series                      (4) Balmer series

18. Two identical photocathodes receive the light of frequencies  $f_1$  and  $f_2$  respectively. If the velocities of the photo-electrons coming out are  $v_1$  and  $v_2$  respectively, then

(1)  $v_1^2 - v_2^2 = \frac{2h}{m} [f_1 - f_2]$


(2)  $v_1^2 + v_2^2 = \frac{2h}{m} [f_1 + f_2]$

(3)  $v_1 + v_2 = \left[ \frac{2h}{m} (f_1 + f_2) \right]^{\frac{1}{2}}$

(4)  $v_1 - v_2 = \left[ \frac{2h}{m} (f_1 - f_2) \right]^{1/2}$

19. What happens to the inductive reactance and the current in a purely inductive circuit if the frequency is halved ?
- (1) Both, inductive reactance and current will be halved.  
 (2) Inductive reactance will be halved and current will be doubled.  
 (3) Inductive reactance will be doubled and current will be halved.  
 (4) Both, inducting reactance and current will be doubled.

20. A sphere of mass 2kg and radius 0.5 m is rolling with an initial speed of  $1 \text{ ms}^{-1}$  goes up an inclined plane which makes an angle of  $30^\circ$  with the horizontal plane, without slipping. How low will the sphere take to return to the starting point A ?



- (1) 0.60 s                      (2) 0.52 s  
(3) 0.57 s                      (4) 0.80 s

### SECTION-B


1. The electric field intensity produced by the radiation coming from a 100 W bulb at a distance of 3m is E. The electric field intensity produced by the radiation coming from 60 W

at the same distance is  $\sqrt{\frac{x}{5}}E$ . Where the value of x = \_\_\_\_\_.

2. A body of mass 1 kg rests on a horizontal floor with which it has a coefficient of static friction  $\frac{1}{\sqrt{3}}$ . It is desired to make the body move by applying the minimum possible force F N. The value of F will be \_\_\_\_\_. (Round off to the Nearest Integer)

[Take  $g = 10 \text{ ms}^{-2}$ ]

3. A boy of mass 4 kg is standing on a piece of wood having mass 5kg . If the coefficient of friction between the wood and the floor is 0.5, the maximum force that the boy can exert on the rope so that the piece of wood does not move from its place is \_\_\_\_\_N.(Round off to the Nearest Integer)



[Take  $g = 10 \text{ ms}^{-2}$ ]

4. Suppose you have taken a dilute solution of oleic acid in such a way that its concentration becomes  $0.01 \text{ cm}^3$  of oleic acid per  $\text{cm}^3$  of the solution. Then you make a thin film of this solution (monomolecular thickness) of area  $4 \text{ cm}^2$  by considering 100 spherical drops of radius  $\left(\frac{3}{40\pi}\right)^{\frac{1}{3}} \times 10^{-3} \text{ cm}$ . Then the thickness of oleic acid layer will be  $x \times 10^{-14} \text{ m}$ .

Where  $x$  is \_\_\_\_\_.

5. A particle of mass  $m$  moves in a circular orbit in a central potential field  $U(r) = U_0 r^4$ . If Bohr's quantization conditions are applied, radii of possible orbitals  $r_n$  vary with  $n^{1/\alpha}$ , where  $\alpha$  is \_\_\_\_\_.

6. The electric field in a region is given by

$$\vec{E} = \frac{2}{5} E_0 \hat{i} + \frac{3}{5} E_0 \hat{j} \text{ with } E_0 = 4.0 \times 10^3 \frac{\text{N}}{\text{C}}.$$


The flux of this field through a rectangular surface area  $0.4 \text{ m}^2$  parallel to the  $Y - Z$  plane is \_\_\_\_\_  $\text{Nm}^2\text{C}^{-1}$ .

7. The disc of mass  $M$  with uniform surface mass density  $\sigma$  is shown in the figure. The centre of mass of the quarter disc (the shaded area) is at

the position  $\frac{x}{3} \frac{a}{\pi}, \frac{x}{3} \frac{a}{\pi}$  where  $x$  is \_\_\_\_\_.


(Round off to the Nearest Integer)

[ $a$  is an area as shown in the figure]



8. The image of an object placed in air formed by a convex refracting surface is at a distance of 10 m behind the surface. The image is real and is at  $\frac{2^{\text{rd}}}{3}$  of the distance of the object from the surface. The wavelength of light inside the surface is  $\frac{2}{3}$  times the wavelength in air. The radius of the curved surface is  $\frac{x}{13}$  m. the value of 'x' is\_\_\_\_\_.

9. A  $2\ \mu\text{F}$  capacitor  $C_1$  is first charged to a potential difference of 10 V using a battery. Then the battery is removed and the capacitor is connected to an uncharged capacitor  $C_2$  of  $8\ \mu\text{F}$ . The charge in  $C_2$  on equilibrium condition is\_\_\_\_\_  $\mu\text{C}$ . (Round off to the Nearest Integer)



10. Seawater at a frequency  $f = 9 \times 10^2$  Hz, has permittivity  $\epsilon = 80\epsilon_0$  and resistivity  $\rho = 0.25\ \Omega\text{m}$ . Imagine a parallel plate capacitor is immersed in seawater and is driven by an alternating voltage source  $V(t) = V_0 \sin(2\pi ft)$ . Then the conduction current density becomes  $10^x$  times the displacement current density after time  $t = \frac{1}{800}$  s. The value of x is \_\_\_\_\_

(Given :  $\frac{1}{4\pi\epsilon_0} = 9 \times 10^9 \text{ Nm}^2\text{C}^{-2}$ )



# FINAL JEE-MAIN EXAMINATION – MARCH, 2021

(Held On Wednesday 17<sup>th</sup> March, 2021) TIME : 3 : 00 PM to 6 : 00 PM

CHEMISTRY	TEST PAPER WITH ANSWER & SOLUTION																				
<p><b>SECTION-A</b></p> <p>1. Fructose is an example of :-                      (1) Pyranose                      (2) Ketohexose                      (3) Aldohexose                      (4) Heptose</p> <p>2. The set of elements that differ in mutual relationship from those of the other sets is :                      (1) Li – Mg                      (2) B – Si                      (3) Be – Al                      (4) Li – Na</p> <p>3. The functional groups that are responsible for the ion-exchange property of cation and anion exchange resins, respectively, are :                      (1) <math>-\text{SO}_3\text{H}</math> and <math>-\text{NH}_2</math>                      (2) <math>-\text{SO}_3\text{H}</math> and <math>-\text{COOH}</math>                      (3) <math>-\text{NH}_2</math> and <math>-\text{COOH}</math>                      (4) <math>-\text{NH}_2</math> and <math>-\text{SO}_3\text{H}</math></p> <p>4. Match List-I and List-II :  <table> <tr> <th>List-I</th><th>List-II</th></tr> <tr> <td>(a) Haematite</td><td>(i) <math>\text{Al}_2\text{O}_3 \cdot x\text{H}_2\text{O}</math></td></tr> <tr> <td>(b) Bauxite</td><td>(ii) <math>\text{Fe}_2\text{O}_3</math></td></tr> <tr> <td>(c) Magnetite</td><td>(iii) <math>\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2</math></td></tr> <tr> <td>(d) Malachite</td><td>(iv) <math>\text{Fe}_3\text{O}_4</math></td></tr> </table>                     Choose the correct answer from the options given below :</p>	List-I	List-II	(a) Haematite	(i) $\text{Al}_2\text{O}_3 \cdot x\text{H}_2\text{O}$	(b) Bauxite	(ii) $\text{Fe}_2\text{O}_3$	(c) Magnetite	(iii) $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$	(d) Malachite	(iv) $\text{Fe}_3\text{O}_4$	<p>(1) (a)-(ii), (b)-(iii), (c)-(i), (d)-(iv)                      (2) (a)-(iv), (b)-(i), (c)-(ii), (d)-(iii)                      (3) (a)-(i), (b)-(iii), (c)-(ii), (d)-(iv)                      (4) (a)-(ii), (b)-(i), (c)-(iv), (d)-(iii)</p> <p>5. The correct pair(s) of the ambident nucleophiles is (are) :                      (A) <math>\text{AgCN/KCN}</math>                      (B) <math>\text{RCOOAg/RCOOK}</math>                      (C) <math>\text{AgNO}_2/\text{KNO}_2</math>                      (D) <math>\text{AgI/KI}</math>                      (1) (B) and (C) only                      (2) (A) only                      (3) (A) and (C) only                      (4) (B) only</p> <p>6. The set that represents the pair of neutral oxides of nitrogen is :                      (1) <math>\text{NO}</math> and <math>\text{N}_2\text{O}</math>                      (2) <math>\text{N}_2\text{O}</math> and <math>\text{N}_2\text{O}_3</math>                      (3) <math>\text{N}_2\text{O}</math> and <math>\text{NO}_2</math>                      (4) <math>\text{NO}</math> and <math>\text{NO}_2</math></p> <p>7. Match List-I with List-II :  <table> <tr> <th>List-I</th><th>List-II</th></tr> <tr> <td>(a) <math>[\text{Co}(\text{NH}_3)_6] [\text{Cr}(\text{CN})_6]</math></td><td>(i) Linkage isomerism</td></tr> <tr> <td>(b) <math>[\text{Co}(\text{NH}_3)_3 (\text{NO}_2)_3]</math></td><td>(ii) Solvate isomerism</td></tr> <tr> <td>(c) <math>[\text{Cr}(\text{H}_2\text{O})_6]\text{Cl}_3</math></td><td>(iii) Co-ordination isomerism</td></tr> <tr> <td>(d) <math>\text{cis-}[\text{CrCl}_2(\text{ox})_2]^{3-}</math></td><td>(iv) Optical isomerism</td></tr> </table>                     Choose the correct answer from the options given below :                      (1) (a)-(iii), (b)-(i), (c)-(ii), (d)-(iv)                      (2) (a)-(iv), (b)-(ii), (c)-(iii), (d)-(i)                      (3) (a)-(ii), (b)-(i), (c)-(iii), (d)-(iv)                      (4) (a)-(i), (b)-(ii), (c)-(iii), (d)-(iv)</p>	List-I	List-II	(a) $[\text{Co}(\text{NH}_3)_6] [\text{Cr}(\text{CN})_6]$	(i) Linkage isomerism	(b) $[\text{Co}(\text{NH}_3)_3 (\text{NO}_2)_3]$	(ii) Solvate isomerism	(c) $[\text{Cr}(\text{H}_2\text{O})_6]\text{Cl}_3$	(iii) Co-ordination isomerism	(d) $\text{cis-}[\text{CrCl}_2(\text{ox})_2]^{3-}$	(iv) Optical isomerism
List-I	List-II																				
(a) Haematite	(i) $\text{Al}_2\text{O}_3 \cdot x\text{H}_2\text{O}$																				
(b) Bauxite	(ii) $\text{Fe}_2\text{O}_3$																				
(c) Magnetite	(iii) $\text{CuCO}_3 \cdot \text{Cu}(\text{OH})_2$																				
(d) Malachite	(iv) $\text{Fe}_3\text{O}_4$																				
List-I	List-II																				
(a) $[\text{Co}(\text{NH}_3)_6] [\text{Cr}(\text{CN})_6]$	(i) Linkage isomerism																				
(b) $[\text{Co}(\text{NH}_3)_3 (\text{NO}_2)_3]$	(ii) Solvate isomerism																				
(c) $[\text{Cr}(\text{H}_2\text{O})_6]\text{Cl}_3$	(iii) Co-ordination isomerism																				
(d) $\text{cis-}[\text{CrCl}_2(\text{ox})_2]^{3-}$	(iv) Optical isomerism																				

8. Primary, secondary and tertiary amines can be separated using :-  
 (1) Para-Toluene sulphonyl chloride  
 (2) Chloroform and KOH  
 (3) Benzene sulphonic acid  
 (4) Acetyl amide

9. The common positive oxidation states for an element with atomic number 24, are :  
 (1) +2 to +6 (2) +1 and +3 to +6  
 (3) +1 and +3 (4) +1 to +6

10. Match List-I with List-II :

**List-I  
Chemical  
Compound**

**List-II  
Used as**

- |                                    |                           |
|------------------------------------|---------------------------|
| (a) Sucralose                      | (i) Synthetic detergent   |
| (b) Glyceryl ester of stearic acid | (ii) Artificial sweetener |
| (c) Sodium benzoate                | (iii) Antiseptic          |
| (d) Bithionol                      | (iv) Food preservative    |

Choose the correct match :

- (1) (a)-(iv), (b)-(iii), (c)-(ii), (d)-(i)  
 (2) (a)-(ii), (b)-(i), (c)-(iv), (d)-(iii)  
 (3) (a)-(iii), (b)-(ii), (c)-(iv), (d)-(i)  
 (4) (a)-(i), (b)-(ii), (c)-(iv), (d)-(iii)

11. Given below are two statements :





**Statement-I** : 2-methylbutane on oxidation with  $\text{KMnO}_4$  gives 2-methylbutan-2-ol.

**Statement-II** : n-alkanes can be easily oxidised to corresponding alcohol with  $\text{KMnO}_4$ .

Choose the correct option :



- (1) Both statement I and statement II are correct  
 (2) Both statement I and statement II are incorrect  
 (3) Statement I is correct but Statement II is incorrect  
 (4) Statement I is incorrect but Statement II is correct

12. Nitrogen can be estimated by Kjeldahl's method for which of the following compound ?



13. Amongst the following, the linear species is :


- (1)  $\text{NO}_2$  (2)  $\text{Cl}_2\text{O}$   
 (3)  $\text{O}_3$  (4)  $\text{N}_3^-$







In the above reactions, the enzyme A and enzyme B respectively are :-

- (1) Amylase and Invertase
- (2) Invertase and Amylase
- (3) Invertase and Zymase
- (4) Zymase and Invertase

- 15.** One of the by-products formed during the recovery of  $\text{NH}_3$  from Solvay process is :
- (1)  $\text{Ca}(\text{OH})_2$                       (2)  $\text{NaHCO}_3$   
(3)  $\text{CaCl}_2$                           (4)  $\text{NH}_4\text{Cl}$




In the above reaction, the structural formula of (A), "X" and "Y" respectively are :



17. For the coagulation of a negative sol, the species below, that has the highest flocculating power is :
- (1)  $\text{SO}_4^{2-}$     (2)  $\text{Ba}^{2+}$     (3)  $\text{Na}^+$     (4)  $\text{PO}_4^{3-}$

- 18.** Which of the following statement(s) is (are) incorrect reason for eutrophication ?
- (A) excess usage of fertilisers
  - (B) excess usage of detergents
  - (C) dense plant population in water bodies
  - (D) lack of nutrients in water bodies that prevent plant growth
- Choose the most appropriate answer from the options given below :
- (1) (A) only
  - (2) (C) only
  - (3) (B) and (D) only
  - (4) (D) only

- 19.** Choose the correct statement regarding the formation of carbocations A and B given :-



- (1) Carbocation B is more stable and formed relatively at faster rate
- (2) Carbocation A is more stable and formed relatively at slow rate
- (3) Carbocation B is more stable and formed relatively at slow rate
- (4) Carbocation A is more stable and formed relatively at faster rate

20. During which of the following processes, does entropy decrease ?
- (A) Freezing of water to ice at  $0^{\circ}\text{C}$   
 (B) Freezing of water to ice at  $-10^{\circ}\text{C}$   
 (C)  $\text{N}_2(\text{g}) + 3\text{H}_2(\text{g}) \rightarrow 2\text{NH}_3(\text{g})$   
 (D) Adsorption of  $\text{CO}(\text{g})$  and lead surface  
 (E) Dissolution of  $\text{NaCl}$  in water

#### SECTION-B

1. A  $\text{KCl}$  solution of conductivity  $0.14 \text{ S m}^{-1}$  shows a resistance of  $4.19 \Omega$  in a conductivity cell. If the same cell is filled with an  $\text{HCl}$  solution, the resistance drops to  $1.03 \Omega$ . The conductivity of the  $\text{HCl}$  solution is  $\text{---} \times 10^{-2} \text{ S m}^{-1}$ . (Round off to the Nearest Integer).

2. On complete reaction of  $\text{FeCl}_3$  with oxalic acid in aqueous solution containing  $\text{KOH}$ , resulted in the formation of product A. The secondary valency of  $\text{Fe}$  in the product A is \_\_\_\_.

(Round off to the Nearest Integer).

3. The reaction  $2\text{A} + \text{B}_2 \rightarrow 2\text{AB}$  is an elementary reaction.

For a certain quantity of reactants, if the volume of the reaction vessel is reduced by a factor of 3, the rate of the reaction increases by a factor of \_\_\_\_\_. (Round off to the Nearest Integer).

4. The total number of  $\text{C}-\text{C}$  sigma bond/s in mesityl oxide ( $\text{C}_6\text{H}_{10}\text{O}$ ) is \_\_\_\_\_. (Round off to the Nearest Integer).

5. A 1 molal  $\text{K}_4\text{Fe}(\text{CN})_6$  solution has a degree of dissociation of 0.4. Its boiling point is equal to that of another solution which contains 18.1 weight percent of a non electrolytic solute A. The molar mass of A is \_\_\_\_ u. (Round off to the Nearest Integer).

[Density of water =  $1.0 \text{ g cm}^{-3}$ ]

6. In the ground state of atomic Fe ( $Z = 26$ ), the spin-only magnetic moment is \_\_\_\_\_  $\times 10^{-1}$  BM. (Round off to the Nearest Integer).

[Given :  $\sqrt{3} = 1.73$ ,  $\sqrt{2} = 1.41$ ]

7. The number of chlorine atoms in 20 mL of chlorine gas at STP is \_\_\_\_\_  $10^{21}$ . (Round off to the Nearest Integer).

[Assume chlorine is an ideal gas at STP

$R = 0.083 \text{ L bar mol}^{-1} \text{ K}^{-1}$ ,  $N_A = 6.023 \times 10^{23}$ ]

8. KBr is doped with  $10^{-5}$  mole percent of  $\text{SrBr}_2$ . The number of cationic vacancies in 1 g of KBr crystal is \_\_\_\_\_  $10^{14}$ . (Round off to the Nearest Integer).


[Atomic Mass : K : 39.1 u, Br : 79.9 u,

$N_A = 6.023 \times 10^{23}$ ]

9. Consider the reaction  $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$ . The temperature at which  $K_C = 20.4$  and  $K_P = 600.1$ , is \_\_\_\_\_ K. (Round off to the Nearest Integer).

[Assume all gases are ideal and  $R = 0.0831 \text{ L bar K}^{-1} \text{ mol}^{-1}$ ]

10.



Consider the above reaction. The percentage yield of amide product is \_\_\_\_\_. (Round off to the Nearest Integer).

(Given : Atomic mass : C : 12.0 u, H : 1.0u, N : 14.0 u, O : 16.0 u, Cl : 35.5 u)

**FINAL JEE–MAIN EXAMINATION – MARCH, 2021**

**(Held On Wednesday 17<sup>th</sup> March, 2021) TIME : 3 : 00 PM to 6 : 00 PM**

**MATHEMATICS**

**TEST PAPER WITH SOLUTION**

**SECTION-A**

1. Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be defined as  $f(x) = e^{-x} \sin x$ . If  $F : [0, 1] \rightarrow \mathbb{R}$  is a differentiable function

such that  $F(x) = \int_0^x f(t) dt$ , then the value of

$\int_0^1 (F'(x) + f(x)) e^x dx$  lies in the interval

(1)  $\left[ \frac{327}{360}, \frac{329}{360} \right]$  (2)  $\left[ \frac{330}{360}, \frac{331}{360} \right]$

(3)  $\left[ \frac{331}{360}, \frac{334}{360} \right]$  (4)  $\left[ \frac{335}{360}, \frac{336}{360} \right]$

2. If the integral  $\int_0^{10} \frac{[\sin 2\pi x]}{e^{x-[x]}} dx = \alpha e^{-1} + \beta e^{-\frac{1}{2}} + \gamma$ ,

where  $\alpha, \beta, \gamma$  are integers and  $[x]$  denotes the greatest integer less than or equal to  $x$ , then the value of  $\alpha + \beta + \gamma$  is equal to :

- (1) 0 (2) 20 (3) 25 (4) 10

3. Let  $y = y(x)$  be the solution of the differential equation  $\cos x (3\sin x + \cos x + 3)dy = (1 + y \sin x (3\sin x + \cos x + 3))dx$ ,

$0 \leq x \leq \frac{\pi}{2}$ ,  $y(0) = 0$ . Then,  $y\left(\frac{\pi}{3}\right)$  is equal to:

- (1)  $2\log_e\left(\frac{2\sqrt{3}+9}{6}\right)$       (2)  $2\log_e\left(\frac{2\sqrt{3}+10}{11}\right)$   
(3)  $2\log_e\left(\frac{\sqrt{3}+7}{2}\right)$       (4)  $2\log_e\left(\frac{3\sqrt{3}-8}{4}\right)$

4. The value of  $\sum_{r=0}^6 ({}^6C_r \cdot {}^6C_{6-r})$  is equal to :

- (1) 1124      (2) 1324      (3) 1024      (4) 924

5. The value of  $\lim_{n \rightarrow \infty} \frac{[r] + [2r] + \dots + [nr]}{n^2}$ , where  $r$  is non-zero real number and  $[r]$  denotes the greatest integer less than or equal to  $r$ , is equal to :
- (1)  $\frac{r}{2}$       (2)  $r$       (3)  $2r$       (4)  $0$

6. The number of solutions of the equation  $\sin^{-1}\left[x^2 + \frac{1}{3}\right] + \cos^{-1}\left[x^2 - \frac{2}{3}\right] = x^2$ , for  $x \in [-1, 1]$ , and  $[x]$  denotes the greatest integer less than or equal to  $x$ , is :
- (1) 2      (2) 0  
(3) 4      (4) Infinite

7. Let a computer program generate only the digits 0 and 1 to form a string of binary numbers with probability of occurrence of 0 at even places be  $\frac{1}{2}$  and probability of occurrence of 0 at the odd place be  $\frac{1}{3}$ . Then the probability that '10' is followed by '01' is equal to :
- (1)  $\frac{1}{18}$       (2)  $\frac{1}{3}$       (3)  $\frac{1}{6}$       (4)  $\frac{1}{9}$



8. The number of solutions of the equation

$$x + 2 \tan x = \frac{\pi}{2} \text{ in the interval } [0, 2\pi] \text{ is :}$$

- (1) 3      (2) 4      (3) 2      (4) 5

9. Let  $S_1$ ,  $S_2$  and  $S_3$  be three sets defined as

$$S_1 = \{z \in \mathbb{C} : |z - 1| \leq \sqrt{2}\}$$

$$S_2 = \{z \in \mathbb{C} : \operatorname{Re}((1 - i)z) \geq 1\}$$

$$S_3 = \{z \in \mathbb{C} : \operatorname{Im}(z) \leq 1\}$$

Then the set  $S_1 \cap S_2 \cap S_3$

- (1) is a singleton  
 (2) has exactly two elements  
 (3) has infinitely many elements  
 (4) has exactly three elements

10. If the curve  $y = y(x)$  is the solution of the differential equation

$$2(x^2 + x^{5/4})dy - y(x + x^{1/4})dx = 2x^{9/4} dx, \quad x > 0$$

which passes through the point

$$\left(1, 1 - \frac{4}{3} \log_e 2\right), \text{ then the value of } y(16) \text{ is equal}$$

to :

(1)  $4\left(\frac{31}{3} + \frac{8}{3} \log_e 3\right)$       (2)  $\left(\frac{31}{3} + \frac{8}{3} \log_e 3\right)$

(3)  $4\left(\frac{31}{3} - \frac{8}{3} \log_e 3\right)$       (4)  $\left(\frac{31}{3} - \frac{8}{3} \log_e 3\right)$

11. If the sides AB, BC and CA of a triangle ABC have 3, 5 and 6 interior points respectively, then the total number of triangles that can be constructed using these points as vertices, is equal to :  
 (1) 364      (2) 240      (3) 333      (4) 360
12. If  $x, y, z$  are in arithmetic progression with common difference  $d$ ,  $x \neq 3d$ , and the determinant of the matrix  $\begin{bmatrix} 3 & 4\sqrt{2} & x \\ 4 & 5\sqrt{2} & y \\ 5 & k & z \end{bmatrix}$  is zero, then the value of  $k^2$  is  
 (1) 72      (2) 12      (3) 36      (4) 6
13. Let  $O$  be the origin. Let  $\overrightarrow{OP} = x\hat{i} + y\hat{j} - \hat{k}$  and  $\overrightarrow{OQ} = -\hat{i} + 2\hat{j} + 3x\hat{k}$ ,  $x, y \in \mathbb{R}$ ,  $x > 0$ , be such that  $|\overrightarrow{PQ}| = \sqrt{20}$  and the vector  $\overrightarrow{OP}$  is perpendicular to  $\overrightarrow{OQ}$ . If  $\overrightarrow{OR} = 3\hat{i} + z\hat{j} - 7\hat{k}$ ,  $z \in \mathbb{R}$ , is coplanar with  $\overrightarrow{OP}$  and  $\overrightarrow{OQ}$ , then the value of  $x^2 + y^2 + z^2$  is equal to  
 (1) 7      (2) 9      (3) 2      (4) 1

14. Two tangents are drawn from a point P to the circle  $x^2 + y^2 - 2x - 4y + 4 = 0$ , such that the angle between these tangents is  $\tan^{-1}\left(\frac{12}{5}\right)$ , where  $\tan^{-1}\left(\frac{12}{5}\right) \in (0, \pi)$ . If the centre of the circle is denoted by C and these tangents touch the circle at points A and B, then the ratio of the areas of  $\Delta PAB$  and  $\Delta CAB$  is :
- (1) 11 : 4    (2) 9 : 4    (3) 3 : 1    (4) 2 : 1

16. Let L be a tangent line to the parabola  $y^2 = 4x - 20$  at (6, 2). If L is also a tangent to the ellipse  $\frac{x^2}{2} + \frac{y^2}{b} = 1$ , then the value of b is equal to :
- (1) 11    (2) 14    (3) 16    (4) 20

17. The value of the limit  $\lim_{\theta \rightarrow 0} \frac{\tan(\pi \cos^2 \theta)}{\sin(2\pi \sin^2 \theta)}$  is equal to :
- (1)  $-\frac{1}{2}$     (2)  $-\frac{1}{4}$     (3) 0    (4)  $\frac{1}{4}$

15. Consider the function  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined by
- $$f(x) = \begin{cases} \left(2 - \sin\left(\frac{1}{x}\right)\right)|x|, & x \neq 0 \\ 0, & x = 0 \end{cases}.$$
- Then f is :
- (1) monotonic on  $(-\infty, 0) \cup (0, \infty)$   
 (2) not monotonic on  $(-\infty, 0)$  and  $(0, \infty)$   
 (3) monotonic on  $(0, \infty)$  only  
 (4) monotonic on  $(-\infty, 0)$  only

18. Let the tangent to the circle  $x^2 + y^2 = 25$  at the point  $R(3, 4)$  meet x-axis and y-axis at point P and Q, respectively. If r is the radius of the circle passing through the origin O and having centre at the incentre of the triangle OPQ, then  $r^2$  is equal to

(1)  $\frac{529}{64}$       (2)  $\frac{125}{72}$       (3)  $\frac{625}{72}$       (4)  $\frac{585}{66}$

19. If the Boolean expression  $(p \wedge q) \otimes (p \otimes q)$  is a tautology, then  $\otimes$  and  $\otimes$  are respectively given by  
 (1)  $\rightarrow, \rightarrow$     (2)  $\wedge, \vee$     (3)  $\vee, \rightarrow$     (4)  $\wedge, \rightarrow$

20. If the equation of plane passing through the mirror image of a point  $(2, 3, 1)$  with respect to line  $\frac{x+1}{2} = \frac{y-3}{1} = \frac{z+2}{-1}$  and containing the line  $\frac{x-2}{3} = \frac{1-y}{2} = \frac{z+1}{1}$  is  $\alpha x + \beta y + \gamma z = 24$ , then  $\alpha + \beta + \gamma$  is equal to :

(1) 20      (2) 19      (3) 18      (4) 21

### SECTION-B

1. If  $1, \log_{10}(4^x - 2)$  and  $\log_{10}\left(4^x + \frac{18}{5}\right)$  are in arithmetic progression for a real number x, then the value of the determinant

$$\begin{vmatrix} 2\left(x - \frac{1}{2}\right) & x-1 & x^2 \\ 1 & 0 & x \\ x & 1 & 0 \end{vmatrix} \text{ is equal to :}$$

2. Let  $f : [-1, 1] \rightarrow \mathbb{R}$  be defined as  $f(x) = ax^2 + bx + c$  for all  $x \in [-1, 1]$ , where  $a, b, c \in \mathbb{R}$  such that  $f(-1) = 2$ ,  $f'(-1) = 1$  and for  $x \in (-1, 1)$  the maximum value of  $f''(x)$  is  $\frac{1}{2}$ . If  $f(x) \leq \alpha$ ,  $x \in [-1, 1]$ , then the least value of  $\alpha$  is equal to \_\_\_\_\_.

3. Let  $f : [-3, 1] \rightarrow \mathbb{R}$  be given as

$$f(x) = \begin{cases} \min \{(x+6), x^2\}, & -3 \leq x \leq 0 \\ \max \{\sqrt{x}, x^2\}, & 0 \leq x \leq 1. \end{cases}$$

If the area bounded by  $y = f(x)$  and  $x$ -axis is  $A$ , then the value of  $6A$  is equal to \_\_\_\_\_.

4. Let  $\tan \alpha$ ,  $\tan \beta$  and  $\tan \gamma$ ;  $\alpha, \beta, \gamma \neq \frac{(2n-1)\pi}{2}$ ,  $n \in \mathbb{N}$  be the slopes of three line segments  $OA$ ,  $OB$  and  $OC$ , respectively, where  $O$  is origin. If circumcentre of  $\triangle ABC$  coincides with origin and its orthocentre lies on  $y$ -axis, then the value

of  $\left( \frac{\cos 3\alpha + \cos 3\beta + \cos 3\gamma}{\cos \alpha \cos \beta \cos \gamma} \right)^2$  is equal to :

5. Consider a set of  $3n$  numbers having variance 4. In this set, the mean of first  $2n$  numbers is 6 and the mean of the remaining  $n$  numbers is 3. A new set is constructed by adding 1 into each of first  $2n$  numbers, and subtracting 1 from each of the remaining  $n$  numbers. If the variance of the new set is  $k$ , then  $9k$  is equal to \_\_\_\_\_.

6. Let the coefficients of third, fourth and fifth terms in the expansion of  $\left(x + \frac{a}{x^2}\right)^n$ ,  $x \neq 0$ , be in the ratio  $12 : 8 : 3$ . Then the term independent of  $x$  in the expansion, is equal to \_\_\_\_\_.

7. Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$  and  $B = \begin{bmatrix} \alpha \\ \beta \end{bmatrix} \neq \begin{bmatrix} 0 \\ 0 \end{bmatrix}$  such that  $AB = B$  and  $a + d = 2021$ , then the value of  $ad - bc$  is equal to \_\_\_\_\_.

8. Let  $\vec{x}$  be a vector in the plane containing vectors  $\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$  and  $\vec{b} = \hat{i} + 2\hat{j} - \hat{k}$ . If the vector  $\vec{x}$  is perpendicular to  $(3\hat{i} + 2\hat{j} - \hat{k})$  and its projection on  $\vec{a}$  is  $\frac{17\sqrt{6}}{2}$ , then the value of  $|\vec{x}|^2$  is equal to \_\_\_\_\_.

9. Let  $I_n = \int_1^e x^{19} (\log|x|)^n dx$ , where  $n \in \mathbb{N}$ . If

$(20)I_{10} = \alpha I_9 + \beta I_8$ , for natural numbers  $\alpha$  and  $\beta$ , then  $\alpha - \beta$  equal to \_\_\_\_\_.

10. Let P be an arbitrary point having sum of the squares of the distance from the planes  $x + y + z = 0$ ,  $lx - nz = 0$  and  $x - 2y + z = 0$ , equal to 9. If the locus of the point P is  $x^2 + y^2 + z^2 = 9$ , then the value of  $l - n$  is equal to \_\_\_\_\_.