

## **AIPMT - 2002**

**Q.1** A mass is suspended separately by two different springs in successive order then time period is t<sub>1</sub> and t<sub>2</sub> respectively. If it is connected by both spring as shown in figure then time period is t<sub>0</sub>, the correct relation is: -



- (3)  $t_0^{-1} = t_1^{-1} + t_2^{-1}$  (4)  $t_0 = t_1 + t_2$
- When an oscillator completes 100 oscillation its 0.2 amplitude reduced to  $\frac{1}{2}$  of initial value. What will be its amplitude, when it completes 200 oscillation: -
  - $(1) \frac{1}{8}$   $(2) \frac{2}{3}$   $(3) \frac{1}{6}$   $(4) \frac{1}{9}$
- 0.3 A circular disc is to be made by using iron and aluminium so that it acquired maximum moment of inertia about geometrical axis. It is possible with:-
  - (1) Aluminium at interior and iron surround to it
  - (2) Iron at interior and aluminium surround to it
  - (3) Using iron and aluminium layers in alternate order
  - (4) Sheet of iron is used at both external surface and aluminium sheet as internal layers
- **Q.4** For the given incident ray as shown in figure, the condition of total internal reflection of this ray the minimum refractive index of prism will be: -



- (1)  $\frac{\sqrt{3}+1}{2}$  (2)  $\frac{\sqrt{2}+1}{2}$  (3)  $\sqrt{\frac{3}{2}}$  (4)  $\sqrt{\frac{7}{6}}$
- Q.5 The value of plank's constant is: -
  - (1)  $6.63 \times 10^{-34} \text{ J/s}$
  - (2)  $6.63 \times 10^{-34} \text{ kg-m}^2/\text{s}$
  - (3)  $6.63 \times 10^{-34} \text{ kg-m}^2$
  - (4)  $6.63 \times 10^{-34} \text{ J-s}^{-1}$

- Displacement between max. P.E. position and **Q.6** max. K.E. position for a particle excuting simple harmonic motion is: -
  - $(1) \pm \frac{a}{2}$
- (2) + a
- $(3) \pm a$
- (4) 1
- Q.7 A disc is rotating with angular speed ω. If a child sits on it, what is conserved: -
  - (1) Linear momentum
  - (2) Angular momentum
  - (3) Kinetic energy
  - (4) Potential energy
- **Q.8** Which is having minimum wavelength: -
  - (1) X-rays
- (2) Ultra violet rays
- (3)  $\gamma$ -rays
- (4) Cosmic rays
- Q.9 If particles are moving with same velocity, then De-Broglie wavelength is maximum for : -
  - (1) Proton
- (2) α-particle
- (3) Neutron
- (4) β-particle
- Q.10 When ultraviolet rays incident on metal plate then photoelectric effect does not occur, it occurs by incidence of: -
  - (1) Infrared rays
- (2) X-rays
- (3) Radio wave
- (4) Light wave
- Q.11 What is the cause of "Green house effect": -
  - (1) Infra-red rays
- (2) Ultra violet rays
- (3) X-rays
- (4) Radio waves
- Which of the following is not the property of Q.12 cathode rays: -
  - (1) It produces heating effect
  - (2) It does not deflecte in electric field
  - (3) It casts shadow
  - (4) It produces flurosence
- Q.13 A solid sphere of radius R is placed on smooth horizontal surface. A horizontal force 'F' is applied at height 'h' from the lowest point. For the maximum, acceleration of centre of mass, which is correct: -
  - (1) h = R
  - (2) h = 2R
  - (3) h = 0
  - (4) No relation between h and R

- Q.14 Diameter of human eye lens is 2 mm. What will be the minimum distance between two points to resolve them, which are situated at a distance of 50 meter from eye. The wavelength of light is 5000 Å :-
  - (1) 2.32 m
- (2) 4.28 mm
- (3) 1.25 cm
- (4) 12.48 cm
- A bulb is located on a wall. Its image is to be Q.15 obtained on a parallel wall with the help of convex lens. If the distance between parallel walls is 'd' then required focal length of lens placed in between the walls is: -
  - (1) Only  $\frac{d}{4}$
  - (2) Only  $\frac{d}{2}$
  - (3) More than  $\frac{d}{4}$  but less than  $\frac{d}{2}$
  - (4) Less than or equal to  $\frac{d}{4}$
- The Wien's displacement law express relation Q.16 between: -
  - (1) Wavelength corresponding to maximum energy and temperature
  - (2) Radiation energy and wavelength
  - (3) Temperature and wavelength
  - (4) Colour of light and temperature
- 0.17 Which of the following is best close to an ideal black body: -
  - (1) Black lamp
  - (2) Cavity maintained at constant temperature
  - (3) Platinum black
  - (4) A lump of charcoal heated to high temp.
- For a black body at temperature 727°C, its Q.18 radiating power is 60 watt and temperature of surrounding is 227°C. If temperature of black body is changed to 1227°C then its radiating power will be: -
  - (1) 304 W
- (2) 320 W
- (3) 240 W
- (4) 120 W
- Q.19 Consider two rods of same length and different specific heats  $(S_1, S_2)$ , conductivities  $(K_1, K_2)$  and area of cross-sections (A1, A2) and both having temperature T<sub>1</sub> and T<sub>2</sub> at their ends. If rate of loss of heat due to conduction is equal, then :-

$$(1) K_1 A_1 = K_2 A$$

(1) 
$$K_1A_1 = K_2A_2$$
 (2)  $\frac{K_1A_1}{S_1} = \frac{K_2A_2}{S_2}$ 

$$(3) K_2 A_1 = K_1 A_2$$

(3) 
$$K_2A_1 = K_1A_2$$
 (4)  $\frac{K_2A_1}{S_2} = \frac{K_1A_2}{S_1}$ 

- Q.20 The efficiency of carnot engine is 50% and temperature of sink is 500K. If temperature of source is kept constant and its efficiency raised to 60%, then the required temperature of the sink will be: -
  - (1) 100 K
- (2) 600 K
- (3) 400 K
- (4) 500 K
- Q.21 Unit of Stefan's constant is: -
  - (1) Watt-m<sup>2</sup>-K<sup>4</sup>
- (2) Watt- $m^2/K^4$
- (3) Watt/m<sup>2</sup>–K
- (4) Watt/m<sup>2</sup>K<sup>4</sup>
- Q.22 Number of atom per unit cell in B.C.C.: -
  - (1)9
- (2)4
- (3)2
- (4) 1
- Q.23 An object of mass 3kg is at rest. Now a force of  $\vec{F} = 6t^2\hat{i} + 4t\hat{j}$  is applied on the object then velocity of object at t = 3 second is:
  - (1)  $18\hat{i} + 3\hat{j}$
- (2)  $18\hat{i} + 6\hat{j}$
- (3)  $3\hat{i} + 18\hat{i}$
- (4)  $18\hat{i} + 4\hat{j}$
- 0.24 A body of mass m is placed on earth surface which is taken from earth surface to a height of h = 3R then change in gravitational potential energy is: -

  - $(1) \frac{\text{mgR}}{4} \qquad (2) \frac{2}{3} \text{mgR}$
  - (3)  $\frac{3}{4}$  mgR (4)  $\frac{\text{mgR}}{2}$
- Q.25 A point P consider at contact point of a wheel on ground which rolls on ground without sliping then value of displacement of point P when wheel completes half of rotation (If radius of wheel is 1m): -
  - (1) 2m
- (2)  $\sqrt{\pi^2 + 4}$  m
- $(3) \pi m$
- (4)  $\sqrt{\pi^2 + 2}$  m
- Q.26 A block of mass 10 kg placed on rough horizontal surface having coefficient of friction µ = 0.5, if a horizontal force of 100 N acting on it then acceleration of the block will be: -
  - (1)  $10 \text{ m/s}^2$
- $(2) 5 \text{ m/s}^2$
- $(3) 15 \text{ m/s}^2$
- $(4) 0.5 \text{ m/s}^2$

- Q.27 A lift of mass 1000 Kg which is moving with acceleration of 1 m/s<sup>2</sup> in upward direction, then the tension developed in string which is connected to lift is:
  - (1) 9800 N
- (2) 10, 800 N
- (3) 11000 N
- (4) 10, 000 N
- Q.28 A particle (A) is droped from a height and another particle (B) is projected in horizontal direction with speed of 5 m/s from the same height then correct statement is:
  - (1) Particle (A) will reach at ground first with respect to particle (B)
  - (2) Particle (B) will reach at ground first with respect to particle (A)
  - (3) Both particles will reach at ground simultaneously
  - (4) Both particles will reach at ground with same speed
- Q.29 A rod of length is 3m and its mass acting per unit length is driectly proportional to distance x from one of its end then its centre of gravity from that end will be at:
  - (1) 1.5 m
- (2) 2 m
- (3) 2.5 m
- (4) 3.0 m
- Q.30 If kinetic energy of a body is increased by 300% than percentage change in momentum will be
  - (1) 100%
- (2) 150%
- (3) 265%
- (4) 73.2%

(3)48

Q.31 For a transistor  $\frac{I_C}{I_E}$  = 0.96, then current gain for

common emitter configuration : -

- (1) 12
- (2) 6
- (4) 24
- Q.32 A wave travelling in positive X-direction with A = 0.2 m velocity = 360 m/s and  $\lambda = 60$  m, then correct expression for the wave is : -

(1) 
$$y = 0.2 \sin \left[2\pi \left(6t + \frac{x}{60}\right)\right]$$

(2) 
$$y = 0.2 \sin \left[\pi \left(6t + \frac{x}{60}\right)\right]$$


(3) 
$$y = 0.2 \sin \left[2\pi \left(6t - \frac{x}{60}\right)\right]$$

(4) 
$$y = 0.2 \sin \left[\pi (6t - \frac{x}{60})\right]$$

- Q.33 A whistle revolves in a circle with angular speed  $\omega = 20$  rad/sec using a string of length 50 cm. If the frequency of sound from the whistle is 385 Hz, then what is the minimum frequency heard by an observer which is far away from the centre:  $-(V_{sound} = 340 \text{ m/s})$ 
  - (1) 385 Hz
- (2) 374 Hz
- (3) 394 Hz
- (4) 333 Hz
- Q.34 In a PN junction: -
  - (1) High potential at N side and low potential at P side
  - (2) High potential at P side and low potential at N side
  - (3) P and N both are at same potential
  - (4) Undetermined
- Q.35 The given truth table is for which logic gate: -

A	В	Y
1	1	0
0	1	1
1	0	1
0	0	1

- (1) NAND (2) XOR (3) NOR (4) OR
- **Q.36** For the given circuit of P-N junction diode which is correct:



- (1) In F.B. the voltage across R is V
- (2) In R.B. the voltage across R is V
- (3) In F.B. the voltage across R is 2 V
- (4) In R.B. the voltage across R is 2 V
- **Q.37** Specific resistance of a conductor increases with:
  - (1) Increase in temperature
  - (2) Increase in cross section area
  - (3) Increase in cross section and decrease in length
  - (4) Decrease in cross section area
- **Q.38** For a series LCR circuit the power loss at resonance is:
  - $(1) \frac{V^2}{\left[\omega L \frac{1}{\omega C}\right]}$
- (2)  $I^2L\omega$
- (3) I<sup>2</sup>R
- (4)  $\frac{V^2}{C\omega}$

- Q.39 Some charge is being given to a conductor. Then its potential: -
  - (1) Is maximum at surface
  - (2) Is maximum at centre
  - (3) Is remain same throughout the conductor
  - (4) Is maximum somewhere between surface and
- For a cell terminal P.D. is 2.2V when circuit is Q.40 open and reduces to 1.8V when cell is connected to a resistance of  $R = 5\Omega$ . Determine internal resistance of cell (r) is then: -
  - $(1) \frac{10}{9} \Omega$
- $(2) \frac{9}{10} \Omega$
- $(3) \frac{11}{9} \Omega$ 
  - $(4) \frac{5}{9} \Omega$
- To convert a galvanometer into a voltmeter one Q.41 should connect a: -
  - (1) High resistance in series with galvanometer
  - (2) Low resistance in series with galvanometer
  - (3) High resistance in parallel with galvanometer
  - (4) Low resistance in parallel with galvanometer
- Q.42 A capacitor of capacity C<sub>1</sub> charged upto V volt and then connected to an uncharged capacitor C<sub>2</sub>. Then final P.D. across each will be

  - (1)  $\frac{C_2 V}{C_1 + C_2}$  (2)  $\frac{C_1 V}{C_1 + C_2}$

  - $(3)\left(1+\frac{C_2}{C_1}\right) \qquad (4)\left(1-\frac{C_2}{C_1}\right)V$
- Q.43 Identical charges (-q) are placed at each corner of a cube of side 'b' then electrical potential energy of charge (+q) which is placed at centre of cube will be
  - (1)  $\frac{-4\sqrt{2}q^2}{\pi \in_0 b}$  (2)  $\frac{-8\sqrt{2}q^2}{\pi \in_0 b}$
  - (3)  $\frac{-4q^2}{\sqrt{3}\pi \in_0 b}$  (4)  $\frac{8\sqrt{2}q^2}{4\pi \in_0 b}$
- Which of the following are suitable for the **Q.44** fusion process: -
  - (1) Light nuclei
  - (2) heavy nuclei
  - (3) Element must be lying in the middle of the periodic table
  - (4) Middle elements, which are lying on binding energy curve

- The magnetic field of given length of wire for Q.45 single turn coil at its centre is 'B' then its value for two turns coil for the same wire is: -
  - (1)  $\frac{B}{4}$
- (2)  $\frac{B}{2}$
- (3) 4B
- (4) 2B
- 0.46 A charge 'q' moves in a region where electric field and magnetic field both exist, then force on it is: -
  - $(1) a (\vec{V} \times \vec{B})$
- (2)  $\vec{a} \cdot \vec{E} + \vec{a} \cdot (\vec{V} \times \vec{B})$
- $(3) q \overrightarrow{E} + q (\overrightarrow{B} \times \overrightarrow{V}) (4) q \overrightarrow{B} + q (\overrightarrow{E} \times \overrightarrow{V})$
- Two bar magnets having same geometry with 0.47 magnetic moments M and 2M, are firstly placed in such a way that their similer poles are same side then its time period of oscillation is  $T_1$ . Now the polarity of one of the magnet is reversed then time period of oscillation is  $T_2$ ,
  - (1)  $T_1 < T_2$
- (2)  $T_1 = T_2$
- (3)  $T_1 > T_2$
- (4)  $T_2 = \infty$
- The velocity of electromagnetic wave is parallel Q.48 to:-
  - (1)  $\vec{B} \times \vec{E}$
- (2)  $\vec{E} \times \vec{B}$
- $(3) \vec{E}$
- (4)  $\vec{B}$
- Q.49 A sample of radioactive element containing  $4 \times 10^{16}$  active nuclei. Half life of element is 10 days, then number of decayed nuclei after 30 days : -(2)  $2 \times 10^{16}$ 
  - $(1) 0.5 \times 10^{16}$
- $(3) 3.5 \times 10^{16}$
- (4)  $1 \times 10^{16}$
- A deuteron is bombarded on 8O16 nucleus then O.50  $\alpha$ -particle is emitted then product nucleus is -
  - $(1)_{7}N^{13}$
- $(2)_{5}B^{10}$
- $(3)_{4}Be^{9}$
- $(4) _{7}N^{14}$
- <sub>92</sub>U<sup>235</sup>, nucleus absorb a neutron and disintegrate Q.51 in  $_{54}Xe^{139}$ ,  $_{38}Sr^{94}$  and x So, What will be the product x:-
  - (1) 3 neutrons
- (2) 2 neutrons
- (3)  $\alpha$  partical
- (4) β partical
- Q.52 In Hydrozen atom, energy of first excited state is - 3.4 eV. Then find out KE of same orbit of Hydrogen atom: -
  - (1) + 3.4 eV
- (2) + 6.8 eV
- (3) 13.6 eV
- (4) + 13.6 eV

- Reaction BaO<sub>2</sub> (s)  $\Longrightarrow$  BaO (s) + O<sub>2</sub>(g); Q.53
  - $\Delta H = +$  ve. In equilibrium condition. Pressure of O<sub>2</sub> is depens on : -
  - (1) Increase mass of BaO<sub>2</sub>
  - (2) Increase mass of BaO
  - (3) Increase temp. on Eq<sup>m</sup>.
  - (4) Increase mass of BaO2 and BaO both
- Q.54 Solubility of MX<sub>2</sub> - type electrolytes is  $0.5 \times 10^{-4}$  Mole/lit. then find out  $K_{sp}$  of elctrolytes: -
  - $(1) 5 \times 10^{-12}$
- (2)  $25 \times 10^{-10}$
- (3)  $1 \times 10^{-13}$
- $(4) 5 \times 10^{-13}$
- Q.55 1 M and 2.5 litre NaOH solution mixed with another 0.5 M and 3 litre NaOH solution. Then find out molarity of resultant solution : -
  - (1) 0.80 M
- (2) 1.0 M
- (3) 0.73 M
- (4) 0.50 M
- Q.56 Which has highest pH: -
  - (1) CH<sub>3</sub>COOK
- (2) Na<sub>2</sub>CO<sub>3</sub>
- (3) NH<sub>4</sub>Cl
- (4) NaNO<sub>3</sub>
- Solution of 0.1 N NH<sub>4</sub>OH and 0.1 N NH<sub>4</sub>Cl has Q.57 pH 9.25, Then find out pkb of NH<sub>4</sub>OH: -
  - (1)9.25
- (2)4.75
- (3) 3.75
- (4) 8.25
- Q.58 Vander waal's real gas, act as a ideal gas, at which conditions: -
  - (1) High temp., Low pressure
  - (2) Low temp., High pressure
  - (3) High temp., High pressure
  - (4) Low temp., Low pressure
- Q.59 Unit of entropy is: -
  - $(1) \text{ JK}^{-1} \text{ mol}^{-1}$
- (2) J mol<sup>-1</sup>
- (3)  $J^{-1}K^{-1} \text{ mol}^{-1}$
- (4) JK mol<sup>-1</sup>
- **Q.60** In a closed insulated container a liquid is stirred with a paddle to increase the temperature which of the following is true: -
  - (1)  $\Delta E = W \neq 0$ , q = 0
  - (2)  $\Delta E = W = q \neq 0$
  - (3)  $\Delta E = 0$ ,  $W = q \neq 0$
  - (4)  $W = 0 \Delta E = q \neq 0$
- Q.61 2 mole of ideal gas at 27°C temp. is expanded reversibly from 2 lit. to 20 lit. Find entropy change (R = 2 cal/mol K): -
  - (1)92.1
- (2) 0
- (3)4
- (4)9.2

- Heat of combustion  $\Delta H^{\circ}$  for C(s), H<sub>2</sub>(g) and Q.62  $CH_4(g)$  are -94, -68 and -213 Kcal/mol. then  $\Delta H^{\circ}$  for  $C(s) + 2H_2(g) \rightarrow CH_4(g)$  is : -
  - (1) 17 Kcal
- (2) 111 Kcal
- (3) 170 Kcal
- (4) 85 Kcal
- $3A \rightarrow 2B$ , rate of reaction  $\frac{+d[B]}{dt}$  is equals to : -Q.63
  - $(1) \frac{3}{2} \frac{d[A]}{dt}$   $(2) \frac{2}{3} \frac{d[A]}{dt}$
  - (3)  $-\frac{1}{3}\frac{d[A]}{dt}$  (4)  $+2\frac{d[A]}{dt}$
- $2A \rightarrow B + C$ Q.64

It would be a zero order reaction when: -

- (1) The rate of reaction is proportional to square of conc. of A
- (2) The rate of reaction remains same at any conc. of A
- (3) The rate remains unchanged at any conc. of B and C
- (4) The rate of reaction doubles if conc. of B is increased to double
- Q.65 Which has maximum molecules: -
  - (1)  $7 \text{ gm N}_2$
- $(2) 2 gm H_2$
- (3) 16 gm NO<sub>2</sub>
- (4)  $16 \text{ gm } O_2$
- **Q.66** A solution contains non volatile solute of molecular mass M2. Which of the following can be used to calculate the molecular mass of solute in terms of osmotic pressure : -

(1) 
$$M_2 = \left(\frac{m_2}{\pi}\right) VRT$$
 (2)  $M_2 = \left(\frac{m_2}{V}\right) \frac{RT}{\pi}$ 

(3) 
$$M_2 = \left(\frac{m_2}{V}\right) \pi R T$$
 (4)  $M_2 = \left(\frac{m_2}{V}\right) \frac{\pi}{R T}$ 

**Note:**  $m_2 \rightarrow$  mass of solute

 $V \rightarrow Volume of solution$ 

 $p \rightarrow Osmotic pressure$ 

- A solution containing components A and B **O.67** follows Raoult's law: -
  - (1) A B attraction force is greater than A Aand B - B
  - (2) A B attraction force is less than A Aand B - B
  - (3) Attraction force remains same in A A and
  - (4) Volume of solution is different from sum of volume of solute and solvent

- **Q.68** Which reaction is not feasible: -
  - (1)  $2 KI + Br_2 \rightarrow 2KBr + I_2$
  - (2) 2 KBr +  $I_2 \rightarrow 2KI + Br_2$
  - (3) 2 KBr + Cl<sub>2</sub>  $\rightarrow$  2KCl + Br<sub>2</sub>
  - (4)  $2H_2O + 2F_2 \rightarrow 4HF + O_2$
- Q.69 In electrolysis of NaCl when Pt electrode is taken then  $H_2$  is liberated at cathode while with Hg cathode it forms sodium amalgam: -
  - (1) Hg is more inert than Pt
  - (2) More voltage is required to reduce H<sup>+</sup> at Hg than at Pt
  - (3) Na is dissolved in Hg while it does not dissolve in Pt
  - (4) Conc. of H<sup>+</sup> ions is larger when Pt electrode is taken
- Q.70 Which of the following statement is true: -
  - (1) Silicon exhibits 4 coordination number in its compound
  - (2) Bond energy of  $F_2$  is less than  $Cl_2$
  - (3) Mn(III) oxidation state is more stable than Mn (II) in aqueous state
  - (4) Elements of 15<sup>th</sup> gp shows only + 3 and + 5 oxidation states
- **Q.71** Which of the following order is wrong: -
  - (1)  $NH_3 < PH_3 < AsH_3 Acidic$
  - (2)  $Li < Be < B < C I^{st}$  IP
  - (3)  $Al_2O_3 < MgO < Na_2O < K_2O$  Basic
  - (4)  $Li^+ < Na^+ < K^+ < Cs^+$  Ionic radius
- Q.72 General electronic configuration of lanthanides is
  - $(1)\,(n-2)\;f^{\,1-14}\,(n-1)\;s^2p^6d^{0-1}\,ns^2$
  - (2)  $(n-2) f^{10-14} (n-1) d^{0-1} ns^2$
  - (3) (n-2) f<sup>0-14</sup>(n-1) d<sup>10</sup> ns<sup>2</sup>
  - (4)  $(n-2) d^{0-1} (n-1) f^{1-14} ns^2$
- Q.73 An atom has electronic configuration  $1s^2 2s^2 2p^6$   $3s^2 3p^6 3d^3 4s^2$ , you will place it in which group:
  - (1) Fifth
- (2) Fifteenth
- (3) Second
- (4) Third
- **Q.74** Which of the following is iso-electronic:
  - $(1)\ CO_2\ ,\ NO_2$
- (2)  $NO_2^-$  ,  $CO_2$
- (3) CN<sup>-</sup>, CO
- (4) SO<sub>2</sub>, CO<sub>2</sub>
- **Q.75** Which of the following has  $p_{\pi} d_{\pi}$  bonding
  - (1)  $NO_3^-$
- (2)  $SO_3^{-2}$
- (3)  $BO_3^{-3}$
- (4)  $CO_3^{-2}$

- Q.76 In NO<sub>3</sub><sup>-</sup> ion number of bond pair and lonepair of electrons on nitrogen atom are: -
  - (1) 2, 2
- (2) 3, 1
- (3) 1, 3
- (4) 4, 0
- Q.77 Which of the following shows maximum number of oxidation states:-
  - (1) Cr
- (2) Fe
- (3) Mn
- (4) V
- Q.78 Atomic number of Cr and Fe are respectively 24 and 26, which of the following is paramagnetic with the spin of electron:
  - (1)  $[Cr(CO)_6]$
- (2) [Fe(CO)<sub>5</sub>]
- (3)  $[Fe(CN)_6]^{-4}$
- (4)  $\left[ \text{Cr}(\text{NH}_3)_6 \right]^{+3}$
- Q.79 The hypothetical complex chloro diaquatriammine cobalt (III) chloride can be represented as:-
  - (1)  $[CoCl(NH_3)_3(H_2O)_2]Cl_2$
  - (2)  $[Co(NH_3)_3(H_2O)Cl_3]$
  - (3)  $[Co(NH_2)_3(H_2O)_2Cl]$
  - (4) [Co(NH<sub>3</sub>)<sub>3</sub>(H<sub>2</sub>O)<sub>3</sub>Cl<sub>3</sub>]
- **Q.80** In the silver plating of copper,  $K[Ag(CN)_2]$  is used instead of  $AgNO_3$ . The reason is:
  - (1) A thin layer of Ag is formed on Cu
  - (2) More voltage is required
  - (3) Ag<sup>+</sup> ions are completely removed from solution
  - (4) Less availability of Ag<sup>+</sup> ions, as Cu can not displace Ag from [Ag(CN)<sub>2</sub>]<sup>-</sup> ion
- Q.81 CuSO<sub>4</sub> when reacts with KCN forms CuCN, which is insoluble in water. It is soluble in excess of KCN, due to formation of the following complex:
  - $(1) K_2[Cu(CN)_4]$
- (2)  $K_3[Cu(CN)_4]$
- (3) CuCN<sub>2</sub>
- $(4) Cu[KCu(CN)_4]$
- Q.82 Position of non polar & polar part in miscell
  - (1) Polar at outer surface but non polar at inner surface
  - (2) Polar at inner surface non polar at outer surface
  - (3) Distributed over all the surface
  - (4) Are present in the surface only
- Q.83 In borex bead test which compound is formed
  - (1) Ortho borate
  - (2) Meta borate
  - (3) Double oxide
  - (4) Tetra borate

- (1) Zn act as oxidising agent when react with HNO2
- (2) HNO<sub>3</sub> is weaker acid then H<sub>2</sub>SO<sub>4</sub> & HCl
- (3) In electrochemical series Zn is above hydrogen
- (4) NO<sub>3</sub><sup>Θ</sup> is reduced in prefference to hydronium ion
- **Q.85** IUPAC name of the following is

$$CH_2 = CH - CH_2 - CH_2 - C \equiv CH$$

- (1) 1, 5-hexenyne
- (2) 1-hexene-5-yne
- (3) 1-hexyne-5-ene
- (4) 1, 5-hexynene

Q.86 
$$C \equiv N$$

$$+ CH_3MgBr \xrightarrow{H_3O^{\oplus}} P$$
OCH<sub>3</sub>

$$\begin{array}{ccc}
OH & O \\
CH-CH_3 & C-CH_3
\end{array}$$

$$\begin{array}{ccc}
OCH_3
\end{array}$$

- Q.87 n-propyl alcohol and isopropyl alcohol can be chemically distinguished by which reagent: -
  - (1) PCl<sub>5</sub>
  - (2) Reduction
  - (3) Oxidation with Potassium dichromate
  - (4) Oznolysis
- Q.88 In the following reaction product 'P' is:-

$$R - C - Cl \xrightarrow{H_2} Pd-BaSO_4 P$$

- (1) RCH<sub>2</sub>OH
- (2) RCOOH
- (3) RCHO
- (4) RCH<sub>3</sub>

Q.89 
$$\overset{\circ}{CH_2}$$
 - C - CH<sub>3</sub> and CH<sub>2</sub> - C - CH<sub>3</sub> are  $\overset{\circ}{CH_2}$  O  $\overset{\circ}{CH_3}$ 

- (1) Resonating structures
- (2) Tautomers
- (3) Geometrical isomers
- (4) Optical isomers

- Q.90 Reactivity order of halides for dehydrohalogenation is:-
  - (1) R F > R Cl > R Br > R I
  - (2) R I > R Br > R Cl > R F
  - (3) R I > R Cl > R Br > R F
  - (4) R F > R I > R Br > R Cl

Q.91 Monomer of 
$$\begin{bmatrix} CH_3 \\ -C-CH_2 - \\ CH_3 \end{bmatrix}$$
 is:

- (1) 2-methyl propene (2) Styrene
- (3) Propylene
- (4) Ethene

$$\mathbf{Q.92} \quad \bigcirc \stackrel{\text{MgBr}}{\bigodot} \xrightarrow{\text{(i)CO}_2} \mathbf{P}$$

In the above reaction product 'P' is: -

CHO COOH
$$(1) \bigcirc O \qquad (2) \bigcirc O$$

$$OH \qquad O \qquad II$$

$$(3) \bigcirc O \qquad (4) C_6H_5 - C - C_6H$$

- Q.93 Cellulose is polymer of: -
  - (1) Glucose
- (2) Fructose
- (3) Ribose
- (4) Sucrose


Q.94 
$$CH_3CH_2CI \xrightarrow{NaCN} X \xrightarrow{Ni/H_2} Y$$

$$Z \xrightarrow{Acetic anhydride}$$

Z in the above reaction sequence is: -

- (1) CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>NHCOCH<sub>3</sub>
- (2) CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>NH<sub>2</sub>
- (3) CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CONHCH<sub>3</sub>
- (4) CH<sub>3</sub>CH<sub>2</sub>CH<sub>2</sub>CONHCOCH<sub>3</sub>
- Q.95 When phenol is treated with CHCl<sub>3</sub> and NaOH, the product formed is:-
  - (1) Benzaldehyde
- (2) Salicylaldehyde
- (3) Salicylic acid
- (4) Benzoic acid
- Q.96 The percentage of C, H and N in an organic compound are 40%, 13.3% and 46.7% respectively then emprirical formula is:
  - $(1) C_3H_{13}N_3$
- (2) CH<sub>2</sub>N
- (3) CH<sub>4</sub>N
- (4) CH<sub>6</sub>N
- **Q.97** Enzymes are made up of : -
  - (1) Edible proteins
  - (2) Proteins with specific structure
  - (3) Nitrogen containing carbohydrates
  - (4) Carbohydrates

- 0.98 Geometrical isomers are differ in : -
  - (1) Poisition of functional group
  - (2) Position of atoms
  - (3) Spatial arrangement of atoms
  - (4) Length of carbon chain
- 0.99 When CH<sub>3</sub>CH<sub>2</sub>CHCl<sub>2</sub> is treated with NaNH<sub>2</sub>, the product formed is: -
  - $(1) CH_3 CH = CH_2$
  - (2)  $CH_3 C \equiv CH$



- O.100 Which is not true statement: -
  - (1)  $\alpha$ -carbon of  $\alpha$ -amino acid is asymmetric
  - (2) All proteins are found in L-form
  - (3) Human body can synthesize all proteins they
  - (4) At pH = 7 both amino and carboxylic groups exist in ionised form
- Q.101 Which is a reducing sugar : -
  - (1) Galactose
  - (2) Gluconic acid
  - (3) β-methyl galactoside
  - (4) Sucrose
- Q.102 Significance of mimicry is -
  - (1) Attack (Offance)
  - (2) Protection (Defence)
  - (3) Both (1) & (2)
  - (4) Isolation
- Q.103 Which of the following is correct match -
  - (1) Down Syndrome =  $21^{st}$  Chromosome
  - (2) Sickel cell anaemia = X Chromosome
  - (3) Haemophilia = Y Chromosome
  - (4) Parkinson Disease = X & Y Chromosome
- Q.104 Some bacteria able to grow in Streptomycin containing medium due to -
  - (1) Natural selection
  - (2) Induced mutation
  - (3) Reproductive isolation
  - (4) Genetic drift

- Q.105 In a population, unrestricted reproductive capacity is called as -
  - (1) Biotic potential
- (2) Fertility
- (3) Carring capacity (4) Birth rate
- Q.106 Change in sequence of nucleotide in DNA is called as -
  - (1) Mutagen
- (2) Mutation
- (3) Recombination
- (4) Translation
- Q.107 Reason of fast speciation in present day crop plants is -
  - (1) Mutation
- (2) Isolation
- (3) Polyploidy
- (4) Sexual Reproduction
- Which of the following is important for Q.108 speciation: -
  - (1) Seasonal isolation
  - (2) Reproductive isolation
  - (3) Behavioural isolation
  - (4) Tropical isolation
- Q.109 Which of the following are homologous organs:-
  - (1) Wings of birds & Locust
  - (2) Wings of birds (Sparrow) & Pectoral fins of fish
  - (3) Wings of bat & Butterfly
  - (4) Legs of frog & Cockroch
- Q.110 Genetic drift oparates in : -
  - (1) Small isolated population
  - (2) Large isolated population
  - (3) Fast reproductive population
  - (4) Slow reproductive population
- **O.111** There is no life on moon due to the absence of -
  - $(1) O_2$
- (2) Water
- (3) Light
- (4) Temperature
- Q.112 According to fossils which discovered up to present time, origin and evolution of man was started from which country -
  - (1) France
- (2) Java
- (3) Africa
- (4) China
- Q.113 Impulse of heart beat originates from: -
  - (1) S.A. Node
- (2) A. V. Node
- (3) Vagus Nerve
- (4) Cardiac Nerve
- Q.114 Which cartilage is present on the end of long bones -
  - (1) Calcified cartilage
  - (2) Hyaline cartilage
  - (3) Elastic cartilage
  - (4) Fibrous cartilage

in human beings: -(1) FSH (2) LH

(3) Ascending straight line

- (3) FSH, LH, Estrogen (4) Progesteron
- Q.122 When both ovary of rat are removed then which hormone is decreased in blood: -
  - (1) Oxytocin
- (2) Prolactin
- (3) Estrogen

(1) Sigmoid

Q.115

Q.119

Melanin protect from :-

(1) U.V. rays

(3) Infrared rays

(1) Vitamin -A

(3) Vitamin - K

Lymph: -

(1) WBC and serum

some proteins

curve for Bacteria: -

(2) No movement at joint

- (4) Gonadotrophic releasing factor
- 0.123Which of the following statement is correct for node of Ranvier of nerve: -
  - (1) Neurilemma is discontinuous
  - (2) Myelin sheath is discontinuous
  - (3) Both neurilemma & Myelin sheath are discontinuous
  - (4) Covered by myelin sheath

- (3) Mutation
- (4) Gene flow
- Q.132 Which of the following occurs more than one and less than five in a chromosome: -
  - (1) Chromatid
- (2) Chromomere
- (3) Centromere
- (4) Telomere
- Q.133 Ribosomes are produced in:
  - (1) Nucleolus
- (2) Cytoplasm
- (3) Mitochondria
- (4) Golgibody

			AIPMT - 2002	
Q.134	Mitotic spindle is mainly composed of which	Q.144	During the formation of bread it becomes	
	protein:-		porous due to release of CO <sub>2</sub> by the action of : -	
	(1) Actin (2) Tubulin		(1) Yeast (2) Bacteria	
	(3) Actomyosin (4) Myoglobin		(3) Virus (4) Protozoans	
Q.135	Cancerous cells can easily be destroyed by	Q.145	In protozoa like Amoeba and Paramecium, a	
	radiations due to : -		organ is found for osmoregulation which is : -	
	(1) Rapid cell division(2) Lack of nutrition		(1) Contractile vacuole (2) Mitochondria	
	(3) Fast mutation (4) Lack of oxygen		(3) Nucleus (4) Food vacuole	
Q.136	Which fungal disease spreads by seed and	Q.146	Which of the following is absent in polluted	
	flowers : -		water : -	
	(1) Loose smut of Wheat		(1) Hydrilla (2) Water hyacinth	
	(2) Corn stunt		(3) Larva of stone fly (4) Blue green algae	
	(3) Covered smut of Barley	Q.147	What is true for individuals of same species	
	(4) Soft rot of Potato		(1) Live in same niche	
Q.137	Sequence of which of the following is used to		(2) Live in same habitat	
	know the phylogeny : -		(3) Interbreeding	
	$(1) m-RNA \qquad \qquad (2) r-RNA$		(4) Live in different habitat	
	$(3) t-RNA \qquad \qquad (4) DNA$	Q.148	In which era reptiles were dominated: -	
Q.138	Which of the following secretes toxins during		(1) Coenozoic era (2) Mesozoic era	
	storage conditions of crop plants : -		(3) Paleozoic era (4) Archaeozoic era	
	(1) Aspergillus (2) Penicillium	Q.149	Number of wild life is continuously decreasing.	
	(3) Fusarium (4) Colletotrichum		What is the main reason of this:-	
Q.139	Which of the following plants produces seeds		<ul><li>(1) Predation</li><li>(2) Cutting down of forest</li></ul>	
	but not flowers : -			
			(3) Destruction of habitat	
Q.140	Best material for the study of mitosis in		(4) Hunting In Angiosperms pollen tube liberate their male	
	laboratory : -	Q.150		
	(1) Anther (2) Root tip		gametes into the : -	
	(3) Leaf tip (4) Ovary		(1) Central cell (2) Antipodal cells	
Q.141	In five kingdom system, the main basis of		(3) Egg cell (4) Synergids	
	classification : -	Q.151	Maximum green house gas released by which	
	(1) Structure of nucleus		country:	
	<ul><li>(2) Nutrition</li><li>(3) Structure of cell wall</li></ul>		(1) India (2) France	
			(3) U.S.A. (4) Britain	
0.142	(4) Asexual reproduction Which of the following is without exception in	Q.152	What is the direction of micropyle in anatropous	
Q.142	Angiosperms : -		ovule:-	
	(1) Presence of vessels		(1) Upward (2) Downward	
		0.150	(3) Right (4) Left	
	(2) Secondary growth	Q.153	Which type of association is found in between	
	(3) Secondary growth		entomophilous flower and pollinating agent : -	
0.142	(4) Autotrophic nutrition		(1) Mutualism (2) Commonsalism	
Q.143	Which bacteria is utilized in Gober gas plant:	0.45	(3) Coperation (4) Co-evolution	
	(1) Methanogens	Q.154	In which of the following notochord is present	
	(2) Nitrifying bacteria	in embryonic stage : -		
	(3) Ammonifying bacteria		(1) All chordates (2) Some chordates	
	(4) Denitrifying bacteria		(3) Vertebrates (4) Non chordates	

				AIPMT - 2002	
Q.155	In Angiosperm all the four microspores of tetrad	Q.164	In Photosynthesis	energy from light reaction to	
	are covered by a layer which is formed by:-		dark reaction is transferred in the form of : -		
	(1) Pectocellulose (2) Callose		(1) ADP	(2) ATP	
	(3) Cellulose (4) Sporopollenin		(3) RUDP	(4) Chlorophyll	
Q.156	In which of the animal dimorphic nucleus is	Q.165	Which of the following absorb light energy for		
	found : -		photosynthesis: -		
	(1) Amoeba proteus		(1) Chlorophyll	(2) Water molecule	
	(2) Trypanosoma gambiens		(3) $O_2$	(4) RUBP	
	(3) Plasmodium vivax		Seed dormancy is	due to the : -	
	(4) Paramecium caudatum		(1) Ethylene	(2) Abscissic acid	
Q.157	Two different species can not live for long duration		(3) IAA	(4) Starch	
	in the same niche or habitat. This law is: -	Q.167	Edible part in man	go is : -	
	(1) Allen's law		(1) Mesocarp	(2) Epicarp	
	(2) Gause's law		(3) Endocarp	(4) Epidermis	
	(3) Competitive exclusion principal	Q.168	What is true for cle	eavage : -	
	(4) Weiseman's theory		(1) Size of embryo increase		
Q.158	Which of the following is a correct pair: -		(2) Size of cells de	crease	
	(1) Cuscuta – parasite		(3) Size of cells increase		
	(2) Dischidia – insectivorous		(4) Size of embryo decrease		
	(3) Opuntia – predator	Q.169	Geocarpic fruit is : -		
	(4) Capsella – hydrophyte		(1) Potato	(2) Peanut	
Q.159	Bamboo plant is growing in a far forest then		(3) Onion	(4) Garlic	
	what will be the trophic level of it: -	Q.170	In which animal nerve cell is present but bra		
	(1) First trophic level $(T_1)$		is absent : -		
	(2) Second trophic level (T <sub>2</sub> )		(1) Sponge	(2) Earthworm	
	(3) Third trophic level (T <sub>3</sub> )		(3) Cockroach	(4) Hydra	
	(4) Fourth trophic level (T <sub>4</sub> )	Q.171	In bacteria, plasmid is : -		
Q.160	Which pigment absorbs the red and farred light		<ol> <li>Extra chromosomal material</li> <li>Main DNA</li> <li>Non functional DNA</li> </ol>		
	(1) Cytochrome (2) Phytochrome				
	(3) Carotenoids (4) Chlorophyll				
Q.161	Opening and closing of stomata is due to the :-		(4) Repetative gene	e	
	<ul> <li>(1) Hormonal change in guard cells</li> <li>(2) Change in Turgor pressure of guard cells</li> <li>(3) Gaseous exchange</li> <li>(4) Respiration</li> </ul>		Transformation experiment was first performed		
			on which bacteria:-		
			(1) E. coli		
			(2) Diplococcus pneumoniae		
Q.162	How many ATP molecules produced by Aerobic		(3) Salmonella		
	oxidation of one molecule of glucose: -		(4) Pasteurella pestis		
	(1) 2 (2) 4	Q.173	Which statement	is correct for bacterial	
	(3) 38 (4) 34		transduction : -		
Q.163	Choose the correct match		(1) Transfer of some genes from one bacteria another bacteria through virus		
	Bladderwert, sundew, venus flytrap: -				
	(1) Nepanthese, Dionea, Drosera		(2) Transfer of some genes from one bacteria to		
	(2) Nepanthese, Utricularia, Vanda		another bacteria by conjugation		
	(3) Utricularia, Drosera, Dionea		(3) Bacteria obtained its DNA directly		
	(4) Dionea, Trapa, Vanda		(4) Bacteria obtain	ned DNA from other external	
	-		source		

## **AIPMT - 2002**

- 0.174 Which steroid is used for transformation: -
  - (1) Cortisol
- (2) Cholesterol
- (3) Testosteron
- (4) Progesteron
- Q.175 Main function of lenticel is: -
  - (1) Transpiration
  - (2) Guttation
  - (3) Gaseous exchange
  - (4) Bleeding
- Q.176 Which of the following is the example of sex linked disease: -
  - (1) AIDS
  - (2) Colour blindness
  - (3) Syphilis
  - (4) Gonorrhoea
- Q.177 Vessels are found in : -
  - (1) All angiosperms and some gymnosperm
  - (2) Most of the angiosperm and gymnosperms
  - (3) All angiosperms, all gymnosperms and some pteridophyta
  - (4) All pteridophyta
- Q.178 In E. Coli, during lactose metabolism repressor binds to:-
  - (1) Regulator gene (2) Operator gene
  - (3) Structural gene (4) Promoter gene
- Q.179 Four radial V.B. are found in: -
  - (1) Dicot root
- (2) Monocot root
- (3) Dicot stem
- (4) Monocot stem
- Q.180 Which of the following is the example of pleiotropic gene; -
  - (1) Haemophilia
- (2) Thalassemea
- (3) Sickle cell anaemia (4) Colour blindness
- **Q.181** A gene said to be dominant if: -
  - (1) It express it's effect only in homozygous
  - (2) It expressed only in heterozygous condition
  - (3) It expressed both in homozygous and heterozygous condition
  - (4) It never expressed in any condition
- 0.182Axillary bud and terminal bud derived from the activity of: -
  - (1) Lateral meristem
  - (2) Intercalary meristem
  - (3) Apical meristem
  - (4) Parenchyma

- Q.183 If a diploid cell is treated with colchicine then it becomes: -
  - (1) Triploid
- (2) Tetraploid
- (3) Diploid
- (4) Monoploid
- Q.184 What is the reason of formation of embryoid from pollen grain in tissue culture medium
  - (1) Cellular totipotency
  - (2) Organogenesis
  - (3) Double fertilization
  - (4) Test tube culture
- A plant of  $F_1$ -generation with genotype Q.185 "AABbCC". On selfing of this plant what is the phenotypic ratio in F<sub>2</sub>-generation : -
  - (1) 3 : 1
  - (2) 1 : 1
  - (3)9:3:3:1
  - (4) 27 : 9 : 9 : 9 : 3 : 3 : 3 : 1
- O.186 In a DNA percentage of thymine is 20% then what is the percentage of guanine: -
  - (1) 20%
- (2) 40%
- (3) 30%
- (4) 60%
- A diseased man marries a normal woman. They Q.187 get three daughter and five sons. All the daughter were diseased and sons were normal. The gene of this disease is: -

  - (1) Sex linked dominant
  - (2) Sex linked recessive
  - (3) Sex limited character
  - (4) Autosomal dominant
- Out of 64 codons, 61 codons code for 20 types Q.188 of amino acid it is called: -
  - (1) Degeneracy of genetic code
  - (2) Overlapping of gene
  - (3) Wobbling of codon
  - (4) Universility of codons
- Q.189 Jacob and Monad studied lactose metabolism in E.Coli and proposed operon concept. Operon concept applicable for : -
  - (1)All prokaryotes
  - (2) All prokaryotes and some eukaryotes
  - (3) All prokaryotes and all eukaryotes
  - (4) All prokaryotes and some protozoanes
- Q.190 Collagen is: -
  - (1) Fibrous protein (2) Globular protein
  - (3) Lipid
- (4) Carbohydrate

## **AIPMT - 2002**

- Q.191 Lipids are insoluble in water because lipids molecules are: -
  - (1) Hydrophilic
- (2) Hydrophobic
- (3) Neutral
- (4) Zwitter ions
- Q.192 Exon part of m-RNAs have code for : -
  - (1) Protein
- (2) Lipid
- (3) Carbohydrate
- (4) Phospholipid
- Q.193 Which of the following statement is true: -
  - (1) Vessels are multicellular and with wide lumen
  - (2) Tracheids are multicellular and with narrow lumen
  - (3) Vessels are unicellular and with narrow lumen
  - (4) Tracheids are unicellular and with wide lumen
- Q.194 Which of the following enzymes are used to join bits of DNA: -
  - (1) Ligase
  - (2) Primase
  - (3) DNA polymerase
  - (4) Endonuclease
- **Q.195** Which of the following crops have been brought to India from New world:
  - (1) Cashewnut, potato, rubber
  - (2) Mango, tea
  - (3) Tea, rubber, mango
  - (4) Coffee
- Q.196 Manipulation of DNA in genetic engineering became possible due to the discovery of: -
  - (1) Restriction endonuclease
  - (2) DNA ligase
  - (3) Transcriptase
  - (4) Primase
- Q.197 There are three genes a, b and c. The percentage of crossing over between a and b is 20%, b and c is 28% and a and c is 8%. What is the sequence of genes on chromosome
  - (1) b, a, c
- (2) a, b, c
- (3) a, c, b
- (4) None
- Q.198 Which of the following reunites the exon segments after RNA splicing: -
  - (1) RNA polymerase
  - (2) RNA primase
  - (3) RNA ligase
  - (4) RNA proteases

- Q.199 Introduction of food plants developed by genetic engineering is not desirable because -
  - (1) Economy of developing countries may suffer
  - (2) These products are less tasty as compared to the already existing products
  - (3) This method is costly
  - (4) There is danger of coming viruses, allergens and toxins with introduced crop
- Q.200 Nucleus of a donor embryonal cell/somatic cell is transferred to an enucleated egg cell. Then after the formation of organism, what shell be true: -
  - (1) Organism will have extranuclear genes of the donor cell
  - (2) Organism will have extra nuclear genes of recipient cell
  - (3) Organism will have extra nuclear genes of both donor and recipient cell
  - (4) Organism will have nuclear genes of recipient cell