

CONNECTING STAKEHOLDERS TO ECOSYSTEM CHANGE WITH ECOLOGICAL FORECAST MODELS IN THE CALIFORNIA CURRENT SYSTEM

Clarissa Anderson (SCCOOS)


*Southern California Coastal Ocean Observing System
Scripps Institution of Oceanography*

Henry Ruhl (CeNCOOS)

*Central and Northern California Ocean Observing System
Monterey Bay Aquarium Research Institute*

Jan Newton (NANOOS)

*Northwest Association of Networked Ocean Observing Systems
University of Washington*




NOAA Integrated Ocean Observing System (IOOS) Coastal Ocean Modeling Testbed: ROMS domains on the west coast: **WCOFS**, **UCSC**, **CA**, **OSU**, **LiveOcean (UW)**. Despite considerable domain overlap, important differences exist between models in terms of resolution, forcing, biogeochemistry and data assimilation

Global Ocean Observing System (GOOS)

GOOS Regional Alliances (GRAs)


GRAs are coalitions of nations and/or institutions which share GOOS principles and goals, but are mostly concerned with local priorities and organized around regional seas or coastal environments. Thirteen GRAs represent different regions of the globe, emphasizing regional priorities, differing by need, resources and culture.



GOOS utilizes the Framework for Ocean Observing to guide its implementation of an integrated and sustained ocean observing system. This systems approach, designed to be flexible and to adapt to evolving scientific, technological and societal needs, helps deliver an ocean observing system with maximized user base and societal impact.



Transforming data for the public and decision-makers



CONSISTENT NATIONAL CAPABILITY

Observations

Data Management

Forecasts/Modeling

User Products

**Outreach and
Education**

Leverage and Link

assure

Regional
Associations


engage

DIVERSE LOCAL STAKEHOLDERS




Each IOOS region supports sustained ocean observing programs
Regional data portals are the gateway to data, ecological forecasts, and derived products


NANOOS Visualization System





CeNCOOS Data Portal




SCCOOS Data Portal



...and all are interoperable with U.S. IOOS,
thus filling regional and national needs



- NVS
- Products
- Education
- Resources
- Merchandise
- Log In
- New Account



Observing extreme events with autonomous underwater vehicles

operational Spray glider lines in California – SCCOOS + CeNCOOS



California Underwater Glider Network

Longest sustained glider lines in the world

PI: Dan Rudnick



Spray Glider





FY18 “Fill the Gaps” Funds

-added new alongshore glider line

<https://spraydata.ucsd.edu/SoCal-index/>


MBON eDNA: The drive for automation

- Fleets of Long Range AUVs with Environmental Sample Processors (ESPs)
- A new window for observing life in the sea



- 2,000 kilometers at one meter per second with primary batteries
- 300 m depth rating
- ANDe™ system that allows high-throughput eDNA sampling with minimal contamination
- 60 cartridges housing filters,
- 3G-ESP long-term, large scale *in situ* eDNA sample processing

BOEM
BUREAU OF OCEAN ENERGY MANAGEMENT




Ecosystem Impacts of Warm Blob on West Coast in 2015-2016


McCabe et al. *GRL* 2016

Widespread Ecosystem Impacts!

Things got Blobular




Temperature (NDBC)



Dungeness Crab Closure
\$100 M Loss
West Coast fishery



CONSISTENT NATIONAL CAPABILITY



Challenge for Global Models: Resolving Coastal and Shelf Sea Physics




Figure 2. Cumulative distribution of the fraction of global (top) and coastal (bottom) ocean, resolving L_1 , L_r , and L_{bt} for different global model resolutions.

L_1 = front/frontal jet, coastal upwelling

L_r = topographic-steered barotropic current

L_{bt} = barotropic tide





Figure 3. Cumulative distribution of number of rivers where scale L_r is resolved at a particular level (e). Based on flow data from the 925 largest ocean-flowing rivers globally (Dai et al. 2009).

Holt et al. *Geosci. Model Dev.* (2017)

High(ish)-Resolution (Nested) Coupled Models



Pacific Northwest & Salish Sea *Live Ocean Model (ROMS)*

Parker MacCready, University of Washington



LiveOcean Forcing and System Overview

3-Day forecast
appears daily on
NANOOS NVS






WRF Winds & Heating


HYCOM Ocean Fields

USGS Rivers

TPXO Tides



ROMS



<http://faculty.washington.edu/pmac/LiveOcean.html>
http://nvs.nanoos.org/Explorer?action=overlay:liveocean_temp

J-SCOPE produces 6-9 month seasonal forecasts of physical conditions, Chl-*a*, O₂, pH, MLD, plankton, and Ω (PI Samantha Seidlecki)



Home

Forecasts

Year in Review

About the Model

Climatology

Model Performance

People

Partners

Disclaimer

Contact



Forecast Origin Dates


Jan 2019 Apr 2018 Jan 2018 Apr 2017 Jan 2017 **Apr 2016** Jan 2016 Apr 2015 Jan 2015 Apr 2014 Apr 2013 Jan 2013

Overview Chlorophyll Sea Surface Temperature Sardines Oxygen Ω CA Current Indicators

Overview

The J-SCOPE forecast system for Washington and Oregon coastal waters presents preliminary results for the ocean acidification conditions during the 2016 upwelling season. The forecast for 2016 is composed of three model runs that make up an ensemble. Each model run is initialized at a different time (April 5, April 15, April 25), and has complementary forcing files from the large scale model CFS.

The forecasts simulate conditions in 2016 with a full carbon model (included DIC and TA). The TA and DIC fields are then used to calculate Ω using CO2sys.




The movie above shows the J-SCOPE forecast for 2016, from ensemble model run #2 initialized on April 15. The 8-day upwelling index is calculated using the method described in Austin and Barth (2002) and can also be found under the California Current

LiveOcean Bottom Oxygen in the NANOOS NVS

The “Comparator” allows real-time comparison with observations

nvs.nanoos.org/Explorer

Seidlecki et al. 2015





APPLICATION to STAKEHOLDERS:

Forecast of corrosive water due to
Ocean Acidification that harms
shellfish aquaculture (WOAC)

The annual value of the shellfish industry in
Washington State is \$108 million.

One in eight oysters consumed in the US comes
from Willapa Bay.






Model forecast of **surface aragonite saturation**
state – corrosive waters from the Columbia River
plume are a key feature.

Often larval oysters in Willapa Bay do not
survive – due to Ocean Acidification.

APPLICATION to STAKEHOLDERS:

Short-term forecasts of phytoplankton blooms and surface water advection from known *Pseudo-nitzschia* HAB hotspots.

-customized for razor clam recreational harvests




Pacific Northwest Harmful Algal Blooms Bulletin

Apr 13, 2018 HAB risk =

HAB risk key:

- = low
- = medium
- = high





PNW HAB Bulletin

Short-term risk



- Beach *Pseudo-nitzschia* abundance & pDA
- Small boat at hotspots (PN, DA)
- ESP moored off La Push, WA
- Ocean currents, Columbia River discharge, satellite chlorophyll
- Marine weather
- Cumulative upwelling index
- LiveOcean forecast model




Steph Moore w/ ESP Partnership with NANOOS



Long-term forecast

- Pacific Ocean Indices

(warmer T years associated with increased DA risk)



Slide courtesy Vera Trainer, NWFSC


Near Real-Time and Research Models for the California Current System

regional downscaling of the physics

Operational 3-km ROMS
with 3DVAR data assimilation
Yi Chao, UCLA
-used for HAB forecasts

Operational 9-km ROMS
with 4DVAR data assimilation
Chris Edwards & Andy Moore, UCSC
-used for EcoCast

REGIONAL OCEAN MODEL SYSTEM (ROMS)



Research-mode ROMS
nests down to 500m


Jim McWilliams, UCLA

-used to test next-gen ecological forecast models
(D. Bianchi, C. Deutsch)




uses nested grids to move across scales
(e.g. Shchepetkin and McWilliams, 2003)
-atmospheric boundary conditions from WRF






California Harmful Algae Mapping (C-HARM) System

Anderson et al., *Harmful Algae* (2009), *GRL* (2011), *Harmful Algae* (2016)



Extensive collaboration with all partners on creation of a monthly **CA HAB Bulletin** distributed via listserv and SCCOOS & HABMAP




Stakeholder engagement is done via web surveys and continual outreach to super end-users

California HAB Bulletin


sccoos.org/california-hab-bulletin/

What is the CA HAB Bulletin?


The purpose of this *experimental* product is to give the public and resource managers a quick outlook of recent toxic (marine) algal blooms in coastal California from models and aggregate data sets. **Monthly reports synthesize model output, near real-time observations, animal strandings, and public health alerts** to provide a more complete picture of the regional variability in harmful algal blooms for stakeholders.




California HAB Bulletin



HABMAP monitoring at 7 stations for HAB species and



C-HARM



The Marine Mammal Center (TMMC) – Sea lion strandings due to domoic acid toxicosis




California Department of Public Health (CDPH)

Lessons Learned & Next Steps

The Future of Coupled Modeling at National Weather Service

- Stakeholders mostly need the higher-resolution, granular predictions, preferably with seasonal outlooks; **must have an iterative feedback loop**
- Will we ever be able to move seamlessly from global to nested coupled models **at nearshore scales relevant to stakeholders?**
 - Requires innovations in physical coupling schemes
 - Requires more progress predicting HABs, OA, Hypoxia impacts in the food web and at nearshore-estuarine scales
- Can one model actually help (or run) them all?
 - Testing the **West Coast Ocean Forecast System (WCOFS)** as a universal backbone for various ecological forecasting efforts on the U.S. West Coast (IOOS COMT)



Thanks to for this opportunity

Collaborators:



NASA Applied Sciences,
Ocean Biology and
Biogeochemistry,
Energy and Water Cycle



Sea Grant
California



NOAA NCCOS CRP
MERHAB & ECOHAB
NOAA OAP

