

Deriving The Speed Of Light In a Vacuum

Antonius Torode

February 15, 2017

We begin with Maxwell's equations which follow in the MKS system of units [1] as

$$\nabla \cdot \vec{E} = \frac{\rho}{\epsilon_0} \quad (1) \quad \nabla \cdot \vec{B} = 0 \quad (3)$$

$$\nabla \times \vec{E} = -\frac{\partial \vec{B}}{\partial t} \quad (2) \quad \nabla \times \vec{B} = \mu_0 \vec{J} + \epsilon_0 \mu_0 \frac{\partial \vec{E}}{\partial t} \quad (4)$$

In the above equations, t is time, ϵ_0 is the permittivity of free space, μ_0 is the permeability of free space, \vec{E} is the electric field, \vec{B} is the magnetic field, and \vec{J} is the current density. The definition of current is given by $\vec{I} \equiv \frac{dQ}{dt} \hat{I}$ which allows us to represent the current density as

$$\vec{J} = \frac{d\vec{I}}{da_{\perp}} = \frac{d}{da_{\perp}} \left(\frac{dQ}{dt} \right) \hat{I}. \quad (5)$$

Now, we can take the curl of equation (2) which gives

$$\nabla \times (\nabla \times \vec{E}) = \nabla \times \left(-\frac{\partial \vec{B}}{\partial t} \right). \quad (6)$$

The left-hand side of this can be manipulated using the vector identity $\nabla \times (\nabla \times \vec{A}) = \nabla(\nabla \cdot \vec{A}) - \nabla^2 \vec{A}$. The right-hand side can also be rearranged since ∇ is not an operation with respect to time which gives

$$\nabla \times (\nabla \times \vec{E}) = \nabla(\nabla \cdot \vec{E}) - \nabla^2 \vec{E} = \nabla \times \left(-\frac{\partial \vec{B}}{\partial t} \right) = -\frac{\partial}{\partial t} (\nabla \times \vec{B}). \quad (7)$$

We can use equation (1) to write this as

$$\nabla \left(\frac{\rho}{\epsilon} \right) - \nabla^2 \vec{E} = -\frac{\partial}{\partial t} (\nabla \times \vec{B}). \quad (8)$$

If we assume we are in a perfect vacuum, then there would contain no matter and thus no charge. Therefore the charge density would be $\rho = 0$. This implies the total charge is zero and thus $\frac{dQ}{dt} = 0$. Hence, we can clearly see from equation (5) that within a vacuum $\vec{J} = 0$. Using these results as well as (4), we can write equation (8) as

$$-\nabla^2 \vec{E} = -\frac{\partial}{\partial t} \left(\epsilon_0 \mu_0 \frac{\partial \vec{E}}{\partial t} \right) \implies \nabla^2 \vec{E} = \epsilon_0 \mu_0 \frac{\partial^2 \vec{E}}{\partial t^2}. \quad (9)$$

This result in equation (9) can be recognized as the wave equation which is generally in the form

$$\nabla^2 \vec{\psi} = \frac{1}{v^2} \frac{\partial^2 \vec{\psi}}{\partial t^2}, \quad (10)$$

where v is the velocity of a wave. Finally, if we think of the electric field \vec{E} as a wave moving through a vacuum, then we can determine its velocity by comparing equations (9) and (10). This is the speed of an electromagnetic wave (the speed of light) which gives

$$v = c = \frac{1}{\sqrt{\epsilon_0 \mu_0}}. \quad (11)$$

We can approximate this based on the values of $\epsilon_0 = 8.85418782 \times 10^{-12} \text{ s}^4 \text{ A}^2 / (\text{m}^3 \text{ kg})$ and $\mu_0 = 4\pi \times 10^{-7} \text{ Wm}$ which gives

$$c = \frac{1}{\sqrt{\epsilon_0 \mu_0}} \approx \frac{1}{\sqrt{[8.85418782 \times 10^{-12} \text{ s}^4 \text{ A}^2 / (\text{m}^3 \text{ kg})][4\pi \times 10^{-7} \text{ Wm}]}} \approx \boxed{299,792,458 \frac{\text{m}}{\text{s}}}. \quad (12)$$

References

- [1] “Wolfram MathWorld: The Web’s Most Extensive Mathematics Resource.” Wolfram MathWorld. N.p., n.d. Web.