Chapter 1

Conclusion and Further Research

1.1 Quasi-linear Differential Problems

Throughout this theis, it has been possible to either use techniques from Applied Mathematics to construct solution strategies on a case by case basis, or use Statistical methods to find a semi-parametric fit. As differential equations become more complex, both approaches begin to rapidly become non-viable. In this section, quasi-linear differential equation models will be touched upon.

The difference between a quasi-linear and a linear differential equation is that the coefficients in a quasi-linear equation are allowed to depend on the unknown function. Instead of an ODE such as $y' = \beta(t)y$, one would have an ODE such as $y' = \beta(y,t)y$. Though quasi-linear problems tend to be reminiscent of linear ones, they are nonetheless substanially more complicated, and require more technical knowledge, and even ingenuity to tackle.

For a quasi-linear variation of a linear ODE, consider the Van Der Pol Equation:

$$y''(t) + \beta(1 - y(t))^2 y'(t) + y(t)$$

This ODE has no obvious solution.

1.1.1 Inviscid Burger's Equation

Even if a solution exists, an estimation strategy might be difficult to derive. Consider the inviscid Burger's Equation:

$$\frac{\partial u(x,t)}{\partial t} + \beta u(x,t) \frac{\partial u(x,t)}{\partial x} = 0$$

This equation is identical to the Transport Equation except that the rate term is equal to $\beta u(x,t)$. The solution is given by:

$$u(x,t) = f(s)$$

Here $f(\cdot)$ is some arbitary function as before, and s is implicitly defined as the solution of the equation $x = \beta f(s)t + s$. Since $s = x - \beta ut$, this can be written as:

$$u(x,t) = f(x - \beta ut)$$

Fitting this model is substantially trickier than the Transport Equation. There is no clean separation between the problem of estimating $f(\cdot)$ and β since u(x,t) appears on the righthand side and scales β .

A further complication is that u(x,t) might only define a *relation*, instead of a function. There might be multiple values of u associated with a given (x,t) that satisfy the solution equation. Physically speaking, multiple values correspond to shock waves.

1.1.2 Discussion

We see that the level of knowledge required to devise fitting strategies can increase substantially even with seemingly modest increases in the complexity of the associated differential equation.

Consider the following quasi-linear model of genetic drift in a population proposed¹ by R.A. Fisher:[8]

$$\frac{\partial u(x,t)}{\partial t} + \beta_1 \frac{\partial u(x,t)}{\partial x} = \beta_2 u(x,t) (1 - u(x,t)) \tag{1.1}$$

This problem is similar to the previous PDEs we discussed, it even admits travelling wave solutions of the form f(x+ct) as Fisher himself noted. Nonetheless, it is a much more difficult problem despite the apparently modest increase in complexity. One would likely have to consult a textbook that covers non-linear PDEs that can generate waves in fair degree of detail to be able to devise a fitting strategy. This is quite a specialised subject!

The overall result is that as the complexity of the differential equation increases, more and more time will be needed to model it correctly.

¹In the Annals of Eugenics...

Bibliography

- [1] John P Boyd. Chebyshev and Fourier spectral methods. Courier Corporation, 2001.
- [2] Stephen Boyd and Lieven Vandenberghe. *Convex optimization*. Cambridge university press, 2004.
- [3] Richard P Brent. Algorithms for minimization without derivatives. Courier Corporation, 2013.
- [4] Jiguo Cao and James O Ramsay. Parameter cascades and profiling in functional data analysis. *Computational Statistics*, 22(3):335–351, 2007.
- [5] Kwun Chuen Gary Chan. Acceleration of expectation-maximization algorithm for length-biased right-censored data. *Lifetime data analysis*, 23(1):102–112, 2017.
- [6] Edwin KP Chong and Stanislaw H Zak. An introduction to optimization, volume 76. John Wiley & Sons, 2013.
- [7] Arthur P Dempster, Nan M Laird, and Donald B Rubin. Maximum likelihood from incomplete data via the em algorithm. *Journal of the Royal Statistical Society:* Series B (Methodological), 39(1):1–22, 1977.
- [8] Ronald Aylmer Fisher. The wave of advance of advantageous genes. *Annals of eugenics*, 7(4):355–369, 1937.
- [9] Bengt Fornberg. Generation of finite difference formulas on arbitrarily spaced grids. *Mathematics of computation*, 51(184):699–706, 1988.
- [10] PR Graves-Morris, DE Roberts, and A Salam. The epsilon algorithm and related topics. *Journal of Computational and Applied Mathematics*, 122(1-2):51–80, 2000.
- [11] David R Hunter and Kenneth Lange. A tutorial on MM algorithms. *The American Statistician*, 58(1):30–37, 2004.
- [12] Eugene Isaacson and Herbert Bishop Keller. Analysis of numerical methods. Courier Corporation, 2012.
- [13] Carl T Kelley. *Iterative methods for linear and nonlinear equations*, volume 16. Siam, 1995.

- [14] Carl T Kelley. Implicit filtering, volume 23. SIAM, 2011.
- [15] C.T. Kelley. A brief introduction to implicit filtering. https://projects.ncsu.edu/crsc/reports/ftp/pdf/crsc-tr02-28.pdf, 2002. [Online; accessed 12-October-2019].
- [16] Jack Kiefer. Sequential minimax search for a maximum. Proceedings of the American mathematical society, 4(3):502–506, 1953.
- [17] Kenneth Lange. Optimization. Springer, 2004.
- [18] Kenneth Lange. Numerical analysis for statisticians. Springer Science & Business Media, 2010.
- [19] Kenneth Lange. The MM algorithm. https://www.stat.berkeley.edu/~aldous/Colloq/lange-talk.pdf, April 2007. [Online, accessed 18-September-2019].
- [20] Randall J LeVeque. Finite difference methods for ordinary and partial differential equations: steady-state and time-dependent problems, volume 98. Siam, 2007.
- [21] Steve McConnell. Code complete. Pearson Education, 2004.
- [22] Geoffrey McLachlan and Thriyambakam Krishnan. The EM algorithm and extensions, volume 382. John Wiley & Sons, 2007.
- [23] J Nocedal and SJ Wright. Numerical Optimisation. Springer verlag, 1999.
- [24] Naoki Osada. Acceleration methods for slowly convergent sequences and their applications. PhD thesis, PhD thesis, Nagoya University, 1993.
- [25] Yudi Pawitan. In all likelihood: statistical modelling and inference using likelihood. Oxford University Press, 2001.
- [26] R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria, 2013. ISBN 3-900051-07-0.
- [27] James Ramsay. Functional data analysis. Encyclopedia of Statistics in Behavioral Science, 2005.
- [28] Jim O Ramsay, Giles Hooker, David Campbell, and Jiguo Cao. Parameter estimation for differential equations: a generalized smoothing approach. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 69(5):741–796, 2007.
- [29] Walter Rudin et al. *Principles of mathematical analysis*, volume 3. McGraw-hill New York, 1964.

- [30] Larry Schumaker. Spline functions: basic theory. Cambridge University Press, 2007.
- [31] Christopher G Small. A survey of multidimensional medians. *International Statistical Review/Revue Internationale de Statistique*, pages 263–277, 1990.
- [32] Karline Soetaert, Thomas Petzoldt, and R. Woodrow Setzer. Solving differential equations in R: Package deSolve. *Journal of Statistical Software*, 33(9):1–25, 2010.
- [33] Gerald Teschl. Ordinary differential equations and dynamical systems, volume 140. American Mathematical Soc., 2012.
- [34] Keller Vandebogart. Method of quadratic interpolation. http://people.math.sc.edu/kellerlv/Quadratic_Interpolation.pdf, September 2017. [Online; accessed 13-September-2019].
- [35] Jet Wimp. Sequence transformations and their applications. Elsevier, 1981.
- [36] Stephen Wright. Optimization for data analysis. In Michael W. Mahoney, John C. Duchi, and John C. Duchi, editors, *The Mathematics of Data*, chapter 2, pages 49–98. American Mathematical Society and IAS/Park City Mathematics Institute and Society for Industrial and Applied Mathematics, 2018.
- [37] Tong Tong Wu, Kenneth Lange, et al. The MM alternative to EM. Statistical Science, 25(4):492–505, 2010.