

Attenuating Neural Threat Expression with Imagination

Highlights

- Imagined extinction reduces neural and physiological conditioned threat responses
- Ventromedial prefrontal cortex is central to both real and imagined extinction
- Nucleus accumbens uniquely predicts the success of imagined extinction

Authors

Marianne Cumella Reddan,
Tor Dessart Wager, Daniela Schiller

Correspondence

tor.wager@colorado.edu (T.D.W.),
daniela.schiller@mssm.edu (D.S.)

In Brief

Reddan et al. demonstrate that threat responses can be extinguished through imagined simulations of the conditioned stimuli. Like real extinction, imagined extinction engages the ventromedial prefrontal cortex, amygdala, and related perceptual cortices. Nucleus accumbens activity predicts an individual's ability to successfully extinguish via imagination.



Attenuating Neural Threat Expression with Imagination

Marianne Cumella Reddan,¹ Tor Dessart Wager,^{1,4,*} and Daniela Schiller^{2,3,4,5,*}

¹Department of Psychology and Neuroscience, University of Colorado Boulder, Boulder, CO 80303, USA

²Departments of Psychiatry and Neuroscience, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

³Friedman Brain Institute, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA

⁴These authors contributed equally

⁵Lead Contact

*Correspondence: tor.wager@colorado.edu (T.D.W.), daniela.schiller@mssm.edu (D.S.)

<https://doi.org/10.1016/j.neuron.2018.10.047>

SUMMARY

Imagination is an internal simulation of real-life events and a common treatment tool for anxiety disorders; however, the neural processes by which imagination exerts behavioral control are unclear. This investigation tests whether and how imagined exposures to a threatening stimulus, conditioned in the real world, influence neural and physiological manifestations of threat. We found that imagined and real extinction are equally effective in the reduction of threat-related neural patterns and physiological responses elicited upon re-exposure to real-world threatening cues. Network connectivity during the extinction phase showed that imagined, like real, extinction engaged the ventromedial prefrontal cortex (vmPFC) as a central hub. vmPFC, primary auditory cortex, and amygdala activation during imagined and real extinction were predictive of individual differences in extinction success. The nucleus accumbens, however, predicted extinction success in the imagined extinction group alone. We conclude that deliberate imagination can attenuate reactions to threat through perceptual and associative learning mechanisms.

INTRODUCTION

Threat likelihoods are rapidly acquired across diverse and dynamic environments. The diminution of learned threat, however, is a slower and more mutable process (LeDoux, 2000). Fear, therefore, often persists into times of safety (Vervliet et al., 2013). Maladaptive defensive reactions negatively impact quality of life, and underlie emotional disorders, such as post-traumatic stress disorder (PTSD), phobias, and anxiety. Extinction learning, the repeated presentation of a conditioned stimulus without its aversive consequence, is one of the most effective ways of reducing learned threat in the laboratory (Graham and Milad, 2011). Extinction is theoretically a core process of exposure therapy, the most prescribed treatment for fear-related dis-

orders (Taylor et al., 2003). Exposure therapy is impractical in some cases because the recreation of cues associated with a traumatic event may be difficult or unethical to reconstruct (e.g., a war zone), or because the intensity of re-exposure is overwhelming for the patient (Storch and McKay, 2013). Imagination creates a unique opportunity to simulate re-exposure in a controlled way so that a patient may immerse themselves at their own pace. Simulation, in theory, provides a method by which persistent and maladaptive threat memories can be targeted for treatment, independent of external stimuli (Arbuthnott et al., 2001). This investigation demonstrates the utility of imagination as a threat-extinction tool and proposes a neural mechanism for imagined extinction that includes a network of brain regions known to support the real extinction learning. These findings expand our understanding of how the human brain modifies threat representations and, in turn, our ability to treat threat-related disorders using mental action.

Imagination, also called mental imagery or mental practice, is a conscious simulation of a stimulus or an event that can impact perception, cognition, and emotion (Holmes and Mathews, 2005; Kosslyn et al., 2006). There is growing evidence that imagination induces neural plasticity (Pascual-Leone et al., 2005), activates perceptual and motor cortices (Kosslyn et al., 2001; Zatorre and Halpern, 2005), enhances real-world performance (Shepard and Metzler, 1971), and supports the prediction of future events (Schacter et al., 2007; Mendelsohn et al., 2014), but its role in emotion regulation remains unclear (Holmes and Mathews, 2010). When effective, imagination holds a unique advantage as an emotion regulation tool because it is changeable, internally directed, and dissociated from immediate sensory input (Taylor et al., 1998).

The clinical advantages of imagination are harnessed by imaginal exposure therapy, where patients imagine triggering events until threat sensations subside (van Minnen and Foa, 2006). Despite imagination's long-standing recognition in the clinic, it has received little attention from the neuroscientific learning community (Mendelsohn et al., 2014; Agren et al., 2017; Soeter and Kindt, 2012; Dadds et al., 1997). This is due, in part, to the clinic's reliance on behavior and self-reports in non-neurotypical populations. When considering behavioral and self-report evidence alone, it is difficult to separate the unique effects of imagination from expectancy or other complex cognitive phenomena that shape emotion and learning. It is also related to the field's



focus on translational research: animal models of imagination are difficult to construct, though they have been attempted (Johnson et al., 2009; Lucantonio et al., 2015; Redish, 2016; Takahashi et al., 2013; Holland, 1990). The current study bridges the gap between the clinic and the laboratory and, in doing so, demonstrates that a scientific interrogation of imagination's role in fear regulation may provide novel insight into how threat memories are represented, accessed, and modified by mental action.

Extinction is a distributed neural process involving the ventromedial prefrontal cortex (vmPFC), hippocampus, and amygdala (Quirk and Mueller, 2008). More recent evidence extends this network to include the nucleus accumbens (NAc) and related perceptual circuitry (Apergis-Schoute et al., 2014; Levita et al., 2012). Learning and plasticity in these regions underlie the creation and sustainment of extinction memories. Imagination may influence emotion processing either by directly modifying affective substrates downstream of sensory cortices, such as the amygdala, or by activating and altering the representation of existing emotional memories via interactions among the PFC, hippocampus, and NAc (Holmes and Mathews, 2010). Behavioral evidence suggests an interaction; vividness of imagination is positively correlated with both memory strength and physiological arousal (Acosta and Vila, 1990; Miller et al., 2008). Furthermore, the vmPFC and hippocampus are part of an imagination network that supports mental time travel (Buckner and Carroll, 2007; Schacter et al., 2007), and the orbitofrontal cortex drives learning related to imagined outcomes (Takahashi et al., 2013). In cases of deliberation, the NAc is recruited to simulate the outcomes of all possible options (Stott and Redish, 2014), and its size is related to individual differences in imaginal ability (Jung et al., 2016).

Drawing across this wide range of evidence, we propose the following mechanism: imagination of the conditioned threat stimulus will activate stimulus-specific perceptual representations that will, in turn, engage the neurocircuitry that underlies threat acquisition: the vmPFC, hippocampus, amygdala, and NAc. In the absence of any danger, repeated imaginings will have the same effect as actual exposures—neural and physiological responses to the conditioned threat stimulus will diminish.

To test the effectiveness of imagined extinction and its neural mechanism, our participants underwent auditory threat conditioning and then were randomized into three groups (Figure 1). The first group performed imagined extinction, that is, they were directed to “play” the conditioned tones “in their head” to the best of their ability. The second underwent real extinction, which consisted of actual exposures to the conditioned auditory stimuli. The third underwent no extinction and instead was directed to imagine two neutral sounds from nature (“birds singing” and “rain falling”), as a control for the general effects of imagination on arousal. The threat memory was then reinstated in all participants through four unsignaled shocks, after which all participants were re-exposed to the conditioned auditory stimuli. Functional magnetic resonance imaging (fMRI) data were collected for each phase, and skin conductance responses (SCRs) were recorded continuously. We hypothesized that both real and imagined extinction would attenuate the neural and bodily expression of threat memory following reinstatement. We expected that

defensive responding would persevere through re-exposure only in the third group, which had no extinction at all.

RESULTS

Imagined Extinction Reduces Whole-Brain Threat Expression to Conditioned Stimuli

We developed a novel whole-brain fMRI predictive pattern, sensitive and specific to threat, in order to test how effectively imagined extinction reduces threat responses in the brain (Figure 2A). This threat-predictive pattern is a multivariate pattern classifier that can be applied to novel activation maps to assess threat expression in subsequent neural activity. It was developed using a linear support vector machine (SVM) trained on subject-wise ($n = 68$) univariate brain activation to the unreinforced CS^+ (threatening) and CS^- (non-threatening) stimuli during threat acquisition. It yielded a classification accuracy of 77% (± 3.6 STE, $p < 0.0001$) in a modified leave-three-subjects-out cross-validation (CV) procedure (Figure 2B), where one subject from each of the three groups was left out of each fold to reduce potential bias from group assignment. The threat-predictive pattern's sensitivity and specificity to threat were 71% (CI: 61%–79%) and 84% (CI: 76%–91%), respectively. It had a positive predictive value (PPV) of 81% (CI: 72%–90%), and an effect size of 0.82, estimated by nonparametric area under the curve. When thresholded (bootstrapped 5,000 samples) and corrected for multiple comparisons (false discovery rate [FDR] corrected, $p < 0.05$), the neural threat-predictive pattern revealed a distributed network of threat representation including the PFC, amygdala, periaqueductal gray (PAG), insula, globus pallidus, and dorsal anterior cingulate cortex (dACC; see Table S1 for complete network). These regions are consistent with those reported in the existing threat conditioning literature, and with the univariate map contrasting the CS^+ and CS^- stimuli in this study (see Figure S1 and Table S2; Critchley and Garfinkel, 2014, Fullana et al., 2016).

Next, we used the neural threat-predictive pattern to evaluate the effectiveness of imagined extinction relative to the control conditions. The pattern was applied to participant-specific brain maps during the re-extinction phase. Reinstatement is often followed by the immediate recovery of defensive reactions, even if extinction learning was successful (Vervliet et al., 2013). However, reinstatement effects are transient in neurotypical humans, lasting only a few trials. (Haaker et al., 2014). We therefore examined whether threat representation was preserved during the late phase of re-extinction (last 5 trials). If extinction learning were less effective or absent, defensive reactions should be maintained during this later phase (Craske et al., 2008). The pattern was applied by taking the dot product between the unthresholded threat-predictive pattern (all voxels) and participant-specific brain maps. This amounts to taking a weighted average over the test data, where the pattern defines the weights, yielding a single value for a given data image. The resulting metric is known as “pattern expression” (or “pattern response”) and is used to assess the magnitude of the activation in a pre-defined pattern (i.e., here defined to differentiate CS^+ versus CS^- and thus serve as a measure of neural threat response). The pattern expression is influenced by both the




Figure 1. Schematic of the Protocol and Methods

This experiment was divided into four phases. (1) Acquisition: all participants ($n = 68$) underwent an auditory threat-conditioning paradigm. The CS⁺ tone was paired with an electric shock 33% of the time. The CS⁻ tone was never paired with shock. A whole-brain multivariate threat-predictive pattern was trained on these trials using support vector machines. (2) Extinction: participants were randomly divided into one of three “extinction” sessions: imagined extinction ($n = 20$), real extinction ($n = 22$), and no extinction (or none, $n = 24$). Trial structure was conserved across groups, and imagined extinction trial order was yoked to real extinction stimulus presentation. (3) Reinstatement: all participants underwent reinstatement, where they received four unsignaled shocks presented 4–12 s apart. (4) Re-extinction: participants were then re-exposed to the conditioned stimuli to assess the effectiveness of the extinction sessions. The threat-predictive pattern was applied to participant-specific brain maps during the threat recovery test period, which was comprised of the last 5 trials of re-extinction. The threat-predictive pattern was applied by taking the dot product between the unthresholded pattern and participant-specific brain maps. The dot product reflects the magnitude of similarity between two vectors.

spatial similarity between the threat-predictive pattern and test data, as well as the magnitude of activation (Woo et al., 2017).

A positive pattern expression value indicates that the tested brain activity is classified as “threatening,” while a negative value indicates that the brain activity is classified as “non-threatening.” Group differences in the magnitude of the neural threat-predictive pattern expression were analyzed, and the effectiveness of the three experimental conditions was compared. Data from two participants were lost during this phase due to a technical error during data collection; therefore, only 66 participants were included in this and the following fMRI analyses.

In this case, as is standard in the field, we trained and tested the threat-predictive pattern in a cross-validated manner (leave-3-participants-out), so that the pattern responses estimated for each individual were always obtained from patterns trained on *other participants’* data, avoiding circularity when testing CS⁺ versus CS⁻ effects on pattern expression (see STAR Methods for more detail; Varoquaux et al., 2017). Group (intervention) effects were always estimated on independent contrasts from the CS⁺ versus CS⁻ contrast used to train the pattern, and tests of intervention effects are also non-circular, with unbiased estimates of effect sizes.

There was a main effect of group as indicated by a one-way ANOVA ($F(2,63) = 3.54$, $p = 0.03$, $\eta^2 = 0.10$; Figure 3A; see Figure S2A for data distribution). As hypothesized, only the no-extinction group exhibited a distributed neural threat response during the late re-exposure (recovery test) period. The threat-predictive pattern expression was positive on average in the no-extinction group ($\bar{x} = 0.55$, $n = 24$), and significantly greater than both imagined extinction ($\bar{x} = -0.20$, $CI = [-1.46, -0.04]$, $n = 20$, $t(37.9) = 2.14$, $p = 0.04$, Hedges $g = 0.64$) and real extinction ($\bar{x} = -0.11$, $CI = [-1.22, -0.11]$, $n = 22$, $t(42.2) = 2.40$, $p = 0.02$, Hedges $g = -0.69$) groups as indicated by two-sample t tests with unequal variance. There was no evidence for a difference between the real and imagined extinction group ($t(31.8) = 0.27$, $p = 0.79$; Bayes factor in favor of the null = 7.62), indicating no detectable differences in the reduction of neural threat expression between these groups and that the null hypothesis is favored 7.62 times more than the alternative.

Because the amygdala is a ROI, we performed this analysis again only in this region, bilaterally. The neural threat-predictive pattern was masked and then applied to the same test phase, late re-extinction. Similar to the results of the whole-brain threat expression, there was a non-significant trend where only the




Figure 2. Multivariate Neural Threat-Predictive Pattern Trained on Acquisition

For 3D neuroimaging files, see <https://doi.org/10.1016/j.neuron.2018.10.047>.

(A) Neural threat-predictive pattern. A distributed pattern of threat was developed using a linear support vector machine trained on subject-wise ($n = 68$) univariate brain activation to the unreinforced CS^+ (threatening) and CS^- (non-threatening) stimuli during threat acquisition. When thresholded (bootstrapped 5,000 samples) and corrected for multiple comparisons (FDR $p < 0.05$), the threat-predictive pattern revealed a distributed network of threat representation including the PFC, periaqueductal gray (PAG), amygdala, insula, subregions of the basal ganglia, and dorsal anterior cingulate (dACC).

(B) ROC plot. The neural threat-predictive pattern yielded a classification accuracy of 77% (± 3.6 STE, $p < 0.0001$, area under the curve [AUC]: 0.82) in a modified leave-three-subjects-out cross-validation (CV) procedure. One subject from each of the three groups was left out of each fold to reduce potential bias from group assignment.

no-extinction group exhibited positive threat pattern expression (see Figure S2D for details).

Together, these findings show that only the no-extinction group expressed the threat pattern following the (no extinction) general imagery session. The results suggest that imagined and real extinction were both effective in diminishing neural pattern reinstatement of threat.

For consistency with prior research, a univariate analysis of the threat recovery period was also performed. Similar, but weaker, results were found in the amygdala and hippocampus, whereas both regions activated more to the CS^+ relative to the CS^- in the no-extinction group than in the other two groups. Indeed, the univariate whole-brain maps contrasting $CS^+ > CS^-$ revealed threat-related responses in the amygdala, insula, and hippocampus of only the no-extinction group (see Figure S3 and Table S3 for details).

Imagined Extinction Reduces Bodily Threat Expression to Conditioned Stimuli

The physiological SCR results in the participants who showed reliable physiological threat acquisition ($n = 42$) complemented the neural findings: relative to no extinction ($\bar{x} = 0.03$, $n = 13$), imagined extinction ($\bar{x} = -0.0004$, $n = 12$, $t(20.1) = 2.31$, $p = 0.03$, $CI = [-0.07, -0.003]$, Hedges $g = -0.88$), and real extinction ($\bar{x} = -0.005$, $n = 17$, $t(26.2) = 2.31$, $p = 0.03$, $CI = [-0.07, -0.004]$, Hedges $g = -0.82$) showed a reduction in threat-related physiological arousal ($F(2,39) = 3.61$, $p = 0.036$, $\eta^2 = 0.16$, Figure 3B; see Figure S2B for data distribution) during the late re-extinction phase. There was no difference between the real and imagined

extinction groups during this period ($t(26.5) = 0.32$, $p = 0.75$, Bayes factor in favor of the null = 2.75). Furthermore, participants' SCR positively correlated with expression of the neural threat-predictive pattern (Spearman's $\rho = 0.34$, $p = 0.03$, Figure 3C), signifying a link between the neural and physiological expressions of threat. Participants ($n = 24$) who did not demonstrate a discriminatory SCR during acquisition, defined as greater SCR to the CS^+ relative to the CS^- during either the first or last half of threat-acquisition on average, were excluded from this analysis because we were unable to determine whether their SCRs were representative of threat-related arousal (see Figure S4 for SCR analysis without this criteria). Importantly, this failure to produce discriminatory SCRs was not related to the order by which stimuli were presented or the counterbalancing of two different Hz tones (see Table S4 for an analysis of order effects in SCR).

To demonstrate that the inclusion of the neural data from SCR non-responders was valid, we retested the recovery of the neural threat pattern expression in the participants whose SCR was not excluded ($n = 42$). The findings yield the same conclusions as this analysis with the complete sample ($F(2,39) = 4.45$, $p = 0.02$, $\eta^2 = 0.10$). The no-extinction group ($\bar{x} = 0.47$, $n = 13$) was significantly larger than both imagined ($\bar{x} = -0.31$, $n = 12$, $t(21.4) = -2.62$, $p = 0.02$, $CI = [-1.40, -0.16]$, Hedges $g = -1.02$) and real extinction ($\bar{x} = -0.21$, $n = 17$, $t(26.94) = -2.68$, $p = 0.01$, $CI = [-1.20, -0.16]$, Hedges $g = -0.94$) groups as indicated by two-sample t tests with unequal variance. Real extinction did not differ from imagined ($t(22.1) = -0.33$, $p = 0.74$, $CI = [-0.70, 0.51]$, Hedges $g = -0.12$).




Figure 3. Imagined Extinction Reduces Neural and Physiological Threat Expression

(A) Neural threat-predictive pattern expression during the recovery test. The threat-predictive pattern was applied to brain activity during the late re-extinction recovery test. Only the no-extinction group (none, $n = 24$) demonstrated threat recovery during late re-extinction, indicated by a comparison of average threat-predictive pattern expression values between groups. This may indicate that both imagined ($n = 20$) and real ($n = 22$) extinction successfully generated an extinction memory. Data are represented as mean $\pm$ SEM.

(B) SCRs during the recovery test. Physiological findings, indicated by SCR, complemented the neural findings. Relative to no extinction ($n = 13$), imagined ($n = 12$) and real extinction ($n = 17$) showed a reduction in threat-related physiological arousal during the recovery test. Data are represented as mean $\pm$ SEM. Participants ($n = 24$) who did not demonstrate a discriminatory SCR during acquisition, defined as greater SCR to the CS+ relative to the CS- during either the first or last half of threat-acquisition on average, were excluded from SCR analysis because we were unable to determine whether their SCRs were representative of threat-related arousal.

(C) Correlation between SCRs and threat-predictive pattern expression during the recovery test. SCRs were positively correlated with expression of the threat-predictive pattern ($\rho = 0.34$, $p = 0.03$), indicating a link between the neural and physiological expressions of threat.

Neural Network Supporting Imagined Extinction

To examine whether brain regions involved in real extinction are also engaged during imagined extinction, we performed a network connectivity analysis among known extinction circuitry (Herry et al., 2010) including vmPFC, hippocampus (CA1), laterobasal amygdala (Amg-LB), central nucleus of the amygdala (Amg-CM), NAc, PAG, and the primary auditory cortex. Anatomical masks of these regions were taken from the SPM anatomy toolbox, except for the vmPFC, NAc, and PAG, which were derived from Neurosynth (Yarkoni et al., 2011). Average signal from the specified ROIs was extracted across the entire extinction blood-oxygen-level-dependent (BOLD) time course without contrast. A contrast was not used because it would not be meaningful to contrast the imagined stimuli in the no-extinction group (imagined sounds of birds singing and rain falling), and contrasting imagined stimuli may mask important activity related to imaginal processes.

This analysis was performed on each group. While each group yielded discriminable networks, which can be accurately classified as either imagined extinction (Figure 4A), real extinction (Figure 4B), or no extinction (Figure 4C) using SVM (Figure 4E), imagined and real extinction have a strikingly similar construction where the vmPFC is a central hub, indicated by a comparison of the vmPFC's betweenness centrality, calculated using the number of shortest paths that pass through the node, across groups (Figure 4D), connected to the primary auditory cortex, CA1, and NAc. The nodes with the highest betweenness centrality in the imagined extinction group, which were also significantly higher than either real or no extinction, were the right NAc,

vmPFC, and right primary auditory cortex, indicating that imagined extinction may engage perceptual and learning circuitry to update the value of the conditioned stimuli (see Table S5 for the complete statistical report on the betweenness centrality comparisons).

Imagined and Real Extinction Recruits vmPFC

Given the centrality of vmPFC as a hub of network connectivity during both imagined and real extinction, we further examined the univariate time course activation of this region. We applied the vmPFC mask to the univariate beta weights, and, using repeated-measures ANOVA, compared differential activation in the vmPFC (CS+ minus CS-) across groups and time (in bins). All acquisition trials were averaged into one bin. The 15 extinction trials were divided into four bins, two early and two late, by averaging four trials in the first three bins and three trials in the last bin. This was decided post hoc in order to better analyze the finer time course without sacrificing power. Early re-extinction was comprised of the first five trials, and late re-extinction was the last five trials. There was a significant main effect of time ($F(6,378) = 3.72$, $p = 0.001$) and a significant interaction between group and time ($F(12,378) = 1.83$, $p = 0.04$; Figure 5).

Descriptively, vmPFC activation during imagined and real extinction was nearly synchronous: it rose slowly and peaked toward the end of extinction training. Pairwise t tests of unequal variance at peak activation (in the third bin) indicated no difference between imagined and real extinction, ($t(39.01) = -0.04$, $p = 0.96$, $CI = [-0.37, 0.35]$, Hedges $g = -0.01$), but vmPFC activation during no extinction was significantly lower than both




Figure 4. Neural Networks Supporting Extinction Learning

Functional connectivity between *a priori* regions of interest (vmPFC, hippocampus (CA1), laterobasal amygdala (Amyg-LB), central nucleus of the amygdala (Amyg-CM), NAc, PAG, and the primary auditory cortex) across the entire extinction session was investigated without a contrast. Links are plotted if the partial correlation test between two nodes was significant after FDR correction within each group. The link length between each pair of nodes is related to the absolute value of the *t*-statistic associated with the partial correlation between each pair; however, link length is a graphical representation of this value, adjusted to exist in a 2D space. Here, shorter links represent greater functional connectivity. The size of each node represents betweenness centrality, which is the number of shortest paths that pass through the node, rescaled for plotting purposes with a sigmoid function. Node colors indicate clusters determined by ward linkage. (A) Imagined extinction connectivity network. The most central nodes (i.e., nodes most connected with other nodes) of the imagined extinction network included the right NAc, vmPFC, left Amyg-CM, and right primary auditory cortex.

(B) Real extinction connectivity network. The most central nodes of real extinction included the left CA1, vmPFC, and right NAc.

(C) No-extinction connectivity network. The most central nodes during the no extinction, unrelated imagination included left NAc, left Amyg-CM, and right NAc. (D) Comparison of nodal centrality across networks. 10 out of 12 nodes yielded group significant differences in betweenness centrality in a one-way ANOVA test, FDR corrected for multiple comparisons ($p < 0.05$). Data are represented as mean $\pm$ SEM. Asterisks represent pairwise significant difference via a post hoc *t* test. The most central nodes to imagined extinction are indicated by a dashed box.

(E) Each network is unique. In order to test the separability of each matrix and thereby assess their relative similarity, we trained three different binary linear support vector machine classifiers on participant connectivity data (the partial correlations between each pair of nodes). Leave-one-out cross-validated accuracy scores revealed that each group was linearly separable from one another with high accuracy (real versus none: 91% $\pm$ 4.2% STE; imagined versus none: 89% $\pm$ 4.8% STE; imagined versus real: 88% $\pm$ 5.0% STE).

imagined ($t(35.93) = 2.18$, $p = 0.035$, $CI = [0.03, 0.69]$, Hedges $g = 0.66$) and real extinction ($t(41.26) = 2.39$, $p = 0.02$, $CI = [0.06, 0.67]$, Hedges $g = 0.70$).

Additionally, correlations of voxel-wise activity patterns at peak activation suggest that activations in the vmPFC are spatially analogous in imagined and real extinction ($R = 0.51$,




Figure 5. Imagined and Real Extinction Recruits the vmPFC

(Left) Depiction of vmPFC anatomical mask used in this analysis. Due to the role of the vmPFC during extinction, it was an *a priori* region of interest. This bilateral mask was created with Neurosynth. (Right) Average differential ($CS^+ > CS^-$) BOLD signal in the vmPFC rose over the course of extinction, peaked in the third bin, and then decreased in the imagined and real extinction groups. The no-extinction group exhibited little to no change in activation across binned trials. The spatial voxel patterns during the third extinction time point are displayed, revealing (nonsignificant)

similarity in the activity patterns between the imagined and real groups. During late re-extinction, vmPFC activity increased in the imagined and real groups, but not in the no-extinction group, which demonstrated threat responses during this phase. Data are represented as mean $\pm$ SEM.

$p = 0.25$; Figure 5), but more data are required to assess significance (see also Figure S5). These results are consistent with previous reports indicating vmPFC activity increases during extinction learning (Schiller and Delgado, 2010; Schiller et al., 2013) and further indicate the ability of imagined extinction to recruit vmPFC in a manner similar to real extinction.

Brain Regions that Predict the Success of Imagined and Real Extinction

Finally, to better understand individual differences in the activation of extinction networks related to extinction success (as defined by the expression of the neural threat-predictive pattern during the recovery test period), we ran a post hoc univariate ordinary least-squares regression, controlling for nuisance signal in the ventricles and white matter, on whole-brain maps of differential activation (CS^+ minus CS^-) during extinction at peak activation (third time bin) with subject threat-predictive pattern expression scores as the predictor within each group (Figures 6A and 6B). When the analysis was thresholded and corrected for multiple comparisons (FDR corrected, $p < 0.05$, $k = 5$), no significant correlates of extinction success were found in the no-extinction group. Therefore, the no-extinction group is not displayed in Figure 6 or discussed here.

To compare the imagined and real extinction FDR corrected thresholded maps, we plotted the conjunction of the FDR thresholded maps (Figure 6C). We report significant activity anywhere within the bilateral anatomical masks of the *a priori* ROIs used in the connectivity analysis. Here, we included the entire amygdala, instead of subnuclei, to provide a more general overview. In this descriptive analysis, individual voxels need not overlap between the two maps in order to conclude that there is activity in the same ROI. Instead, significant activity in *any* voxel within the ROI was used to construct the Venn diagram descriptive summary of the two maps (Figure 6C). Because we were most interested in brain regions that predicted extinction success, we focused on the negatively correlated results alone, as a negative correlation meant that the threat-predictive pattern expression had decreased. The max effect size (partial η^2) was estimated within each ROI to compare the importance of the effect between groups. Activity in vmPFC (imagined: partial $\eta^2 = 0.42$; real: partial $\eta^2 = 0.48$), primary auditory cortex (imagined: partial $\eta^2 = 0.62$; real: partial

$\eta^2 = 0.37$), and amygdala (imagined: partial $\eta^2 = 0.50$; real: partial $\eta^2 = 0.49$) were negatively correlated with threat recovery in both the imagined and real extinction groups (refer to Figure S6 for the distribution of the data supporting these effects). The NAc (partial $\eta^2 = 0.42$) predicted extinction success uniquely in the imagined extinction group, while CA1 (partial $\eta^2 = 0.33$) predicted extinction success uniquely in the real extinction group. It is possible that imagined extinction does not influence learning in the hippocampus the same way as real extinction does but instead changes representations of stimulus value via the NAc.

DISCUSSION

Imagination is an important cognitive and emotion regulation tool that allows simulation of the real world. This investigation demonstrates that imagined extinction can reduce learned responses to threat. Using a novel neural threat-predictive pattern, we found that imagined extinction reduces threat-related brain activity, and, using SCR, we found that imagined extinction reduces threat-related physiology upon re-exposure. Furthermore, we demonstrate that the neural mechanisms underlying imagined and real extinction share a “common core” network of brain regions, wherein the vmPFC acts as a central hub of connectivity among the hippocampus, NAc, and auditory cortex. By contrast, nonspecific imagination (no extinction) depends upon a different network of brain activity and is not effective against the recovery of threat-related neural activation and arousal.

People differ in both their imaginal ability and threat response magnitudes (Miller et al., 2008). We therefore expected individual differences in the efficacy of imagined extinction. When exploring these individual differences, we found that activation strength in the vmPFC, auditory cortex, and amygdala were related to extinction efficacy in both the imagined and real extinction groups. Activity in the NAc, however, predicted extinction efficacy in the imagined extinction group alone, while activity in the hippocampus was uniquely related to the efficacy of real extinction. Together with the connectivity network analysis, this evidence suggests that imagined extinction alters learned threat responses by updating value representations associated with the conditioned stimuli, through interactions




Figure 6. Brain Regions that Predict the Success of Imagined and Real Extinction

A linear regression was performed on whole-brain maps of differential activation ($CS^+ > CS^-$) during extinction (time point 3) with participants' threat-predictive pattern expression scores during the recovery test as the predictor. Because we were most interested in brain regions that predicted extinction success, we focused on the negatively correlated results (see Table S6 for a complete table of activation).

(A and B) When thresholded and corrected for multiple comparisons (FDR corrected, $p < 0.05$), both imagined (A) and real extinction (B) revealed activity in the vmPFC, primary auditory cortex, and amygdala were related to extinction success. No significant correlates of extinction success were found in the no-extinction group.

(C) Conjunction. Signal in *a priori* regions of interest (ROIs) that survived correction in the whole-brain analysis was assessed. The NAc predicted extinction success uniquely in the imagined extinction group, while CA1 predicted extinction success uniquely in the real extinction group. Effect sizes in the ROIs were estimated via max partial η^2 , which ranges from 0 to 1.

among the auditory cortex, vmPFC, amygdala, and NAc. How does imagined extinction change the value of a threat representation learned through experience with the real world? There are several possible ways by which imagination may induce neural processes that change one's responses to learned threat. Here, we discuss two possible learning mechanisms.

Imagined Extinction as New Learning

Our findings provide strong evidence that imagined extinction engages the perceptual circuitry active during the encoding of real-world threat contingencies. One interpretation of this finding is that imagined extinction is a "true" simulation of real extinction learning. Simulation of real exposures may activate the extinction circuitry to form a new stimulus-outcome association that competes with the original learned threat association for expression. Like real extinction, vmPFC activity during imagined extinction may reflect integration of information distributed across the amygdala, NAc, and hippocampus (Nieuwenhuis and Takashima, 2011; Hartley and Phelps, 2010). As this information is integrated in a safe context, the vmPFC may suppress representations in the limbic system, which are no longer accurate or relevant.

This interpretation views imagined extinction as real extinction without the external stimulus. It follows then, that imagined extinction, like real extinction, would be marked by a gradual reduction in physiological responses to the conditioned threat stimulus (Verviet et al., 2013). We did not find evidence for this; as real extinction SCR responses decreased through time, imagined extinction responses, which were initially lower, slightly rose (Figure S4). However, the efficacy of exposure therapy is not well predicted by threat expression during or at the end of exposure (Craske et al., 2008), and SCR results in this investigation are possibly underpowered to fully examine this hypothesis. We also found differences in the degree to which the two processes involved the hippocampus, a region that is central to formation and recall of an extinction memory. Therefore, we maintain the position that though imagined and real extinction share core processes and accomplish comparable reductions in threat responding upon re-exposure, as a whole, they are functionally distinct.

Another type of new learning that imagination may influence is belief formation. This type of learning is more similar to cognitive approaches to emotion regulation (McRae, 2016). For example,

instead of learning something new about reinforcement contingencies in their environment, a person engaging in imagined extinction training may learn something new about their agency within it. Clinical reports indicate that it is not just the vividness of one's imagination that predicts therapeutic outcomes, but the interaction between vividness and one's sense of control over the imagery (Richardson, 1969; Singer and Pope, 1978). Hence, imagination can be used to increase one's internal locus of control, which may alter—not existing memories—but the way one approaches future threats. Thus, imagination may provide a stepping stone toward real exposure therapy. Alternatively, imagined exposures can yield adverse effects. Anxiety-driven imagination that strengthens the belief that one is unsafe is unlikely to have positive therapeutic outcomes (Jones and Davey, 1990). Indeed, emotional imagery is a hallmark of PTSD re-experiencing (Holmes and Mathews, 2010). In this way, imagination can exasperate a fear and promote avoidance behaviors. Drawing across this disparate evidence, we conclude that if imagination is well directed, so that the person feels safe and in control, one can reconceptualize learned threat contingencies in a wider context and change the value of the threatening stimulus via expectancy violation, which is a key component of extinction learning (Craske et al., 2014; Rescorla and Wagner, 1972).

Imagined Extinction as “Unlearning”

Instead of engendering new learning, imagined extinction may alter the existing threat association itself, whereby one unlearns the threat association. Repeated imagination of actions or events that never actually happened can lead to their being falsely remembered as having occurred in reality (Thomas et al., 2007), especially if the imagined events are consistent with existing knowledge. Intrusions of imagination on existing memories are thought to occur through a reconsolidation process (Schiller and Phelps, 2011; Schacter et al., 2011). Reconsolidation occurs after a consolidated memory re-enters a labile state following reactivation. The memory may be “updated” with new information and then reconsolidated. During imagined extinction, the perceptual simulation of real exposures may recreate the original learning context and reactivate the original threat memory in the amygdala, thereby assuming the state of initial learning. Similar to the theory behind gradual extinction learning (Gershman et al., 2015), which posits that a gradual reduction in reinforcement reduces the magnitude of prediction errors allowing extinction to become an extended state of acquisition learning; imagined extinction may, while recreating the original learning state, diminish or even omit prediction errors due to the lack of external input. Despite the lack of input, percepts continue to be experienced without external reinforcement. This may cause the memory trace to weaken its associations with threat-related physiology. Reconsolidation, of course, can only take place after the memory has first been consolidated. This experiment occurred in a single visit, so we are unable to assess “reconsolidation.” However, in this time frame, imagined extinction may become an extension of acquisition, so that only one stimulus–outcome memory is formed.

One way to disentangle these two learning hypotheses would be to test the durability of imagined extinction. Real extinction

suppresses the expression of a fear memory. It does not modify nor erase the original memory; this is why extinguished fear responses spontaneously recover over time (Quirk and Mueller, 2008). Reconsolidation modifies the original threat memory and therefore prevents the return of fear both spontaneously over time and after reinstatement (Schacter et al., 2011). Evidence from the clinic suggests that the effects of imaginal processes endure through time (Zoellner et al., 2017); however, a controlled investigation of imagined extinction assessing more than one visit would be necessary in order to determine whether imagined extinction is vulnerable to spontaneous recovery.

When interpreting the finding that imagined extinction engages perceptual circuitry active during acquisition, it is important to note that this experiment used a unimodal auditory conditioning paradigm. There is growing evidence that auditory processing has a direct impact on the learning structures supporting emotional behaviors. For example, amygdala-dependent threat-learning induces long-lasting changes in the sensitivity of the auditory cortex, in both the mouse and the human, and these changes persist after extinction (Bakin and Weinberger, 1990; Weinberger, 1998; Wigstrand et al., 2016). Indeed, Apergis-Schoute et al. (2014) found auditory association cortex BOLD signal is resistant to extinction. This suggests that sensory-cortical processing of threatening stimuli can be long lasting, even when immediate threat and corresponding amygdala activation have subsided. It is possible that this grounding of the CS in the sensory region where it was learned may allow for a quicker response to the threatening stimulus in the future. It is also possible that auditory sensation itself can gate negative arousal: music has been shown to alleviate pain during surgery recovery, even when the music is played while a person is under anesthesia (Hole et al., 2015; Kahloul et al., 2017). This evidence also suggests that the choice of sounds imagined by the no-extinction control group could have affected their performance. In this case, if the sounds of birds or rain were calming, this might produce extinction-like effects, which would work against the group differences (more powerful effects of both real and imagined extinction) that we observed here. To better understand the role of perceptual processes in imagined extinction, future studies could test for such unconditioned effects of affective qualities of multimodal imagined stimuli.

Nothing in the real world can be *perfectly* represented by the brain, but a “good enough” approximation allows one to predict and navigate their environment. Imagination is a process by which information about one's environment can be simulated and reorganized in order to improve predictions and learn under reduced risk. Imagination, when reduced to this basic functionality, is not unique to humans. Animals demonstrate deliberation when making choices, and this deliberation is thought to be analogous to imagined simulations of possible outcomes (Redish, 2016). Moser and Moser (2011) demonstrate that the resting mouse will both “replay” past actions and “preplay” future ones in a maze task. Furthermore, Takahashi et al. (2013) show that rats imagine outcomes when inferring paths through a mechanism that requires the integration of reinforcement histories of environmental cues via the orbitofrontal cortex. The human brain, however, may use imagination to draw upon richer experiences and cognitive frameworks in order to influence

memory, learning, emotion, decision making, expectations, and beliefs. In this way, imagination is a process that can inform the study of cognition and behavior in both human and non-human animals.

This investigation has strong implications for the treatment of anxiety and threat-related disorders. While the integration of imagination with exposure therapy is not new, our approach to threat simulation is. Clinical applications of imagination are not pure exposure tools: patients have expectations of recovery, they sometimes are trained to control their breath, and imagination may be combined with cognitive restructuring techniques, or other talk-based therapies (Craske et al., 2014). In this experiment, the only difference between the real and imagined extinction procedure is the existence of the external stimulus. This allowed us to directly test whether stimulus re-exposure is critical to extinction learning, or whether it can be internally simulated. We conclude that an internal simulation of a real-world experience can alter the way one responds to that situation in the future. Indeed, imagined exposures to threatening stimuli are effective in the reduction of learned threat responses and evoke a network of brain activation similar to real extinction. These novel findings bridge a long-standing gap between clinical practice and cognitive neuroscience. Once a topic reserved only for poets and philosophers, imagination is now being regarded by psychologists as an important cognitive tool for both decision-making and emotion regulation.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- CONTACT FOR REAGENT AND RESOURCE SHARING
- EXPERIMENTAL MODEL AND SUBJECT DETAILS
- METHOD DETAILS
 - Experimental Procedures
- DATA COLLECTION
 - Skin Conductance Signal Acquisition
 - Neuroimaging Acquisition
- QUANTIFICATION AND STATISTICAL ANALYSIS
 - SCR Preprocessing and Signal Decomposition
 - SCR Data Normalization
 - Outlier Removal
 - Imaging Preprocessing
 - Univariate Analysis
 - Classification analysis
 - Network Analysis Preprocessing
- DATA AND SOFTWARE AVAILABILITY

SUPPLEMENTAL INFORMATION

Supplemental Information includes six figures and six tables and can be found with this article online at <https://doi.org/10.1016/j.neuron.2018.10.047>.

ACKNOWLEDGMENTS

The authors would like to thank Dino Levy for discussions on the conceptualization of this study. Funding was provided by NIDA R03 DA035305, Klingenstein-Simons Fellowship Award in the Neurosciences to D.S., and NIDA R01 DA035484 to T.D.W.

AUTHOR CONTRIBUTIONS

Conceptualization, M.C.R., T.D.W., and D.S.; Methodology, M.C.R., T.D.W., and D.S.; Formal Analysis, M.R.; Investigation, M.C.R.; Data Curation, M.C.R.; Writing – Original Draft, M.C.R.; Writing – Review and Editing, T.D.W. and D.S.; Funding Acquisition, D.S., Resources, D.S.; Visualization, M.C.R.; Supervision, T.D.W. and D.S.

DECLARATION OF INTERESTS

The authors declare no competing interests.

Received: July 27, 2018

Revised: October 6, 2018

Accepted: October 24, 2018

Published: November 21, 2018

REFERENCES

- Acosta, A., and Vila, J. (1990). Emotional imagery: Effect of autonomic response information on physiological arousal. *Cogn. Emotion* 4, 145–160.
- Agren, T., Björkstrand, J., and Fredrikson, M. (2017). Disruption of human fear reconsolidation using imaginal and in vivo extinction. *Behav. Brain Res.* 319, 9–15.
- Apergis-Schoute, A.M., Schiller, D., LeDoux, J.E., and Phelps, E.A. (2014). Extinction resistant changes in the human auditory association cortex following threat learning. *Neurobiol. Learn. Mem.* 113, 109–114.
- Arbuthnott, K.D., Arbuthnott, D.W., and Rossiter, L. (2001). Guided imagery and memory: Implications for psychotherapists. *J. Couns. Psychol.* 48, 123–132.
- Atlas, L.Y., Bolger, N., Lindquist, M.A., and Wager, T.D. (2010). Brain mediators of predictive cue effects on perceived pain. *J. Neurosci.* 30, 12964–12977.
- Bakin, J.S., and Weinberger, N.M. (1990). Classical conditioning induces CS-specific receptive field plasticity in the auditory cortex of the guinea pig. *Brain Res.* 536, 271–286.
- Ben-Shakhar, G. (1985). Standardization within individuals: A simple method to neutralize individual differences in skin conductance. *Psychophysiology* 22, 292–299.
- Benedek, M., and Kaernbach, C. (2010). A continuous measure of phasic electrodermal activity. *J. Neurosci. Methods* 190, 80–91.
- Buckner, R.L., and Carroll, D.C. (2007). Self-projection and the brain. *Trends Cogn. Sci.* 11, 49–57.
- Craske, M.G., Kircanski, K., Zelikowsky, M., Mystkowski, J., Chowdhury, N., and Baker, A. (2008). Optimizing inhibitory learning during exposure therapy. *Behav. Res. Ther.* 46, 5–27.
- Craske, M.G., Treanor, M., Conway, C.C., Zbozinek, T., and Vervliet, B. (2014). Maximizing exposure therapy: An inhibitory learning approach. *Behav. Res. Ther.* 58, 10–23.
- Critchley, H., and Garfinkel, S. (2014). Neural correlates of fear: Insights from neuroimaging. *Neurosci. Biobehav. Rev.* 3, 111–125.
- Dadds, M.R., Bovbjerg, D.H., Redd, W.H., and Cutmore, T.R.H. (1997). Imagery in human classical conditioning. *Psychol. Bull.* 122, 89–103.
- Fullana, M.A., Harrison, B.J., Soriano-Mas, C., Vervliet, B., Cardoner, N., Àvila-Parcet, A., and Radua, J. (2016). Neural signatures of human fear conditioning: An updated and extended meta-analysis of fMRI studies. *Mol. Psychiatry* 21, 500–508.
- Gershman, S.J., Norman, K.A., and Niv, Y. (2015). Discovering latent causes in reinforcement learning. *Curr. Opin. Behav. Sci.* 5, 43–50.
- Graham, B.M., and Milad, M.R. (2011). The study of fear extinction: Implications for anxiety disorders. *Am. J. Psychiatry* 168, 1255–1265.

- Haaker, J., Golkar, A., Hermans, D., and Lonsdorf, T.B. (2014). A review on human reinstatement studies: An overview and methodological challenges. *Learn. Mem.* 21, 424–440.
- Hartley, C.A., and Phelps, E.A. (2010). Changing fear: The neurocircuitry of emotion regulation. *Neuropsychopharmacology* 35, 136–146.
- Herry, C., Ferraguti, F., Singewald, N., Letzkus, J.J., Ehrlich, I., and Lüthi, A. (2010). Neuronal circuits of fear extinction. *Eur. J. Neurosci.* 31, 599–612.
- Hole, J., Hirsch, M., Ball, E., and Meads, C. (2015). Music as an aid for post-operative recovery in adults: A systematic review and meta-analysis. *Lancet* 386, 1659–1671.
- Holland, P.C. (1990). Event representation in Pavlovian conditioning: Image and action. *Cognition* 37, 105–131.
- Holmes, E.A., and Mathews, A. (2005). Mental imagery and emotion: A special relationship? *Emotion* 5, 489–497.
- Holmes, E.A., and Mathews, A. (2010). Mental imagery in emotion and emotional disorders. *Clin. Psychol. Rev.* 30, 349–362.
- Johnson, A., Fenton, A.A., Kentros, C., and Redish, A.D. (2009). Looking for cognition in the structure within the noise. *Trends Cogn. Sci.* 13, 55–64.
- Jones, T., and Davey, G.C. (1990). The effects of cued UCS rehearsal on the retention of differential 'fear' conditioning: An experimental analogue of the "worry" process. *Behav. Res. Ther.* 28, 159–164.
- Jung, R.E., Flores, R.A., and Hunter, D. (2016). A new measure of imagination ability: Anatomical brain imaging correlates. *Front. Psychol.* 7, 496.
- Kahloul, M., Mhamdi, S., Nakhli, M.S., Sfehyi, A.N., Azzaza, M., Chaouch, A., and Naija, W. (2017). Effects of music therapy under general anesthesia in patients undergoing abdominal surgery. *Libyan J. Med.* 12, 1260886.
- Kosslyn, S.M., Ganis, G., and Thompson, W.L. (2001). Neural foundations of imagery. *Nat. Rev. Neurosci.* 2, 635–642.
- Kosslyn, S., Thompson, W., and Ganis, G. (2006). *The Case for Mental Imagery* (Oxford University Press).
- Lee, Y., Nelder, J.A., and Pawitan, Y. (2006). *Generalized Linear Models with Random Effects* (Chapman & Hall/CRC Press).
- LeDoux, J.E. (2000). Emotion circuits in the brain. *Annu. Rev. Neurosci.* 23, 155–184.
- Levita, L., Hoskin, R., and Champi, S. (2012). Avoidance of harm and anxiety: A role for the nucleus accumbens. *Neuroimage* 62, 189–198.
- Lucantonio, F., Gardner, M.P.H., Mirens, A., Newman, L.E., Takahashi, Y.K., and Schoenbaum, G. (2015). Neural estimates of imagined outcomes in basolateral amygdala depend on orbitofrontal cortex. *J. Neurosci.* 35, 16521–16530.
- McRae, K. (2016). Cognitive emotion regulation: A review of theory and scientific findings. *Curr. Opin. Behav. Sci.* 10, 119–124.
- Mendelsohn, A., Pine, A., and Schiller, D. (2014). Between thoughts and actions: Motivationally salient cues invigorate mental action in the human brain. *Neuron* 81, 207–217.
- Miller, G.A., Levin, D.N., Kozak, M.J., Cook, E.W., McLean, A., and Lang, P.J. (2008). Individual differences in imagery and the psychophysiology of emotion. *Cogn. Emotion* 1, 367–390.
- Morey, R.D., Rouder, J.N., Pratte, M.S., and Speckman, P.L. (2011). Using MCMC chain outputs to efficiently estimate Bayes factors. *J. Math. Psychol.* 55, 368–378.
- Moser, E.I., and Moser, M.B. (2011). Neuroscience: Seeing into the future. *Nature* 469, 303–304.
- Mumford, J.A., Turner, B.O., Ashby, F.G., and Poldrack, R.A. (2012). Deconvolving BOLD activation in event-related designs for multivoxel pattern classification analyses. *Neuroimage* 59, 2636–2643.
- Naveteur, J., and Bague, E.F. (1987). Individual differences in electrodermal activity as a function of subjects' anxiety. *Pers. Individ. Dif.* 8, 615–626.
- Nieuwenhuis, I.L., and Takashima, A. (2011). The role of the ventromedial prefrontal cortex in memory consolidation. *Behav. Brain Res.* 218, 325–334.
- Pascual-Leone, A., Amedi, A., Fregni, F., and Merabet, L.B. (2005). The plastic human brain cortex. *Annu. Rev. Neurosci.* 28, 377–401.
- Penny, W., Friston, K., Ashburner, J., Kiebel, S., and Nichols, T. (2007). *Statistical Parametric Mapping: The Analysis of Functional Brain Images* (Academic Press).
- Quirk, G.J., and Mueller, D. (2008). Neural mechanisms of extinction learning and retrieval. *Neuropsychopharmacology* 33, 56–72.
- Reddan, M.C., Lindquist, M.A., and Wager, T.D. (2017). Effect size estimation in neuroimaging. *JAMA Psychiatry* 74, 207–208.
- Redish, A.D. (2016). Vicarious trial and error. *Nat. Rev. Neurosci.* 17, 147–159.
- Rescorla, R.A., and Wagner, A.R. (1972). A theory of Pavlovian conditioning: Variations in the effectiveness of reinforcement and non-reinforcement. In *Classical Conditioning II: Current Research and Theory*, A.H. Prokasy, ed., ed. (Appleton-Century-Croft), pp. 64–99.
- Richardson, A. (1969). *Mental Imagery* (Springer).
- Rissman, J., Gazzaley, A., and D'Esposito, M. (2004). Measuring functional connectivity during distinct stages of a cognitive task. *Neuroimage* 23, 752–763.
- Schacter, D.L., Addis, D.R., and Buckner, R.L. (2007). Remembering the past to imagine the future: The prospective brain. *Nat. Rev. Neurosci.* 8, 657–661.
- Schacter, D.L., Guerin, S.A., and St Jacques, P.L. (2011). Memory distortion: An adaptive perspective. *Trends Cogn. Sci.* 15, 467–474.
- Schiller, D., and Delgado, M.R. (2010). Overlapping neural systems mediating extinction, reversal and regulation of fear. *Trends Cogn. Sci.* 14, 268–276.
- Schiller, D., and Phelps, E.A. (2011). Does reconsolidation occur in humans? *Front. Behav. Neurosci.* 5, 24.
- Schiller, D., Kanen, J.W., LeDoux, J.E., Monfils, M.H., and Phelps, E.A. (2013). Extinction during reconsolidation of threat memory diminishes prefrontal cortex involvement. *Proc. Natl. Acad. Sci. USA* 110, 20040–20045.
- Shepard, R.N., and Metzler, J. (1971). Mental rotation of three-dimensional objects. *Science* 171, 701–703.
- Singer, J.L., and Pope, K.S. (1978). *The Power of Human Imagination: New Methods in Psychotherapy* (Plenum Press).
- Soeter, M., and Kindt, M. (2012). Erasing fear for an imagined threat event. *Psychoneuroendocrinology* 37, 1769–1779.
- Storch, E., and McKay, D. (2013). Therapist barriers to the dissemination of exposure therapy. In *Handbook of Treating Variants and Complications in Anxiety Disorders*, J. Brand, J.M. Reid, and D. McKay, eds. (Springer), pp. 363–373.
- Stott, J.J., and Redish, A.D. (2014). A functional difference in information processing between orbitofrontal cortex and ventral striatum during decision-making behaviour. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 369, 1–12.
- Takahashi, Y.K., Chang, C.Y., Lucantonio, F., Haney, R.Z., Berg, B.A., Yau, H.J., Bonci, A., and Schoenbaum, G. (2013). Neural estimates of imagined outcomes in the orbitofrontal cortex drive behavior and learning. *Neuron* 80, 507–518.
- Taylor, S.E., Pham, L.B., Rivkin, I.D., and Armor, D.A. (1998). Harnessing the imagination. *Mental simulation, self-regulation, and coping*. *Am. Psychol.* 53, 429–439.
- Taylor, S., Thordarson, D.S., Maxfield, L., Fedoroff, I.C., Lovell, K., and Ogrodniczuk, J. (2003). Comparative efficacy, speed, and adverse effects of three PTSD treatments: Exposure therapy, EMDR, and relaxation training. *J. Consult. Clin. Psychol.* 71, 330–338.
- Thomas, A.K., Hannula, D.E., and Loftus, E.F. (2007). How self-relevant imagination affects memory for behaviour. *Appl. Cogn. Psychol.* 21, 69–86.
- van Minnen, A., and Foa, E.B. (2006). The effect of imaginal exposure length on outcome of treatment for PTSD. *J. Trauma. Stress* 19, 427–438.
- Varoquaux, G., Raamana, P.R., Engemann, D.A., Hoyos-Idrobo, A., Schwartz, Y., and Thirion, B. (2017). Assessing and tuning brain decoders: Cross-validation, caveats, and guidelines. *Neuroimage* 145 (Pt B), 166–179.

- Vervliet, B., Craske, M.G., and Hermans, D. (2013). Fear extinction and relapse: State of the art. *Annu. Rev. Clin. Psychol.* 9, 215–248.
- Weinberger, N.M. (1998). Physiological memory in primary auditory cortex: Characteristics and mechanisms. *Neurobiol. Learn. Mem.* 70, 226–251.
- Wigstrand, M.B., Schiff, H.C., Fyhn, M., LeDoux, J.E., and Sears, R.M. (2016). Primary auditory cortex regulates threat memory specificity. *Learn. Mem.* 24, 55–58.
- Woo, C.-W., Chang, L.J., Lindquist, M.A., and Wager, T.D. (2017). Building better biomarkers: Brain models in translational neuroimaging. *Nat. Neurosci.* 20, 365–377.
- Yarkoni, T., Poldrack, R.A., Nichols, T.E., Van Essen, D.C., and Wager, T.D. (2011). Large-scale automated synthesis of human functional neuroimaging data. *Nat. Methods* 8, 665–670.
- Zatorre, R.J., and Halpern, A.R. (2005). Mental concerts: Musical imagery and auditory cortex. *Neuron* 47, 9–12.
- Zoellner, L.A., Telch, M., Foa, E.B., Farach, F.J., McLean, C.P., Gallop, R., Bluett, E.J., Cobb, A., and Gonzalez-Lima, F. (2017). Enhancing extinction learning in posttraumatic stress disorder with brief daily imaginal exposure and methylene blue: A randomized controlled trial. *J. Clin. Psychiatry* 78, e782–e789.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Deposited Data		
Source code and analysis scripts for all figure data	Mendeley	https://doi.org/10.17632/46np3y2zg.1
Software and Algorithms		
CanlabRepository, MATLAB	In this paper	https://github.com/canlab/
SPM8, MATLAB	Penny et al., 2007	https://www.fil.ion.ucl.ac.uk/spm/software/spm8/
BayesFactor, R package	Morey et al., 2011	https://cran.r-project.org/web/packages/BayesFactor/BayesFactor.pdf
Ledlab, MATLAB package	Benedek and Kaernbach, 2010	http://www.ledlab.de/
Lme4, R package	Lee et al., 2006	https://cran.r-project.org/web/packages/lme4/lme4.pdf

CONTACT FOR REAGENT AND RESOURCE SHARING

Further information and requests for resources and code should be directed to and will be fulfilled by the Lead Contact, Daniela Schiller (daniela.schiller@mssm.edu).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Sixty-eight (45 females, average age = 29.64 STD 15.89, 45 participants were white, 5 African American, 9 Hispanic/Latino, 8 Asian, and 1 Native American) adult healthy members of the New York community, who had not participated in a shock-related experiment in at least six months prior to recruitment participated in this experiment. The gender identity of the participants is not known. Participants were only asked to provide their biological sex as male or female.

Participants were recruited through internal and surrounding communities near Icahn School of Medicine at Mount Sinai and New York University. All participants gave informed consent and were paid for their participation. Participants were not aware that there were multiple experimental conditions. This study was approved by the Icahn School of Medicine at Mount Sinai Institutional Review Board.

METHOD DETAILS

Experimental Procedures

Eligibility Criteria

Participants were required to be between the ages of 18 to 65. Participants who met criteria for disorders under the DSM-IV were excluded from this study. Eligibility was determined by the SCID (structured clinical interview for DSM-IV), which was conducted by the experimenter prior to the scheduling of the MRI experiment.

Randomization

Participants were randomly assigned to an experimental condition (imagined, real, or no extinction) and order (A or B) by the experimenter upon arrival. A power analysis was not calculated for this study. Sample sizes were determined based on previously published studies of threat conditioning and extinction. At least 20 participants were decided to be necessary in each of the three groups.

Shock Work-Up Procedure

Mild electric shocks were delivered through a bar electrode attached to the right wrist. A Grass Instruments stimulator was used with cable leads that were magnetically shielded and grounded through an RF filter. Shock strength was determined by a work-up procedure. In this procedure, the subject was first given a mild shock (200 ms duration, 50 pulses/s) which was gradually increased to a level the subject indicated as maximally “uncomfortable, but not painful” (the maximum shock that could given was 50 V). The average shock strength of participants in this experiment was 24.28 V (STD 10.19).

Stimuli

Several auditory stimuli were used in this experiment: (1) a 800 Hz pure “high” tone, (2) a 170 Hz pure “low” tone, (3) a sound clip of birds singing, and (4) a sound clip of rain falling. The tones were used because they had been previously used in threat conditioning

experiments (Apergis-Schoute et al., 2014). The nature sounds were selected for the no extinction control session because the sounds of birds singing and rain falling are quite commonly associated with relaxation, and therefore, they control for the aspect of imagination which we intended to control for: its relaxing, internally-directed properties.

Practice Session

All 68 participants completed a “practice session” before the experiment in which they were asked to listen to and imagine four types of auditory stimuli. This practice allowed participants to familiarize themselves with the cued auditory imagination procedure. The following auditory stimuli were presented for 4 s two times each: (1) a 800 Hz pure “high” tone, (2) a 170 Hz pure “low” tone, (3) a sound clip of birds singing, and (4) a sound clip of rain falling. Participants were then told the cue words which will indicate to them which sound to imagine and when. The cue words were: high, low, birds, and rain, respectively. Participants were then cued to imagine each sound twice to practice the cued-imagination process. Participants were given the cue word ‘stop,’ to indicate when they should stop imagining the sound after each 4 s cued-imagination trial.

Phase 1: Threat Acquisition

All 68 participants underwent an auditory threat-conditioning paradigm inside of the fMRI environment. The CSs were two easily distinguishable pure tones (800 and 170 Hz). They were delivered through MR-designed headphones, and were outside of the range of scanner noise. The US was a mild electric shock to the right wrist. All CSs were presented for 4 s, with a 10 s fixed inter-trial-interval (ITI). One of the tones was designated as the CS+ and was paired with shock on 33% of the trials. The other tone was the CS- and was never paired with shock. Assignment of the high and low tones to the CS+ and CS- conditions was counterbalanced across participants. Participants were not informed of the reinforcement contingency. They were told that they might receive shocks during the experiment, but that no further information could be given. Acquisition consisted of 8 randomized repetitions of CS+ tones paired with shock, 8 randomized repetitions of unreinforced CS+ tones, and 8 randomized repetitions of CS- tones.

Phase 2: Extinction Training

Participants then underwent one of three “extinction training sessions.” The first control group underwent real extinction training, which consisted of 15 pseudorandomized unreinforced presentations of each CS+ and CS- tone. The second control group was cued to imagine two neutral sounds from nature, which they heard previously during the practice session: birds singing and rain falling. There were 15 pseudorandomized cues to imagine each the birds and the rain. The third group, which was of main theoretical interest, was cued to imagine the CS+ and CS-, 15 times each, in a pseudorandomized order yoked to the presentation pattern of the real extinction-training group. No shocks were delivered during any of the extinction training sessions.

Phase 3: Reinstatement

All 68 participants underwent reinstatement of the threat memory: Four unsignaled shocks were delivered in temporally random pattern, whereby shocks were delivered between 4 to 12 s apart.

Phase 4: Re-Extinction

All 68 participants were exposed again to the CS+ and CS- tones. They received 10 presentations of each, in a randomized order.

DATA COLLECTION

Skin Conductance Signal Acquisition

Skin conductance was assessed with shielded Ag–AgCl electrodes attached to the middle phalanges of the second and third fingers of the left hand using BIOPAC systems EDA module. The electrode cables were grounded through an RF filter panel.

Neuroimaging Acquisition

The study was conducted at the NYU Center for Brain Imaging using a 3T Siemens Allegra scanner and a 32 channel Siemens head coil. Functional scans used a gradient echo sequence, TR = 2 s, TE = 20, flip ANGLE = 90, 183 FOV = 192, 3 mm slice thickness. A total of 39 axial slices were sampled for whole brain coverage. The in-plane resolution was 3 mm x 3 mm. Functional image acquisition was divided into four runs, one for each session, however the run for phase 3, reinstatement was not analyzed due to its short duration (approximately 1 minute). Between runs there was a break of approximately 15–30 s where the experimenter checked with the participant to make sure they were comfortable and alert. The scanning session ended with MPRAGE anatomical scans to obtain a 3D volume for slice selection.

QUANTIFICATION AND STATISTICAL ANALYSIS

SCR Preprocessing and Signal Decomposition

Skin conductance analysis was performed using the MATLAB package, Ledalab (Benedek and Kaernbach, 2010). Raw traces were downsampled to 100 Hz, then preprocessed with the software’s adaptive smoothing function, which employs a Gaussian low-pass filter to the data. Finally, the signal was deconvolved using a continuous decomposition analysis (CDA). This analysis separates the traces into continuous signals of tonic and phasic activity. The ‘phasic’ driving component of the signal can then be analyzed while controlling for slow, tonic changes. Event-related responses to stimuli were assessed by extracting the average phasic activity of SCRs occurring within a half a second after stimulus onset until a half a second after stimulus offset for each trial. The SCR analysis included only trials that did not coterminate with a presentation of the US.

SCR Data Normalization

Event-related phasic SCRs were next range normalized and log transformed within subjects, across all trials. This procedure was done so that individual variability in the signal would not bias further analyses.

Outlier Removal

Participants who did not demonstrate greater SCRs to the CS+ relative to the CS- on average across all acquisition trials ($n = 24$) were removed from analysis because it would be meaningless to investigate threat recovery in the signal of participants who did not demonstrate initial learning. It is important to note that individual differences in SCR are known to exist and that individuals vary in the reliability of their responding due to the equipment, properties of the skin, or psychological traits; Ben-Shakhar, 1985; Naveteur and Baque, 1987). A lack of a conditioned SCR does not mean the person did not learn cognitively or autonomically, only that the learning is not evident in their SCR signal.

Imaging Preprocessing

Imaging data were preprocessed and analyzed using SPM8 (<http://www.fil.ion.ucl.ac.uk/spm>) and custom MATLAB (MATLAB, The MathWorks, Inc., Natick, MA) code available from the authors' website (<http://canlab.colorado.edu>). Images were first unwarped and realigned to the first image of the series using a six-parameter, rigid-body transformation. The realigned images were then coregistered to each participant's T2-weighted structural image and normalized to the 152 MNI template using a 12-parameter nonlinear and affine transformation. The images were resliced during preprocessing from 3x3x3 voxels to 2x2x2 which is standard in SPM. Spatially normalized images were smoothed with a 6 mm full-width-at-half-maximum (FWHM) Gaussian kernel and filtered with a high pass filter (180 s cutoff).

Univariate Analysis

A univariate general linear model (GLM) was used to create images for the prediction analysis (Figure 2). For Phase 1: Acquisition, the model included three boxcar regressors, one for each stimulus presentation type: reinforced CS+, unreinforced CS+, and CS-. For the remaining phases, a single trial model, where each trial is modeled by one regressor was applied (Rissman et al., 2004; Mumford et al., 2012; Atlas et al., 2010). All regressors were convolved with a double gamma HRF function, and an additional 24 covariate regressors modeled movement effects (6 realignment parameters demeaned, their 1st derivatives, and the squares of these 12 regressors). Univariate analyses are reported in Figures 5 and 6 as well as Figures S1, S3, and S6.

Classification analysis

Using a cross-validated pattern classification procedure, we developed a whole brain neural pattern predictive of threat. This combination of multivariate pattern analysis and machine learning allows for the creation of a single integrated model of threat experiencing that accounts for activity distributed across networks. This approach has greater power and less bias than mass univariate approaches (Reddan et al., 2017). First, a support vector machine classifier was trained on single subject first-level GLM contrast images to distinguish between responses to the CS+ > baseline and responses to the CS- > baseline during Phase 1: Acquisition. In order to validate the neural threat-predictive pattern and reduce the chance of overfitting, we used a modified leave-one-subject out cross validation procedure which splits the data into "training" and "testing" data. Typically, a leave-one-subject-out procedure uses $N-1$ subjects for training and the remaining subject for testing. Because our subjects were divided into three groups for the extinction phase, we used a stricter $N-3$ procedure, where one subject from each group was held out. This efficient procedure allows every data point to serve as both training and test data. The resulting cross-validated neural threat-predictive pattern consists of the weights of each voxel in predicting the unreinforced CS+ or CS- stimulus presentation plus the intercept. However, to determine the voxels that made the most reliable contributions to the classification, we performed a bootstrap test. This involved taking 5,000 samples with replacement from the training dataset and repeating the prediction process with each bootstrap sample. This distribution was then converted into a z-value at each voxel and thresholded based on a voxel's corresponding p value, FDR corrected $p < 0.05$.

All pattern expression analyses using the test data were performed with the full set of non-thresholded weights, which included all nonzero voxels, unless otherwise specified (i.e., partial pattern expression looking at just the amygdala). We use the thresholded map to elucidate which voxels were most important for the classification. The non-thresholded weight map was applied to test data in an independent, leave-k-subjects-out fashion so that the participant whose data were being tested was not included in the neural-threat-predictive pattern (Varoquaux et al., 2017). That is, the weight map from the cross-validation fold that held out the participant to be tested was used to assess threat pattern expression in that subject. In this way, the pattern expression estimates for each individual were obtained from patterns trained on *other participants'* data, avoiding circularity. Pattern expression is calculated by taking the dot product between the neural threat pattern (classifier map) and a 'test' activation map, in this case, activation to the CS+ and CS- during the 'recovery test period.' A positive pattern expression is produced when positive data values are found in regions that are positive (CS+ > CS-) in the classifier map and negative data values are found in regions negative (CS- > CS+) in the classifier map.

Network Analysis Preprocessing

Preprocessing for the extinction network connectivity analysis (Figure 4) was completed using in-house MATLAB software (canlab_connectivity_preproc, <https://www.github.com/canlab>). Time course data were bandpass filtered (.008 - 0.25 window) and linear trend was removed. Nuisance regressors included the 24 motion parameters, spikes, and the top five principle components of signal from the white matter and ventricles.

Additional details about the specific analyses supporting the figures in this paper can be found in both the Results sections of the main paper and in the figure legends.

DATA AND SOFTWARE AVAILABILITY

Data and analysis scripts for this paper can be found at:

Mendeley: <https://dx.doi.org/10.17632/46npy3y2zg.1>

Open Science Framework: <https://osf.io/68yww/>

https://github.com/canlab/Attenuating_neural_threat_expression_with_imagination_2018/

Neuron, Volume 100

Supplemental Information

**Attenuating Neural Threat Expression
with Imagination**

Marianne Cumella Reddan, Tor Dessart Wager, and Daniela Schiller

Supplementary Tables and Figures

Supplementary Table 1. Significant clusters within the neural threat-predictive pattern. Related to Figure 2A

Region		MNI Coordinates			voxels	mm3	max stat
		x	y	z			
Cortex							
Frontal Lobe							
R	Inferior Frontal Gyrus	48	26	-18	1	8	-3.73
L	Middle Frontal Gyrus	-38	52	-10	256	2048	-5.73
R	Middle Frontal Gyrus	32	36	46	8	64	-3.92
R	Superior Frontal Gyrus	38	48	20	5	40	-3.90
R	Superior Frontal Gyrus	40	34	30	18	144	-4.56
L	Anterior Cingulate	-14	42	6	1	8	3.70
L	Cingulate Gyrus	-4	8	40	1	8	3.73
Temporal Lobe							
R	Insula	42	-16	14	4	32	3.80
L	Fusiform Gyrus	-44	-54	-8	12	96	4.17
R	Inferior Temporal Gyrus	28	-12	-42	7	56	3.94
		64	-14	-20	2	16	-3.84
R	Inferior Temporal Gyrus	64	-52	-18	7	56	4.27
L	Middle Temporal Gyrus	-52	-68	24	20	160	-4.13
		-62	-32	-12	2	16	-3.71
R	Middle Temporal Gyrus	54	-2	-8	9	72	-4.13
		66	-26	-4	4	32	-3.81
L	Superior Temporal Gyrus	-50	-2	6	1	8	3.70
		-48	-48	10	1	8	3.70
R	Parahippocampal Gyrus	12	-16	-16	2	16	3.96
L	Parahippocampal Gyrus	-24	-46	-4	4	32	4.23
R	Amygdala	24	-4	-6	11	88	4.03
Parietal Lobe							
L	Inferior Parietal Lobule	-48	-66	40	106	848	-4.85
R	Postcentral Gyrus	54	-24	22	13	104	4.41
L	Precuneus	-32	-64	36	1	8	-3.83
R	Precuneus	14	-58	42	1	8	-3.79
R	Superior Parietal Lobule	14	-56	64	47	376	-4.35
Occipital Lobe							

R	Cuneus	16	-76	10	4	32	4.02
SubCortex							
L	Basal Ganglia:Globus Pallidus	-12	0	-6	14	112	4.36
L	Thalamus: Ventral Anterior Nucleus	-8	-4	8	7	56	4.28
Cerebellum							
L	Cerebellum, Declive	-6	-74	-18	99	792	5.91
		-8	-60	-18	32	256	4.12
B Midbrain							
L	Periacqueductal Gray	-2	-26	-6	204	1632	6.04
R	Substantia Nigra	12	-18	-12	3	24	3.93

* Thresholded (bootstrapped 5,000 samples) and corrected for multiple comparisons (FDR $p < 0.05$, $k = 1$)

Supplementary Table 2. Significant clusters in univariate acquisition (CS+>CS-) map. Related to STAR METHODS QUANTIFICATION AND STATISTICAL ANALYSIS *Univariate Analysis* and Figure S1

Region		MNI Coordinates			voxels	mm3	maxstat
		x	y	z			
Cortex							
Frontal Lobe							
L	Medial Frontal Gyrus	-10	56	-14	75	600	-4.08
		-38	54	-8	753	6024	-5.01
L	Middle Frontal Gyrus	-42	18	42	51	408	-3.76
		-58	26	28	8	64	-3.41
L	Premotor Cortex	-10	-26	46	12	96	3.89
L	Orbitofrontal Cortex	-6	42	-28	17	136	-3.6
		-4	30	18	63	504	4.48
L	Anterior Cingulate Gyrus	-2	24	24	356	2848	4.92
		-20	22	24	33	264	4.37
		-4	8	40	427	3416	6.2
R	Middle Frontal Gyrus	58	44	-8	6	48	-3.38
R	Inferior Frontal Gyrus	36	20	-6	206	1648	4.2
R	Superior Frontal Gyrus	20	48	38	6	48	-3.43
R	Precentral Gyrus	34	-22	48	26	208	3.73
		48	-12	48	19	152	3.43
R	Mid Cingulate Gyrus	6	-20	28	9	72	3.51
R	Anterior Cingulate Gyrus	8	-22	44	109	872	4.15
		6	2	42	461	3688	6.53
Temporal Lobe							
L	Middle Temporal Gyrus	-60	-30	-12	685	5480	-5.73
		-38	-20	16	44	352	3.95
		-36	10	2	285	2280	4.94
L	Insula	-36	0	-6	116	928	4.1
		-30	20	8	417	3336	6.05
		-40	0	10	616	4928	6.65
L	Superior Temporal Gyrus	-56	2	6	146	1168	5.48
L	Transverse Temporal Gyrus	-52	-22	12	290	2320	5.15
L	Fusiform Gyrus	-40	-52	-6	14	112	7.82
R	Middle Temporal Gyrus	66	-20	-18	276	2208	-5.29
		58	-36	20	119	952	4.42
R	Insula	36	-22	18	153	1224	5.12
		48	-18	16	250	2000	5.24

	38	-10	-2	81	648	4.23
	36	-16	10	104	832	4.21
R Superior Temporal Gyrus	48	-58	28	90	720	-4.19
	56	-2	4	61	488	4.61
R Amygdala	22	-14	-8	69	552	4.43
Parietal Lobe						
	-46	-66	40	1337	10696	-5.98
L Inferior Parietal Lobule	-56	-28	24	627	5016	6.72
	-50	-38	22	123	984	4.48
L Precuneus	0	-60	36	99	792	-3.75
	56	-26	22	303	2424	6.46
R Inferior Parietal Lobule	50	-32	24	86	688	4.62
R Superior Parietal Lobule	40	-68	52	36	288	-3.91
R Angular Gyrus	38	-62	34	14	112	-3.69
Occipital Lobe						
L Middle Occipital Gyrus	-38	-92	14	30	240	-4.13
SubCortex						
L Caudate	-14	-12	20	6	48	3.37
L Lentiform Nucleus	-14	4	-6	476	3808	7.82
R Caudate	10	10	2	232	1856	4.16
R Lentiform Nucleus	24	2	-4	251	2008	5.34
R Nucleus Accumbens	22	12	-12	232	1856	4.32
R Claustrum	32	6	8	391	3128	5.72
R Thalamus: Medial Dorsal Nucleus	6	-18	10	83	664	3.97
Cerebellum						
L Cerebellum, Culmen	-28	-50	-28	147	1176	5.01
	-6	-56	-14	171	1368	4.83
L Cerebellum, Declive	-6	-70	-18	77	616	4.39
	4	-58	-28	27	216	3.66
R Cerebellum, Culmen	40	-50	-32	26	208	3.7
	2	-46	-16	5	40	3.34
R Cerebellum, Anterior Lobe	2	-48	-12	16	128	3.62
Brainstem						
L Midbrain, Periaqueductal Gray	-2	-14	-12	356	2848	5.41
R Pons	8	-34	-42	40	320	4.27
R Midbrain, Substantia Nigra	10	-20	-12	317	2536	5.79

* Thresholded and corrected for multiple comparisons (FDR $p < 0.05$, $k=5$)

Supplementary Table 3. Univariate activation during recovery test (CS + > CS-). Related to STAR METHODS QUANTIFICATION AND STATISTICAL ANALYSIS *Univariate Analysis* and Figure S4.

Imagined	Region	MNI Coordinates			voxels	mm3	max stat
		x	y	z			
Cortex							
Temporal Lobe							
R	'Temporal_Inf_R (aal)'	46	-70	-4	284	2272	-5.07
Occipital Lobe							
L	'Calcarine_L (aal)'	-10	-98	-6	111	888	-4.69
L	'Cuneus_L (aal)'	-6	-86	16	234	1872	-5.09
L	Occipital_Mid'	-26	-86	34	191	1528	-4.8
R	Occipital_Mid'	32	-74	32	151	1208	-4.59
Real	Region	MNI Coordinates			voxels	mm3	max stat
		x	y	z			
Cortex							
Frontal Lobe							
R	'Inferior Frontal Gyrus'	48	38	2	43	344	-4.57
R	'Inferior Frontal Gyrus'	58	8	36	80	640	-4.59
R	'Middle Frontal Gyrus'	24	28	-18	108	864	-4.29
R	'Middle Frontal Gyrus'	24	-2	48	98	784	-4.47
Temporal Lobe							
L	Amygdala	-16	-8	-32	120	960	-5.01
R	'Insula'	46	8	12	50	400	-4.43
R	Insula	42	4	24	71	568	-4.73
R	'Fusiform Gyrus'	56	-50	-24	68	544	-4.37
L	'Superior Temporal Gyrus'	-36	0	-20	66	528	-4.32
L	'Superior Temporal Gyrus'	-54	-42	8	76	608	-4.32
R	'Superior Temporal Gyrus'	40	6	-22	75	600	-4.33
L	Hippocampus	-40	-22	-14	52	416	-4.41
Parietal Lobe							
R	'Inferior Parietal Lobule'	56	-38	24	254	2032	-4.85
R	'Parietal_Inf_R (aal)'	32	-46	50	634	5072	-5.01
L	'Postcentral Gyrus'	-58	-24	40	205	1640	-4.52
Occipital Lobe							
L	'Middle Occipital Gyrus'	-32	-82	8	1349	10792	-6.02

R 'Precuneus'			20	-72	28	2978	23824	-6.19
Cerebellum								
L 'Culmen'			-44	-44	-30	294	2352	-5.56
L 'Declive'			-28	-84	-24	96	768	-5.37
Brainstem								
L 'Pons'			-8	-12	-32	240	1920	-5.01
None	Region	MNI Coordinates			voxels	mm3	max stat	
		x	y	z				
Cortex								
Frontal Lobe								
L 'Cingulate Gyrus'			-6	-12	30	60	480	5.11
R 'Middle Frontal Gyrus'			32	28	36	50	400	-4.14
L 'Inferior Frontal Gyrus'			-30	32	-4	117	936	5.45
Temporal Lobe								
L Amygdala			-10	-8	-14	68	544	4.21
L Hippocampus			-36	-28	-12	80	640	4.62
R Insula			46	4	20	77	616	-4.19

* Thresholded uncorrected ($p < 0.05$, $k = 5$)

**Supplementary Table 4. There was no effect of order or stimulus counterbalancing on SCR,
Related to Figure 3 and Figure S4**

Type III ANOVA with Satterthwaite's method

Source	<i>df</i>	<i>SS</i>	<i>MS</i>	<i>F</i>	<i>P</i>
Group	2, 60	0.003	0.002	0.325	0.724
Order	1, 60	0.002	0.002	0.355	0.554
Group:Order	2, 60	0.001	0.001	0.128	0.880

Post hoc within group contrasts (all phases): Order A – Order B

Group	<i>df</i>	<i>Estimate</i>	<i>SE</i>	<i>t.ratio</i>	<i>P</i>
Imagined Extinction	60	-0.001	0.016	-0.042	0.967
Standard Extinction	60	0.010	0.014	0.735	0.465
No Extinction	60	0.006	0.014	0.412	0.682

Within Acquisition Phase (all groups) ANOVA Contrast: Order A – B

	<i>df</i>	<i>SS</i>	<i>MS</i>	<i>F</i>	<i>P</i>
Order	1	0.004	0.004	0.256	0.614
Residuals	64	0.885	0.014		

Supplementary Table 5. Post hoc analyses of the neural network supporting imagined extinction. Related to Figure 4D

10 out of 12 nodes yielded group significant differences in betweenness centrality in a one-way ANOVA test, FDR corrected for multiple comparisons.

One-way Analysis of Variance of the Betweenness Centrality of the L NAc Node by Group

Source	df	SS	MS	F	P	η^2
Between groups	2	18008.8	9004.42	35.17	5.54e-11	0.53
Within groups	63	16131.1	256.05			
Total	65	34139.9				

Significant pair-wise two-sample t-tests with unequal variance:

None > Imagined $t(23) = 8.68, P < 0.0001, CI = [30.92, 50.26], \text{Hedges } g = 2.35$
None > Real $t(38.39) = 3.03, P = 0.004, CI = [5.56, 27.97], \text{Hedges } g = -0.86$
Real > Imagined $t(21) = 8.04, P < 0.0001, CI = [17.66, 29.98], \text{Hedges } g = 2.32$

One-way Analysis of Variance of the Betweenness Centrality of the R NAc Node by Group

Source	df	SS	MS	F	P	η^2
Between groups	2	12522.6	6261.31	21.03	1.01e-07	0.40
Within groups	63	18757.9	297.74			
Total	65	31280.5				

Significant pair-wise two-sample t-tests with unequal variance:

Imagined > None $t(25.42) = 5.79, P < 0.0001, CI = [17.57, 36.93], \text{Hedges } g = 1.58$
Imagined > Real $t(24.17) = 8.13, P < 0.0001, CI = [24.05, 40.40], \text{Hedges } g = 2.36$

One-way Analysis of Variance of the Betweenness Centrality of the L Amygdala-CM Node by Group

Source	df	SS	MS	F	P	η^2
Between groups	2	20386.3	10193.2	58.51	4.33e-15	0.65
Within groups	63	10975.5	174.2			
Total	65	31361.8				

Significant pair-wise two-sample t-tests with unequal variance:

Imagined > Real $t(25.8) = 13.59, P < 0.0001, CI = [32.08, 43.53], \text{Hedges } g = 3.96$
None > Real $t(40.87) = 8.14, P < 0.0002, CI = [27.69, 45.96], \text{Hedges } g = 2.32$

One-way Analysis of Variance of the Betweenness Centrality of the R Amygdala-CM Node by Group

Source	df	SS	MS	F	P	η^2
Between groups	2	492.1	246.038	1.59	0.211 n.s.	0.04
Within groups	63	9725.2	154.368			
Total	65	10217.3				

* ANOVA not significant; pair-wise t-tests not reported

One-way Analysis of Variance of the Betweenness Centrality of the L Amygdala-LB Node by Group

Source	df	SS	MS	F	P	η^2
Between groups	2	225.12	112.56	3.49	0.039 n.s.	0.10
Within groups	63	2031.91	32.25			
Total	65	2257.03				

* ANOVA not significant; pair-wise t-tests not reported

One-way Analysis of Variance of the Betweenness Centrality of the R Amygdala-LB Node by Group

Source	df	SS	MS	F	P	η^2
Between groups	2	5558	2779.00	17.6	8.44e-07	0.36
Within groups	63	9945.03	157.858			
Total	65	15503.03				

Significant pair-wise two-sample t-tests with unequal variance:

Real > Imagined $t(39.42) = 4.21, P = 0.0001, CI = [9.26, 26.34], \text{Hedges } g = 1.26$
Real > None $t(35.89) = 5.38, P < 0.0001, CI = [12.83, 28.34], \text{Hedges } g = 1.59$

One-way Analysis of Variance of the Betweenness Centrality of the L Auditory (Te1.0) Node by Group

Source	df	SS	MS	F	P	η^2
Between groups	2	6524.12	3262.06	761.66	7.36e-45	0.96
Within groups	63	269.82	4.28			
Total	65	6793.94				

Significant pair-wise two-sample t-tests with unequal variance:

Real > Imagined $t(21) = 27.60, P < 0.0001, CI = [19.50, 22.68], \text{Hedges } g = 7.97$
Real > None $t(21) = 27.59, P < 0.0001, CI = [19.50, 22.68], \text{Hedges } g = 8.37$

One-way Analysis of Variance of the Betweenness Centrality of the R Auditory (Te1.0) Node by Group

Source	df	SS	MS	F	P	η^2
Between groups	2	5551.62	2775.81	45619.84	2.65e-100	1.00
Within groups	63	3.83	0.06			
Total	65	5555.45				


Significant pair-wise two-sample t-tests with unequal variance:

Imagined > Real $t(23) = \text{Inf}, P < 0.0001, CI = [20, 20], \text{Hedges } g = \text{Inf}$
Imagined > None $t(23) = 239.00, P < 0.0001, CI = [19.74, 20.01], \text{Hedges } g = 64.74$


One-way Analysis of Variance of the Betweenness Centrality of the L CA1 Node by Group

Source	df	SS	MS	F	P	η^2
Between groups	2	31964.4	15982.2	251.05	9.71e-31	0.89
Within groups	63	4010.6	63.7			

Total	65	35975				
<i>Significant pair-wise two-sample t-tests with unequal variance:</i>						
Real > Imagined	$t(21.84) = 15.73, P < 0.0001, CI = [40.28, 52.52], \text{Hedges } g = 4.55$					
Real > None	$t(21.03) = 16.06, P < 0.0001, CI = [40.84, 52.99], \text{Hedges } g = 4.87$					
<i>One-way Analysis of Variance of the Betweenness Centrality of the R CA1 Node by Group</i>						
Source	df	SS	MS	F	P	η^2
Between groups	2	2978.95	1489.48	78.75	7.27e-18	0.71
Within groups	63	1191.53	18.91			
Total	65	4170.48				
<i>Significant pair-wise two-sample t-tests with unequal variance:</i>						
None > Real	$t(23) = 11.48, P < 0.0001, CI = [12.01, 17.31], \text{Hedges } g = 3.18$					
Imagined > Real	$t(19) = 15.68, P < 0.0001, CI = [11.87, 15.53], \text{Hedges } g = 4.99$					
<i>One-way Analysis of Variance of the Betweenness Centrality of the PAG Node by Group</i>						
Source	df	SS	MS	F	P	η^2
Between groups	2	3931.62	1965.81	46.95	3.29e-13	0.60
Within groups	63	2637.65	41.87			
Total	65	6569.27				
<i>Significant pair-wise two-sample t-tests with unequal variance:</i>						
None > Imagined	$t(29.74) = 2.84, P = 0.008, CI = [1.80, 11.04], \text{Hedges } g = 0.88$					
None > Real	$t(44) = 12.26, P < 0.0001, CI = [15.31, 21.34], \text{Hedges } g = 3.54$					
Imagined > Real	$t(28.77) = 5.32, p < 0.0001, CI = [7.33, 16.49], \text{Hedges } g = 1.65$					
<i>One-way Analysis of Variance of the Betweenness Centrality of the vmPFC Node by Group</i>						
Source	df	SS	MS	F	P	η^2
Between groups	2	19691.5	9845.73	30.79	4.72e-10	0.49
Within groups	63	20148.3	319.81			
Total	65	39839.8				
<i>Significant pair-wise two-sample t-tests with unequal variance:</i>						
Imagined > Real	$t(25.48) = 4.33, P = 0.0002, CI = [11.05, 31.04], \text{Hedges } g = 1.26$					
Imagined > None	$t(29) = 9.63, P < 0.0001, CI = [33.41, 51.42], \text{Hedges } g = 2.66$					
Real > None	$t(42.96) = 3.45, P = 0.001, CI = [8.88, 33.87], \text{Hedges } g = 1.00$					




Supplementary Figure 1. Univariate activation during threat Acquisition (CS+ > CS-), Related to Figure 2. This is the FDR corrected contrast map (CS+ > CS-) resulting from a univariate general linear model applied to all subjects ($n = 68$) during the Threat Acquisition phase. This map is similar to the multivariate Neural Threat-Predictive Pattern in the main text. Regions with positive activations include the dACC, PAG, amygdala and insula. Regions with negative activations include the precuneus and orbitofrontal cortex. See Table S2 for a detailed list of regions.




Supplementary Figure 2. Imagined and real extinction reduce neural and physiological threat expression, Related to Figures 2 and 3. A-B. Violin plot version of Figure 3 panels A and B in the main text. See Figure 3 in main text for details. **C.** The unthresholded classification weights in the amygdala extracted from the whole brain threat predictive pattern in Figure 2 of the main text. The neural threat-predictive pattern was masked with a bilateral anatomical mask of the amygdala and then applied to the threat recovery test phase (late re-extinction) in order to determine if threat expression was altered in this region of interest. **D.** The partial threat pattern expression of the amygdala was assessed in the three groups (analogous to Figure 3A). Group differences were not significant, but there is a trend consistent with the findings in this paper: The no extinction group increased threat expression in the amygdala during the recovery test but the imagined and real extinction groups did not ($F(2,63) = 2.35$, $p = 0.11$, $\eta^2 = 0.07$). The no extinction group ($\bar{x} = 0.02$, $n = 24$) is greater than real ($\bar{x} = -0.02$, $n = 22$; $t(38.68) = 2.00$, $p = 0.05$, Hedges $g = 0.59$), and is non-significantly greater than imagined ($\bar{x} = -0.0005$, $n = 20$, $t(41.04) = 1.46$, $p = 0.15$, Hedges $g = 0.43$). No difference was found between the imagined and real ($t(38.10) = 0.75$, $p = 0.46$, Hedges $g = 0.22$) extinction groups.



a. Imagined Extinction




b. Real Extinction



c. No Extinction




d. Comparison of average signal in a priori ROIs




Supplementary Figure 3. Univariate Activation during Recovery Test (CS + > CS-), Related to Figure 3. A general linear model contrasting activation to the last 5 trials of re-extinction CS+ > CS- was applied across the whole brain within each group. Uncorrected t-statistics are plotted for each map ($P < 0.05$, cluster size $k = 5$; see Table S3 for complete list of activations). **A. Imagined Extinction.** Amongst the significant clusters was the cuneus, which was negatively activated to the CS+ relative to CS-. This map demonstrates no evidence for threat recovery in the Imagined Extinction group. **B. Real Extinction.** Activations were distributed and largely negative. Notably, there were decreased responses to the CS+ relative to the CS- in the amygdala and the Inferior Parietal Lobule (IPL). **C. No Extinction.** The no extinction group demonstrated distributed positive activations, that is, greater responding to the threatening relative to the safety stimulus, in several regions known to be involved in threat expression, including the amygdala, anterior insula, hippocampus (CA1), and cingulate cortex. **D. Comparison of average signal in a priori ROIs.**

Unthresholded beta-weights in bilateral a priori ROIs, the amygdala and CA1, were extracted and averaged from the maps in A-C and then compared across groups. A one-way ANOVA revealed a non-significant trend of group effects in the amygdala ($F(2,63) = 2.23$, $p = 0.12$, $\eta^2 = 0.07$). Pairwise differences, revealed by two-sample t -tests, were not significant, but demonstrate a trend that activation in the amygdala is greater in the no extinction group ($\bar{x} = 0.09$, $n = 24$) than in the imagined ($t(35.64) = 1.95$, $p = 0.06$, $CI = [0.36, -0.01]$, Hedges $g = 0.59$) or real extinction groups ($t(39.77) = 1.734$, $p = 0.09$, $CI = [0.33, -0.03]$, Hedges $g = 0.51$). There was no difference between the imagined ($\bar{x} = -0.09$, $n = 20$) and real extinction ($\bar{x} = -0.06$, $n = 22$) groups ($t(39.61) = -0.25$, $p = 0.81$, $CI = [-0.23, 0.18]$, Hedges $g = -0.07$). A one-way ANOVA revealed group effects in the CA1 ($F(2,63) = 3.61$, $p = 0.03$, $\eta^2 = 0.10$). Pairwise t -tests revealed that activation in CA1 is greater in the no extinction group ($\bar{x} = 0.09$) than in the imagined ($t(38.86) = -2.38$, $p = 0.02$, $CI = [-0.27, -0.02]$, Hedges $g = -0.71$) or real extinction groups ($t(43.20) = -2.24$, $p = 0.03$, $CI = [-0.25, -0.01]$, Hedges $g = -0.65$). There was no difference between the imagined ($\bar{x} = -0.06$) and real extinction ($\bar{x} = -0.04$) groups ($t(39.13) = -0.25$, $p = 0.80$, $CI = [-0.15, 0.11]$, Hedges $g = -0.08$).




Supplementary Figure 4. Average skin conductance responses across all phases, Related to Figure 3. *A. Average Differential SCR Across Each Phase.* A linear mixed effects model including group and phase as predictors, indicated a significant effect of phase across the entire experiment ($F(4,156) = 9.19, P = 1.08e-06$). There was a trend towards a significant group by phase interaction ($F(8,156) = 1.78, P = 0.08$). The planned group comparison of threat-related SCRs during late re-extinction was significant and included in the main text. Post hoc analysis of group differences during early and late extinction phases revealed a significant group by phase interaction ($F(2,39) = 5.01, P = 0.01$) and a significant pairwise difference between imagined and real extinction groups during early extinction ($t(77.96) = -2.770, P = 0.02$; P value adjustment via tukey method for comparing a family of 3 estimates) Participants who did not demonstrate greater SCRs to the CS+ relative to the CS- on average across all acquisition trials ($n = 24$) were removed from analysis because it would be

meaningless to investigate threat recovery in the signal of participants who did not demonstrate initial learning. *B. Average Differential SCR Across Each Phase for All Subjects ($n = 66$)*. When participants who did not show a discriminatory SCR are not removed from the analysis, effects are diminished, but the trend remains. Additionally, this plot shows why it was necessary to remove participants who did not demonstrate a discriminatory SCR during learning; if there is no acquisition baseline it is difficult to assess the importance of recovery during late re-extinction.



Supplementary Figure 5. Imagined and real extinction yield similar patterns in the vmPFC during extinction, Related to Figure 5. To assess the similarity of the representational content in the vmPFC across ‘extinction’ sessions, voxel-wise activations were tested for correlations between groups during extinction (the third time bin; Figure 4B). The third time bin was selected *post hoc*, based on the average temporal activation data in order to focus on the peak activation during extinction. Imagined and real extinction had the highest correlation coefficient ($R = 0.51$, $P = 0.25$). Imagined and no extinction were weakly positively correlated ($R = 0.20$, $P = 0.76$) and real and no extinction were also weakly positively correlated ($R = 0.27$, $P = 0.66$). These correlations were not significant according to a nonparametric bootstrap test (10,000 samples). These results suggest that activations in the vmPFC are spatially analogous in imagined and real extinction, but require further data.



Supplementary Figure 6. Distribution of data supporting the effects in ROIs that predict the success of imagined and real extinction, Related to Figure 6. Scatterplots are shown for descriptive purposes. These plots illustrate the distribution of individual data values in Figure 6, but should not be taken as indicative of the true effect sizes (Reddan, Lindquist, & Wager, 2017).