

Fast Algorithm of General Quadratic Estimator Normalization

Toshiya Namikawa

Latest revision : November 21, 2016

Contents

1 Preliminaries	1
1.1 CMB	2
1.2 Lensing	2
1.3 Rotation	2
1.4 Inhomogeneous Reionization	2
1.5 Spherical Harmonics and Wigner-3j	3
1.6 Derivatives of Spherical Harmonics	3
1.7 Map derivatives	3
2 Distortion of CMB anisotropies	4
2.1 Lensing distortion	4
2.1.1 Lens distortion in harmonic space: Temperature	4
2.1.2 Lens distortion in harmonic space: Polarization	5
2.2 Tau distortion	6
2.3 Rotation distortion	7
2.4 Summary	7
3 Quadratic estimator	8
3.1 Distortion induced anisotropies	8
3.2 Weight Function	8
3.3 Weight Function: Derivations	8
4 General Fast formalism of Normalization	10
4.1 Normalization and Kernel function	10
4.2 Noise covariance and kernel function	11
5 Explicit Kernel Functions	13
5.1 Kernel Functions: Lensing	13
5.2 Kernel Functions: Rotation	16

1 Preliminaries

Here I will generalize the fast modal algorithm of [1, 2] to the case of rotation and others. In the followings, for multipoles of the CMB anisotropies, we use small letters (e.g., ℓ), while large letters are used for multipoles of the distortion fields (lensing, rotation, etc).

1.1 CMB

The CMB temperature fluctuations are denoted as Θ and the CMB linear polarization is expressed by the Stokes parameters, Q and U . The harmonic coefficients of the temperature anisotropies (and, in general, any scalar quantities x) are given by

$$x_{LM} = \int d^2\hat{n} Y_{LM}^*(\hat{n}) x(\hat{n}). \quad (1)$$

where Y_{LM} is the spin-0 spherical harmonics. On the other hand, the values of Q and U are changed by the rotation of the sphere. These Stokes parameters are therefore usually transformed into the rotational invariant quantities, the E and B modes, as

$$[E \pm iB]_{\ell m} = \int d^2\hat{n} (Y_{\ell m}^{\pm 2})^*(\hat{n}) [Q \pm iU](\hat{n}). \quad (2)$$

Here, $Y_{\ell m}^{\pm 2}$ is the spin-2 spherical harmonics. For short notation, we also use

$$\begin{aligned} \Xi^\pm &= E \pm iB, \\ P^\pm &= Q \pm iU \end{aligned} \quad (3)$$

1.2 Lensing

The lensing effect on CMB anisotropies is described as remapping of the unlensed CMB anisotropies by the deflection angle,

$$X(\hat{n}) = X(\hat{n} + \mathbf{d}), \quad (4)$$

where X is Θ or P^\pm .

The deflection angle of the CMB lensing is decomposed into the lensing potential, ϕ , and curl mode, ϖ , as

$$\mathbf{d} = \nabla\phi + \Delta\varpi, \quad (5)$$

where the operator $\Delta = \star\nabla$ denotes the derivatives with 90° rotation counterclockwise on the plane perpendicular to the line-of-sight direction and then operation. The harmonic coefficients of ϕ and ϖ are given by Eq. (1). The remapping of the CMB anisotropies is then given by

$$X(\hat{n}) = X(\hat{n}) + [\nabla\phi + \Delta\varpi] \cdot \nabla X + \mathcal{O}(\phi^2, \varpi^2). \quad (6)$$

1.3 Rotation

If the rotation angle is small, the modulation of polarization after rotation by an angle α is given by (e.g. [3])

$$\delta P^\pm = \pm 2\alpha P^\pm. \quad (7)$$

The harmonic coefficients of α is given by Eq. (1).

1.4 Inhomogeneous Reionization

The inhomogeneities of the reionization could vary the optical depth τ across the CMB sky. If the spatial variation of τ is very small, this leads to the modulation in CMB temperature and polarization as (e.g. [1, 4])

$$\Theta \rightarrow \Theta + \tau\Theta, P^\pm \rightarrow P^\pm + \tau P^\pm. \quad (8)$$

The harmonic coefficients of τ is given by Eq. (1).

1.5 Spherical Harmonics and Wigner-3j

The spherical harmonics is related to the Wigner-3j symbols as [5]

$$\int d^2\hat{n} Y_{\ell_1 m_1}^{s_1} Y_{\ell_2 m_2}^{s_2} Y_{\ell_3 m_3}^{s_3} = \sqrt{\frac{(2\ell_1+1)(2\ell_2+1)(2\ell_3+1)}{4\pi}} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ -s_1 & -s_2 & -s_3 \end{pmatrix} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ m_1 & m_2 & m_3 \end{pmatrix}. \quad (9)$$

1.6 Derivatives of Spherical Harmonics

In general, denoting $a_\ell^s = \sqrt{(\ell-s)(\ell+s)/2}$, the derivative of the spherical harmonics is given by

$$\nabla Y_{\ell m}^s = a_\ell^s Y_{\ell m}^{s+1} \mathbf{e}^* - a_\ell^{-s} Y_{\ell m}^{s-1} \mathbf{e}. \quad (10)$$

Here, we introduce the polarization vector \mathbf{e} which are defined

$$\mathbf{e} = \frac{\mathbf{e}_1 + i\mathbf{e}_2}{\sqrt{2}} \quad (11)$$

with \mathbf{e}_i denoting the basis vectors orthogonal to the radial vector. The polarization vector satisfies $\mathbf{e} \cdot \mathbf{e} = 0$, $\mathbf{e} \cdot \mathbf{e}^* = 1$, $\star \mathbf{e} = -i\mathbf{e}$. In particular, for $s = 0$,

$$\nabla Y_{\ell m} = a_\ell^0 (Y_{\ell m}^1 \mathbf{e}^* - Y_{\ell m}^{-1} \mathbf{e}), \quad (12)$$

and, for $s = \pm 2$, denoting $a_\ell^\pm = a_\ell^{\pm 2}$,

$$\begin{aligned} \nabla Y_{\ell m}^2 &= a_\ell^+ Y_{\ell m}^3 \mathbf{e}^* - a_\ell^- Y_{\ell m}^1 \mathbf{e}, \\ \nabla Y_{\ell m}^{-2} &= a_\ell^- Y_{\ell m}^{-1} \mathbf{e}^* - a_\ell^+ Y_{\ell m}^{-3} \mathbf{e}. \end{aligned} \quad (13)$$

1.7 Map derivatives

Derivative of scalar quantities such as the CMB temperature fluctuations and lensing potential is

$$\nabla x = \sum_{LM} x_{LM} \nabla Y_{LM} = \sum_{LM} x_{LM} a_L^0 (Y_{LM}^1 \mathbf{e}^* - Y_{LM}^{-1} \mathbf{e}) = x^+ \mathbf{e}^* - x^- \mathbf{e}. \quad (14)$$

where we define

$$x^\pm \equiv \sum_{LM} x_{LM} a_L^0 Y_{LM}^{\pm 1}, \quad (15)$$

and $(x^+)^* = -x^-$. The rotation of a pseudo-scalar quantity is given by

$$\Delta \varpi = \sum_{LM} \varpi_{LM} \Delta Y_{LM} = \sum_{LM} \varpi_{LM} a_L^0 i (Y_{LM}^1 \mathbf{e}^* + Y_{LM}^{-1} \mathbf{e}) = i(\varpi^+ \mathbf{e}^* + \varpi^- \mathbf{e}), \quad (16)$$

and $(\varpi^+)^* = -\varpi^-$. Spin-2 fields such as the CMB linear polarization is given by

$$\nabla P^+ = \sum_{\ell m} \Xi_{\ell m}^+ \nabla Y_{\ell m}^2 = \sum_{\ell m} \Xi_{\ell m}^+ (a_\ell^+ Y_{\ell m}^3 \mathbf{e}^* - a_\ell^- Y_{\ell m}^1 \mathbf{e}) = \Xi^{++} \mathbf{e}^* - \Xi^{+-} \mathbf{e}, \quad (17)$$

$$\nabla P^- = (\nabla P^+)^* = \sum_{\ell m} \Xi_{\ell m}^- \nabla Y_{\ell m}^{-2} = \sum_{\ell m} \Xi_{\ell m}^- (a_\ell^- Y_{\ell m}^{-1} \mathbf{e}^* - a_\ell^+ Y_{\ell m}^{-3} \mathbf{e}) = \Xi^{-+} \mathbf{e}^* - \Xi^{--} \mathbf{e}. \quad (18)$$

Note that $(\Xi^{++})^* = -\Xi^{--}$ and $(\Xi^{+-})^* = -\Xi^{-+}$.

2 Distortion of CMB anisotropies

In the following, we first define useful quantities to compute the distortion effect. The parity symmetry indicator is given by

$$p_{\ell_1 \ell_2 \ell_3}^{\pm} \equiv \frac{1 \pm (-1)^{\ell_1 + \ell_2 + \ell_3}}{2}, \quad (19)$$

$$(20)$$

An even (odd) parity quantity contains p^+ (p^-). A multipole factor is defined as

$$\gamma_{\ell_1 \ell_2 \ell_3} \equiv \sqrt{\frac{(2\ell_1 + 1)(2\ell_2 + 1)(2\ell_3 + 1)}{4\pi}}. \quad (21)$$

The convolutuon operator in full sky is defined as

$$\widetilde{\sum}_{LM\ell'm'}^{(\ell m)} \equiv \sum_{LM\ell'm'} (-1)^m \begin{pmatrix} \ell & L & \ell' \\ -m & M & m' \end{pmatrix}. \quad (22)$$

2.1 Lensing distortion

The lensing contributions in the position space become

$$\begin{aligned} \delta^{\phi} \Theta &= \nabla \phi \cdot \nabla \Theta = -\phi^- \Theta^+ - \phi^+ \Theta^-, \\ \delta^{\varpi} \Theta &= \Delta \varpi \cdot \nabla \Theta = i(\varpi^- \Theta^+ - \varpi^+ \Theta^-), \\ \delta^{\phi} P^{\pm} &= \nabla \phi \cdot \nabla P^{\pm} = -\phi^- \Xi^{\pm+} - \phi^+ \Xi^{\pm-}, \\ \delta^{\varpi} P^{\pm} &= \Delta \varpi \cdot \nabla P^{\pm} = i(\varpi^- \Xi^{\pm+} - \varpi^+ \Xi^{\pm-}). \end{aligned} \quad (23)$$

2.1.1 Lens distortion in harmonic space: Temperature

The harmonics transform of the lensing contributions is

$$\begin{aligned} \delta^{\phi} \Theta_{\ell m} &= - \int d^2 \hat{n} Y_{\ell m}^* [\phi^- \Theta^+ + \phi^+ \Theta^-] \\ &= - \sum_{LM\ell'm'} \phi_{LM} \Theta_{\ell'm'} a_L^0 a_{\ell'}^0 \int d^2 \hat{n} (-1)^m Y_{\ell, -m} [Y_{LM}^{-1} Y_{\ell'm'}^1 + Y_{LM}^1 Y_{\ell'm'}^{-1}] \\ &= - \sum_{LM\ell'm'} \phi_{LM} \Theta_{\ell'm'} 2a_L^0 a_{\ell'}^0 p_{\ell L \ell'}^+ \gamma_{\ell L \ell'} (-1)^m \begin{pmatrix} \ell & L & \ell' \\ -m & M & m' \end{pmatrix} \begin{pmatrix} \ell & L & \ell' \\ 0 & 1 & -1 \end{pmatrix} \\ &= - \widetilde{\sum}_{LM\ell'm'}^{(\ell m)} \phi_{LM} \Theta_{\ell'm'} 2a_L^0 a_{\ell'}^0 p_{\ell L \ell'}^+ \gamma_{\ell L \ell'} \begin{pmatrix} \ell & L & \ell' \\ 0 & 1 & -1 \end{pmatrix} \\ &= \widetilde{\sum}_{LM\ell'm'}^{(\ell m)} \phi_{LM} \Theta_{\ell'm'} W_{\ell L \ell'}^{\phi, 0}. \end{aligned} \quad (24)$$

Here we introduce coefficients $c_{\phi} = 1$ and $c_{\varpi} = -i$, and denote

$$W_{\ell_1 \ell_2 \ell_3}^{\phi, 0} = -2c_{\phi} a_{\ell_2}^0 a_{\ell_3}^0 p_{\ell_1 \ell_2 \ell_3}^+ \gamma_{\ell_1 \ell_2 \ell_3} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ 0 & 1 & -1 \end{pmatrix}. \quad (25)$$

Note that $(W_{\ell_1 \ell_2 \ell_3}^{\phi, 0})^* = W_{\ell_1 \ell_2 \ell_3}^{\phi, 0}$.

On the other hand, for curl mode,

$$\begin{aligned}
\delta^\varpi \Theta_{\ell m} &= i \int d^2 \hat{\mathbf{n}} Y_{\ell m}^* [\varpi^- \Theta^+ - \varpi^+ \Theta^-] \\
&= \sum_{LM\ell'm'} \varpi_{LM} \Theta_{\ell'm'} 2i a_L^0 a_{\ell'}^0 p_{\ell L \ell'}^- \gamma_{\ell L \ell'} (-1)^m \begin{pmatrix} \ell & L & \ell' \\ -m & M & m' \end{pmatrix} \begin{pmatrix} \ell & L & \ell' \\ 0 & 1 & -1 \end{pmatrix} \\
&= \widetilde{\sum_{LM\ell'm'}^{(\ell m)}} \varpi_{LM} \Theta_{\ell'm'} W_{\ell L \ell'}^{\varpi, 0},
\end{aligned} \tag{26}$$

with

$$W_{\ell_1 \ell_2 \ell_3}^{\varpi, 0} = -2c_\varpi a_{\ell_2}^0 a_{\ell_3}^0 p_{\ell_1 \ell_2 \ell_3}^- \gamma_{\ell_1 \ell_2 \ell_3} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ 0 & 1 & -1 \end{pmatrix}. \tag{27}$$

Note that the above quantity is consistent with Ref. [6] and also $(W_{\ell_1 \ell_2 \ell_3}^{\varpi, 0})^* = (-1)^{\ell_1 + \ell_2 + \ell_3} W_{\ell_1 \ell_2 \ell_3}^{\varpi, 0}$ which is consistent with (27).

2.1.2 Lens distortion in harmonic space: Polarization

The lensed anisotropies for polarizations are given by

$$\begin{aligned}
\delta^\phi \Xi_{\ell m}^\pm &= - \int d^2 \hat{\mathbf{n}} (Y_{\ell m}^{\pm 2})^* [\phi^- \Xi^{\pm+} + \phi^+ \Xi^{\pm-}] \\
&= - \sum_{LM\ell'm'} \phi_{LM} \Xi_{\ell'm'}^\pm a_L^0 \int d^2 \hat{\mathbf{n}} (Y_{\ell m}^{\pm 2})^* [a_{\ell'}^+ Y_{LM}^{\mp 1} Y_{\ell'm'}^{\pm 3} + a_{\ell'}^- Y_{LM}^{\pm 1} Y_{\ell'm'}^{\pm 1}] \\
&= - \sum_{LM\ell'm'} (-1)^m \begin{pmatrix} \ell & L & \ell' \\ -m & M & m' \end{pmatrix} \phi_{LM} \Xi_{\ell'm'}^\pm \gamma_{\ell L \ell'} a_L^0 \left[a_{\ell'}^+ \begin{pmatrix} \ell & L & \ell' \\ \mp 2 & \mp 1 & \pm 3 \end{pmatrix} + a_{\ell'}^- \begin{pmatrix} \ell & L & \ell' \\ \mp 2 & \pm 1 & \pm 1 \end{pmatrix} \right] \\
&= \widetilde{\sum_{LM\ell'm'}^{(\ell m)}} \phi_{LM} \Xi_{\ell'm'}^\pm W_{\ell L \ell'}^{\phi, \pm 2},
\end{aligned} \tag{28}$$

with

$$S_{\ell_1 \ell_2 \ell_3}^{\phi, 2} = (-1)^{\ell_1 + \ell_2 + \ell_3} S_{\ell_1 \ell_2 \ell_3}^{\phi, -2} = -c_\phi \gamma_{\ell_1 \ell_2 \ell_3} a_{\ell_2}^0 \left[a_{\ell_3}^+ \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ -2 & -1 & 3 \end{pmatrix} + a_{\ell_3}^- \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ -2 & 1 & 1 \end{pmatrix} \right]. \tag{29}$$

For curl mode,

$$\begin{aligned}
\delta^\varpi \Xi_{\ell m}^\pm &= i \int d^2 \hat{\mathbf{n}} (Y_{\ell m}^{\pm 2})^* [\varpi^- \Xi^{\pm+} - \varpi^+ \Xi^{\pm-}] \\
&= \pm i \sum_{LM\ell'm'} (-1)^m \begin{pmatrix} \ell & L & \ell' \\ -m & M & m' \end{pmatrix} \varpi_{LM} \Xi_{\ell'm'}^\pm a_L^0 \gamma_{\ell L \ell'} \left[a_{\ell'}^+ \begin{pmatrix} \ell & L & \ell' \\ \mp 2 & \mp 1 & \pm 3 \end{pmatrix} - a_{\ell'}^- \begin{pmatrix} \ell & L & \ell' \\ \mp 2 & \pm 1 & \pm 1 \end{pmatrix} \right] \\
&= \widetilde{\sum_{LM\ell'm'}^{(\ell m)}} \varpi_{LM} \Xi_{\ell'm'}^\pm W_{\ell L \ell'}^{\varpi, \pm 2},
\end{aligned} \tag{30}$$

with

$$W_{\ell_1 \ell_2 \ell_3}^{\varpi, 2} = -(-1)^{\ell_1 + \ell_2 + \ell_3} W_{\ell_1 \ell_2 \ell_3}^{\varpi, -2} = -c_\varpi \gamma_{\ell_1 \ell_2 \ell_3} a_{\ell_2}^0 \left[a_{\ell_3}^+ \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ -2 & -1 & 3 \end{pmatrix} - a_{\ell_3}^- \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ -2 & 1 & 1 \end{pmatrix} \right]. \tag{31}$$

Now we consider the lensed E/B modes separately. In general, for $X^\pm = A \pm iB = (a \pm ib)c^{(\pm)}$,

$$A = \frac{X^+ + X^-}{2} = \left(a \frac{c^{(+)} + c^{(-)}}{2} + ib \frac{c^{(+)} - c^{(-)}}{2} \right), \tag{32}$$

$$B = \frac{X^+ - X^-}{2i} = \left(-ai \frac{c^{(+)} - c^{(-)}}{2} + b \frac{c^{(+)} + c^{(-)}}{2} \right) \tag{33}$$

The lensing correction terms for E/B modes are then given by

$$\delta^x E_{\ell m} = \widetilde{\sum_{LM\ell'm'}}^{(\ell m)} \phi_{LM} [W_{\ell L\ell'}^{x,+} E_{\ell'm'} + W_{\ell L\ell'}^{x,-} B_{\ell'm'}] , \quad (34)$$

$$\delta^x B_{\ell m} = \widetilde{\sum_{LM\ell'm'}}^{(\ell m)} \phi_{LM} [-W_{\ell L\ell'}^{x,-} E_{\ell'm'} + W_{\ell L\ell'}^{x,+} B_{\ell'm'}] . \quad (35)$$

Here we define

$$\begin{aligned} W_{\ell_1\ell_2\ell_3}^{x,+} &\equiv \frac{W_{\ell_1\ell_2\ell_3}^{x,2} + W_{\ell_1\ell_2\ell_3}^{x,-2}}{2} = \frac{1 + c_x^2(-1)^{\ell_1+\ell_2+\ell_3}}{2} W_{\ell_1\ell_2\ell_3}^{x,2} \\ &= -\wp_{\ell_1\ell_2\ell_3}^{x,+} \gamma_{\ell_1\ell_2\ell_3} a_{\ell_2}^0 \left[a_{\ell_3}^+ \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ -2 & -1 & 3 \end{pmatrix} + c_x^2 a_{\ell_3}^- \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ -2 & 1 & 1 \end{pmatrix} \right] , \\ W_{\ell_1\ell_2\ell_3}^{x,-} &\equiv i \frac{W_{\ell_1\ell_2\ell_3}^{x,2} - W_{\ell_1\ell_2\ell_3}^{x,-2}}{2} = i \frac{1 - c_x^2(-1)^{\ell_1+\ell_2+\ell_3}}{2} W_{\ell_1\ell_2\ell_3}^{x,2} \\ &= -\wp_{\ell_1\ell_2\ell_3}^{x,-} \gamma_{\ell_1\ell_2\ell_3} a_{\ell_2}^0 \left[a_{\ell_3}^+ \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ -2 & -1 & 3 \end{pmatrix} + c_x^2 a_{\ell_3}^- \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ -2 & 1 & 1 \end{pmatrix} \right] , \end{aligned} \quad (36)$$

where we reintroduce a parity indicator as

$$\begin{aligned} \wp_{\ell_1\ell_2\ell_3}^{x,+} &= c_x \frac{1 + c_x^2(-1)^{\ell_1+\ell_2+\ell_3}}{2} , \\ \wp_{\ell_1\ell_2\ell_3}^{x,-} &= i c_x \frac{1 - c_x^2(-1)^{\ell_1+\ell_2+\ell_3}}{2} . \end{aligned} \quad (37)$$

2.2 Tau distortion

The harmonics transform of $\tau(\hat{n})\Theta(\hat{n})$ is

$$\begin{aligned} \delta\Theta_{\ell m} &= \int d^2\hat{n} Y_{\ell m}^* \tau(\hat{n})\Theta(\hat{n}) \\ &= \sum_{LM\ell'm'} \tau_{LM} \Theta_{\ell'm'} \int d^2\hat{n} Y_{\ell m}^* Y_{LM} Y_{\ell'm'} \\ &= \sum_{LM\ell'm'} \tau_{LM} \Theta_{\ell'm'} p_{\ell L\ell'}^+ \gamma_{\ell L\ell'} (-1)^m \begin{pmatrix} \ell & L & \ell' \\ -m & M & m' \end{pmatrix} \begin{pmatrix} \ell & L & \ell' \\ 0 & 0 & 0 \end{pmatrix} \\ &= \widetilde{\sum_{LM\ell'm'}}^{(\ell m)} \tau_{LM} \Theta_{\ell'm'} W_{\ell L\ell'}^{\tau,0} , \end{aligned} \quad (38)$$

where

$$W_{\ell L\ell'}^{\tau,0} = p_{\ell L\ell'}^+ \gamma_{\ell L\ell'} \begin{pmatrix} \ell & L & \ell' \\ 0 & 0 & 0 \end{pmatrix} . \quad (39)$$

2.3 Rotation distortion

The E and B modes after the rotation are given by

$$\begin{aligned}
\delta\Xi^\pm &= \pm 2 \int d^2\hat{\mathbf{n}} (Y_{\ell m}^{\pm 2})^* \alpha P^\pm \\
&= \pm 2 \sum_{LM\ell'm'} \alpha_{LM} \Xi_{\ell'm'}^\pm \int d^2\hat{\mathbf{n}} (Y_{\ell m}^{\pm 2})^* Y_{LM} Y_{\ell'm'}^{\pm 2} \\
&= \pm 2 \sum_{LM\ell'm'} (-1)^m \begin{pmatrix} \ell & L & \ell' \\ -m & M & m' \end{pmatrix} \alpha_{LM} \Xi_{\ell'm'}^\pm \gamma_{\ell L \ell'} \begin{pmatrix} \ell & L & \ell' \\ \mp 2 & 0 & \pm 2 \end{pmatrix} \\
&= \widetilde{\sum_{LM\ell'm'}^{(\ell m)}} \alpha_{LM} \Xi_{\ell'm'}^\pm W_{\ell L \ell'}^{\alpha, \pm 2},
\end{aligned} \tag{40}$$

with

$$W_{\ell_1 \ell_2 \ell_3}^{\alpha, \pm 2} = \pm 2 \gamma_{\ell_1 \ell_2 \ell_3} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ \mp 2 & 0 & \pm 2 \end{pmatrix}. \tag{41}$$

The distorted E and B modes are then described as

$$\delta E_{\ell m} = \widetilde{\sum_{LM\ell'm'}^{(\ell m)}} \alpha_{LM} (E_{\ell'm'} W_{\ell L \ell'}^{\alpha, +} + B_{\ell'm'} W_{\ell L \ell'}^{\alpha, -}), \tag{42}$$

$$\delta B_{\ell m} = \widetilde{\sum_{LM\ell'm'}^{(\ell m)}} \alpha_{LM} (-E_{\ell'm'} W_{\ell L \ell'}^{\alpha, -} + B_{\ell'm'} W_{\ell L \ell'}^{\alpha, +}) \tag{43}$$

where we define $c_\alpha = 1$ and

$$W_{\ell_1 \ell_2 \ell_3}^{\alpha, \pm} = 2 \varphi_{\ell_1 \ell_2 \ell_3}^{\alpha, \mp} \gamma_{\ell_1 \ell_2 \ell_3} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ -2 & 0 & 2 \end{pmatrix} \tag{44}$$

2.4 Summary

The above all distortions are described in the following form:

$$\delta\Theta_{\ell m} = \widetilde{\sum_{LM\ell'm'}^{(\ell m)}} x_{LM} \Theta_{\ell'm'} W_{\ell L \ell'}^{x, 0}, \tag{45}$$

$$\delta E_{\ell m} = \widetilde{\sum_{LM\ell'm'}^{(\ell m)}} x_{LM} (E_{\ell'm'} W_{\ell L \ell'}^{x, +} + B_{\ell'm'} W_{\ell L \ell'}^{x, -}), \tag{46}$$

$$\delta B_{\ell m} = \widetilde{\sum_{LM\ell'm'}^{(\ell m)}} x_{LM} (-E_{\ell'm'} W_{\ell L \ell'}^{x, -} + B_{\ell'm'} W_{\ell L \ell'}^{x, +}) \tag{47}$$

where x is a distortion field. The functional form of W is given above.

3 Quadratic estimator

3.1 Distortion induced anisotropies

The distortion fields x described above induce the off-diagonal elements of the covariance ($\ell \neq \ell'$ or $m \neq m'$),

$$\langle \tilde{X}_{\ell m} \tilde{Y}_{\ell' m'} \rangle_{\text{CMB}} = \sum_{LM} \begin{pmatrix} \ell & \ell' & L \\ m & m' & M \end{pmatrix} f_{\ell L \ell'}^{x, XY} x_{LM}^*, \quad (48)$$

where $\langle \cdots \rangle_{\text{CMB}}$ denotes the ensemble average over the primary CMB anisotropies with a fixed realization of the distortion fields. We ignore the higher-order terms of the distortion fields. The functional form of the weight functions f are discussed later.

With a quadratic combination of observed CMB anisotropies, \hat{X} and \hat{Y} , the general quadratic estimators are formed as (e.g., [7]),

$$[\hat{x}_{LM}^{XY}]^* = A_L^{x, XY} \sum_{\ell \ell' m m'} \begin{pmatrix} \ell & \ell' & L \\ m & m' & M \end{pmatrix} g_{\ell L \ell'}^{x, XY} \hat{X}_{\ell m} \hat{Y}_{\ell' m'}. \quad (49)$$

Here we define

$$g_{\ell L \ell'}^{x, XY} = \frac{[f_{\ell L \ell'}^{x, XY}]^*}{\Delta^{XY} \hat{C}_{\ell}^{XX} \hat{C}_{\ell'}^{YY}} \quad (50)$$

$$A_L^{x, XY} = \frac{1}{2L+1} \sum_{\ell \ell'} f_{\ell L \ell'}^{x, XY} g_{\ell L \ell'}^{x, XY}, \quad (51)$$

where $\Delta^{XX} = 2$, $\Delta^{EB} = \Delta^{TB} = 1$, and \hat{C}_{ℓ}^{XX} (\hat{C}_{ℓ}^{YY}) is the observed power spectrum.

3.2 Weight Function

The weight functions are, in general, given as

$$f_{\ell L \ell'}^{x, (\Theta\Theta)} = W_{\ell L \ell'}^{x, 0} C_{\ell'}^{\Theta\Theta} + p_x W_{\ell' L \ell}^{x, 0} C_{\ell}^{\Theta\Theta}, \quad (52)$$

$$f_{\ell L \ell'}^{x, (\Theta E)} = W_{\ell L \ell'}^{x, 0} C_{\ell'}^{\Theta E} + p_x W_{\ell' L \ell}^{x, +} C_{\ell}^{\Theta E}, \quad (53)$$

$$f_{\ell L \ell'}^{x, (\Theta B)} = p_x W_{\ell' L \ell}^{x, -} C_{\ell}^{\Theta E}, \quad (54)$$

$$f_{\ell L \ell'}^{x, (EE)} = W_{\ell L \ell'}^{x, +} C_{\ell'}^{EE} + p_x W_{\ell' L \ell}^{x, +} C_{\ell}^{EE}, \quad (55)$$

$$f_{\ell L \ell'}^{x, (EB)} = W_{\ell L \ell'}^{x, -} C_{\ell'}^{BB} + p_x W_{\ell' L \ell}^{x, -} C_{\ell}^{EE}, \quad (56)$$

$$f_{\ell L \ell'}^{x, (BB)} = W_{\ell L \ell'}^{x, +} C_{\ell'}^{BB} + p_x W_{\ell' L \ell}^{x, +} C_{\ell}^{BB}. \quad (57)$$

Here, the parity index is $p_{\phi} = p_{\epsilon} = 1$ and $p_{\varpi} = p_{\alpha} = -1$. Strictly speaking, p_x should be $(-1)^{\ell' + L + \ell}$. However, W is only non-zero when $\ell' + L + \ell$ is even, and vice versa. The parity even quantities are $x = \phi$ and ϵ . The odd parity quantities are $x = \varpi$ and α . Note that the above weight functions are consistent with Ref. [6] ($W_{\ell L \ell'}^{x, -} = -_{\Theta} S_{\ell L \ell'}^x$) for the lensing case.

3.3 Weight Function: Derivations

Let us first consider the temperature case. There are two contributions to the temperature quadratic estimator, and the one is given as

$$\begin{aligned} \langle \Theta_{\ell'' m''} \delta \Theta_{\ell m} \rangle &= \sum_{LM \ell' m'}^{(\ell m)} x_{LM} \langle \Theta_{\ell'' m''} \Theta_{\ell' m'} \rangle W_{\ell L \ell'}^{x, 0} \\ &= \sum_{LM \ell' m'}^{(\ell m)} x_{LM} \delta_{\ell'' \ell'} \delta_{m'', -m'} (-1)^{m'} C_{\ell'}^{\Theta\Theta} W_{\ell L \ell'}^{x, 0} \\ &= \sum_{LM} \begin{pmatrix} \ell & \ell'' & L \\ m & m'' & M \end{pmatrix} x_{LM}^* C_{\ell''}^{\Theta\Theta} W_{\ell L \ell''}^{x, 0}. \end{aligned} \quad (58)$$

Here, we use

$$\begin{aligned}
\widetilde{\sum_{LM\ell'm'}}^{(\ell m)} \delta_{\ell''\ell'} \delta_{m'',-m'} (-1)^{m'} x_{LM} &= \sum_{LM} (-1)^{m+m'} \begin{pmatrix} \ell & L & \ell'' \\ -m & M & -m'' \end{pmatrix} x_{LM} \\
&= \sum_{LM} (-1)^{-M} \begin{pmatrix} \ell & \ell'' & L \\ m & m'' & -M \end{pmatrix} x_{LM} \\
&= \sum_{LM} \begin{pmatrix} \ell & \ell'' & L \\ m & m'' & M \end{pmatrix} x_{LM}^* \tag{59}
\end{aligned}$$

The other term is obtained by exchanging $(\ell'', m'') \leftrightarrow (\ell, m)$ and is given by

$$\langle \Theta_{\ell m} \delta \Theta_{\ell'' m''} \rangle = \sum_{LM} (-1)^{\ell+\ell''+L} \begin{pmatrix} \ell & \ell'' & L \\ m & m'' & M \end{pmatrix} x_{LM}^* C_{\ell}^{\Theta\Theta} W_{\ell' L \ell}^{x,0}. \tag{60}$$

The sign $(-1)^{\ell+\ell''+L}$ depends on the parity of W .

For EB estimator, the two contributions are

$$\begin{aligned}
E_{\ell'' m''} \delta B_{\ell m} &= - \widetilde{\sum_{LM\ell'm'}}^{(\ell m)} x_{LM} \langle E_{\ell'' m''} E_{\ell' m'} \rangle W_{\ell L \ell'}^{x,-} \\
&= - \widetilde{\sum_{LM\ell'm'}}^{(\ell m)} x_{LM} (-1)^{m''} \delta_{\ell''\ell'} \delta_{m'',-m'} C_{\ell''}^{\text{EE}} W_{\ell L \ell'}^{x,-} \\
&= - \sum_{LM} \begin{pmatrix} \ell & \ell'' & L \\ m & m'' & M \end{pmatrix} x_{LM}^* C_{\ell''}^{\text{EE}} W_{\ell L \ell'}^{x,-}, \tag{61}
\end{aligned}$$

and

$$\begin{aligned}
B_{\ell'' m''} \delta E_{\ell m} &= \widetilde{\sum_{LM\ell'm'}}^{(\ell m)} x_{LM} \langle E_{\ell'' m''} E_{\ell' m'} \rangle W_{\ell L \ell'}^{x,-} \\
&= \sum_{LM} (-1)^{\ell+\ell''+L} \begin{pmatrix} \ell & \ell'' & L \\ m & m'' & M \end{pmatrix} x_{LM}^* C_{\ell}^{\text{EE}} W_{\ell' L \ell}^{x,-}. \tag{62}
\end{aligned}$$

4 General Fast formalism of Normalization

4.1 Normalization and Kernel function

The normalization of the $\Theta\Theta$ quadratic estimator is

$$\begin{aligned} \frac{1}{A_L^{x,(\Theta\Theta)}} &= \frac{1}{2L+1} \sum_{\ell\ell'} \frac{[W_{\ell L\ell'}^{x,0} C_{\ell'}^{\Theta\Theta} + p_x W_{\ell' L\ell}^{x,0} C_{\ell}^{\Theta\Theta}]^2}{2\widehat{C}_{\ell}^{\Theta\Theta} \widehat{C}_{\ell'}^{\Theta\Theta}} \\ &= \frac{1}{2} \Sigma_L^{(0),x} \left[\frac{1}{\widehat{C}^{\Theta\Theta}}, \frac{(C^{\Theta\Theta})^2}{\widehat{C}^{\Theta\Theta}} \right] + p_x \Gamma_L^{(0),x} \left[\frac{C^{\Theta\Theta}}{\widehat{C}^{\Theta\Theta}}, \frac{C^{\Theta\Theta}}{\widehat{C}^{\Theta\Theta}} \right] + \frac{1}{2} \Sigma_L^{(0),x} \left[\frac{(C^{\Theta\Theta})^2}{\widehat{C}^{\Theta\Theta}}, \frac{1}{\widehat{C}^{\Theta\Theta}} \right], \end{aligned} \quad (63)$$

where we define kernel functions as

$$\Sigma_L^{(0),x}[A, B] = \frac{1}{2L+1} \sum_{\ell\ell'} (W_{\ell L\ell'}^{x,0})^2 A_{\ell} B_{\ell'}, \quad (64)$$

$$\Gamma_L^{(0),x}[A, B] = \frac{1}{2L+1} \sum_{\ell\ell'} W_{\ell L\ell'}^{x,0} W_{\ell' L\ell}^{x,0} A_{\ell} B_{\ell'}. \quad (65)$$

For ΘE ,

$$\begin{aligned} \frac{1}{A_L^{x,(\Theta E)}} &= \frac{1}{2L+1} \sum_{\ell\ell'} \frac{|W_{\ell L\ell'}^{x,0} C_{\ell'}^{\Theta E} + p_x W_{\ell' L\ell}^{x,+} C_{\ell}^{\Theta E}|^2}{\widehat{C}_{\ell}^{\Theta\Theta} \widehat{C}_{\ell'}^{\Theta E}} \\ &= \frac{1}{2L+1} \sum_{\ell\ell'} \left[(W_{\ell L\ell'}^{x,0})^2 \frac{(C_{\ell'}^{\Theta E})^2}{\widehat{C}_{\ell}^{\Theta\Theta} \widehat{C}_{\ell'}^{\Theta E}} + 2p_x W_{\ell L\ell'}^{x,0} W_{\ell' L\ell}^{x,+} \frac{C_{\ell'}^{\Theta E} C_{\ell}^{\Theta E}}{\widehat{C}_{\ell}^{\Theta\Theta} \widehat{C}_{\ell'}^{\Theta E}} + (W_{\ell' L\ell}^{x,+})^2 \frac{(C_{\ell}^{\Theta E})^2}{\widehat{C}_{\ell}^{\Theta\Theta} \widehat{C}_{\ell'}^{\Theta E}} \right] \\ &= \Sigma_L^{(0),x} \left[\frac{1}{\widehat{C}^{\Theta\Theta}}, \frac{(C^{\Theta E})^2}{\widehat{C}^{\Theta E}} \right] + 2p_x \Gamma_L^{(\times),x} \left[\frac{C^{\Theta E}}{\widehat{C}^{\Theta\Theta}}, \frac{C^{\Theta E}}{\widehat{C}^{\Theta E}} \right] + \Sigma_L^{(+),x} \left[\frac{1}{\widehat{C}^{\Theta E}}, \frac{(C^{\Theta E})^2}{\widehat{C}^{\Theta\Theta}} \right], \end{aligned} \quad (66)$$

where kernel functions are defined as

$$\begin{aligned} \Gamma_L^{(\times),x}[A, B] &= \frac{1}{2L+1} \sum_{\ell\ell'} W_{\ell L\ell'}^{x,0} W_{\ell' L\ell}^{x,+} A_{\ell} B_{\ell'}, \\ \Sigma_L^{(+),x}[A, B] &= \frac{1}{2L+1} \sum_{\ell\ell'} (W_{\ell L\ell'}^{x,+})^2 A_{\ell} B_{\ell'}. \end{aligned} \quad (67)$$

For ΘB ,

$$\begin{aligned} \frac{1}{A_L^{x,(\Theta B)}} &= \frac{1}{2L+1} \sum_{\ell\ell'} \frac{|W_{\ell' L\ell}^{x,-} C_{\ell}^{\Theta E}|^2}{\widehat{C}_{\ell}^{\Theta\Theta} \widehat{C}_{\ell'}^{\Theta B}} \\ &= \Sigma_L^{(-),x} \left[\frac{1}{\widehat{C}^{\Theta B}}, \frac{(C^{\Theta E})^2}{\widehat{C}^{\Theta\Theta}} \right], \end{aligned} \quad (68)$$

where

$$\Sigma_L^{(-),x}[A, B] = \frac{1}{2L+1} \sum_{\ell\ell'} |W_{\ell L\ell'}^{x,-}|^2 A_{\ell} B_{\ell'}. \quad (69)$$

For EE (and for BB by replacing $EE \rightarrow BB$),

$$\begin{aligned} \frac{1}{A_L^x} &= \frac{1}{2L+1} \sum_{\ell\ell'} \frac{|W_{\ell L\ell'}^{x,+} C_{\ell'}^{\Theta E} + p_x W_{\ell' L\ell}^{x,+} C_{\ell}^{\Theta E}|^2}{2\widehat{C}_{\ell}^{\Theta E} \widehat{C}_{\ell'}^{\Theta E}} \\ &= \Sigma_L^{(+),x} \left[\frac{1}{\widehat{C}^{\Theta E}}, \frac{(C^{\Theta E})^2}{\widehat{C}^{\Theta E}} \right] + p_x \Gamma_L^{(+),x} \left[\frac{C^{\Theta E}}{\widehat{C}^{\Theta E}}, \frac{C^{\Theta E}}{\widehat{C}^{\Theta E}} \right], \end{aligned} \quad (70)$$

where

$$\Gamma_L^{(+),x}[A, B] = \frac{1}{2L+1} \sum_{\ell\ell'} W_{\ell'L\ell}^{x,+} W_{\ell L\ell'}^{x,+} A_\ell B_{\ell'} = \Gamma_L^{(+),x}[B, A]. \quad (71)$$

For EB ,

$$\begin{aligned} \frac{1}{A_L^{x,(EB)}} &= \frac{1}{2L+1} \sum_{\ell\ell'} \frac{|W_{\ell L\ell'}^{x,-} C_{\ell'}^{\text{BB}} + p_x W_{\ell'L\ell}^{x,-} C_\ell^{\text{EE}}|^2}{\widehat{C}_\ell^{\text{EE}} \widehat{C}_{\ell'}^{\text{BB}}} \\ &= \Sigma_L^{(-),x} \left[\frac{1}{\widehat{C}^{\text{EE}}}, \frac{(C^{\text{BB}})^2}{\widehat{C}^{\text{BB}}} \right] + 2p_x \Gamma_L^{(-),x} \left[\frac{C^{\text{EE}}}{\widehat{C}^{\text{EE}}}, \frac{C^{\text{BB}}}{\widehat{C}^{\text{BB}}} \right] + \Sigma_L^{(-),x} \left[\frac{1}{\widehat{C}^{\text{BB}}}, \frac{(C^{\text{EE}})^2}{\widehat{C}^{\text{EE}}} \right], \end{aligned} \quad (72)$$

where

$$\Gamma_L^{(-),x}[A, B] = \frac{1}{2L+1} \sum_{\ell\ell'} [W_{\ell L\ell'}^{x,-}]^* W_{\ell'L\ell}^{x,-} A_\ell B_{\ell'} = \Gamma_L^{(-),x}[B, A]. \quad (73)$$

4.2 Noise covariance and kernel function

For $\Theta\Theta\Theta E$,

$$\begin{aligned} \frac{A_L^{x,(\Theta\Theta)} A_L^{x,(\Theta E)}}{N_L^{x,(\Theta\Theta\Theta E)}} &= \frac{1}{2L+1} \sum_{\ell\ell'} \left[\frac{W_{\ell L\ell'}^{x,0} C_{\ell'}^{\Theta\Theta}}{2\widehat{C}_\ell^{\Theta\Theta} \widehat{C}_{\ell'}^{\Theta\Theta}} + p_x(\ell \leftrightarrow \ell') \right] \left[\frac{(W_{\ell L\ell'}^{x,0} C_{\ell'}^{\Theta E} + p_x W_{\ell'L\ell}^{x,+} C_\ell^{\Theta E}) \widehat{C}_{\ell'}^{\Theta E}}{\widehat{C}_\ell^{\text{EE}}} + p_x(\ell \leftrightarrow \ell') \right] \\ &= \frac{1}{2L+1} \sum_{\ell\ell'} \left[\frac{W_{\ell L\ell'}^{x,0} C_{\ell'}^{\Theta\Theta}}{\widehat{C}_\ell^{\Theta\Theta} \widehat{C}_{\ell'}^{\Theta\Theta}} \frac{(W_{\ell L\ell'}^{x,0} C_{\ell'}^{\Theta E} + p_x W_{\ell'L\ell}^{x,+} C_\ell^{\Theta E}) \widehat{C}_{\ell'}^{\Theta E}}{\widehat{C}_\ell^{\text{EE}}} \right. \\ &\quad \left. + p_x \frac{W_{\ell L\ell'}^{x,0} C_{\ell'}^{\Theta\Theta}}{\widehat{C}_\ell^{\Theta\Theta} \widehat{C}_{\ell'}^{\Theta\Theta}} \frac{(W_{\ell'L\ell}^{x,0} C_\ell^{\Theta E} + p_x W_{\ell L\ell'}^{x,+} C_{\ell'}^{\Theta E}) \widehat{C}_\ell^{\Theta E}}{\widehat{C}_{\ell'}^{\text{EE}}} \right] \\ &= \Sigma_L^{(0),x} \left[\frac{1}{\widehat{C}^{\Theta\Theta}}, \frac{C^{\Theta\Theta} C^{\Theta E} \widehat{C}^{\Theta E}}{\widehat{C}^{\Theta\Theta} \widehat{C}^{\text{EE}}} \right] + p_x \Gamma_L^{(\times),x} \left[\frac{C^{\Theta E}}{\widehat{C}^{\Theta\Theta}}, \frac{C^{\Theta\Theta} \widehat{C}^{\Theta E}}{\widehat{C}^{\Theta\Theta} \widehat{C}^{\text{EE}}} \right] \\ &\quad + p_x \Gamma_L^{(0),x} \left[\frac{C^{\Theta E} \widehat{C}^{\Theta E}}{\widehat{C}^{\Theta\Theta} \widehat{C}^{\text{EE}}}, \frac{C^{\Theta\Theta}}{\widehat{C}^{\Theta\Theta}} \right] + \Sigma_L^{(\times),x} \left[\frac{\widehat{C}^{\Theta E}}{\widehat{C}^{\Theta\Theta} \widehat{C}^{\text{EE}}}, \frac{C^{\Theta E} C^{\Theta\Theta}}{\widehat{C}^{\Theta\Theta}} \right], \end{aligned} \quad (74)$$

where

$$\Sigma_L^{(\times),x}[A, B] = \frac{1}{2L+1} \sum_{\ell\ell'} W_{\ell L\ell'}^{x,0} W_{\ell L\ell'}^{x,+} A_\ell B_{\ell'}. \quad (75)$$

For $\Theta\Theta EE$,

$$\begin{aligned} \frac{A_L^{x,(\Theta\Theta)} A_L^{x,(EE)}}{N_L^{x,(\Theta\Theta EE)}} &= \frac{1}{2L+1} \sum_{\ell\ell'} \left[\frac{W_{\ell L\ell'}^{x,0} C_{\ell'}^{\Theta\Theta}}{2\widehat{C}_\ell^{\Theta\Theta} \widehat{C}_{\ell'}^{\Theta\Theta}} + p_x(\ell \leftrightarrow \ell') \right] \left[\frac{(W_{\ell L\ell'}^{x,+} C_{\ell'}^{\text{EE}} + p_x W_{\ell'L\ell}^{x,+} C_\ell^{\text{EE}}) \widehat{C}_\ell^{\Theta E} \widehat{C}_{\ell'}^{\Theta E}}{2\widehat{C}_\ell^{\text{EE}} \widehat{C}_{\ell'}^{\text{EE}}} + p_x(\ell \leftrightarrow \ell') \right] \\ &= \frac{1}{2L+1} \sum_{\ell\ell'} \frac{W_{\ell L\ell'}^{x,0} C_{\ell'}^{\Theta\Theta}}{\widehat{C}_\ell^{\Theta\Theta} \widehat{C}_{\ell'}^{\Theta\Theta}} \left[\frac{(W_{\ell L\ell'}^{x,+} C_{\ell'}^{\text{EE}} + p_x W_{\ell'L\ell}^{x,+} C_\ell^{\text{EE}}) \widehat{C}_\ell^{\Theta E} \widehat{C}_{\ell'}^{\Theta E}}{2\widehat{C}_\ell^{\text{EE}} \widehat{C}_{\ell'}^{\text{EE}}} + p_x(\ell \leftrightarrow \ell') \right] \\ &= \frac{1}{2L+1} \sum_{\ell\ell'} \frac{W_{\ell L\ell'}^{x,0} C_{\ell'}^{\Theta\Theta}}{\widehat{C}_\ell^{\Theta\Theta} \widehat{C}_{\ell'}^{\Theta\Theta}} \frac{(W_{\ell L\ell'}^{x,+} C_{\ell'}^{\text{EE}} + p_x W_{\ell'L\ell}^{x,+} C_\ell^{\text{EE}}) \widehat{C}_\ell^{\Theta E} \widehat{C}_{\ell'}^{\Theta E}}{\widehat{C}_\ell^{\text{EE}} \widehat{C}_{\ell'}^{\text{EE}}} \\ &= \Sigma_L^{(0),x} \left[\frac{\widehat{C}^{\Theta E}}{\widehat{C}^{\Theta\Theta} \widehat{C}^{\text{EE}}}, \frac{C^{\Theta\Theta} C^{\text{EE}} \widehat{C}^{\Theta E}}{\widehat{C}^{\Theta\Theta} \widehat{C}^{\text{EE}}} \right] + p_x \Gamma_L^{(\times),x} \left[\frac{\widehat{C}^{\Theta E} C^{\text{EE}}}{\widehat{C}^{\Theta\Theta} \widehat{C}^{\text{EE}}}, \frac{C^{\Theta\Theta} \widehat{C}^{\Theta E}}{\widehat{C}^{\Theta\Theta} \widehat{C}^{\text{EE}}} \right]. \end{aligned} \quad (76)$$

For ΘEEE ,

$$\begin{aligned}
\frac{A_L^{x,(\Theta E)} A_L^{x,(EE)}}{N_L^{x,(\Theta EEE)}} &= \frac{1}{2L+1} \sum_{\ell\ell'} \left[\frac{W_{\ell L\ell'}^{x,+} C_{\ell'}^{EE}}{2\hat{C}_{\ell}^{EE} \hat{C}_{\ell'}^{EE}} + p_x(\ell \leftrightarrow \ell') \right] \left[\frac{(W_{\ell L\ell'}^{x,0} C_{\ell'}^{\Theta E} + p_x W_{\ell' L\ell}^{x,+} C_{\ell}^{\Theta E}) \hat{C}_{\ell}^{\Theta E}}{\hat{C}_{\ell}^{\Theta\Theta}} + p_x(\ell \leftrightarrow \ell') \right] \\
&= \frac{1}{2L+1} \sum_{\ell\ell'} \left[\frac{W_{\ell L\ell'}^{x,+} C_{\ell'}^{EE}}{\hat{C}_{\ell}^{EE} \hat{C}_{\ell'}^{EE}} + p_x \frac{W_{\ell' L\ell}^{x,+} C_{\ell}^{EE}}{\hat{C}_{\ell}^{EE} \hat{C}_{\ell'}^{EE}} \right] \left[\frac{(W_{\ell L\ell'}^{x,0} C_{\ell'}^{\Theta E} + p_x W_{\ell' L\ell}^{x,+} C_{\ell}^{\Theta E}) \hat{C}_{\ell}^{\Theta E}}{\hat{C}_{\ell}^{\Theta\Theta}} \right] \\
&= \Sigma_L^{(\times),x} \left[\frac{\hat{C}^{\Theta E}}{\hat{C}^{\Theta\Theta} \hat{C}^{EE}}, \frac{C^{\Theta E} C^{EE}}{\hat{C}^{EE}} \right] + p_x \Gamma_L^{(+),x} \left[\frac{C^{\Theta E} \hat{C}^{\Theta E}}{\hat{C}^{\Theta\Theta} \hat{C}^{EE}}, \frac{C^{EE}}{\hat{C}^{EE}} \right] \\
&\quad + p_x \Gamma_L^{(\times),x} \left[\frac{\hat{C}^{\Theta E} C^{EE}}{\hat{C}^{\Theta\Theta} \hat{C}^{EE}}, \frac{C^{\Theta E}}{\hat{C}^{EE}} \right] + \Sigma_L^{(+),x} \left[\frac{C^{\Theta E} \hat{C}^{\Theta E} C^{EE}}{\hat{C}^{\Theta\Theta} \hat{C}^{EE}}, \frac{1}{\hat{C}^{EE}} \right]. \tag{77}
\end{aligned}$$

For ΘBEB ,

$$\begin{aligned}
\frac{A_L^{x,(\Theta B)} A_L^{x,(EB)}}{N_L^{x,(\Theta BEB)}} &= \frac{1}{2L+1} \sum_{\ell\ell'} \left[\frac{(W_{\ell L\ell'}^{x,-})^* C_{\ell'}^{BB} - p_x (W_{\ell' L\ell}^{x,-})^* C_{\ell}^{EE}}{\hat{C}_{\ell}^{EE} \hat{C}_{\ell'}^{BB}} \right] \left[\frac{-p_x W_{\ell' L\ell}^{x,-} C_{\ell}^{\Theta E} \hat{C}_{\ell}^{\Theta E}}{\hat{C}_{\ell}^{\Theta\Theta}} \right] \\
&= -p_x \Gamma_L^{(-),x} \left[\frac{C^{\Theta E} \hat{C}^{\Theta E}}{\hat{C}^{\Theta\Theta} \hat{C}^{EE}}, \frac{C^{BB}}{\hat{C}^{BB}} \right] + \Sigma_L^{(-),x} \left[\frac{C^{\Theta E} \hat{C}^{\Theta E} C^{EE}}{\hat{C}^{\Theta\Theta} \hat{C}^{EE}}, \frac{1}{\hat{C}^{BB}} \right]. \tag{78}
\end{aligned}$$

5 Explicit Kernel Functions

Here we consider expression for the Kernel functions in terms of the Wigner d-functions. In the following calculations, we frequently use

$$\int_{-1}^1 d\mu d_{s_1, s'_1}^{\ell_1}(\beta) d_{s_2, s'_2}^{\ell_2}(\beta) d_{s_3, s'_3}^{\ell_3}(\beta) = 2 \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ s_1 & s_2 & s_3 \end{pmatrix} \begin{pmatrix} \ell_1 & \ell_2 & \ell_3 \\ s'_1 & s'_2 & s'_3 \end{pmatrix}, \quad (79)$$

with $s_1 + s_2 + s_3 = s'_1 + s'_2 + s'_3 = 0$ and $\mu = \cos \beta$, and the symmetric property:

$$d_{mm'}^{\ell}(\beta) = (-1)^{m-m'} d_{-m, -m'}^{\ell}(\beta) = (-1)^{m-m'} d_{m'm}^{\ell}(\beta) \quad (80)$$

$$d_{mm'}^{\ell}(\beta) = (-1)^{\ell+m} d_{m, -m'}^{\ell}(\pi - \beta). \quad (81)$$

We also define

$$X^{p \dots q} = a_{\ell}^p \dots a_{\ell}^q X_{\ell}. \quad (82)$$

5.1 Kernel Functions: Lensing

We obtain

$$\begin{aligned} \Sigma_L^{(0),x}[A, B] &= \frac{1}{2L+1} \sum_{\ell \ell'} |W_{\ell L \ell'}^{x,0}|^2 A_{\ell} B_{\ell'} \\ &= \sum_{\ell \ell'} 4\pi L(L+1) \frac{2\ell+1}{4\pi} A_{\ell} \frac{2\ell'+1}{4\pi} B_{\ell'} \ell'(\ell'+1) \frac{1+c_x^2(-1)^{\ell+L+\ell'}}{2} \begin{pmatrix} \ell & L & \ell' \\ 0 & 1 & -1 \end{pmatrix}^2 \\ &= \pi L(L+1) \sum_{\ell \ell'} \frac{2\ell+1}{4\pi} A_{\ell} \frac{2\ell'+1}{4\pi} B_{\ell'} 2\ell'(\ell'+1) \left[\begin{pmatrix} \ell & L & \ell' \\ 0 & 1 & -1 \end{pmatrix}^2 + c_x^2 \begin{pmatrix} \ell & L & \ell' \\ 0 & -1 & 1 \end{pmatrix} \begin{pmatrix} \ell & L & \ell' \\ 0 & 1 & -1 \end{pmatrix} \right] \\ &= \int_{-1}^1 d\mu \pi L(L+1) \sum_{\ell \ell'} \frac{2\ell+1}{4\pi} A_{\ell} \frac{2\ell'+1}{4\pi} B_{\ell'} \ell'(\ell'+1) [d_{00}^{\ell} d_{11}^L d_{11}^{\ell'} + c_x^2 d_{00}^{\ell} d_{1,-1}^L d_{1,-1}^{\ell'}] \\ &= \int_{-1}^1 d\mu \pi L(L+1) \{ \xi_{00}[A] \xi_{11}[B^{00}] d_{11}^L + c_x^2 \xi_{00}[A] \xi_{1,-1}[B^{00}] d_{1,-1}^L \}, \end{aligned} \quad (83)$$

where

$$\xi_{mm'}[A] = \sum_{\ell} \frac{2\ell+1}{4\pi} A_{\ell} d_{mm'}^{\ell}. \quad (84)$$

The cross-term is

$$\begin{aligned} \Gamma_L^{(0),x}[A, B] &= \frac{1}{2L+1} \sum_{\ell \ell'} (W_{\ell L \ell'}^{x,0})^* W_{\ell' L \ell}^{x,0} A_{\ell} B_{\ell'} \\ &= \sum_{\ell \ell'} 4\pi L(L+1) \frac{2\ell+1}{4\pi} A_{\ell} \frac{2\ell'+1}{4\pi} B_{\ell'} a_{\ell}^0 a_{\ell'}^0 \frac{1+c_x^2(-1)^{\ell+L+\ell'}}{2} \begin{pmatrix} \ell & L & \ell' \\ 0 & 1 & -1 \end{pmatrix} \begin{pmatrix} \ell' & L & \ell \\ 0 & 1 & -1 \end{pmatrix} \\ &= \pi L(L+1) \sum_{\ell \ell'} \frac{2\ell+1}{4\pi} A_{\ell} \frac{2\ell'+1}{4\pi} B_{\ell'}^0 2 \left[\begin{pmatrix} \ell & L & \ell' \\ 0 & 1 & -1 \end{pmatrix} \begin{pmatrix} \ell & L & \ell' \\ 1 & -1 & 0 \end{pmatrix} + c_x^2 \begin{pmatrix} \ell & L & \ell' \\ 0 & 1 & -1 \end{pmatrix} \begin{pmatrix} \ell & L & \ell' \\ -1 & 1 & 0 \end{pmatrix} \right] \\ &= \int_{-1}^1 d\mu \pi L(L+1) \sum_{\ell \ell'} \frac{2\ell+1}{4\pi} A_{\ell} \frac{2\ell'+1}{4\pi} B_{\ell'}^0 [d_{01}^{\ell} d_{1,-1}^L d_{-1,0}^{\ell'} + c_x^2 d_{0,-1}^{\ell} d_{11}^L d_{-1,0}^{\ell'}] \\ &= - \int_{-1}^1 d\mu \pi L(L+1) \{ \xi_{01}[A^0] \xi_{0,-1}[B^0] d_{1,-1}^L + c_x^2 \xi_{01}[A^0] \xi_{01}[B^0] d_{11}^L \}. \end{aligned} \quad (85)$$

Denoting $p = \pm$ and $x = \phi, \varpi$, the polarization auto kernel is

$$\begin{aligned}
\Sigma_L^{(p),x}[A, B] &= \frac{1}{2L+1} \sum_{\ell\ell'} |W_{\ell L \ell'}^{x,p}|^2 A_\ell B_{\ell'} \\
&= \pi L(L+1) \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} 2[1 + pc_x^2(-1)^{\ell+L+\ell'}] \left[a_{\ell'}^+ \begin{pmatrix} \ell & L & \ell' \\ -2 & -1 & 3 \end{pmatrix} + c_x^2 a_{\ell'}^- \begin{pmatrix} \ell & L & \ell' \\ -2 & 1 & 1 \end{pmatrix} \right]^2 \\
&= \pi L(L+1) \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} [1 + pc_x^2(-1)^{\ell+L+\ell'}] \\
&\quad \times 2 \left[(a_{\ell'}^+)^2 \begin{pmatrix} \ell & L & \ell' \\ -2 & -1 & 3 \end{pmatrix}^2 + (a_{\ell'}^-)^2 \begin{pmatrix} \ell & L & \ell' \\ -2 & 1 & 1 \end{pmatrix}^2 + 2c_x^2 a_{\ell'}^+ a_{\ell'}^- \begin{pmatrix} \ell & L & \ell' \\ -2 & 1 & 1 \end{pmatrix} \begin{pmatrix} \ell & L & \ell' \\ -2 & -1 & 3 \end{pmatrix} \right] \\
&= \pi L(L+1) \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} \\
&\quad \times 2 \left[(a_{\ell'}^+)^2 \begin{pmatrix} \ell & L & \ell' \\ -2 & -1 & 3 \end{pmatrix}^2 + (a_{\ell'}^-)^2 \begin{pmatrix} \ell & L & \ell' \\ -2 & 1 & 1 \end{pmatrix}^2 + 2c_x^2 a_{\ell'}^+ a_{\ell'}^- \begin{pmatrix} \ell & L & \ell' \\ -2 & 1 & 1 \end{pmatrix} \begin{pmatrix} \ell & L & \ell' \\ -2 & -1 & 3 \end{pmatrix} \right. \\
&\quad \left. + pc_x^2 (a_{\ell'}^+)^2 \begin{pmatrix} \ell & L & \ell' \\ -2 & -1 & 3 \end{pmatrix} \begin{pmatrix} \ell & L & \ell' \\ 2 & 1 & -3 \end{pmatrix} + pc_x^2 (a_{\ell'}^-)^2 \begin{pmatrix} \ell & L & \ell' \\ -2 & 1 & 1 \end{pmatrix} \begin{pmatrix} \ell & L & \ell' \\ 2 & -1 & -1 \end{pmatrix} + 2pa_{\ell'}^+ a_{\ell'}^- \begin{pmatrix} \ell & L & \ell' \\ -2 & 1 & 1 \end{pmatrix} \begin{pmatrix} \ell & L & \ell' \\ 2 & 1 & -3 \end{pmatrix} \right] \\
&= \int_{-1}^1 d\mu \pi L(L+1) \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} [(a_{\ell'}^+)^2 d_{22}^\ell d_{11}^L d_{33}^{\ell'} + (a_{\ell'}^-)^2 d_{22}^\ell d_{11}^L d_{11}^{\ell'} \\
&\quad + 2c_x^2 a_{\ell'}^+ a_{\ell'}^- d_{22}^\ell d_{1,-1}^L d_{13}^{\ell'} + pc_x^2 (a_{\ell'}^+)^2 d_{-2,2}^\ell d_{-1,1}^L d_{3,-3}^{\ell'} + pc_x^2 (a_{\ell'}^-)^2 d_{-2,2}^\ell d_{1,-1}^L d_{1,-1}^{\ell'} + 2pa_{\ell'}^+ a_{\ell'}^- d_{-2,2}^\ell d_{11}^L d_{1,-3}^{\ell'}] \\
&= \int_{-1}^1 d\mu \pi L(L+1) [(\xi_{22}[A]\xi_{33}[B^{++}] + \xi_{22}[A]\xi_{11}[B^{--}] + 2p\xi_{2,-2}[A]\xi_{3,-1}[B^{+-}])d_{11}^L \\
&\quad + c_x^2(p\xi_{2,-2}[A]\xi_{3,-3}[B^{++}] + p\xi_{2,-2}[A]\xi_{1,-1}[B^{--}] + 2\xi_{22}[A]\xi_{31}[B^{+-}])d_{1,-1}^L]. \tag{86}
\end{aligned}$$

The polarization cross kernel is

$$\begin{aligned}
\Gamma_L^{(p),x}[A, B] &= \frac{1}{2L+1} \sum_{\ell\ell'} (W_{\ell L \ell'}^{x,p})^* W_{\ell' L \ell}^{x,p} A_\ell B_{\ell'} \\
&= \pi L(L+1) \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} 2[1 + pc_x^2(-1)^{\ell+L+\ell'}] \\
&\quad \times \left[a_{\ell'}^+ \begin{pmatrix} \ell & L & \ell' \\ -2 & -1 & 3 \end{pmatrix} + c_x^2 a_{\ell'}^- \begin{pmatrix} \ell & L & \ell' \\ -2 & 1 & 1 \end{pmatrix} \right] \left[a_{\ell'}^+ \begin{pmatrix} \ell' & L & \ell \\ -2 & -1 & 3 \end{pmatrix} + c_x^2 a_{\ell'}^- \begin{pmatrix} \ell' & L & \ell \\ -2 & 1 & 1 \end{pmatrix} \right] \\
&= \pi L(L+1) \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} 2[(-1)^{\ell+L+\ell'} + pc_x^2] \\
&\quad \times \left[a_{\ell'}^+ \begin{pmatrix} \ell & L & \ell' \\ -2 & -1 & 3 \end{pmatrix} + c_x^2 a_{\ell'}^- \begin{pmatrix} \ell & L & \ell' \\ -2 & 1 & 1 \end{pmatrix} \right] \left[a_{\ell'}^+ \begin{pmatrix} \ell & L & \ell' \\ 3 & -1 & -2 \end{pmatrix} + c_x^2 a_{\ell'}^- \begin{pmatrix} \ell & L & \ell' \\ 1 & 1 & -2 \end{pmatrix} \right] \\
&= \pi L(L+1) \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} \\
&\quad \times 2 \left\{ \left[a_{\ell'}^+ \begin{pmatrix} \ell & L & \ell' \\ 2 & 1 & -3 \end{pmatrix} + c_x^2 a_{\ell'}^- \begin{pmatrix} \ell & L & \ell' \\ 2 & -1 & -1 \end{pmatrix} \right] \left[a_{\ell'}^+ \begin{pmatrix} \ell & L & \ell' \\ 3 & -1 & -2 \end{pmatrix} + c_x^2 a_{\ell'}^- \begin{pmatrix} \ell & L & \ell' \\ 1 & 1 & -2 \end{pmatrix} \right] \right. \\
&\quad \left. + p \left[c_x^2 a_{\ell'}^+ \begin{pmatrix} \ell & L & \ell' \\ -2 & -1 & 3 \end{pmatrix} + a_{\ell'}^- \begin{pmatrix} \ell & L & \ell' \\ -2 & 1 & 1 \end{pmatrix} \right] \left[a_{\ell'}^+ \begin{pmatrix} \ell & L & \ell' \\ 3 & -1 & -2 \end{pmatrix} + c_x^2 a_{\ell'}^- \begin{pmatrix} \ell & L & \ell' \\ 1 & 1 & -2 \end{pmatrix} \right] \right\} \\
&= \int_{-1}^1 d\mu \pi L(L+1) \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} \\
&\quad \times [a_{\ell'}^+ a_{\ell'}^+ d_{23}^L d_{1,-1}^L d_{-3,-2}^{\ell'} + c_x^2 a_{\ell'}^+ a_{\ell'}^- d_{21}^L d_{11}^L d_{-3,-2}^{\ell'} + c_x^2 a_{\ell'}^- a_{\ell'}^+ d_{23}^L d_{11}^L d_{-1,-2}^{\ell'} + a_{\ell'}^- a_{\ell'}^- d_{21}^L d_{1,-1}^L d_{-1,-2}^{\ell'} \\
&\quad + p(c_x^2 a_{\ell'}^+ a_{\ell'}^+ d_{-2,3}^L d_{11}^L d_{3,-2}^{\ell'} + a_{\ell'}^+ a_{\ell'}^- d_{-2,1}^L d_{1,-1}^L d_{3,-2}^{\ell'} + a_{\ell'}^- a_{\ell'}^+ d_{-2,3}^L d_{1,-1}^L d_{1,-2}^{\ell'} + c_x^2 a_{\ell'}^- a_{\ell'}^- d_{-2,1}^L d_{11}^L d_{1,-2}^{\ell'})] \\
&= \int_{-1}^1 d\mu \pi L(L+1) [-c_x^2 (\xi_{21}[A^-] \xi_{32}[B^+] + \xi_{32}[A^+] \xi_{21}[B^-] + p \xi_{3,-2}[A^+] \xi_{3,-2}[B^+] + p \xi_{2,-1}[A^-] \xi_{2,-1}[B^-]) d_{11}^L \\
&\quad + (\xi_{32}[A^+] \xi_{32}[B^+] + \xi_{21}[A^-] \xi_{21}[B^-] - p \xi_{2,-1}[A^-] \xi_{3,-2}[B^+] - p \xi_{3,-2}[A^+] \xi_{2,-1}[B^-]) d_{1,-1}^L]. \quad (87)
\end{aligned}$$

The temperature-polarization kernel is

$$\begin{aligned}
\Sigma_L^{(\times),x}[A, B] &= \frac{1}{2L+1} \sum_{\ell\ell'} (W_{\ell L \ell'}^{x,0})^* W_{\ell' L \ell}^{x,+} A_\ell B_{\ell'} \\
&= \pi L(L+1) \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} a_{\ell'}^0 2[1 + c_x^2(-1)^{\ell+L+\ell'}] \\
&\quad \times \begin{pmatrix} \ell & L & \ell' \\ 0 & 1 & -1 \end{pmatrix} \left[a_{\ell'}^+ \begin{pmatrix} \ell & L & \ell' \\ -2 & -1 & 3 \end{pmatrix} + c_x^2 a_{\ell'}^- \begin{pmatrix} \ell & L & \ell' \\ -2 & 1 & 1 \end{pmatrix} \right] \\
&= \pi L(L+1) \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} a_{\ell'}^0 \\
&\quad \times 2 \left[\begin{pmatrix} \ell & L & \ell' \\ 0 & 1 & -1 \end{pmatrix} + c_x^2 \begin{pmatrix} \ell & L & \ell' \\ 0 & -1 & 1 \end{pmatrix} \right] \left[a_{\ell'}^+ \begin{pmatrix} \ell & L & \ell' \\ -2 & -1 & 3 \end{pmatrix} + c_x^2 a_{\ell'}^- \begin{pmatrix} \ell & L & \ell' \\ -2 & 1 & 1 \end{pmatrix} \right] \\
&= \int_{-1}^1 d\mu \pi L(L+1) \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} a_{\ell'}^0 \\
&\quad \times \left[a_{\ell'}^+ d_{0,-2}^L d_{1,-1}^L d_{-1,3}^{\ell'} + c_x^2 a_{\ell'}^- d_{0,-2}^L d_{11}^L d_{-1,1}^{\ell'} + c_x^2 a_{\ell'}^+ d_{0,-2}^L d_{11}^L d_{13}^{\ell'} + a_{\ell'}^- d_{0,-2}^L d_{-1,1}^L d_{11}^{\ell'} \right] \\
&= \int_{-1}^1 d\mu \pi L(L+1) \{ c_x^2 (\xi_{20}[A] \xi_{1,-1}[B^{0-}] + \xi_{20}[A] \xi_{31}[B^{0+}]) d_{11}^L \\
&\quad + (\xi_{20}[A] \xi_{3,-1}[B^{0+}] + \xi_{20}[A] \xi_{11}[B^{0-}]) d_{1,-1}^L \}, \quad (88)
\end{aligned}$$

and

$$\begin{aligned}
\Gamma_L^{(\times),x}[A,B] &= \frac{1}{2L+1} \sum_{\ell\ell'} (W_{\ell L \ell'}^{x,0})^* W_{\ell' L \ell}^{x,+} A_\ell B_{\ell'} \\
&= \pi L(L+1) \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} a_{\ell'}^0 2[1+c_x^2(-1)^{\ell+L+\ell'}] \\
&\quad \times \begin{pmatrix} \ell & L & \ell' \\ 0 & 1 & -1 \end{pmatrix} \left[a_\ell^+ \begin{pmatrix} \ell' & L & \ell \\ -2 & -1 & 3 \end{pmatrix} + c_x^2 a_\ell^- \begin{pmatrix} \ell' & L & \ell \\ -2 & 1 & 1 \end{pmatrix} \right] \\
&= \pi L(L+1) \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} a_{\ell'}^0 \\
&\quad \times 2 \left[\begin{pmatrix} \ell & L & \ell' \\ 0 & 1 & -1 \end{pmatrix} + c_x^2 \begin{pmatrix} \ell & L & \ell' \\ 0 & -1 & 1 \end{pmatrix} \right] \left[a_\ell^+ \begin{pmatrix} \ell & L & \ell' \\ -3 & 1 & 2 \end{pmatrix} + c_x^2 a_\ell^- \begin{pmatrix} \ell & L & \ell' \\ -1 & -1 & 2 \end{pmatrix} \right] \\
&= \int_{-1}^1 d\mu \pi L(L+1) \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} a_{\ell'}^0 \\
&\quad \times \left[a_\ell^+ d_{0,-3}^\ell d_{11}^L d_{-1,2}^{\ell'} + c_x^2 a_\ell^- d_{0,-1}^\ell d_{1,-1}^L d_{-1,2}^{\ell'} + c_x^2 a_\ell^+ d_{0,-3}^\ell d_{1,-1}^L d_{12}^{\ell'} + a_\ell^- d_{0,-1}^\ell d_{11}^L d_{12}^{\ell'} \right] \\
&= \int_{-1}^1 d\mu \pi L(L+1) \{ -(\xi_{30}[A^+] \xi_{2,-1}[B^0] + \xi_{10}[A^-] \xi_{12}[B^0]) d_{11}^L \\
&\quad - c_x^2 (\xi_{10}[A^-] \xi_{2,-1}[B^0] + \xi_{30}[A^0] \xi_{21}[B^-]) d_{1,-1}^L \} .
\end{aligned} \tag{89}$$

5.2 Kernel Functions: Rotation

Next we consider the kernel functions for $x = \alpha$. If $p = -$ and $x = \alpha$,

$$\begin{aligned}
\Sigma_L^{(-),\alpha}[A,B] &= \pi \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} 8[1+(-1)^{\ell+L+\ell'}] \begin{pmatrix} \ell & L & \ell' \\ -2 & 0 & 2 \end{pmatrix}^2 \\
&= \pi L(L+1) \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} 8 \left[\begin{pmatrix} \ell & L & \ell' \\ -2 & 0 & 2 \end{pmatrix}^2 + \begin{pmatrix} \ell & L & \ell' \\ -2 & 0 & 2 \end{pmatrix} \begin{pmatrix} \ell & L & \ell' \\ 2 & 0 & -2 \end{pmatrix} \right] \\
&= \int_{-1}^1 d\mu \pi L(L+1) \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} 4(d_{-2,-2}^\ell d_{00}^L d_{22}^{\ell'} + d_{-2,2}^\ell d_{00}^L d_{2,-2}^{\ell'}) \\
&= \int_{-1}^1 d\mu 4\pi (\xi_{-2,-2}[A] \xi_{22}[B] + \xi_{-2,22}[A] \xi_{22}[B]) d_{00}^L ,
\end{aligned} \tag{90}$$

and

$$\begin{aligned}
\Gamma_L^{(-),\alpha}[A,B] &= \pi \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} 8[1+(-1)^{\ell+L+\ell'}] \begin{pmatrix} \ell & L & \ell' \\ -2 & 0 & 2 \end{pmatrix} \begin{pmatrix} \ell' & L & \ell \\ -2 & 0 & 2 \end{pmatrix} \\
&= \pi \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} \\
&\quad \times 8 \begin{pmatrix} \ell & L & \ell' \\ -2 & 0 & 2 \end{pmatrix} \left[\begin{pmatrix} \ell & L & \ell' \\ -2 & 0 & 2 \end{pmatrix} + \begin{pmatrix} \ell & L & \ell' \\ 2 & 0 & -2 \end{pmatrix} \right] \\
&= \int_{-1}^1 d\mu \pi \sum_{\ell\ell'} \frac{2\ell+1}{4\pi} A_\ell \frac{2\ell'+1}{4\pi} B_{\ell'} 4[d_{-2,-2}^\ell d_{00}^L d_{22}^{\ell'} + d_{-2,2}^\ell d_{00}^L d_{2,-2}^{\ell'}] \\
&= \int_{-1}^1 d\mu 4\pi (\xi_{-2,-2}[A] \xi_{22}[B] + \xi_{-2,2}[A] \xi_{2,-2}[B]) d_{00}^L .
\end{aligned} \tag{91}$$

References

- [1] C. Dvorkin, W. Hu, and K. M. Smith, “*B-mode CMB Polarization from Patchy Screening during Reionization*”, *Phys. Rev. D* **79** (2009) 107302, [arXiv:0902.4413].
- [2] K. M. Smith *et al.*, “*Delensing CMB Polarization with External Datasets*”, *J. Cosmol. Astropart. Phys.* **1206** (2012) 014, [arXiv:1010.0048].
- [3] V. Gluscevic, M. Kamionkowski, and A. Cooray, “*Derotation of the cosmic microwave background polarization: Full-sky formalism*”, *Phys. Rev. D* **80** (2009) 023510.
- [4] V. Gluscevic, M. Kamionkowski, and D. Hanson, “*Patchy Screening of the Cosmic Microwave Background by Inhomogeneous Reionization*”, *Phys. Rev. D* **87** (2013) 047303, [arXiv:1210.5507].
- [5] D. Varshalovich, A. Moskalev, and V. Kersonskii. World Scientific, Singapore.
- [6] T. Namikawa, D. Yamauchi, and A. Taruya, “*Full-sky lensing reconstruction of gradient and curl modes from CMB maps*”, *J. Cosmol. Astropart. Phys.* **1201** (2012) 007, [arXiv:1110.1718].
- [7] T. Okamoto and W. Hu, “*CMB Lensing Reconstruction on the Full Sky*”, *Phys. Rev. D* **67** (2003) 083002, [astro-ph/0301031].