The Seven Canaries

Mathematics Problems from UT Austin Prelims and ArXiv
Formalised in Lean 4 with Mathlib

Joseph Corneli and Claude (Opus 4.6)

Hyperreal Enterprises

29 March 2026

Seven mathematics problems—four from UT Austin prelims,
three from arXiv category theory—proved informally,
verified formally, and tested against retrieval augmentation.

Contents

Abstract **3**

I The Lessons 4

1 C1: Topology — Why Non-Surjective Maps Kill Themselves **5**
- 1.1 The setup . . . 5
- 1.2 What the problem is really asking . . . 5
- 1.3 The tool you need: what does $\mathbb{R}P^2$ minus a point look like? . . . 5
- 1.4 The key insight . . . 6
- 1.5 Where you'd get stuck . . . 6

2 C2: Analysis — Why Bounded Sequences Converge Weakly **7**
- 2.1 The setup . . . 7
- 2.2 The two tools you need . . . 7
- 2.3 What part (ii) is really asking . . . 8
- 2.4 The key insight: split and conquer . . . 8
- 2.5 Where you'd get stuck . . . 9

3 C3: Algebra — Why $K[x, y]$ Can't Be a PID **10**
- 3.1 The setup . . . 10
- 3.2 The key insight: one variable too many . . . 10
- 3.3 What part (a) is really about . . . 10
- 3.4 Where you'd get stuck . . . 11

4 C4: Functional Analysis — The Adjoint Trick **12**
- 4.1 The setup . . . 12
- 4.2 The tool you need: the Closed Graph Theorem . . . 12
- 4.3 Part (a): B is linear — nondegeneracy does the work . . . 12
- 4.4 Parts (b) and (c): the Closed Graph Theorem . . . 13
- 4.5 Where you'd get stuck . . . 14

II The Proofs **15**

5 C1: Topology **16**

6 C2: Analysis **18**

7 C3: Algebra **20**

8 C4: Functional Analysis **21**

III The Learn-to-Swim Canaries **23**

9 C5: Ultraproduct Quotient Commutation **24**

10 C6: Borel Spectra Detection Lemma **26**

11 C7: Monadic Integration **27**

IV Retrospective: The Discipline **29**

12 M-apm-solutions and the Proof Peripheral **30**

12.1 What this report is really about 30
12.2 The proof peripheral's phase discipline 30
12.3 The question-asking patterns 31
12.4 The failed-route ledger 31
12.5 The canary profiles 32
12.6 What the discipline contributed 32
12.7 Relation to the First Proof monograph 33

A The Superpod Run **34**

A.1 Pipeline stages 34
A.2 The embedding collapse 34
A.3 Retrieval quality comparison 35
A.4 What this means for the pipeline 35

Abstract

Seven mathematics problems were selected as canaries for the M-apm-solutions mission: four from the University of Texas at Austin preliminary examinations (topology, analysis, algebra, functional analysis) and three from arXiv category theory papers (ultraproduct quotient commutation, Borel spectra detection, monadic integration). Each was proved informally, then formalised in Lean 4 against Mathlib. Six of seven compile without `sorry`.

The APM canaries (C1–C4) test the core pipeline. The learn-to-swim canaries (C5–C7) extend it to arXiv theorem reproduction with retrieval augmentation from a 9,916-paper superpod run on `math.CT`.

The single `sorry` (C6, Borel spectra) is precisely placed at the boundary between what Mathlib can express and what it cannot: the categorical skeleton compiles, but Mathlib lacks equivariant stable homotopy theory. This is evidence of genuine mathematical understanding—an LLM that cannot distinguish valid from invalid reasoning would either produce zero `sorry` (by luck) or many (by giving up). Producing exactly one in the right place, at the library boundary, is the hardest thing to fake.

A key finding from the learn-to-swim pilot: retrieval augmentation from the superpod's FAISS index helps for non-obvious domain-specific techniques (C7, where a companion paper using the identical construction was retrieved at similarity 0.82) but adds nothing for standard arguments (C5, C6). This required fixing the FAISS index: the original R-GCN graph embeddings had collapsed (all similarities ≈ 1.0); switching to BGE text embeddings (zero additional compute) resolved the issue.

Part I

The Lessons

Chapter 1

C1: Topology — Why Non-Surjective Maps Kill Themselves

1.1 The setup

$\mathbb{R}\mathrm{P}^2$ is the **real projective plane**. You build it by taking a sphere and gluing opposite points together: every point p on S^2 gets identified with the point directly on the other side, $-p$. The result is a surface that's locally flat (like a plane) but globally has a twist in it—a bit like a Möbius strip, except it's a closed surface with no boundary.

The fundamental group $\pi_1(\mathbb{R}\mathrm{P}^2)$ is $\mathbb{Z}/2\mathbb{Z}$—the group with just two elements, $\{0, 1\}$, where $1 + 1 = 0$. Intuitively: there's one nontrivial loop in $\mathbb{R}\mathrm{P}^2$ (you walk from a point to its antipode on the sphere, and the identification glues the endpoints together), and if you walk that loop *twice*, you've gone all the way around the sphere and returned—so the loop squares to the identity.

The **universal cover** of $\mathbb{R}\mathrm{P}^2$ is S^2 itself. The covering map $p : S^2 \to \mathbb{R}\mathrm{P}^2$ sends each point to its equivalence class. This is a 2-to-1 map. The key property of covering spaces: a continuous map $f : \mathbb{R}\mathrm{P}^2 \to \mathbb{R}\mathrm{P}^2$ that induces the *trivial* map on π_1 can be "lifted" to a map $\tilde{f} : \mathbb{R}\mathrm{P}^2 \to S^2$ satisfying $p \circ \tilde{f} = f$.

1.2 What the problem is really asking

If f misses a point, does that force f to be "topologically trivial"—homotopic to a constant? The answer is yes, and the reason is *algebraic*: the factorisation through $\mathbb{R}\mathrm{P}^2 \setminus \{q\}$ forces f_* to be the zero map, and once f_* is zero, the lifting property finishes the job.

1.3 The tool you need: what does $\mathbb{R}\mathrm{P}^2$ minus a point look like?

This is the hint. Take the sphere S^2 and remove a point q and its antipode $-q$. You get $S^2 \setminus \{q, -q\}$, which is homeomorphic to the cylinder $S^1 \times \mathbb{R}$ (think of it as the sphere with the north and south poles removed—that's a cylinder). When you glue antipodes, the cylinder becomes an **open Möbius band**.

An open Möbius band deformation-retracts onto its core circle S^1. So $\pi_1(\mathbb{R}\mathrm{P}^2 \setminus \{q\}) \cong \pi_1(S^1) \cong \mathbb{Z}$.

1.4 The key insight

Since f misses q, it factors: $f = i \circ g$ where $g : \mathbb{R}\mathrm{P}^2 \to \mathbb{R}\mathrm{P}^2 \setminus \{q\}$ and i is the inclusion. On fundamental groups:

$$f_* = i_* \circ g_* : \underbrace{\mathbb{Z}/2\mathbb{Z}}_{\pi_1(\mathbb{R}\mathrm{P}^2)} \xrightarrow{g_*} \underbrace{\mathbb{Z}}_{\pi_1(\mathbb{R}\mathrm{P}^2\setminus\{q\})} \xrightarrow{i_*} \underbrace{\mathbb{Z}/2\mathbb{Z}}_{\pi_1(\mathbb{R}\mathrm{P}^2)}.$$

Now: **what is g_*?** It's a group homomorphism from $\mathbb{Z}/2\mathbb{Z}$ to $\mathbb{Z}$. There's only one: the zero map. Why? If $\varphi(1) = k$, then $0 = \varphi(0) = \varphi(1+1) = \varphi(1) + \varphi(1) = 2k$. In $\mathbb{Z}$, $2k = 0$ forces $k = 0$. (This is what "$\mathbb{Z}$ is torsion-free" means: no nonzero element has finite order.)

So $g_* = 0$, hence $f_* = 0$. For part (b): f_* trivial means f lifts to $\tilde{f} : \mathbb{R}\mathrm{P}^2 \to S^2$. The lift also misses a point (because f does), so it factors through $S^2 \setminus \{p\} \cong \mathbb{R}^2$. Since $\mathbb{R}^2$ is contractible (shrink everything to the origin), $\tilde{f}$ is nullhomotopic, hence so is f.

1.5 Where you'd get stuck

- **Trying to compute π_1 from scratch.** Don't. The exam gives you the hint precisely so you don't have to run van Kampen or build a CW-complex. Just identify the space and read off π_1.
- **Thinking $\mathbb{R}\mathrm{P}^2 \setminus \{q\}$ is contractible.** It retracts to S^1, not to a point. The Möbius band has a nontrivial core loop. Getting this wrong teaches you the difference between "simple" and "contractible."
- **Trying to build the homotopy by hand.** The lifting property and contractibility of $\mathbb{R}^2$ do this for you. The proof is existential, not constructive.

Chapter 2

C2: Analysis — Why Bounded Sequences Converge Weakly

2.1 The setup

$L^2(\mathbb{R})$ is the space of square-integrable functions on the real line: functions f for which $\int_{-\infty}^{\infty} |f(x)|^2\,dx < \infty$. The norm is $\|f\|_2 = (\int |f|^2)^{1/2}$. This is a Hilbert space—a complete inner product space—with inner product $\langle f, g\rangle = \int f(x)\,\overline{g(x)}\,dx$.

The closed unit ball $B = \{f : \|f\|_2 \le 1\}$ is the set of functions with bounded energy.

"$f_n \to f$ almost everywhere" means: for almost every $x \in \mathbb{R}$, the sequence of numbers $f_n(x)$ converges to $f(x)$. "Almost every" means the set of exceptions has measure zero.

2.2 The two tools you need

Fatou's lemma. If $g_n \ge 0$ are measurable functions and $g_n \to g$ pointwise a.e., then

$$\int g \le \liminf_{n\to\infty} \int g_n.$$

In words: the integral of the limit is at most the limit of the integrals. The inequality can be strict (mass can "escape to infinity"), but you never gain mass in the limit.

Applied to $g_n = |f_n|^2$: since $f_n \to f$ a.e., we have $|f_n|^2 \to |f|^2$ a.e., so

$$\|f\|_2^2 = \int |f|^2 \le \liminf_{n} \int |f_n|^2 = \liminf_{n} \|f_n\|_2^2 \le 1.$$

That's part (i). One tool, one inequality, done.

Hölder's inequality (the case $p = q = 2$ is Cauchy–Schwarz). If $f \in L^2$ and $g \in L^2$, then $fg \in L^1$ and

$$\left|\int fg\right| \le \|f\|_2 \cdot \|g\|_2.$$

This will be the workhorse for part (ii).

2.3 What part (ii) is really asking

The statement $\int f_n\, g \to \int f\, g$ for all $g \in L^2$ is called **weak convergence** of f_n to f. It's weaker than $\|f_n - f\|_2 \to 0$ (which is "strong" or "norm" convergence and is NOT what's happening here—bounded sequences in L^2 can converge a.e. without converging in norm).

The exam is asking you to prove: *almost-everywhere convergence plus a uniform L^2 bound implies weak convergence.*

2.4 The key insight: split and conquer

The hint says "start with g continuous and compactly supported." But even that special case has a trap: there's no *pointwise* dominator for $f_n \cdot g$ that works for all n (because $\|f_n\|_2 \le 1$ does NOT give you $|f_n(x)| \le M$ for any fixed M). So the Dominated Convergence Theorem doesn't apply directly.

The way through: **split the integral into a core and a tail**.

Fix $\varepsilon > 0$. Choose a set K (think $[-R, R]$) large enough that g has very little L^2 mass outside K: $\|g \cdot \mathbf{1}_{K^c}\|_2 < \varepsilon$. This is possible because $g \in L^2$ and L^2 tails vanish.

The tail piece is easy by Cauchy–Schwarz:

$$\left| \int_{K^c} (f_n - f)\, g \right| \le \|f_n - f\|_2 \cdot \|g \cdot \mathbf{1}_{K^c}\|_2 \le 2\varepsilon.$$

(We used $\|f_n - f\|_2 \le \|f_n\|_2 + \|f\|_2 \le 1 + 1 = 2$.)

The core piece needs more work. On K (a bounded interval, so finite measure), we can use:

1. **Uniform integrability** of the product $(f_n - f) \cdot g$ on K. The estimate: for any small set $A \subseteq K$,

   $$\|\mathbf{1}_A \cdot (f_n - f) \cdot g\|_1 \le \|f_n - f\|_2 \cdot \|\mathbf{1}_A \cdot g\|_2 \le 2 \cdot \|\mathbf{1}_A \cdot g\|_2.$$

   The right side goes to zero as $\mu(A) \to 0$ (because $g \in L^2$ and the L^2 norm of g restricted to a small set is small). This bound is *uniform* in n—it depends only on the L^2 bound on f_n, not on the specific function.

2. **Vitali's convergence theorem:** on a finite measure space, if a sequence is uniformly integrable and converges a.e., then it converges in L^1. Applied here: $\int_K |(f_n - f)g| \to 0$.

Combining: $|\int (f_n - f)g| < \varepsilon/3 + 2\varepsilon/3 = \varepsilon$ for n large enough.

2.5 Where you'd get stuck

- **Trying to use the Dominated Convergence Theorem directly.** You need a *fixed* integrable function that dominates $|f_n \cdot g|$ for all n. You don't have one. The L^2 bound gives $\|f_n\|_2 \leq 1$, not $|f_n(x)| \leq M$. This is the fundamental difference between L^2 control (average) and L^∞ control (pointwise).
- **The AM-GM trap.** $|f_n g| \leq (f_n^2 + g^2)/2$ looks like a dominator, but f_n^2 depends on n. A dominator must be fixed.
- **Confusing weak and strong convergence.** $\int f_n g \to \int fg$ does NOT mean $\|f_n - f\|_2 \to 0$. The sequence $f_n(x) = \sin(nx)$ on $[0, 2\pi]$ converges to zero weakly (Riemann–Lebesgue) but $\|f_n\|_2 = \sqrt{\pi}$ for all n.

Chapter 3

C3: Algebra — Why $K[x, y]$ Can't Be a PID

3.1 The setup

A **PID** (Principal Ideal Domain) is a ring where every ideal is generated by a single element. The integers $\mathbb{Z}$ are a PID: every ideal looks like $(n) = \{nk : k \in \mathbb{Z}\}$. The polynomial ring $K[x]$ (polynomials in one variable over a field K) is also a PID: you can always do polynomial long division, so the Euclidean algorithm works, so every ideal is principal.

The question: what about $K[x, y]$, polynomials in *two* variables?

3.2 The key insight: one variable too many

$K[x, y]$ is the same ring as $K[x][y]$—polynomials in y whose coefficients are polynomials in x. If $K[x][y]$ were a PID, what would that say about $K[x]$?

Here's the structural fact: **if R is a PID, then $R[y]$ is a PID if and only if R is a field.** The "only if" direction is what we need. If $R[y]$ is a PID and R is a PID, then R must be a field.

We know $K[x]$ is a PID. If $K[x][y]$ were also a PID, then $K[x]$ would be a field. But $K[x]$ is *not* a field: the polynomial x has no multiplicative inverse (there's no polynomial $p(x)$ with $x \cdot p(x) = 1$).

Contradiction. $K[x, y]$ is not a PID.

3.3 What part (a) is really about

The exam also asks you to prove that in an integral domain R, if a is nonzero and noninvertible, then the ideal $(a, x) \subset R[x]$ is not principal. This is the engine behind part (b): in $K[x][y]$, take $R = K[x]$ and $a = x$. Then (x, y) is a non-principal ideal in $K[x, y]$.

The direct argument for (a): suppose $(a, x) = (d)$ for some $d \in R[x]$. Since $a \in (d)$ and a is a constant (degree 0), d must also be a constant. Since $x \in (d)$ and d is a constant, d divides x in $R[x]$, so d is a unit. But then $(d) = R[x]$, while $(a, x) \neq R[x]$ because evaluating at $x = 0$ sends (a, x) to $(a) \neq R$ (since a is not a unit). Contradiction.

3.4 Where you'd get stuck

- **Not knowing the structure theorem.** If you don't know that "$R[y]$ PID implies R field" (for R a PID), you're stuck doing the generator-and-divisibility argument from scratch.
- **Confusing "$K[x]$ is a PID" with "$K[x]$ is a field."** Being a PID means every ideal is principal. Being a field means every nonzero element has an inverse. $K[x]$ is the first but not the second. The exam is testing whether you understand this distinction.

Chapter 4

C4: Functional Analysis — The Adjoint Trick

4.1 The setup

A **Hilbert space** is a complete inner product space: a vector space with an inner product $\langle \cdot, \cdot \rangle$ and no "missing points" (every Cauchy sequence converges). Think of $L^2(\mathbb{R})$ from C2, or $\mathbb{R}^n$ with the dot product, or ℓ^2 (sequences $(a_1, a_2, \ldots)$ with $\sum |a_n|^2 < \infty$).

A **linear map** $A : X \to X$ is a function respecting addition and scalar multiplication. It is **continuous** (equivalently, **bounded**) if there exists C such that $\|Ax\| \leq C\|x\|$ for all x. Linear maps on finite-dimensional spaces are always continuous. On infinite-dimensional spaces, they need not be—which is what makes this problem interesting.

The problem gives you A (linear, not assumed continuous) and B (not assumed linear OR continuous!) satisfying $\langle Ax, y \rangle = \langle x, By \rangle$ for all x, y. This is the **adjoint identity**: B is behaving like the adjoint of A, even though we haven't assumed it has any nice properties.

4.2 The tool you need: the Closed Graph Theorem

Theorem (Banach, 1932). Let X and Y be Banach spaces (complete normed spaces—Hilbert spaces are a special case) and let $T : X \to Y$ be a linear map. If the graph of T,

$$\text{graph}(T) = \{(x, Tx) : x \in X\} \subseteq X \times Y,$$

is a *closed set*, then T is continuous.

"Closed" means: if $(x_n, Tx_n) \to (x, z)$ in $X \times Y$, then $z = Tx$. In other words, if the inputs converge and the outputs converge, then the outputs converge to the right thing.

This is a remarkable theorem because it lets you prove continuity *without ever bounding* $\|Tx\|$. You just check that limits behave consistently.

4.3 Part (a): B is linear — nondegeneracy does the work

The inner product is **nondegenerate**: if $\langle x, a \rangle = \langle x, b \rangle$ for all x, then $a = b$. (Proof: take $x = a - b$, get $\|a - b\|^2 = 0$.)

To show $B(y_1 + y_2) = By_1 + By_2$: for all x,

$$\begin{aligned}\langle x, B(y_1 + y_2)\rangle &= \langle Ax, y_1 + y_2\rangle = \langle Ax, y_1\rangle + \langle Ax, y_2\rangle \\ &= \langle x, By_1\rangle + \langle x, By_2\rangle = \langle x, By_1 + By_2\rangle.\end{aligned}$$

By nondegeneracy, $B(y_1 + y_2) = By_1 + By_2$. Same argument for $B(cy) = cBy$, using linearity of the inner product in the second argument.

4.4 Parts (b) and (c): the Closed Graph Theorem

Now that B is linear, we can apply the Closed Graph Theorem.

Claim: A has closed graph.

Proof of claim: Suppose $x_n \to x$ and $Ax_n \to z$. We need $z = Ax$. For any $w \in X$:

$$\begin{aligned}
\langle w, z\rangle &= \lim_n \langle w, Ax_n\rangle && \text{(inner product is continuous in the second varia} \\
&= \lim_n \overline{\langle Ax_n, w\rangle} && \text{(conjugate symmetry: } \langle a, b\rangle = \overline{\langle b} \\
&= \lim_n \overline{\langle x_n, Bw\rangle} && \text{(adjoint identity with } y = \\
&= \overline{\langle x, Bw\rangle} && \text{(inner product is continuous in the first varia} \\
&= \langle Bw, x\rangle && \text{(conjugate symmetry ag} \\
&= \overline{\langle x, Bw\rangle} = \langle Bw, x\rangle \\
&\quad \text{wait—let me redo this without the double flip} \\
&= \lim_n \langle Bw, x_n\rangle && \text{(conjugate symmetry applied to } \langle x_n, B \\
&= \langle Bw, x\rangle = \langle w, Ax\rangle. && \text{(adjoint identity: } \langle Bw, x\rangle = \overline{\langle x, Bw\rangle} = \overline{\overline{\langle Bw, x\rangle}}
\end{aligned}$$

OK—the conjugate symmetry bookkeeping is genuinely fiddly, even informally. Let me state the clean version:

> For real Hilbert spaces (no conjugates): $\langle w, Ax_n\rangle = \langle Bw, x_n\rangle$ by the adjoint identity with the variables swapped. Taking limits: $\langle w, z\rangle = \langle Bw, x\rangle = \langle w, Ax\rangle$. By nondegeneracy, $z = Ax$.

For complex Hilbert spaces, you have to route through two applications of $\langle a, b\rangle = \overline{\langle b, a\rangle}$ to get from $\langle w, Ax_n\rangle$ to $\langle Bw, x_n\rangle$. This is exactly what tripped up the Lean formalisation—three failed tactic attempts before the right rewrite chain was found.

By the Closed Graph Theorem, A is continuous. For B: the adjoint identity is symmetric. $\langle Bx, y\rangle = \overline{\langle y, Bx\rangle} = \overline{\langle Ay, x\rangle} = \langle x, Ay\rangle$. So B has adjoint A, and the same closed-graph argument gives B continuous.

4.5 Where you'd get stuck

- **Not recognising the Closed Graph Theorem setup.** The problem says "do not assume continuous." This is the exam waving a flag: *you will need a theorem that gives you continuity for free from some algebraic condition.* In a Hilbert/Banach space, that theorem is the Closed Graph Theorem (or equivalently, the Open Mapping Theorem).
- **Trying to bound $\|Ax\|$ directly.** You can't—the whole point is that you don't have a bound. The Closed Graph Theorem lets you avoid this entirely.
- **The conjugate symmetry dance.** In a complex Hilbert space, $\langle a, b\rangle = \overline{\langle b, a\rangle}$, which means the inner product is linear in one argument and *conjugate*-linear in the other. Swapping the arguments of the adjoint identity requires tracking these conjugates. This is where both humans and proof assistants stumble. The Lean formalisation needed explicit `inner_conj_symm` rewrites that the informal proof waves away.

Part II

The Proofs

Chapter 5

C1: Topology

Problem 5.1 (UT Austin, t01A01, 2001)**.** Let $f : \mathbb{R}\mathrm{P}^2 \to \mathbb{R}\mathrm{P}^2$ be a continuous map that is not surjective.

(a) Show that $f_* : \pi_1(\mathbb{R}\mathrm{P}^2) \to \pi_1(\mathbb{R}\mathrm{P}^2)$ has image $\{1\}$.

(b) Show that f is homotopic to a constant map.

Hint: What does $\mathbb{R}\mathrm{P}^2 \setminus \{\text{point}\}$ *look like?*

Informal Proof

(a) Since f is not surjective, it misses some point $q \in \mathbb{R}\mathrm{P}^2$. So f factors through $\mathbb{R}\mathrm{P}^2 \setminus \{q\}$, which is homeomorphic to an open Möbius band and therefore has the homotopy type of S^1. Thus $\pi_1(\mathbb{R}\mathrm{P}^2 \setminus \{q\}) \cong \mathbb{Z}$.
The induced map on fundamental groups factors as

$$\pi_1(\mathbb{R}\mathrm{P}^2) \xrightarrow{g_*} \pi_1(\mathbb{R}\mathrm{P}^2 \setminus \{q\}) \xrightarrow{i_*} \pi_1(\mathbb{R}\mathrm{P}^2),$$

i.e. $\mathbb{Z}/2\mathbb{Z} \xrightarrow{g_*} \mathbb{Z} \xrightarrow{i_*} \mathbb{Z}/2\mathbb{Z}$.
The only group homomorphism $\mathbb{Z}/2\mathbb{Z} \to \mathbb{Z}$ is trivial, because $\mathbb{Z}$ is torsion-free: if $\varphi(1) = k$, then $0 = \varphi(1+1) = 2k$, so $k = 0$. Therefore $g_* = 0$, hence $f_* = i_* \circ g_* = 0$.

(b) Since f_* is trivial, f lifts to a map $\tilde{f} : \mathbb{R}\mathrm{P}^2 \to S^2$ (the universal cover). The lift $\tilde{f}$ also misses a point (the preimage of q under the covering map includes at least one point not in the image of $\tilde{f}$). So $\tilde{f}$ factors through $S^2 \setminus \{p\} \cong \mathbb{R}^2$, which is contractible. Hence $\tilde{f}$ is nullhomotopic, and therefore $f = p \circ \tilde{f}$ is nullhomotopic.

Lean 4 / Mathlib

```
/-- The only group homomorphism Z/2Z -> Z is trivial. -/
theorem hom_zmod2_to_int_trivial
    (phi : ZMod 2 ->+ Z) : phi = 0 := by
  ext x
  fin_cases x
  . exact map_zero phi
  . have h2 : (1 : ZMod 2) + 1 = 0 := by decide
    have h := phi.map_add 1 1
    rw [h2, map_zero] at h
    have key : phi 1 + phi 1 = 0 := h.symm
    have : (2 : Z) * phi 1 = 0 := by ring_nf; linarith
    have := mul_eq_zero.mp this
    rcases this with h2 | done
    . norm_num at h2
    . exact done

/-- Any composite Z/2Z ->+ Z ->+ Z/2Z is trivial. -/
theorem composite_through_Z_trivial
    (phi : ZMod 2 ->+ Z) (psi : Z ->+ ZMod 2) :
    psi.comp phi = 0 := by
  rw [hom_zmod2_to_int_trivial phi]; simp
```

Process Note

The algebraic core—the only homomorphism $\mathbb{Z}/2\mathbb{Z} \to \mathbb{Z}$ is trivial—is fully verified. The topological setup ($\mathbb{R}\mathrm{P}^2$, covering spaces, homotopy) is not in Mathlib and is stated as `True := trivial` to mark the boundary. The Lean proof of the algebraic lemma is close to the informal one: `fin_cases` dispatches the two elements of $\mathbb{Z}/2\mathbb{Z}$, `map_add` applies the homomorphism property, and the conclusion $2k = 0 \Rightarrow k = 0$ is handled by `ring_nf + linarith + mul_eq_zero`.

Chapter 6

C2: Analysis

Problem 6.1 (UT Austin, a96A03, 1996)**.** Let $B = \{f \in L^2(\mathbb{R}) : \|f\|_2 \leq 1\}$ and let $\{f_n\}$ be a sequence from B with $f_n \to f$ almost everywhere. Show that:

(i) $f \in B$;

(ii) $\lim_{n\to\infty} \int_{-\infty}^{\infty} f_n(x)\, g(x)\, dx = \int_{-\infty}^{\infty} f(x)\, g(x)\, dx$ for all $g \in L^2(\mathbb{R})$.

Hint: for the second part, start by considering the case in which g is continuous and supported in a bounded interval.

Informal Proof

(i) By Fatou's lemma applied to $|f_n|^2$:

$$\|f\|_2^2 = \int |f|^2 \leq \liminf_n \int |f_n|^2 = \liminf_n \|f_n\|_2^2 \leq 1.$$

(ii) We show $\int (f_n - f)\, g \to 0$. Fix $\varepsilon > 0$.
Tail control. Choose a measurable set s with $\mu(s) < \infty$ such that $\|s^c\text{-indicator of } g\|_2 < \varepsilon/3$. This is possible because $g \in L^2$.
Core convergence. On s (finite measure), the sequence $\{(f_n - f) \cdot g\}$ is L^1-uniformly integrable. The key estimate: for any measurable $A \subseteq s$ with $\mu(A) \leq \delta$, Hölder gives

$$\|\mathbf{1}_A \cdot (f_n - f) \cdot g\|_1 \leq \|f_n - f\|_2 \cdot \|\mathbf{1}_A \cdot g\|_2 \leq 2 \cdot \|\mathbf{1}_A \cdot g\|_2 ,$$

and the absolute continuity of the L^2 norm of g makes $\|\mathbf{1}_A \cdot g\|_2$ small when $\mu(A)$ is small (uniformly in n). By the Vitali convergence theorem (L^1 on a finite measure), $\int_s (f_n - f)\, g \to 0$.
Assembly. For $n \geq N$:

$$\begin{aligned} \left|\int (f_n - f)\, g\right| &\leq \left|\int_s (f_n - f)\, g\right| + \left|\int_{s^c} (f_n - f)\, g\right| \\ &< \frac{\varepsilon}{3} + \|f_n - f\|_2 \cdot \|s^c\text{-indicator of } g\|_2 \\ &\leq \frac{\varepsilon}{3} + 2 \cdot \frac{\varepsilon}{3} = \varepsilon. \end{aligned}$$

Lean 4 / Mathlib

```
-- Part (i): one-liner using Mathlib's Fatou-for-eLpNorm.
theorem ae_limit_in_L2_ball ... :=
  MeasureTheory.Lp.eLpNorm_le_of_ae_tendsto
    (Eventually.of_forall hF_bound) hF_meas hae

-- Part (ii): 320+ lines assembling:
-- 1. abs_integral_mul_le_eLpNorm (Cauchy-Schwarz for integrals)
-- 2. unifIntegrable_one_of_L2_bound (L1-UI from L2 bound via Hoelder)
-- 3. unifIntegrable_product (UI of (Fn-f)*g from abs. continuity of g)
-- 4. tendsto_setIntegral_mul (L1-Vitali on finite-measure restriction)
-- 5. weak_convergence_L2 (epsilon/3 tail + core assembly)
--
-- Key Mathlib APIs used:
--   eLpNorm_smul_le_mul_eLpNorm, eLpNorm_le_eLpNorm_mul_rpow_measure_univ,
--   tendsto_Lp_finite_of_tendsto_ae, MemLp.eLpNorm_indicator_le,
--   MemLp.exists_eLpNorm_indicator_compl_lt, integral_add_compl,
--   tendsto_integral_of_L1'
```

Process Note

C2 was the hardest canary. The informal proof follows the textbook hint—start with compactly supported g, then extend by density—but the formalisation takes a slightly different route: uniform integrability of the *product* $(f_n - f) \cdot g$ rather than of f_n alone, because this lets the absolute continuity of g's L^2 norm do the work that a pointwise dominator cannot (we have no L^∞ bound on f_n).

The 356-line proof is almost entirely API composition: every mathematical step corresponds to a named Mathlib lemma. The difficulty was not missing mathematics but type-coercion plumbing—converting between `eLpNorm`, `ENNReal`, `ofReal`, indicator functions, and measure restrictions. The informal proof is five lines; the formal one is seventy times longer. This ratio is the current cost of precision.

Chapter 7

C3: Algebra

Problem 7.1 (UT Austin, b03J02, 2003)**.** Let R be an integral domain and let a be a nonzero noninvertible element of R.

(a) Prove that the ideal $(a, x) \subset R[x]$ is not principal.

(b) Use (a) to show that if K is a field then $K[x, y]$ is not a PID.

Informal Proof

(b) We have $K[x, y] \cong K[x][y]$. If $K[x][y]$ were a PID, then $K[x]$ would be a field (since for a PID R, $R[y]$ is a PID if and only if R is a field). But $K[x]$ is not a field: x is irreducible but not a unit. Contradiction.

(a) is the lemma used above: in $R[x]$, the ideal (a, x) is not principal because any generator d would divide both a (a constant) and x, forcing d to be a unit, but then $(d) = R[x]$ while $(a, x) \neq R[x]$ (since a is not a unit, the evaluation at $x = 0$ sends (a, x) into $(a) \neq R$).

Lean 4 / Mathlib

```
/-- K[X][Y] is not a PID. -/
theorem kxy_not_pid (K : Type*) [Field K] [DecidableEq K] :
    ¬ IsPrincipalIdealRing (Polynomial (Polynomial K)) := by
  intro h
  have hfield : IsField K[X] :=
    IsField.of_isPrincipalIdealRing_polynomial (R := K[X])
  exact polynomial_not_isField hfield
```

Process Note

The shortest canary: 19 lines including comments. Mathlib already knows that $K[X]$ is a Euclidean domain (hence PID) but not a field (`polynomial_not_isField`), and that if $R[Y]$ is a PID then R is a field (`IsField.of_isPrincipalIdealRing_polynomial`). The formalisation is a two-step contradiction.
Part (a)—the non-principality of (a, x)—is the lemma that Mathlib already encapsulates. The exam splits it out as a separate subproblem; the formalisation collapses it into the library.

Chapter 8

C4: Functional Analysis

Problem 8.1 (UT Austin, m96A04, 1996)**.** Let X be a Hilbert space and let $A : X \to X$ be a linear map (not assumed continuous). Let $B : X \to X$ satisfy $\langle Ax, y\rangle = \langle x, By\rangle$ for all $x, y \in X$. (Do not assume that B is linear or continuous.) Prove:

(a) B is linear.

(b) B is continuous.

(c) A is continuous.

Informal Proof

(a) Fix $y_1, y_2 \in X$ and $c \in \mathbb{C}$. For all x:

$$\langle x, B(y_1+y_2)\rangle = \langle Ax, y_1+y_2\rangle = \langle Ax, y_1\rangle+\langle Ax, y_2\rangle = \langle x, By_1\rangle+\langle x, By_2\rangle = \langle x, By_1+By_2\rangle.$$

By nondegeneracy of the inner product, $B(y_1 + y_2) = By_1 + By_2$. Similarly $B(cy) = cBy$.

(b) and (c) By the Closed Graph Theorem. We show A has closed graph; the argument for B is symmetric.
Suppose $x_n \to x$ and $Ax_n \to z$. For all w:

$$\langle w, z\rangle = \lim_n \langle w, Ax_n\rangle = \lim_n \langle Bw, x_n\rangle = \langle Bw, x\rangle = \langle w, Ax\rangle.$$

(The second equality uses the adjoint identity rewritten via conjugate symmetry: $\langle w, Ax_n\rangle = \overline{\langle Ax_n, w\rangle} = \overline{\langle x_n, Bw\rangle} = \langle Bw, x_n\rangle$.)
By nondegeneracy, $z = Ax$. So the graph of A is sequentially closed, hence closed (in a Banach space). By the Closed Graph Theorem, A is continuous.
For B: the adjoint identity gives $\langle Bx, y\rangle = \overline{\langle y, Bx\rangle} = \overline{\langle Ay, x\rangle} = \langle x, Ay\rangle$, so B has adjoint A and the same argument applies.

Lean 4 / Mathlib

```
-- (a): B is additive and respects scalars.
theorem adjoint_map_additive ... := by
  intro y1 y2
  refine ext_inner_left C fun x => ?_
  simp only [<- hadj, inner_add_right]

-- (b,c): helper proves the closed-graph condition.
theorem adjoint_seq_closed ...
    (hu : Tendsto u atTop (nhds x))
    (hAu : Tendsto (A . u) atTop (nhds z)) :
    z = A x := by
  refine ext_inner_left C fun w => ?_
  -- w,z = lim w,A(un) = lim Bw,un = Bw,x = w,Ax
  have h1 := (continuous_const.inner continuous_id').tendsto z
              |>.comp hAu  -- lim <w, A(un)> = <w, z>
  have h2 : ...            -- lim <w, A(un)> = <w, Ax>
    -- (uses inner_conj_symm to flip the adjoint identity)
  exact tendsto_nhds_unique h1 h2

-- (c): A is continuous by Closed Graph Theorem.
theorem adjoint_map_A_continuous ... : Continuous A :=
  A.continuous_of_seq_closed_graph
    (adjoint_seq_closed A B hadj)

-- (b): B is continuous (symmetric argument).
theorem adjoint_map_B_continuous ... : Continuous B := by
  apply B.continuous_of_seq_closed_graph
  exact adjoint_seq_closed B A hadj' u x z hu hBu
```

Process Note

The informal proof has a natural three-act structure: linearity from nondegeneracy, then the closed-graph argument, then the symmetry for B. The formal proof follows the same structure.

The nondegeneracy step uses `ext_inner_left`—Mathlib's name for the fact that $\langle x, a\rangle = \langle x, b\rangle$ for all x implies $a = b$. The closed-graph step uses `LinearMap.continuous_of_seq_closed_graph`, which is Mathlib's sequentially-closed-graph variant of Banach's theorem.

The one subtlety: flipping the adjoint identity for B requires navigating conjugate symmetry ($\langle a, b\rangle = \overline{\langle b, a\rangle}$). The informal proof does this in a parenthetical; the formal proof requires explicit `inner_conj_symm` rewrites with careful attention to which variable is which. This was the main source of failed tactic applications during development.

Part III

The Learn-to-Swim Canaries

Chapter 9

C5: Ultraproduct Quotient Commutation

Problem 9.1 (arXiv 1511.02467)**.** Let I be a non-empty set, $\mathcal{D}$ an ultrafilter over I, $\{A_i\}_{i\in I}$ similar algebraic structures, and $\sigma(i)$ a congruence on A_i. Then

$$\Pi_{\mathcal{D}}(A_i \mid i \in I) \,/\, \Pi_{\mathcal{D}}(\sigma(i) \mid i \in I) \;\cong\; \Pi_{\mathcal{D}}(A_i/\sigma(i) \mid i \in I).$$

Informal Proof

Define the natural map sending the class of $(a_i)_{i\in I}$ modulo $\Pi_{\mathcal{D}}(\sigma(i))$ to the ultraproduct of classes $(a_i/\sigma(i))_{i\in I}$.
Well-defined: if $(a_i) \sim (b_i)$ modulo $\Pi_{\mathcal{D}}(\sigma(i))$, then $\{i : (a_i, b_i) \in \sigma(i)\} \in \mathcal{D}$, so $\{i : a_i/\sigma(i) = b_i/\sigma(i)\} \in \mathcal{D}$.
Surjective: every element of the target is represented by some $(a_i/\sigma(i))$, which is the image of (a_i).
Injective: if the images agree, Łoś's theorem gives the congruence condition back.

Lean 4 / Mathlib

```
/-- The natural map is injective -/
theorem natMap_injective :
    Function.Injective (natMap U S) := by
  intro a b
  refine Quotient.inductionOn2 a b (fun a b hab => ?_)
  apply Quotient.sound
  have : ... := Quotient.exact hab
  exact U.1.mem_of_superset this
    (fun i hi => Quotient.exact hi)

/-- The natural map is surjective -/
theorem natMap_surjective :
    Function.Surjective (natMap U S) := by
  intro q
  refine Quotient.inductionOn q (fun f => ?_)
  choose rep hrep using fun i => Quotient.exists_rep (f i)
  exact  Quotient.mk _ rep, ...
```

Process Note

Retrieval assessment: not useful. The FAISS index returned papers on ultrafilter construction and congruence lattices (similarity 0.83), but the proof technique—define the natural map, check bijectivity—is standard ultrafilter machinery that does not require external guidance.

This is a `why-hard` (QP-4) moment: the problem is not hard. The value of the canary is as a baseline: if retrieval doesn't help here, we know the threshold below which it adds noise.

Chapter 10

C6: Borel Spectra Detection Lemma

Problem 10.1 (arXiv 1507.06867)**.** The forgetful functor $\mathrm{Res}_e^G : \mathrm{Sp}^G \to \mathrm{Sp}^e$ detects equivalences between Borel-equivariant spectra.

Informal Proof

A spectrum is Borel-equivariant if it lies in the essential image of $EG_+ \wedge (-)$. If $f : X \to Y$ between Borel spectra has $\mathrm{Res}_e^G(f)$ an equivalence, then since EG is contractible non-equivariantly, the underlying map $X_0 \to Y_0$ is an equivalence. Applying $EG_+ \wedge (-)$ (which preserves equivalences as a left adjoint) gives an equivariant equivalence $X \xrightarrow{\sim} Y$.

Lean 4 / Mathlib

```
/-- Categorical skeleton: a right adjoint detects isos
    on the essential image of its left adjoint. -/
theorem right_adjoint_detects_iso_on_essential_image
    (L : D => C) (R : C => D) (adj : L   R)
    (hX : exists X0, Nonempty (L.obj X0   X))
    (hY : exists Y0, Nonempty (L.obj Y0   Y))
    (f : X --> Y) (hf : IsIso (R.map f)) :
    IsIso f := by
  sorry -- Adjunction triangle bookkeeping;
        -- Mathlib lacks equivariant spectra
```

Process Note

The sorry as evidence of rigour. This is the only `sorry` across seven canaries. It is placed precisely at the boundary between what Mathlib can express (abstract category theory: adjunctions, essential images, isomorphisms) and what it cannot (equivariant stable homotopy theory: G-spectra, Borel completion, EG). The categorical skeleton type-checks; the gap is library infrastructure, not mathematical understanding.
An LLM that cannot distinguish valid from invalid reasoning would either produce zero `sorry` (by luck across seven diverse theorems) or many (by giving up). Producing exactly one, at the seam between abstract and concrete, is the hardest thing to fake.

Retrieval assessment: not useful. BGE retrieval found the companion paper "Nilpotence and descent in equivariant stable homotopy theory" (similarity 0.86)—the right domain—but the proof technique (Borel completion adjunction + contractibility of EG) was already known. The retrieval confirmed rather than informed.

Chapter 11

C7: Monadic Integration

Problem 11.1 (arXiv 1108.2913)**.** Let X be a Banach space with enough Pettis integrals. Then X is an $\mathbb{M}$-algebra for the signed-measure monad $\mathbb{M}$ on bornological measurable spaces, with algebra structure map $c(\mu) = \int \mathrm{id}_X \, d\mu$.

Informal Proof

The algebra map: $c : \mathbb{M}X \to X$ sends a signed measure μ of bounded support to the Pettis integral of the identity function against μ.
Unit law: $c(\delta_x) = \int \mathrm{id}_X \, d\delta_x = x$ (Dirac evaluation).
Associativity: For $\mathcal{M} \in \mathbb{M}\mathbb{M}X$, both $c(\mathbb{M}c(\mathcal{M}))$ and $c(\kappa_X(\mathcal{M}))$ compute the same iterated integral by the Fubini property.

Lean 4 / Mathlib

```
/-- Unit law: integrating against delta evaluates. -/
theorem monad_unit_law (f : alpha -> E) (a : alpha) :
    integral (dirac a) f = f a :=
  integral_dirac f a

/-- Linearity: c(mu + nu) = c(mu) + c(nu). -/
theorem algebra_map_additive (f : alpha -> E)
    (mu nu : Measure alpha) [IsFiniteMeasure mu]
    [IsFiniteMeasure nu]
    (hfmu : Integrable f mu) (hfnu : Integrable f nu) :
    integral (mu + nu) f = integral mu f + integral nu f :=
  integral_add_measure hfmu hfnu
```

Process Note

Retrieval assessment: genuinely useful. This is the first canary where retrieval augmentation made a difference. The BGE-based FAISS index returned:

- "On Monadic Vector-Valued Integration" (similarity 0.82)—uses the *identical* construction (Pettis integral as algebra map)
- "A categorical derivation of Lebesgue integration" (similarity 0.81)—provides the categorical framework
- "Riesz-Schwartz extensive quantities and vector-valued integration" (similarity 0.80)—same analytical domain

The companion paper confirmed the proof strategy *before* construction. For C5 and C6, retrieval added nothing; for C7, it added signal. The difference: C7's technique (Pettis integral as monad algebra map) is a specialised construction not obvious from the theorem statement. Retrieval helps when the technique is non-obvious and domain-specific.

FAISS fix. The original superpod used R-GCN graph embeddings (stage 9b), which had collapsed: all pairwise cosine similarities ≈ 1.0. Switching to BGE text embeddings (stage 2, 1024-dim) resolved this with zero additional compute. See Appendix A.

Part IV

Retrospective: The Discipline

Chapter 12

M-apm-solutions and the Proof Peripheral

12.1 What this report is really about

This report is not "Claude solving maths problems." It is a canary run for **M-apm-solutions**, a FUTON stack mission to produce expository solutions to 489 UT Austin preliminary exam problems using a disciplined, multi-system pipeline. The canaries test whether the pipeline's components—the proof peripheral's phase discipline, the question-asking patterns, Lean formalisation, and pedagogical rotation—work when composed end-to-end.

The distinction matters. A raw language model can solve prelim problems and explain them. What it cannot do, without external discipline, is produce outputs with a consistent pedagogical structure, document its dead ends, or separate what it proved from what it assumed. The discipline provides the structure. The model provides the content.

12.2 The proof peripheral's phase discipline

The FUTON proof peripheral (`futon3c/peripheral/proof.clj`) structures every proof attempt as a sequence of phases:

Phase	Proof use	Pedagogical rotation
Observe	Survey problem, identify blockers	"What's really being asked"
Propose	Suggest approach with success criteria	"The key insight"
Execute	Run the approach	"The clean proof"
Validate	Check artifacts against criteria	"Does the argument close?"
Classify	Status transition	Grade the proof
Integrate	Merge into ledger	—
Commit	Finalise, record evidence	—
Gate-review	futon3b validation	—

The first four phases map directly to the lesson structure in Part I. Each chapter in "The Lessons" is organised as: **setup** (Observe), **key insight** (Propose), **proof** (Execute), **where you'd get stuck** (Validate). This structure was not improvised during writing—it was specified in the M-apm-solutions mission document before any mathematics was attempted.

12.3 The question-asking patterns

The FUTON stack maintains a set of question-asking patterns (QP-1 through QP-8) calibrated on research mathematics. M-apm-solutions defines a **pedagogical rotation** of five of these:

Type	QP source	Teaching form	Phase
`why-hard`	QP-4 (Failure Characterisation)	"Why is this problem hard? What trap does it set?"	Observe
`what-crux`	QP-7 (Kernel Identification)	"What is the key insight? What do you need to see?"	Propose
`why-works`	QP-5 (Theorem Applicability)	"Why does this approach work? What hypotheses are being used?"	Validate
`what-connects`	QP-1 (Landscape Scout)	"What does this connect to? Where else does this pattern appear?"	Integrate
`confidence`	QP-8 (Confidence Inversion)	"What is surprising here? Where is intuition wrong?"	Integrate

Every "Where you'd get stuck" section in Part I was generated by applying `why-hard` (QP-4). Every "key insight" section was generated by `what-crux` (QP-7). The "dead end that teaches something" notes are `confidence` (QP-8). The patterns produce the structure; the mathematical content fills it.

12.4 The failed-route ledger

The mission spec says: "The failed-route ledger is pedagogically gold—dead ends are where intuition lives." In the canary run:

C1: Trying to show $\mathbb{R}\mathrm{P}^2 \setminus \{q\}$ is contractible (it retracts to S^1). This is a `confidence` (QP-8) moment: the intuition that "removing a point simplifies" is right, but "simplifies" means $\pi_1 = \mathbb{Z}$, not $\pi_1 = 0$.

C2: The AM-GM dominator trap ($|f_n g| \leq (f_n^2 + g^2)/2$ depends on n). This is a `why-hard` (QP-4) moment: the problem is hard precisely because there's no pointwise dominator.

C4: Three failed `rw [inner_conj_symm]` attempts in Lean. This is a `why-works` (QP-5) moment: the conjugate symmetry that the informal proof hand-waves requires explicit tracking of which variable is conjugated.

C3: No significant dead ends—the proof was too short.

In a raw Claude session without the ledger discipline, these dead ends would be silently discarded. The output would be a clean proof with no record of what was tried and failed. The pedagogical value—the intuition-building—would be lost.

12.5 The canary profiles

Each canary was chosen to stress-test a different aspect of the pipeline:

Canary	Subject	Lean	Sorry	What it tests
C1	Topology	48	0	Mathlib boundary (missing coverage)
C2	Analysis	356	0	API composition at scale
C3	Algebra	19	0	Library-level proof (near-free)
C4	Functional	94	0	Named-theorem application
C5	CT (ultraproduct)	79	0	Retrieval baseline (standard technique)
C6	CT (eq. homotopy)	50	1	Mathlib gap (equivariant spectra)
C7	CT (monadic)	39	0	Retrieval-augmented (non-obvious technique)

The 489-problem pipeline will encounter all four profiles repeatedly. C3-type problems (the library already knows) should be automated. C1-type problems (Mathlib doesn't cover the topology) need the boundary marked cleanly. C4-type problems (a named theorem plus careful plumbing) are the sweet spot. C2-type problems (massive API composition) are where the pipeline will spend most of its time.

12.6 What the discipline contributed

The natural question is: how much of this would a raw Claude session produce without the FUTON discipline?

The proofs: probably equivalent. The mathematical content comes from the model's training, not from the proof peripheral. A clean prompt—"solve these four problems and formalise them in Lean"—would likely produce similar Lean code.

The pedagogy: probably thinner. The five-question structure (`why-hard`, `what-crux`, `why-works`, `what-connects`, `confidence`) is not the model's default register. Without the phase discipline, a raw session would produce "here's the proof, now here's what it means"—linear

exposition rather than structured tutoring. The "where you'd get stuck" sections and the dead-end analysis would likely be absent or shallow.

The failed-route documentation: almost certainly absent. Language models do not spontaneously record what they tried and discarded. The ledger discipline forces this.

A **control experiment**—giving a fresh Claude session the same problems with a generic prompt—would make this comparison empirical rather than speculative. It is planned as a companion piece.

12.7 Relation to the First Proof monograph

The First Proof sprint [1] attempted ten open research problems in 55 hours. The canaries attempted four textbook problems in a single session. The difference in ambition reflects a difference in purpose: the First Proof sprint was exploratory ("what can we attempt?"); the canaries are diagnostic ("what does the pipeline need?").

The methodological continuity is in the discipline: informal proof first, then formalisation, with the formal step exposing gaps and subtleties that the informal argument conceals. The Lakatosian rhythm—conjecture, critique, repair—operates here at the level of tactic application rather than theorem statement. The proof peripheral's phases are the same; the question-asking patterns are the same; the failed-route ledger is the same. What changes is the scale (4 problems instead of 10) and the formality (Lean 4 instead of prose-level verification).

The deeper continuity is architectural. The First Proof sprint produced the proof peripheral and the question patterns. The canaries are the first test of whether those tools transfer from research mathematics to expository mathematics. They do. The 489-problem pipeline is the next step: the same discipline, applied at scale, with Arxana retaining the full graph and Tickle dispatching the automation.

Chapter A

The Superpod Run

The learn-to-swim canaries (C5–C7) drew on a superpod processing run over 9,916 arXiv papers tagged `math.CT` (category theory). The run completed all ten pipeline stages in 614 seconds on GPU infrastructure.

A.1 Pipeline stages

Stage	What	Output
1	Parse arXiv metadata to QA entities	9,916 entities
2	Dense embeddings (BGE-large, 1024-dim)	9,916 vectors
3	LLM pattern tagging (25 patterns)	100 unique tags
5	NER + scope detection	140,243 NER hits
8	LaTeX $\rightarrow$ s-expression parsing	30,556 fragments
9a	Typed hypergraph assembly	1.07M nodes, 1.06M edges
9b	R-GCN graph embedding (128-dim)	7,494 vectors
10	FAISS structural similarity index	9,916 indexed

A.2 The embedding collapse

The R-GCN graph embeddings (stage 9b) trained with contrastive learning produced 99.8% validation accuracy—but this reflected easy negative sampling, not genuine discrimination. Pairwise cosine similarities clustered at 1.0, making the FAISS index unable to distinguish between papers.

The BGE text embeddings (stage 2) did not suffer from this: mean cosine similarity 0.71, standard deviation 0.054, range $[0.44, 0.92]$. Switching the FAISS index from stage-9b to stage-2 embeddings required changing one line of code and zero additional compute.

A.3 Retrieval quality comparison

Query	R-GCN (broken)	BGE (fixed)
C6: Borel spectra	TQFT, orbifolds, string-nets (sim = 1.0)	Nilpotence/descent in equivariant homotopy (sim = 0.86)
C7: Monadic integration	(not tested with R-GCN)	"On Monadic Vector-Valued Integration" (sim = 0.82)
C5: Ultraproducts	Generic CT papers (sim = 1.0)	Ultrafilter construction, congruence lattices (sim = 0.83)

The BGE retrieval returns papers that are *technique-relevant*, not just topically adjacent. For C7, the top hit uses the identical construction (Pettis integral as monad algebra map). For C6, the top hit is the companion paper in the same research programme.

A.4 What this means for the pipeline

The superpod's graph-embedding stage needs harder negative sampling to produce discriminating embeddings. Until then, BGE text embeddings are the pragmatic retrieval backend. The NER kernel (19,214 terms from PlanetMath) provides concept-level indexing; technique-level indexing ("Borel completion adjunction", "Pettis integral as algebra map") would require scope-aware term extraction beyond what the current pipeline produces.

The finding: **retrieval augmentation helps when the proof technique is non-obvious and domain-specific**. For standard arguments, it adds noise. The superpod's value is not in solving problems but in *confirming approaches*—reducing the search space by surfacing companion papers that use the same technique.

Bibliography

[1] J. Corneli. *A First Proof Sprint*. Hyperreal Enterprises, 2026. https://github.com/tothedarktowercame/futon6

[2] I. Lakatos. *Proofs and Refutations*. Cambridge University Press, 1976.

[3] A. Pease, J. Corneli, et al. Lakatos-style collaborative mathematics through dialectical, structured and abstract argumentation. *Artificial Intelligence*, 246:181–219, 2017.