Centre universitaire de santé McGill

McGill University Health Centre

☐ HME MCH ☒ HGM MGH ☒ HRV RVH
☐ HNM MNH ☒ ITM MCI ☐ CL LC

* F M U - 1 6 0 0 *

CESSATION TABAGIQUE
Ordonnance médicale

SMOKING CESSATION
Physician order

ALLERGIES : _____ **Poids** / *Weight* (kg): _____

Initiales du médecin Pour chaque ordonnance *Physician's Initials for each order*	**ORDONNANCE DU MÉDECIN /** *PHYSICIAN'S ORDERS*	**Initiales de l'infirmier(ère) notées** *Nurse's initials noted*
	See over for titration guidelines	
	Smoking **less than 10 cigarettes per day: (check one)** ☐ Nicotine **Patch** 14 mg daily x _____ weeks **OR 1 gum or lozenge q 1-2 hour PRN (maximum 16 pieces daily)** ☐ Nicotine **Lozenge** 1 mg x _____ weeks **OR** ☐ Nicotine **Gum** 2 mg x _____ weeks **OR** ☐ Nicotine **Lozenge** 2 mg x _____ weeks	
	Smoking **10 to 20 cigarettes per day:** Nicotine **Patch** 21 mg daily x _____ weeks **AND (check one) 1 gum or lozenge q 1-2 hour PRN (maximum 16 pieces daily)** ☐ Nicotine **Lozenge** 1 mg x _____ weeks **OR** ☐ Nicotine **Gum** 2 mg x _____ weeks **OR** ☐ Nicotine **Lozenge** 2 mg x _____ weeks	
	Smoking **20 to 30 cigarettes per day:** Nicotine **Patch** 28 mg daily x _____ weeks **AND (check one) 1 gum or lozenge q 1-2 hour PRN (maximum 16 pieces daily)** ☐ Nicotine **Lozenge** 1 mg x _____ weeks **OR** ☐ Nicotine **Gum** 2 mg x _____ weeks **OR** ☐ Nicotine **Lozenge** 2 mg x _____ weeks	
	Smoking **more than 30 cigarettes per day:** Nicotine **Patch** 35 mg daily x _____ weeks **AND (check one) 1 gum or lozenge q 1-2 hour PRN (maximum 16 pieces daily)** ☐ Nicotine **Lozenge** 1 mg x _____ weeks **OR** ☐ Nicotine **Gum** 2 mg x _____ weeks **OR** ☐ Nicotine **Lozenge** 2 mg x _____ weeks	
	OTHER FIRST LINE MEDICATIONS	
	Bupropion SR (**Zyban®**) 150 mg PO q AM x 3 days, then 150 mg PO BID x _____ weeks **Can be combined with nicotine replacement therapy (NRT)**	
	☐ **This patient is included in the IMPACT program** Varenicline (**Champix®**) 0.5 mg PO q AM x 3 days, 0.5 mg PO BID x 4 days, then 1 mg po BID x _____ weeks **May be combined with nicotine replacement therapy for the first week only**	
	Patient is on varenicline at home: Varenicline (**Champix®**) 1 mg PO BID x _____ weeks	

	Nom en lettres moulées *Name in print*	**Signature**	**N° Permis** *License N°*	**Heure** *Time* 00:00	**Date** AAYY/MM/JD
Médecin *Physician*					

	Nom en lettres moulées et/ou Numéro de permis *Name in print and/or License Number*	**Parapher /** *Initial*	**Heure** *Time* 00:00	**Date** AAYY/MM/JD
Infirmier(ère) *Nurse*				
Pharmacien(ne) *Pharmacist*				

Original-Pharmacie/Pharmacy **Copie Jaune-Dossier Médical** / Yellow Copy-Medical Record

IMPORTANT

If withdrawal or craving symptoms persist, increase patch dose by 7 mg every 24 to 48 h to a maximum of 42 mg and encourage increased use of fast acting nicotine replacement therapy (gum and lozenges) to a maximum of 16 pieces per day.

- Withdrawal & craving symptoms: desire to smoke, anger/irritability/frustration, anxiety/nervousness, sadness/loss of interest, difficulty concentrating, restlessness, increased appetite and insomnia.
- Ensure patient has fast acting NRT at bedside for easy access.
- Always review and assess proper use of nicotine replacement therapy (as outlined below).
  - Eating and drinking anything except water should be avoided for 15 minutes before and during use of fast acting NRT.
  - Nicotine gum should be chewed slowly until the taste becomes strong, then placed between the cheek and the gums for one minute. Repeat and discard after 30 minutes.
- Patch may be removed at bedtime if patient complains of “insomnia and/or nightmares”.
- It may be necessary for some patients to remain on NRT longer than the protocol below. Extension over 12 weeks may not be covered by their insurance.
- If patient continues to have the urge to smoke, please consult smoking cessation practitioner.

Guidelines for the Titration of Nicotine Replacement Therapy

Patient should consult a family doctor or a pharmacist 2 - 4 weeks after discharge

Smoking less than 10 cigarettes per day	Smoking 10 - 20 cigarettes per day	Smoking 20 - 30 cigarettes per day	Smoking more than 30 cigarettes per day
Long acting nicotine therapy (daily)			
Nicotine patch 14 mg **X 6 weeks**	Nicotine patch 21 mg **X 6 weeks**	Nicotine patch 28 mg **X 4 weeks**	Nicotine patch 35 mg **X 4 weeks**
Nicotine patch 7 mg **X 4-6 weeks**	Nicotine patch 14 mg **X 4 weeks**	Nicotine patch 21 mg **X 4 weeks**	Nicotine patch 28 mg **X 2 weeks**
	Nicotine patch 7 mg **X 2 weeks**	Nicotine patch 14 mg **X 2 weeks**	Nicotine patch 21 mg **X 2 weeks**
		Nicotine patch 7 mg **X 2 weeks**	Nicotine patch 14 mg **X 2 weeks**
			Nicotine patch 7 mg **X 2 weeks**
Watch for symptoms of overdose: nausea, excessive salivation, sweating, headache and palpitations			
Fast acting nicotine therapy (week 1 to week 12)			
Nicotine (Thrive®) Lozenge 1 or 2 mg and Nicotine (Nicorette®) Gum 2 mg			
Use fast acting as needed to relieve withdrawal & craving symptoms every 1 - 2 h (maximum 16 pieces per day)			
Other first line medications			
Bupropion (Zyban®) 150 mg daily (day 1 - 3) then BID (day 4 to week 12) • It may be necessary to have the dose lowered temporarily or permanently if patient experiences nausea or other side effects. Contraindications: history of seizures or eating disorder, concomitant use of another form of Bupropion, MAO inhibitor use in the past 14 days or a recent history of discontinuation of alcohol or sedatives.			
Varenicline (Champix®) 0.5 mg daily (day 1 - 3), 0.5mg BID (day 4 - 7) then 1 mg (week 2 to week 12) • May be extended up to 24 weeks if not smoking at 12 weeks. • It may be necessary to decrease the dose temporarily or permanently if patient experiences nausea or other side effects. Contraindications: Severe renal failure and concomitant use of cimetidine. Observe for neuropsychiatric side effects particularly in context of past depression or suicide attempt.			