Centre universitaire de santé McGill

McGill University Health Centre

☒ HME MCH ☒ HGM MGH ☒ HRV RVH
☒ HNM MNH ☒ ITM MCI ☒ HL LH

* F M U - 3 3 3 8 *

Ordonnances Médicales

Medical Orders

ALLERGIES _____

Poids / Weight (kg) _____ **Taille** / Height (cm) _____ **Surface corporelle / BSA (m^2)** _____

DATE-(AAYY/MM/JD) HEURE/TIME (00:00)	**ORDONNANCES DU PRESCRIPTEUR/** PRESCRIBER'S ORDERS **Le prescripteur doit indiquer son nom en lettres moulées, apposer sa signature et son N° de permis** Prescriber must write name in print letters, sign and write license No.	**Initiales de l'infirmier(ère) notées** Nurse's initials **noted**

PHARMACEUTICAL ORDERS – ANALGESIA AND SEDATION – ADULT ICU

GENERAL

• **NOTIFY THE INTENSIVE CARE TEAM OR THE ANESTHESIOLOGIST** if analgesia or sedation targets are not met, or if the maximum rate is reached without success.

• Each rate increase should be preceded by a bolus to ensure a rapid onset of action.

ANALGESIA

Analgesia assessment scale = CPOT (Critical-Care Pain Observation Tool)

1st CHOICE: IF eGFR 50 mL/min OR MORE:

☐ **Morphine** 1 mg/mL

- ☐ **Bolus:** _____ mg IV q _____ min PRN if CPOT 3 or more **OR** before a potentially painful procedure
- ☐ **Variable dose: Initial dose:** _____ mg/h
  Dose range: from _____ mg/h to _____ mg/h for CPOT less than 3
- ☐ **Fixed dose:** _____ mg/h (no rate adjustment without physician approval)

2nd CHOICE: IF eGFR BETWEEN 25 AND 49 mL/min:

☐ **HYDROmorphone** 0.4 mg/mL

- ☐ **Bolus:** _____ mg IV q _____ min PRN if CPOT 3 or more **OR** before a potentially painful procedure
- ☐ **Variable dose: Initial dose:** _____ mg/h
  Dose range: from _____ mg/h to _____ mg/h for CPOT less than 3
- ☐ **Fixed dose:** _____ mg/h (no dose adjustment without physician approval)

3rd CHOICE: eGFR LESS THAN 25 mL/min OR RENAL REPLACEMENT THERAPY OR MORPHINE ALLERGY:

☐ **Fentanyl** 50 mcg/mL

- ☐ **Bolus:** _____ mcg IV q _____ min PRN if CPOT 3 or more **OR** before a potentially painful procedure
- ☐ **Variable dose: Initial dose:** _____ mcg/h
  Dose range: from _____ mcg/h to _____ mcg/h for CPOT less than 3
- ☐ **Fixed dose:** _____ mcg/h (no dose adjustment without physician approval)

Prescriber (print name): _____ **Signature:** _____

License no.: _____ **Date** (yyyy/mm/dd): _____/____/____ **Time** (hh:mm): ____:____

Centre universitaire de santé McGill

McGill University Health Centre

☒ HME MCH ☒ HGM MGH ☒ HRV RVH
☒ HNM MNH ☒ ITM MCI ☒ HL LH

* F M U - 3 3 3 8 *

Ordonnances Médicales

Medical Orders

ALLERGIES _____

Poids / Weight (kg) _____ **Taille** / Height (cm) _____ **Surface corporelle / BSA (m^2)** _____

DATE-(AAYY/MM/JD) HEURE/TIME (00:00)	**ORDONNANCES DU PRESCRIPTEUR/** PRESCRIBER'S ORDERS **Le prescripteur doit indiquer son nom en lettres moulées, apposer sa signature et son N° de permis** Prescriber must write name in print letters, sign and write license No.	**Initiales de l'infirmier(ère) notées** Nurse's initials **noted**

SEDATION

Use the **sedation assessment scale = RASS** (Richmond Agitation-Sedation Scale).

☐ **Midazolam** 1 mg/mL
- ☐ **Bolus:** _____ mg IV q _____ min PRN if RASS 1 or more
- ☐ **Variable dose: Initial dose:** _____ mg/h
  - **Dose range:**
    - ☐ **Standard:** from _____ mg/h to _____ mg/h for RASS 0 to −2 (do not use if patient is receiving neuromuscular blockade)
    - ☐ **Deep sedation:** from _____ mg/h to _____ mg/h for ☐ RASS −3 to −5 or ☐ RASS −5
- ☐ **Fixed dose:** _____ mg/h (no dose adjustment without physician approval)

☐ **Propofol** 10 mg/mL (**USUAL MAXIMUM DOSE: 65 MCG/KG/MIN**)

Bolus: If a bolus is given for agitation and ventilator dyssynchrony, also increase the infusion dose if possible. Notify the intensivist or anesthesiologist if the maximum infusion dose is reached and boluses are needed.
- ☐ **Midazolam** _____ mg IV q _____ min PRN if RASS 1 or more — do not give through the propofol line (physicochemical incompatibility)

OR
- ☐ **Propofol** ☐ 10 mg (1 mL) OR ☐ 20 mg (2 mL) IV q 10 min PRN if RASS 1 or more
- ☐ **Variable dose: Initial dose:** _____ mcg/kg/min
  - **Dose range:**
    - ☐ **Standard:** from _____ to _____ mcg/kg/min for RASS 0 to −2 (do not use if patient is receiving neuromuscular blockade)
    - ☐ **Deep sedation:** from _____ mcg/kg/min to 65 mcg/kg/min for ☐ RASS −3 to −5 or ☐ RASS −5
- ☐ **Fixed dose:** _____ mcg/kg/min (no dose adjustment without physician approval)

GASTROPROTECTION – STRESS ULCER PROPHYLAXIS

Start: ☐ Today **OR** ☐ On day 2 post-intubation; intubation date (yyyy/mm/dd): _____/____/____

☐ **Famotidine** according to renal function:
- eGFR 60 mL/min or more: ☐ 20 mg PO BID ☐ 20 mg IV BID
- eGFR 30 to 59 mL/min: ☐ 20 mg PO daily ☐ 20 mg IV daily
- eGFR less than 30 mL/min or intermittent/continuous renal replacement therapy: **favour a PPI (see below)**

☐ **Lansoprazole orally disintegrating** 30 mg via NG tube daily

☐ **If strict NPO: Pantoprazole** 40 mg IV daily

ORAL CARE – PREVENTION OF VENTILATOR-ASSOCIATED PNEUMONIA

☒ **Chlorhexidine** 0.12%, 15 mL swabbed during mouth care BID

☒ **Lip balm (Glaxal Base®)** topically q4h scheduled and q2h PRN

Prescriber (print name): _____ **Signature:** _____

License no.: _____ **Date** (yyyy/mm/dd): _____/____/____ **Time** (hh:mm): ____:____