Centre universitaire de santé McGill

McGill University Health Centre

X HME MCH X HGM MGH X HRV RVH
X HNM MNH X ITM MCI X HL LH

* F M U - 3 3 3 8 *

Ordonnances Médicales

Medical Orders

ALLERGIES _____

Poids / Weight (kg) _____ **Taille** / Height (cm) _____ **Surface corporelle / BSA (m^2)** _____

DATE-(AAYY/MM/JD) HEURE/TIME (00:00)	**ORDONNANCES DU PRESCRIPTEUR/** PRESCRIBER'S ORDERS **Le prescripteur doit indiquer son nom en lettres moulées, apposer sa signature et son N° de permis** Prescriber must write name in print letters, sign and write license No.	**Initiales de l'infirmier(ère) notées** Nurse's initials **noted**

PREVENTION OF IODINATED CONTRAST-INDUCED NEPHROPATHY – INPATIENT AND EMERGENCY

Exam date (yyyy/mm/dd): _____/____/____

Ensure a creatinine result is available from within the 7 days before the exam for patients with at least one of the risk factors below.

☐ 70 years or older
☐ Diabetes
☐ Heart disease
☐ Chemotherapy (active cancer under treatment or treatment within the last 6 weeks)
☐ Kidney disease
☐ Kidney transplant
☐ Nephrology follow-up recommended

☐ **If no result is available, draw a serum creatinine if risk factors are present**

MEASURES ACCORDING TO eGFR

☐ **Patient with eGFR greater than 60 mL/min/1.73 m²:**
- ✓ Have the patient drink 500 mL of water before the exam.

☐ **Patient with eGFR greater than 30 and up to 60 mL/min/1.73 m²:**
- ✓ Have the patient drink 500 mL of water before the exam.
- ✓ Hold loop diuretics (furosemide, bumetanide, ethacrynic acid) and metolazone on the day of the exam.

☐ **Patient with eGFR 30 mL/min/1.73 m² or less:** the radiologist and the referring physician assess the risks/benefits.

If iodinated contrast injection is confirmed by the radiologist:
- ✓ NaCl 0.9% IV at 3 mL/kg/h (max 300 mL): _____ for one hour before the exam, then 1.5 mL/kg/h (max 150 mL/h): _____ for 4 hours after the patient returns to the care unit after the exam
- ✓ Stop metFORMIN on the day of the exam. Resumption to be assessed by the treating physician (NB: contraindicated if eGFR less than 30 mL/min).
- ✓ Hold loop diuretics (furosemide, bumetanide, ethacrynic acid) and metolazone on the day of the exam.
- ✓ Draw a serum creatinine 48 hours after the exam.

Prescriber (print name): _____ **Signature:** _____

License no.: _____ **Date** (yyyy/mm/dd): _____/____/____ **Time** (hh:mm): ____:____