Centre universitaire de santé McGill

McGill University Health Centre

☒ HME MCH ☒ HGM MGH ☒ HRV RVH
☒ HNM MNH ☒ ITM MCI ☒ HL LH

* F M U - 3 3 3 8 *

Ordonnances Médicales

Medical Orders

ALLERGIES _____

Poids / Weight (kg) _____ **Taille** / Height (cm) _____ **Surface corporelle / BSA (m^2)** _____

DATE-(AAYY/MM/JD) HEURE/TIME (00:00)	**ORDONNANCES DU PRESCRIPTEUR/** PRESCRIBER'S ORDERS **Le prescripteur doit indiquer son nom en lettres moulées, apposer sa signature et son N° de permis** Prescriber must write name in print letters, sign and write license No.	**Initiales de l'infirmier(ère) notées** Nurse's initials **noted**

VASOPRESSORS / INOTROPES / CHRONOTROPES – ADULT

ADMINISTRATION GUIDELINES

- **DO NOT** give boluses of vasopressors, dobutamine or isoproterenol.
- **In case of extravasation:** notify MD to consider intradermal administration of **phentolamine**.

VASOPRESSOR AGENTS

Targets: ☐ SBP _____ mmHg or more ☐ MAP _____ mmHg or more ☐ CPP _____ to _____ mmHg

☐ **Norepinephrine** 32 mcg/mL
- ☐ Start at _____ mcg/kg/min IV infusion; titrate to reach the targets above
  Notify if the dose is _____ mcg/kg/min or more
- ☐ Fixed dose: _____ mcg/kg/min

Suggested initial dose: 0.05 mcg/kg/min
Titration/weaning: steps of 0.02 mcg/kg/min q5min

☐ **EPINEPHrine** 32 mcg/mL
- ☐ Start at _____ mcg/kg/min IV infusion; titrate to reach the targets above
  Notify if the dose is _____ mcg/kg/min or more
- ☐ Fixed dose: _____ mcg/kg/min

Suggested initial dose: 0.05 mcg/kg/min
Titration/weaning: steps of 0.02 mcg/kg/min q5min

☐ **Vasopressin** 0.24 unit/mL — start at:
☐ 0.04 unit/min IV infusion ☐ _____ unit/min IV infusion
Subsequent dose adjustment per physician order
Notify physician if: _____

Suggested dose: 0.01 to 0.04 unit/min
MAXIMUM fixed dose: 0.04 unit/min

☐ **DOPamine** 1600 mcg/mL
- ☐ Start at _____ mcg/kg/min IV infusion; titrate to reach the targets above
  Notify if the dose is _____ mcg/kg/min or more

Suggested initial dose: 2 to 5 mcg/kg/min
Titration/weaning: steps of 2.5 to 5 mcg/kg/min q5–10min
Usual doses: 2 to 25 mcg/kg/min

Vasopressor weaning
- Decrease doses gradually at 5- to 10-minute intervals, according to clinical judgement.
- **Vasopressin:** wean per physician order (longer intervals: decrease every 30 to 60 minutes).
- **If more than one vasopressor is running:** confirm with the physician which agent to wean first, unless specified here: ☐ Wean _____ first.
- **If vasopressin is running:** consider weaning it first unless otherwise specified by the treating physician.

Prescriber (print name): _____ **Signature:** _____

License no.: _____ **Date** (yyyy/mm/dd): _____/____/____ **Time** (hh:mm): ____:____

Centre universitaire de santé McGill

McGill University Health Centre

☒ HME MCH ☒ HGM MGH ☒ HRV RVH
☒ HNM MNH ☒ ITM MCI ☒ HL LH

* F M U - 3 3 3 8 *

Ordonnances Médicales

Medical Orders

ALLERGIES _____

Poids / Weight (kg) _____ **Taille** / Height (cm) _____ **Surface corporelle / BSA (m^2)** _____

DATE-(AAYY/MM/JD) HEURE/TIME (00:00)	**ORDONNANCES DU PRESCRIPTEUR/** PRESCRIBER'S ORDERS **Le prescripteur doit indiquer son nom en lettres moulées, apposer sa signature et son N° de permis** Prescriber must write name in print letters, sign and write license No.	**Initiales de l'infirmier(ère) notées** Nurse's initials **noted**

INOTROPIC AGENTS

☐ **DOBUTamine** 1 mg/mL
Start at: ☐ 2.5 mcg/kg/min ☐ _____ mcg/kg/min IV infusion
Subsequent dose adjustment per physician order. Notify physician if:
_____

Suggested dose: 2.5 to 20 mcg/kg/min

☐ **Milrinone** 200 mcg/mL *Caution in renal impairment*
☐ Loading dose: 50 mcg/kg, i.e. _____ mcg IV over 10 min, given from the infusion
Start at: ☐ 0.375 mcg/kg/min ☐ _____ mcg/kg/min IV infusion
Subsequent dose adjustment per physician order. Notify physician if:
_____

Suggested dose: see table below

***Renal impairment:** increased half-life and risk of accumulation

Creatinine clearance	Recommended starting dose		
	Minimum dose	**Standard dose**	**Maximum dose**
60 mL/min or more	0.375 mcg/kg/min	0.5 mcg/kg/min	0.75 mcg/kg/min
50 to 59 mL/min	0.25 mcg/kg/min	0.375 mcg/kg/min	0.5 mcg/kg/min
40 to 49 mL/min	0.125 mcg/kg/min	0.25 mcg/kg/min	0.375 mcg/kg/min
30 to 39 mL/min	0.0625 mcg/kg/min	0.125 mcg/kg/min	0.25 mcg/kg/min
20 to 29 mL/min	Not recommended	0.0625 mcg/kg/min	0.125 mcg/kg/min
10 to 19 mL/min	Not recommended		0.0625 mcg/kg/min
9 mL/min or less	Not recommended		

Inotrope weaning: per individual physician order.

CHRONOTROPIC AGENT

☐ **Isoproterenol** 10 mcg/mL
☐ Start at _____ mcg/min IV infusion
Titrate for: ☐ HR _____ beats/min or more
Notify physician if: _____

Suggested initial dose: 0.5 to 2 mcg/min
Titration: steps of 1 mcg/min q5min
Usual doses: 2 to 10 mcg/min

Chronotrope weaning: per individual physician order.

Prescriber (print name): _____ **Signature:** _____

License no.: _____ **Date** (yyyy/mm/dd): _____/____/____ **Time** (hh:mm): ____:____