Centre universitaire de santé McGill

McGill University Health Centre

X HME MCH X HGM MGH X HRV RVH
X HNM MNH X ITM MCI X HL LH

* F M U - 3 3 3 8 *

Ordonnances Médicales

Medical Orders

ALLERGIES _____

Poids / Weight (kg) _____ **Taille** / Height (cm) _____ **Surface corporelle / BSA (m^2)** _____

DATE-(AAYY/MM/JD) HEURE/TIME (00:00)	**ORDONNANCES DU PRESCRIPTEUR/** PRESCRIBER'S ORDERS **Le prescripteur doit indiquer son nom en lettres moulées, apposer sa signature et son N° de permis** Prescriber must write name in print letters, sign and write license No.	**Initiales de l'infirmier(ère) notées** Nurse's initials **noted**

PHARMACEUTICAL ORDERS – DISTRESS AND BRONCHIAL CONGESTION – PALLIATIVE CARE ADULT

DISTRESS ORDER

INDICATIONS: major respiratory distress, **or** massive hemorrhage, **or** intolerable pain not relieved by the prescribed medications — **with panic AND agitation**.

MIDAZOLAM (Versed®) – syringe no. 1

☐ Midazolam _____ mg (min. suggested dose: 5 mg)

OPIOID – syringe no. 2

☐ Morphine _____ mg (min. suggested dose: 5 mg)

OR

☐ HYDROmorphone _____ mg (min. suggested dose: 1 mg)

OR

☐ FentaNYL _____ mcg (min. suggested dose: 50 mcg)

SCOPOLAMINE – syringe no. 3

☐ Scopolamine **0.4 mg**

All medications:

☐ **IV** or ☐ **SC**

See suggested doses below

If the IV route is not available, use the SC route. By SC route, scopolamine may be given with the opioid in the second syringe if the total volume is less than 2 mL.

Notify the physician 10 to 20 minutes after each administration of the distress order.

If **distress persists**, repeat all distress order medications **once only**: 20 minutes after SC administration; 10 minutes after IV administration.

BRONCHIAL CONGESTION ORDER (WITHOUT DISTRESS) – check only one

☐ **GLYCOPYRROLATE** (Robinul®) _____ mg SC q _____ h PRN (suggested dose = 0.4 mg q4h PRN)

OR

☐ **Scopolamine** _____ mg SC q _____ h PRN

Prescriber (print name): _____ **Signature:** _____

License no.: _____ **Date** (yyyy/mm/dd): _____/____/____ **Time** (hh:mm): ____:____

SUGGESTED DOSES FOR THE DISTRESS ORDER

MIDAZOLAM (Versed®)	**Suggested dose**
Weight 70 kg or less **AND** equivalent lorazepam dose less than 3 mg/24 h	Midazolam 5 mg
Weight more than 70 kg **OR** equivalent lorazepam dose 3 to 12 mg/24 h	Midazolam 10 mg
Equivalent lorazepam dose more than 12 mg/24 h	Midazolam 15 mg

Benzodiazepine equivalent doses					
Lorazepam (Ativan®)	Alprazolam (Xanax®)	Bromazepam (Lectopam®)	Clonazepam (Rivotril®)	Oxazepam (Serax®)	Temazepam (Restoril®)
1 mg	0.5 mg	3 mg	0.5 mg *	15 mg	15 mg

The proposed clonazepam equivalence is intentionally conservative

Morphine received per 24 h		**per 4 h**	**Suggested dose**
PO	**SC or IV**	**SC or IV**	
0 to 30 mg	0 to 15 mg	0 to 2.5 mg	5 mg
30.1 to 120 mg	15.1 to 60 mg	2.6 to 10 mg	10 mg
120.1 to 360 mg	60.1 to 180 mg	10.1 to 30 mg	15 mg
More than 360 mg	More than 180 mg	More than 30 mg	5% of the PO dose received per 24 h (maximum 50 mg) or 10% of the SC or IV dose received per 24 h (maximum 50 mg)

HYDROmorphone received per 24 h		**per 4 h**	**Suggested dose**
PO	**SC or IV**	**SC or IV**	
0 to 6 mg	0 to 3 mg	0 to 0.5 mg	1 mg
6.1 to 24 mg	3.1 to 12 mg	0.6 to 2 mg	2 mg
24.1 to 72 mg	12.1 to 36 mg	2.1 to 6 mg	3 mg
More than 72 mg	More than 36 mg	More than 6 mg	5% of the PO dose received per 24 h (maximum 10 mg) or 10% of the SC or IV dose received per 24 h (maximum 10 mg)

FentaNYL: transdermal patch dose or fentaNYL infusion rate	**Suggested dose**		
	Morphine	**HYDROmorphone**	**FentaNYL**
0 to 12 mcg/h	5 mg	1 mg	50 mcg
13 to 50 mcg/h	10 mg	2 mg	100 mcg
51 to 150 mcg/h	15 mg	3 mg	150 mcg
151 to 300 mcg/h	25 mg	5 mg	250 mcg
More than 300 mcg/h	50 mg	10 mg	500 mcg