Centre universitaire de santé McGill

McGill University Health Centre

☒ HME MCH ☒ HGM MGH ☒ HRV RVH
☒ HNM MNH ☒ ITM MCI ☒ HL LH

* F M U - 3 3 3 8 *

Ordonnances Médicales

Medical Orders

ALLERGIES _____

Poids / Weight (kg) _____ **Taille** / Height (cm) _____ **Surface corporelle / BSA (m^2)** _____

DATE-(AAYY/MM/JD) HEURE/TIME (00:00)	**ORDONNANCES DU PRESCRIPTEUR/** PRESCRIBER'S ORDERS **Le prescripteur doit indiquer son nom en lettres moulées, apposer sa signature et son N° de permis** Prescriber must write name in print letters, sign and write license No.	**Initiales de l'infirmier(ère) notées** Nurse's initials **noted**

PHARMACEUTICAL ORDERS – SHIVERING CONTROL – TARGETED TEMPERATURE MANAGEMENT

SHIVERING PREVENTION

☒ **Acetaminophen** 975 mg PO or PR q6h SCHEDULED
If weight less than 45 kg: 650 mg PO or PR q6h SCHEDULED
☒ **Magnesium sulfate** 2 g IV infused over 2 hours × 2 consecutive doses (total 4 g) at the start of the cooling phase

TREATMENT OF SHIVERING WHEN BSAS SCORE IS 2 OR 3

- **Give treatments in the order presented**
- **If a treatment fails (BSAS 2 or 3 despite the treatment): proceed to the next treatment**
- **This order set ends once body temperature has returned to 37 °C**

1. **Meperidine** 50 mg IV q4h PRN (maximum 100 mg/24 hours)
2. **IF SHIVERING 15 MINUTES AFTER MEPERIDINE OR IF SHIVERING AND MAXIMUM MEPERIDINE DOSE REACHED:**
   ☒ Stop meperidine
   ☒ **Notify the intensivist or anesthesiologist** to deepen sedation and target RASS −5. Sedation and target must be prescribed by the treating physician on the order set: **Analgesia/Sedation – Mechanical Ventilation – Intensive Care – Adult**
3. **IF SHIVERING DESPITE DEEP SEDATION (RASS −5):**
   ☒ Maintain a sedation target of RASS −5
   ☒ **Rocuronium** 50 mg IV q1h PRN (maximum 3 doses over a 6-hour period)
   If weight more than 100 kg: Rocuronium 100 mg IV q1h PRN (maximum 3 doses over a 6-hour period)
4. **IF SHIVERING AND MAXIMUM ROCURONIUM DOSE REACHED:**
   ☒ Stop rocuronium
   ☒ **Cisatracurium** 0.2 mg/kg, i.e. _____ mg IV bolus × 1, followed by the infusion
   Cisatracurium 2 mg/mL IV infusion: start at 0.5 mcg/kg/min, then titrate to a maximum of 3 mcg/kg/min for **TOF (***train of four***) 2/4**, until body temperature returns to 37 °C, then stop

SHIVERING ASSESSMENT SCALE (BSAS = *Bedside Shivering Assessment Scale*)

0 – None: no shivering
1 – Mild: shivering localized to the neck/thorax; may be detected only as ECG artifact or by palpation
2 – Moderate: intermittent shivering involving the upper extremities ± thorax
3 – Severe: generalized shivering or sustained shivering of the upper or lower extremities

Prescriber (print name): _____ **Signature:** _____

License no.: _____ **Date** (yyyy/mm/dd): _____/____/____ **Time** (hh:mm): ____:____