

Number of quantifier alternations in Huang and nested

1 Preliminaries

For $\sigma = \text{mgu}(\varphi, \psi)$ for two terms or literals φ and ψ , we denote by σ_i for $1 \leq i \leq n$ the i th substitution which is added to σ by the unification algorithm, where $n = |\text{dom}(\sigma)|$. We define $\sigma_0 \stackrel{\text{def}}{=} \text{id}$.

We furthermore denote the composition of all σ_k for $i \leq k \leq j$ by $\sigma_{(i,j)}$. Hence $\sigma = \sigma_{(1,n)} = \sigma_{(0,n)}$.

A literal l is called a Φ -literal if its predicate symbol is Φ -colored.

NB: The notion of single-colored is considered to be deprecated here.

In a literal or term ϕ containing a subterm t , t is said to occur *below* a Φ -symbol s if in the syntax tree representation of ϕ , there is a node labelled s on the path from the root to t . Note that the colored symbol may also be the predicate symbol. Moreover, t is said to occur *directly below* a Φ -symbol if it occurs below the Φ -symbol s and in the syntax tree representation of ϕ on the path from s to t , no nodes with labels with colored symbol occur.

Quantifier alternations in I usually assumes the quantifier-alternation-minimizing arrangement of quantifiers in I . The lemma statements hence talk about the *minimal* number of quantifier alternations, which is indeed easily obtainable, i.e. it's a lower and upper bound at the same time.

In the following, we assume that the maximum \max of an empty sequence is defined to be 0 and constants are treated as function symbols of arity 0. Furthermore \perp is used to denote a color which is not possessed by any symbol.

Definition 1 (Color alternation col-alt). Let Γ and Δ be sets of formulas and t be a term.

$$\text{col-alt}(t) \stackrel{\text{def}}{=} \text{col-alt}_{\perp}(t)$$

$$\text{col-alt}_\Phi(t) \stackrel{\text{def}}{=} \begin{cases} 0 & t \text{ is a variable} \\ \max(\text{col-alt}_\Phi(t_1), \dots, \text{col-alt}_\Phi(t_n)) & t = f(t_1, \dots, t_n) \text{ is grey} \\ \max(\text{col-alt}_\Phi(t_1), \dots, \text{col-alt}_\Phi(t_n)) & t = f(t_1, \dots, t_n) \text{ is of color } \Phi \\ 1 + \max(\text{col-alt}_\Psi(t_1), \dots, \text{col-alt}_\Psi(t_n)) & t = f(t_1, \dots, t_n) \text{ is of color } \Psi, \\ & \Phi \neq \Psi \end{cases}$$

\triangle

Definition 2 (Quantifier alternation quant-alt). Let A be a formula.

$$\text{quant-alt}(A) \stackrel{\text{def}}{=} \text{quant-alt}_\perp(A)$$

$$\text{quant-alt}_Q(A) \stackrel{\text{def}}{=} \begin{cases} 0 & A \text{ is an atom} \\ \text{quant-alt}_Q(B) & A \equiv \neg B \\ \max(\text{quant-alt}_Q(B), \text{quant-alt}_Q(C)) & A \equiv B \circ C, \circ \in \{\wedge, \vee, \supset\} \\ \text{quant-alt}_Q(B) & A \equiv Q'B, Q = Q' \\ 1 + \text{quant-alt}_{Q'}(B) & A \equiv Q'B, Q \neq Q' \end{cases}$$

\triangle

Note that this definition of quantifier alternations handles formulas in prenex and non-prenex form.

Definition 3. We define $\text{PI}_{\text{step}}^\circ$ to coincide with PI_{step} but without applying the substitution σ in each of the cases. Analogously, if $C \equiv D\sigma$, we use C° to denote D . \triangle

Hence $\text{PI}_{\text{step}}^\circ(\cdot)\sigma = \text{PI}_{\text{step}}(\cdot)$.

2 Occurrence of terms in the interpolant

Definition 4 (PI^*). PI^* is defined as PI with the difference that in PI^* , all literals are considered to be grey. \triangle

Hence $\text{PI}_{\text{init}}^*$ coincides with PI_{init} . $\text{PI}_{\text{step}}^*$ coincides with PI_{step} in case of factorisation and paramodulation inferences. For resolution inferences, the first two cases in the definition of PI_{step} do not occur for $\text{PI}_{\text{step}}^*$.

PI^* enjoys the convenient property that it absorbs every literal which occurs some clause:

$\langle \text{prop:every_lit_in_pi_star} \rangle$ **Proposition 5.** , For every literal which occurs in a clause of a resolution refutation π , a respective successor occurs in $\text{PI}^*(\pi)$.

Proof. By structural induction. \square

$\text{ma:grey_lits_of_pi_star_in_pi}$ **Lemma 6.** For every clause C of a resolution refutation, every grey literal, which occurs in $\text{PI}^*(C)$, also occurs in $\text{PI}(C)$.

Proof. Note that PI_{init} and $\text{PI}_{\text{init}}^*$ coincide and PI_{step} and $\text{PI}_{\text{step}}^*$ only differ for resolution inferences. But more specifically, they only differ on resolution inferences, where the resolved literal is colored. However here, no grey literals are lost. \square

Note that in PI^* , we can conveniently reason about the occurrence of terms as no terms are lost throughout the extraction. However Lemma ?? allows us to transfer results about grey literals to PI . We can also give similar results about general literals and equalities occurring in the resolution refutation:

$\langle \text{lemma:grey_lits_all_in_PI} \rangle$ **Lemma 7.** *If there is a grey literal λ in a clause C of a resolution refutation π , then a successor of λ occurs in $\text{PI}(\pi)$.*

Proof. Immediate by the definition of PI . \square

$\langle \text{lemma:equalities_all_in_PI} \rangle$ **Lemma 8.** *For every equality $s = t$ of a clause in a resolution refutation π , a successor of $s = t$ occurs in $\text{PI}(\pi)$.*

Proof. Equalities in clauses are only removed by means of paramodulation and as π derives the empty clause, all equalities are removed eventually. For any paramodulation inference ι using the equality $s = t$, $\text{PI}_{\text{step}}(\iota, I_1, I_2)$ contains $s = t$. \square

We now make some considerations in the form of four lemmata about the construction of terms of certain shapes in the context of interpolant extraction. In the following, we abbreviate $\text{PI}_{\text{step}}^{\circ}(\iota, \text{PI}^*(C_1), \dots, \text{PI}^*(C_n)) \vee C^{\circ}$ by χ .

$\langle \text{lemma:var_below_phi_symbol} \rangle$ **Lemma 9.** *Let ι be a resolution or factorisation inference in a refutation of $\Gamma \cup \Delta$. Suppose that a variable u occurs directly below a Φ -symbol in $\chi\sigma_{(0,i)}$ for $i \geq 1$. Then at least one of the following statements holds:*

- $\langle 14_1 \rangle$ 1. *The variable u occurs directly below a Φ -symbol in $\chi\sigma_{(0,i-1)}$.*
- $\langle 14_5 \rangle$ 2. *The variable u occurs at a grey position in a grey literal or at a grey position in an equality in $\chi\sigma_{(0,i)}$.*
- $\langle 14_2 \rangle$ 3. *There is a variable v such that*
 - u occurs grey in $v\sigma_i$ and*
 - v occurs in $\chi\sigma_{(0,i-1)}$ directly below a Φ -symbol as well as directly below a Ψ -symbol*

Proof. We consider the different situations under which the situation in question is introduced by means of unification:

- There is already a literal in $\chi\sigma_{(0,i-1)}$ where u occurs directly below a Φ -symbol and σ_i does not change this. Then clearly 1 is the case.
- There is a variable v in $\chi\sigma_{(0,i-1)}$ such that $v\sigma_i$ contains u directly below a Φ -symbol. As then v is unified with the term $v\sigma_i$, $v\sigma_i$ must occur in $\chi\sigma_{(0,i-1)}$, which implies that 1 is the case.
- There is a variable v which occurs directly below a Φ -symbol such that u occurs grey in $v\sigma_i$.

Hence in the resolved or factorised literals λ and λ' in $\chi\sigma_{(0,i-1)}$, there is a position p such that without loss of generality $\lambda|_p = v$ and u occurs grey in $\lambda'|_p$. Note that due to the definition of the unification algorithm, λ and λ' must coincide on the path to p .

By Proposition 5, λ and λ' occur in χ irrespective of their coloring.

We distinguish cases based on the position p :

- Suppose that p occurs directly below a Φ -symbol. Then as u occurs grey in $\lambda|_p$, u occurs directly below a Φ -symbol in $\chi\sigma_{(0,i-1)}$ and 1 is the case.
- Suppose that p occurs directly below a Ψ -symbol. Then v occurs directly below a Ψ -symbol in $\lambda|_p$ and clearly 3 is the case.
- Suppose that p does not occur directly below a colored symbol. Then p does not occur below any colored symbol, hence u is contained in a grey literal in a grey position in $\chi\sigma_{(0,i-1)}$. As σ_i is trivial on u , this occurrence of u also is present in $\chi\sigma_{(0,i)}$ and hence 2 is the case. \square

`(var_below_phi_symbol_paramod)` **Lemma 10.** *Let ι be a paramodulation inference in a refutation of $\Gamma \cup \Delta$. Suppose that a variable u occurs directly below a Φ -symbol in $\chi\sigma_{(0,i)}$ for $i \geq 1$. Then at least one of the following statements holds:*

- $\langle 15_1 \rangle$ 1. *The variable u occurs directly below a Φ -symbol in $\chi\sigma_{(0,i-1)}$.*
- $\langle 15_5 \rangle$ 2. *The variable u occurs at a grey position in a grey literal or at a grey position in an equality in $\chi\sigma_{(0,i)}$.*
- $\langle 15_2 \rangle$ 3. *There is a variable v such that*
 - *u occurs grey in $v\sigma_i$ and*
 - *v occurs in $\chi\sigma_{(0,i-1)}$ directly below a Φ -symbol as well as directly below a Ψ -symbol*

Proof. Let ι be a paramodulation inference of the clauses $C_1 : r_1 = r_2 \vee D$ and $C_2 : E[r]_p$ with $\sigma = \text{mgu}(\iota) = \text{mgu}(r_1, r)$ yielding $C : (D \vee E[r_2]_p)\sigma$. We consider the different situations under which the situation in question is introduced by means of unification:

- There is already a literal in $\chi\sigma_{(0,i-1)}$ where u occurs directly below a Φ -symbol and σ_i does not change this. Then 1 is the case.
- There is a variable v in $\chi\sigma_{(0,i-1)}$ such that $v\sigma_i$ contains u directly below a Φ -symbol. As then v is unified with the term $v\sigma_i$, $v\sigma_i$ must occur in $\chi\sigma_{(0,i-1)}$, which implies that 1 is the case.
- The variable u occurs grey in r_2 and p in E is directly below a Φ -symbol. But then u occurs grey in an equality in $\chi\sigma_{(0,i-1)}$ and as σ_i is trivial on u also in $\chi\sigma_{(0,i)}$, hence 2 holds.
- Suppose that some variable v occurs directly below a Φ -symbol in $\chi\sigma_{(0,i-1)}$ such that u occurs grey in $v\sigma_i$. Then by the definition of the unification algorithm, there exists a position q such that one of $r_1|_q$ and $r|_q$ is v and the other one contains a grey occurrence of u .

We distinguish cases based on the position q :

- Suppose that q occurs directly below a Φ -symbol. Then clearly 1 is the case.
- Suppose that q occurs directly below a Ψ -symbol. Then as the variable v also occurs directly below a Φ -symbol and u occurs grey in $v\sigma_i$, 3 is the case.

- Suppose that q is a grey position. Then 2 is the case: Either u occurs grey in r_1 in $\chi\sigma_{(0,i-1)}$ and then also in $\chi\sigma_{(0,i)}$, or otherwise v occurs grey in r_1 in $\chi\sigma_{(0,i-1)}$, but as $v\sigma_i$ contains u grey, then u occurs grey in the successor of r_1 in $\chi\sigma_{(0,i)}$. \square

`<lemma:col_change>` **Lemma 11.** *Let ι be an inference of a resolution refutation of $\Gamma \cup \Delta$. Suppose that a variable u occurs directly below a Φ -symbol as well as directly below a Ψ -symbol in $\chi\sigma_{(0,i)}$. Then u occurs grey in a grey literal or grey in an equality in $\chi\sigma_{(0,i)}$.*

Proof. We proceed by induction over the refutation. As the original clauses each contain symbols of at most one color, the base case is trivially true.

For the induction step, suppose that an inference makes use of the clauses C_1, \dots, C_n and that the lemma holds for $\text{PI}^*(C_j) \vee C_j$ for $1 \leq j \leq n$.

Note that then, the lemma holds for $\chi = \text{PI}_{\text{step}}^{*\circ}(\iota, \text{PI}^*(C_1), \dots, \text{PI}^*(C_n)) \vee C^\circ$. This is because as all clauses are variable-disjoint, no variable can occur χ both directly below a Φ -symbol as well as directly below a Ψ -symbol if this was not the case in $\text{PI}^*(C_j) \vee C_j$ for some j , for which the lemma by assumption holds. Furthermore, by the definition of PI^* , every literal which occurs in $\text{PI}^*(C_j) \vee C_i$ for some j occurs in χ .

Hence it remains to show that the lemma holds for $\chi\sigma = \chi\sigma_0 \dots \sigma_m$, which we do by induction over i for $1 \leq i \leq m$. Suppose that the lemma holds for $\chi\sigma_{(0,i-1)}$ and in $\chi\sigma_{(0,i)}$, the variable u occurs directly below a Φ -symbol as well as directly below a Ψ -term.

Then by the lemmata 9 and 10, we can deduce that one of the following statements holds for $\Omega = \Phi$ as well as $\Omega = \Psi$. We denote case j for $\Omega = \Phi$ by j^Φ and for $\Omega = \Psi$ by j^Ψ .

- (16_1) 1. The variable u occurs directly below a Ω -symbol in $\chi\sigma_{(0,i-1)}$.
- (16_4) 2. The variable u occurs at a grey position in a grey literal or at a grey position in an equality in $\chi\sigma_{(0,i)}$.
- (16_2) 3. There is a variable v such that
 - u occurs grey in $v\sigma_i$ and
 - v occurs in $\chi\sigma_{(0,i-1)}$ directly below a Φ -symbol as well as directly below a Ψ -symbol

If 2^Φ or 2^Ψ is the case, we clearly are done. On the other hand 3^Φ or 3^Ψ is the case, then by the induction hypothesis, v occurs grey in a grey literal or grey in an equality in $\chi\sigma_{(0,i-1)}$. As u occurs grey in $v\sigma_i$, we obtain that then, u occurs grey in a grey literal or grey in an equality in χ .

Hence the only remaining possibility is that both 1^Φ and 1^Ψ hold. But then u occurs directly below a Φ -symbol as well as below a Ψ -symbol in $\chi\sigma_{(0,i-1)}$ and again by the induction hypothesis, we obtain that u occurs grey in a grey literal or grey in an equality in $\chi\sigma_{(0,i-1)}$ and as σ_i is trivial on u , the same occurrence of u is present in $\chi\sigma_{(0,i)}$. \square

`<lemma:subterm_in_grey_lit_star>` **Lemma 12.** *If $\text{PI}^*(C) \vee C$ for a clause C in a resolution refutation of $\Gamma \cup \Delta$ contains a maximal colored occurrence of a Φ -term $t[s]$, which contains a maximal Ψ -colored term s , then s occurs grey in a grey literal or grey in an equality in $\text{PI}^*(C) \vee C$.*

Proof. We proceed by induction over the resolution refutation. As the original clauses each contain symbols of at most one color, the base case is trivially true.

The induction step is laid out similarly as in the proof of Lemma 11. We suppose that an inference makes use of the clauses C_1, \dots, C_n and that the lemma holds for $\text{PI}^*(C_j) \vee C_j$ for $1 \leq j \leq n$. Then the lemma holds for $\chi = \text{PI}_{\text{step}}^{*\circ}(\iota, \text{PI}^*(C_1), \dots, \text{PI}^*(C_n)) \vee C^\circ$ as no new terms are introduced in χ and all literals from $\text{PI}^*(C_j) \vee C_j$ for $1 \leq j \leq n$ occur in χ .

It remains to show that the lemma holds for $\chi\sigma = \chi\sigma_0 \dots \sigma_m$, which we do by induction over i for $0 \leq i \leq m$. We distinguish on the situation under which a unification leads to the term $t[s]$.

- Suppose for some variable u that $u\sigma_i$ contains $t[s]$. Then u is unified with a term which contains $t[s]$ and occurs in $\chi\sigma_{(0, i-1)}$. Hence by the induction hypothesis, s occurs grey in a grey literal or grey in an equality in $\chi\sigma_{(0, i-1)}$ and, as σ_i does not change this, also in $\chi\sigma_{(0, i)}$.
- Otherwise there is a variable u which occurs directly below a Φ -symbol and $v\sigma_i$ contains a grey occurrence of s . We distinguish based on the occurrences of u in $\chi\sigma_{(0, i-1)}$:
 - Suppose that u occurs in $\chi\sigma_{(0, i-1)}$ in grey a grey literal or grey in an equality. Then clearly we are done.
 - Suppose that u occurs in $\chi\sigma_{(0, i-1)}$ directly below a Ψ -symbol. Then by Lemma 11, u occurs grey in a grey literal or grey in an equality in $\chi\sigma_{(0, i-1)}$, whose successor in $\chi\sigma_{(0, i)}$ is an occurrence of s of the same coloring.
 - Suppose that u occurs in $\chi\sigma_{(0, i-1)}$ only directly below a Ψ -symbol. Here, we differentiate between the types of inference of the current induction step:
 - * Suppose that the inference of the current induction step is a resolution or a factorisation inference. As u occurs grey in $v\sigma_i$, there is a position p such that for the resolved or factorised literals λ and λ' it holds without loss of generality that $\lambda|_p = u$ and s occurs grey in $\lambda'|_p$. Note that λ and λ' agree on the path to p , including the predicate symbol.. Now as by assumption u only occurs directly below a Φ -symbol, so must s . But then s occurs directly below a Φ -symbol in $\chi\sigma_{(0, i-1)}$ and we get the result by the induction hypothesis.
 - * Suppose that the inference of the current induction step is a paramodulation inference. Let it use the the clauses $C_1 : r_1 = r_2 \vee D$ and $C_2 : E[r]_p$ with $\sigma = \text{mgu}(\iota) = \text{mgu}(r_1, r)$ to yield $C : (D \vee E[r_2]_p)\sigma$. As u is affected by σ_i , it must occur in r_1 or r . Let \hat{u} refer to this occurrence.
 - Suppose that \hat{u} occurs directly below a Φ -colored function symbol. If \hat{u} is contained in r_1 , then s must be contained in r directly below a Φ -symbol as r_1 and r are unifiable. We then get the result by the induction hypothesis.

If otherwise \hat{u} is contained in r , then there are two possibilities for the occurrence of s in r_1 :

Either \hat{u} occurs in a Φ -colored function symbol in r . Then s occurs in a Φ -colored function symbol in r_1 and we get the result by the induction hypothesis.

Otherwise \hat{u} occurs grey in r , but r occurs directly below a Φ -colored function symbol in E . Then however, as r and r_1 are unifiable, s must occur grey in r_1 and hence grey in an equality.

· Suppose that \hat{u} occurs directly below a Φ -colored predicate symbol.

Then as the equality predicate is not considered to be colored, u must occur grey in r . But then as r_1 and r are unifiable, s must occur grey in r_1 and hence grey in an equality. \square

(lemma:subterm_in_grey_lit) **Lemma 13.** *Same as 12 with PI in place of PI^* .*

Proof. Note that PI and PI^* only differ with respect to colored literals. Let C be a clause of a resolution refutation of $\Gamma \cup \Delta$. As the literals occurring in $\text{PI}(C)$ are a subset of the literals occurring in $\text{PI}^*(C)$, the lemma prerequisites hold true only for terms for which they also hold in $\text{PI}^*(C)$.

Therefore we can deduce that if a maximal colored Φ -term $t[s]$ containing a maximal Ψ -colored term s occurs in $\text{PI}(C) \vee C$, then $t[s]$ also occurs in $\text{PI}^*(C) \vee C$ and by Lemma 12, the term s occurs grey in a grey literal or grey in an equality in $\text{PI}^*(C) \vee C$. However by Lemma 6, all grey literals and equalities of $\text{PI}^*(C)$ also occur in $\text{PI}(C)$. \square

(lt_in_grey_lit_then_quant_alt) **Lemma 14.** *If a term with n color alternations occurs in a grey literal or in an equality in $\text{PI}(C) \vee C$ for a clause C , then the interpolant I produced in Theorem ?? contains at least n quantifier alternations.*

Proof. We perform an induction on n and show the strengthening that the quantification of the lifting variable which replaces a term with n color alternations is required to be in the scope of the quantification of $n - 1$ alternating quantifiers.

For $n = 0$, no colored terms occur in I and hence also no quantifiers. Moreover for $n = 1$, there are terms of one color which evidently require at least one quantifier.

Suppose that the statement holds for $n - 1$ for $n > 1$ and that a term t with $\text{col-alt}(t) = n$ occurs in $\text{PI}(C)$. We assume without loss of generality that t is a Φ -term. Then t contains some Ψ -colored term s with $\text{col-alt}(s) = n - 1$ and by Lemma 13, s occurs grey in a grey literal or an equality in $\text{PI}(C) \vee C$. By Lemma 7 and Lemma 8, a successor of s occurs in $\text{PI}(\pi)$. Note that as s occurs in a grey position, any successor of s also occurs in a grey position.

By the induction hypothesis, the quantification of the lifting variable for s requires $n - 1$ alternated quantifiers. As s is a subterm of t and t is lifted, t must be quantified in the scope of the quantification of s , and as t and s are of different color, their quantifier type is different. Hence the quantification of the lifting variable for t requires n quantifier alternations. \square

$\langle \text{prop:color_alt_eq_quant_alt} \rangle$ **Proposition 15.** *If a term with n color alternations occurs in $\text{PI}(C) \vee C$ for a clause C , then the interpolant I produced in Theorem ?? contains at least $n - 1$ quantifier alternations.*

Proof. By Lemma 13, a term with $n - 1$ color alternations occurs in a grey literal or an equality in $\text{PI}(C) \vee C$. Lemma 14 gives the result. \square

We present an example which illustrates that the occurrence of a term with n color alternations in $\text{PI}(C) \vee C$ for a clause C can lead to an interpolant with $n - 1$ quantifier alternations (but no less as Proposition 15 shows).

Example 16. Let $\Gamma = \{\neg P(a)\}$ and $\Delta = \{P(x) \vee Q(f(x)), \neg Q(y)\}$. Consider the following refutation of $\Gamma \cup \Delta$:

$$\frac{\frac{\neg P(a) \mid \perp \quad P(x) \vee Q(f(x)) \mid \top}{Q(f(a)) \mid \neg P(a)} \text{res}_{x \mapsto a} \quad \neg Q(y) \mid \top}{\square \mid \neg P(a)} \text{res}_{y \mapsto f(a)}$$

In this example, Theorem ?? yields the interpolant $I \equiv \exists y_a \neg P(y_a)$ with $\text{quant-alt}(I) = 1$. The existence of the term $f(a)$ with $\text{col-alt}(f(a)) = 2$ in a clause of the refutation implies that $\text{quant-alt}(I) \geq 1$. \triangle

3 LI

Conjectured Lemma 17. *Let C be a clause of a resolution refutation of $\Gamma \cup \Delta$. Then $\ell[\text{PI}(C)]$ and $\ell[\text{LI}(C)]$ agree if we disregard the quantifiers.*

Proof. Note that $\text{PI}(C)$ and $\text{LI}(C)$ only differ with respect to the lifting conditions of LI. If one of them becomes true for a term t in $\text{LI}(C)$, it is lifted, but this does not affect the lemma statement. \square

the only difference is that it might be t in $\text{PI}(C)$ where it's x_t in $\text{LI}(C)$, but this does not interfere with the lemmata on quantifier alternation and stuff.

MOSTLY WRONG

- a term in $\text{LI}(C)$ is not lifted if a subterm of it is still in C
- If a term is lifted, then all its possible and future subterms are already in $\text{LI}(C)$, hence all of them are actually there.
- all terms are lifted eventually
- at the stage where a term is lifted, all subterms are there

OR:

• WRONG: OTHER WAY AROUND!!!

For a term t in $\text{LI}(C) \vee C$, if a subterm s of t occurs in $\text{LI}(C) \vee C$ which is not lifted, then t is not lifted.

• WRONG: OTHER WAY AROUND!!!

When a term is lifted, then all its subterms in $\text{LI}(C) \vee C$ are already lifted or are lifted right now.

Conjectured Lemma 18. *If a term t in $\text{LI}^\bullet(C)$ is lifted in $\text{LI}(C)$, then no subterms of t are lifted in $\text{LI}^\bullet(C)$ (i.e. have been lifted in previous stages).*

- A subterm s of t is lifted either when t is lifted or later. In any case, the quantifier for t occurs in the scope of the quantifier of s

3.1 Big picture

- if max col term t in $\text{LI}(C)$ not lifted, then t also occurs in $\text{PI}(C)$ (1).
- for every max col term t in $\text{PI}(C)$, either t or $\ell[t]$ occurs in $\text{LI}(C)$ (2)
- if max col term t in $\text{LI}(C)$ not lifted, then all subterms s of t are not lifted as well. (3)
- Hence:

Supp max col t in $\text{LI}(C)$ is lifted in $\text{LI}(C)$ but not in $\text{LI}^\bullet(C)$.

Then t also occurs in $\text{PI}(C)$ by (1).

Then by that lemma, any subterm s of t occurs in $\text{PI}(C)$ grey in grey lit or grey in equality. Hence it occurs also in $\text{LI}(C)$, either lifted or not lifted (2).

By (3), it is not lifted.

It however will eventually be lifted (cf. TODO), so the quantification of t will be in the scope of the quantifier for s .

(lemma:pi_li_1) **Conjectured Lemma 19.** *Let C be a clause in a resolution refutation of $\Gamma \cup \Delta$. If a maximal colored term t occurs in $\text{LI}^\bullet(C)$ in a literal λ at position p , then t also occurs in a corresponding literal at position p in $\text{PI}(C)$.*

Proof. **TODO:**

□

(lemma:pi_li_2) **Conjectured Lemma 20.** *Let C be a clause in a resolution refutation of $\Gamma \cup \Delta$. If a maximal colored term t occurs in $\text{PI}(C)$ in a literal λ at position p , then either t or z_t occurs in a corresponding literal at position p in $\text{LI}(C)$. **TODO: OR $\text{LI}^\bullet(C)$***

Proof. **TODO:**

□

(lemma:pi_li_3) **Conjectured Lemma 21.** *If a maximal colored term t in $\text{LI}^\bullet(C)$ is not lifted, then no subterm s of t , which occurs in $\text{LI}^\bullet(C) \vee C$, is lifted.*

Proof.

□

Conjectured Lemma 22. ***TODO: merge with 5.3. of thesis (lemma:lifting_conditions)***

(lemma:pi_li_3) *Let C be a clause of a resolution refutation such that $\text{LI}^\bullet(C)$ contains a maximal colored Φ -term t which is lifted in $\text{LI}(C)$. Suppose that t contains a Ψ -colored subterm s . Then s occurs as a subterm of t in $\text{LI}^\bullet(C)$.*

Proof. By the construction of LI, this lemma is only violated if the term s or a respective predecessor is lifted due to fulfilling one of the lifting conditions.

For the sake of contradiction, suppose that this is the case in the inference creating the clause C' . Let s' and t' be the respective predecessors of s and t in C'

- Suppose that s' is lifted due to containing a variable which does not occur in C' . Then as s' is a subterm of t' , t' contains this variable as well and therefore is lifted in $\text{LI}(C')$, contradicting the assumption.

ICI

□

(lemma:grey_lits_all_in_LI) **Conjectured Lemma 23.** *Let C be a clause in a resolution refutation π of $\Gamma \cup \Delta$. Then for any grey literal λ in $\text{LI}^\bullet(C) \vee C$, a successor of λ occurs in $\text{LI}^\bullet(C') \vee C'$, where C' is a successor of C .*

Proof. TODO:

□

(lemma:equalities_all_in_LI) **Conjectured Lemma 24.** *Let C be a clause in a resolution refutation π of $\Gamma \cup \Delta$. Then for any equality $s = t$ in $\text{LI}^\bullet(C) \vee C$, a successor of $s = t$ occurs in $\text{LI}^\bullet(C') \vee C'$, where C' is a successor of C .*

Proof. TODO:

□

Conjectured Lemma 25. *If a term t with $\text{col-alt}(t) = n$ occurs grey in a grey literal or grey in an equality $\text{LI}^\bullet(C)$ for some C , then $\text{LI}(\pi)$ has at least n quantifier alternations.*

Proof. looks good

We show that if t occurs in $\text{LI}^\bullet(C)$, then the “successor” of $\text{LI}^\bullet(C)$ which is contained in $\text{LI}(\pi)$ is enclosed by n alternating quantifiers.

By the Lemmata 23 and 24, a successor of t occurs in $\text{LI}^\bullet(C') \vee C'$ for any successor C' of C . Let C' be the successor of C such that $\text{LI}^\bullet(C')$ contains a successor of t which is not a lifting variable and $\text{LI}(C')$ contains a successor of t which is a lifting variable. As all colored terms are lifted eventually, such a clause C' must occur.

By Lemma 19, t occurs at the same position in $\text{PI}(C')$. We assume with loss of generality that t is a Φ -term. Then t contains some Ψ -colored term s with $\text{col-alt}(s) = n - 1$. By Lemma 13, s occurs grey in a grey literal or grey in an equality in $\text{PI}(C') \vee C'$.

By Lemma 20, then either s or z_s occurs in $\text{LI}(C')$ TODO: OR $\text{LI}^\bullet(C')$ at a corresponding literal and position. As t is not lifted in $\text{LI}^\bullet(C') \vee C'$ and s is a subterm of t , by Lemma 22, s is also not lifted in $\text{LI}^\bullet(C') \vee C'$.

As t is lifted in this step, one quantifier is added. Note that even if s is also lifted in this step, then it is lifted after t , or otherwise it is lifted at a later stage.

Let $\text{LI}^\bullet(C)'$ be the “formula right after t has been lifted and quantified, but not the remaining ones of this stage”.

Then s occurs in $\text{LI}^\bullet(C)' \vee C$. Hence by the induction hypothesis, the successor of $\text{LI}^\bullet(C)'$ in $\text{LI}(\pi)$ is enclosed by $n - 1$ quantifier alternations. As s and t are of opposite colores, their quantifier type is different and the lifting

and quantification of t adds one more quantifier alternation to $\text{LI}^\bullet(C)$. Hence the successor of $\text{LI}^\bullet(C)$ in $\text{LI}(\pi)$ is enclosed by n quantifier alternations. \square

Proof. Supp t not lifted in $\text{LI}^\bullet(C)$ but in $\text{LI}(C)$ and $\text{col-alt}(t) = 1$. Then t is colored and $\text{LI}(C)$ (and $\text{LI}(\pi)$) has one color alternation

Supp t not lifted in $\text{LI}^\bullet(C)$ but in $\text{LI}(C)$ and $\text{col-alt}(t) = n$. Then by Lemma 22, no subterm of t is lifted. By TODO, the subterm s of t occurs in $\text{LI}^\bullet(C)$ and by the argument above is not lifted. By the induction hypothesis, $n - 1$ quantifier alternations are added to the “current lifted interpolant”.

as s and t of different color, different quantifier. hence the lifting of t adds one quantifier alternation, plus the $n - 1$ which come later, so we have at least n in the end. \square

Conjectured Lemma 26. *Let π be a refutation of $\Gamma \cup \Delta$ and C a clause in π . Let t be a term with n color alternations such that t is not a lifting variable. If t occurs in a grey literal or in an equality in $\text{LI}^\bullet(C) \vee C$, then $\text{LI}(\pi)$ contains at least n quantifier alternations.*

Proof. ONLY IND STEP FOR NOW:

By the Lemmata 23 and 24, a successor of t occurs in $\text{LI}^\bullet(C') \vee C'$ for any successor C' of C . Let C' be the successor of C such that $\text{LI}^\bullet(C')$ contains a successor of t which is not a lifting variable and $\text{LI}(C')$ contains a successor of t which is a lifting variable. As all colored terms are lifted eventually, such a clause C' must occur.

By Lemma 19, t occurs at the same position in $\text{PI}(C')$. We assume with loss of generality that t is a Φ -term. Then t contains some Ψ -colored term s with $\text{col-alt}(s) = n - 1$. By Lemma 13, s occurs grey in a grey literal or grey in an equality in $\text{PI}(C') \vee C'$.

By Lemma 20, then either s or z_s occurs in $\text{LI}(C')$ **TODO: OR $\text{LI}^\bullet(C')$** at a corresponding literal and position. As t is not lifted in $\text{LI}^\bullet(C') \vee C'$ and s is a subterm of t , by Lemma 22, s is also not lifted in $\text{LI}^\bullet(C') \vee C'$.

As however s is a colored term which occurs in a grey literal or an equality in $\text{LI}^\bullet(C') \vee C'$, by the induction hypothesis, $\text{LI}(\pi)$ contains at least $n - 1$ quantifier alternations. **TODO: weak!**

OTHER STATEMENT: n col alt $\Rightarrow n$ nested liftings still to come

\square

TODO: show that terms in $\text{PI}(C)$ actually occur in $\text{LI}(C)$, hence $\text{PI}(C)$ and $\text{LI}(C)$ have same quant alts