

Interpolation in First-Order Logic with Equality

Masterstudium:
Computational Intelligence


Bernhard Mallinger

Technische Universität Wien
Institut für diskrete Mathematik und Geometrie
Forschungsgruppe: Computational Logic
Betreuer: Ass.Prof. Stefan Hetzl

Craig Interpolation

Theorem (Craig). Let A and B be first-order formulas such that $\models A \supset B$ where A contains red and gray symbols and B contains of blue and gray symbols. Then there is an interpolant I containing of gray symbols for A and B such that:

- ▶ $\models A \supset I$
- ▶ $\models I \supset B$



⇒ Interpolants give a concise logical summary of the implication

Applications of Craig Interpolation

Theoretical:

- ▶ Proof of Beth's Definability Theorem

Practical:

- ▶ Program analysis: Detect loop invariants
- ▶ Model checking: Overapproximate set of reachable states

Aim and Scope of the Thesis

Give comprehensive account of existing techniques and extend them:

- ▶ Model-theoretic proof
- ▶ Reduction to first-order logic without equality
- ▶ Interpolant extraction from resolution proofs

Model-theoretic proof

- ▶ Non-constructive proof:
 - ▶ Let T_A and $T_{\neg B}$ be theories extending A and $\neg B$
 - ▶ Build model from maximal consistent intersection of T_A and $T_{\neg B}$ (assuming the non-existence of interpolants)
 $\Rightarrow A \wedge \neg B$ satisfiable
- ▶ Related to Robinson's Joint Consistency Theorem

Reduction to first-order logic without equality [1]

Translate equality and function symbols:


$$\begin{aligned} (P(c))^* &\equiv \exists x (C(x) \wedge P(x)) \\ (P(f(c)))^* &\equiv \exists x (\exists y (C(y) \wedge F(y, x)) \wedge P(x)) \\ (s = t)^* &\equiv E(s, t) \end{aligned}$$

Add theory of equality:


$$\varphi \rightarrow T_E \supset \varphi^*$$

⇒ Then calculate interpolant in reduced logic

Interpolant extraction from proofs in two phases [2]



Interpolant extraction from proofs in one phase



Contributions

- ▶ We introduced the one phase-approach.
- ▶ We showed that the number of quantifier alternations in the interpolant essentially corresponds to the number of color alternations in terms.

References

- [1] William Craig.
Linear Reasoning. A New Form of the Herbrand-Gentzen Theorem.
Journal of Symbolic Logic, 22(3):250–268, 1957.
- [2] Guoxiang Huang.
Constructing Craig Interpolation Formulas.
In *Proc COCOON '95*, p. 181–190, 1995.