
Contents

Contents	1
1 Introduction	3
1.1 Preliminaries	3
1.1.1 Unification	4
1.2 Craig Interpolation	5
2 Calculi	8
2.1 Resolution	8
2.2 Resolution and Interpolation	9
2.2.1 Interpolation and Skolemisation	10
2.2.2 Interpolation and structure-preserving Normal Form Transformation	11
2.3 Sequent Calculus	14
3 Reduction to First-Order Logic without Equality	17
3.1 Translation of formulas	18
3.2 Computation of interpolants	21
3.3 Proof by reduction	27
4 Interpolant extraction from resolution proofs in two phases	29
4.1 Layout of the proof	29
4.2 Extraction of propositional interpolants	30
4.3 Lifting of colored symbols	31
4.4 Symmetry of the extracted interpolants	33
4.5 Main lemma	35

4.6	Quantifying over lifting variables	39
A	WT: Huang's proof	40
A.1	Propositional interpolants	40
A.2	Propositional refutations	44
A.3	Lifting of colored symbols	46
	Bibliography	53

Introduction

1.1 Preliminaries

this section contains all the required notation but will just be written up nicely in the final version

The language of a first-order formula A is denoted by $L(A)$ and contains all predicate, constant and function symbols that occur in A . These are also referred to as the *non-logical symbols* of A . The *logical symbols* on the other hand include all logical connectives, quantifiers, the equality symbol ($=$) as well as symbols denoting truth (\top) and falsity (\perp).

For formulas A_1, \dots, A_n , $L(A_1, \dots, A_n) = \bigcup_{1 \leq i \leq n} L(A_i)$.

A term s is a subterm of a term t if s occurs in t . s is a strict subterm of t if s is a subterm of t and $s \neq t$.

An occurrence of Φ -term is called *maximal* if it does not occur as subterm of another Φ -term. An occurrence of a colored term t is a maximal colored term if it does not occur as subterm of another colored term.

We denote x_1, \dots, x_n by \bar{x} .

For a set of formulas Φ , $\neg\Phi$ denotes $\{\neg A \mid A \in \Phi\}$.

A substitution is a mapping of variables to terms. It is denoted by $\phi[x/t]$, where ϕ is a formula or term where each occurrence of the variable x is replaced by the term t . A substitution σ is called trivial on x if $x\sigma = x$. Otherwise it is called non-trivial.

An *abstraction* on the other hand is a mapping of terms to variables. It is denoted by $\phi\{t/x\}$, where ϕ is a formula or term where each occurrence of the term t is replaced by the variable x .

colors
are only
defined
later

A term s is an *instance* of a term t if there exists a substitution σ such that $t\sigma = s$. If s is an instance of t , then t is an *abstraction* of s . Note that the abstraction- and instance-relation are reflexive. s is a *proper* instance (abstraction) of t if s is an instance (abstraction) of t and $s \neq t$.

The length of a term or formula ϕ is the number of logical and non-logical symbols in ϕ .

$A[s]_p$ denotes A with an occurrence of s at position p .

$A[s]$ denotes A where the term s occurs on some set of positions Φ . $A[t]$ denotes $A[s]$ where on each position in Φ , s has been replaced by t . Due to its vagueness, this notation is mostly used in order to emphasis that the term s does occur in A in some way.

TODO: define Σ as subformula set; possibly remove definition in chapter 2

TODO: define superterm the same way as subterm

TODO: define \equiv as syntactic equality. Define also \Leftrightarrow , \leftrightarrow .

TODO: define what we mean by model and free variables. (need universal quantification of free vars)

need it for paramodulation proof

proposal: free vars are implicitly universally quantified

this should work out in general, and when needed, we add an explicit interpretation of some free variables

$M \models P(x)$ is short for $M \models \forall \text{FV}(P(x)) P(x)$

$M_\alpha \models P(x)$ is short for M entails $P(x)$ with all variables interpreted by α interpreted like that (basically $P(x)$ with “substitution” α)

TODO: define ground, non-ground

TODO: define set-notation of unifiers

TODO: define infinite-domain unifiers

TODO: define range and domain of substitutions

TODO: define prenex formulas with matrix and prefix

TODO: define prefix of term position: e.g. u in $f(c, g(u, x, h(a)))$ has the prefix $f(., g(., ., .))$, or possible written as sequence of symbols (algo: always go to parent starting at u)

TODO: abstraction has 2 contradictory definitions above

TODO: free vars: FV and possibly LV: free lifting vars. probably FV does not include LV.

1.1.1 Unification

We specify the unification algorithm which is used in this thesis in order enable formal reasoning.

Definition 1 (Unification algorithm). Let id denote the identity function and **fail** be returned by mgu in case the arguments are not unifiable to signify that the mgu of the given arguments is not defined. We treat constants as 0-ary functions. Let s and t denote terms and x a variable.

The most general unifier mgu of two literals $l = A(s_1, \dots, s_n)$ and $l' = A(t_1, \dots, t_n)$ is defined to be $\text{mgu}(\{(s_1, t_1), \dots, (s_n, t_n)\})$.

The mgu for a set of pairs of terms T is defined as follows:

$\text{mgu}(\emptyset) = \text{id}$

$$\text{mgu}(\{t\} \cup T) = \begin{cases} \text{fail} & \text{if } t = (x, s) \text{ or } t = (s, x) \text{ and } x \\ & \text{occurs in } s \text{ but } x \neq s \\ \text{mgu}(T[x/s])[x/s] \cup \{x \mapsto s\} & \text{if } t = (x, s) \text{ or } t = (s, x) \text{ and } x \\ & \text{does not occur in } s \text{ or } x = s \\ \text{fail} & \text{if } t = (f(s_1, \dots, s_n), g(s_1, \dots, s_n)) \\ & \text{with } f \neq g \\ \text{mgu}(T \cup \{(s_1, t_1), \dots, (t_n, s_n)\}) & \text{if } t = (f(s_1, \dots, s_n), f(t_1, \dots, t_n)) \\ \text{mgu}(T) & \text{if } t = (s, s) \end{cases}$$

For $\sigma = \text{mgu}(l, l')$ for two literals l and l' we denote by σ_i the i th substitution, which in the course of the run of the algorithm has been added to σ . We denote composition of substitutions σ_i up to including σ_j by $\sigma_{(i,j)}$. \triangle

Hence for a unifier σ , we have that $\sigma = \sigma_1 \cdots \sigma_n = \sigma_{(i,j)}$, where n is $|\text{dom}(\sigma)|$. Note that $|\text{dom}(\sigma_i)| = 1$ for $1 \leq i \leq n$.

1.2 Craig Interpolation

TODO: write some text about what interpolation means and that we prove more or less only reverse interpolation, but that's fine by the proposition

Definition 2. Let Γ and Δ be sets of first-order formulas. An *interpolant* of Γ and Δ is a first-order formula I such that

1. $\Gamma \models I$
2. $I \models \Delta$
3. $L(I) \subseteq L(\Gamma) \cap L(\Delta)$.

A *reverse interpolant* of Γ and Δ is a first-order formula I such that I meets conditions 1 and 3 of an interpolant as well as:

$$2'. \Delta \models \neg I \quad \triangle$$

Theorem 3 (Interpolation). *Let Γ and Δ be sets of first-order formulas such that $\Gamma \models \Delta$. Then there exists an interpolant for Γ and Δ .*

Theorem 4 (Reverse Interpolation). *Let Γ and Δ be sets of first-order formulas such that $\Gamma \cup \Delta$ is unsatisfiable. Then there exists a reverse interpolant for Γ and Δ .*

Proposition 5. *Theorem 3 and 4 are equivalent.*

Proof. Let Γ and Δ be sets of first-order formulas such that $\Gamma \models \Delta$. Then $\Gamma \cup \neg\Delta$ is unsatisfiable. By Theorem 4, there exists a reverse interpolant I for Γ and $\neg\Delta$. As $\neg\Delta \models \neg I$, we get by contraposition that $I \models \Delta$, hence I is an interpolant for Γ and Δ .

For the other direction, let Γ and Δ be sets of first-order formulas such that $\Gamma \cup \Delta$ is unsatisfiable. Then $\Gamma \models \neg\Delta$, hence by Theorem 3, there exists an interpolant I of Γ and $\neg\Delta$. But as thus $I \models \neg\Delta$, we get by contraposition that $\Delta \models \neg I$, so I is a reverse interpolant for Γ and Δ . \square

As the notions of interpolation and reverse interpolation in this sense coincide, we will in the following only speak of interpolation where will be clear from the context which definition applies.

Lemma 6. *Let $\Gamma, \Gamma', \Delta, \Delta'$ be sets of first order formulas such that $\Gamma \Leftrightarrow \Gamma'$ and $\Delta \Leftrightarrow \Delta'$ and $L(\Gamma) \cap L(\Delta) = L(\Gamma') \cap L(\Delta')$. Then I is an interpolant for Γ and Δ if and only if I is an interpolant for Γ' and Δ' .*

Proof. Clearly $\Gamma \models I$ holds if and only if $\Gamma' \models I$ and similarly $\Delta \models \neg I$ holds if and only if $\Delta' \models \neg I$. As the intersections of the respective languages coincide, the language condition on I is satisfied in both directions. \square

Remark. In Lemma 6, it is not sufficient to require that $\Gamma \Leftrightarrow \Gamma'$ and $\Delta \Leftrightarrow \Delta'$. Consider the example where $\Gamma = \Delta = \{\forall x(x = c)\}$ and $\Gamma' = \Delta' = \{\forall x(x = d)\}$. Then even though Γ and Γ' as well as Δ and Δ' have the same models, $L(\Gamma) \cap L(\Delta) = \{c\}$ whereas $L(\Gamma') \cap L(\Delta') = \{d\}$. Therefore $\forall x(x = c)$ is an interpolant for Γ and Δ but not for Γ' and Δ' . \triangle

In the context of interpolation, every non-logical symbol is assigned a color which indicates its origin(s).

Definition 7 (Coloring). A non-logical symbol is said to be Γ (Δ)-*colored* if it only occurs in Γ (Δ) and *grey* in case it occurs in both Γ and Δ . A symbol is *colored* if it is Γ - or Δ -colored. A term is a Φ -*term* if its outermost symbol is Φ -colored.

A term t is *single-colored* if t is Φ -colored for some Φ and all colored symbols in t are Φ -colored. A term t is *mixed-colored* if t is Φ -colored for some Φ and t contains a term which is colored but not Φ -colored. **TODO: fix either mixed-colored or multi-colored, currently both is used synonymously** Note that a multicolored Φ -term consequently is a term whose outermost symbol is Φ -colored and contains a colored but not Φ -colored subterm.

An occurrence of a term t is called Φ -*colored* if t is contained in a maximal Φ -colored term. t is called *colored* if t is of any color and *grey* otherwise.

A variable is a *color-changing* variable if it occurs both in a single-colored Φ -term and a single-colored Ψ -term in a given context. \triangle

We sometimes use Φ and Ψ as colors to emphasise that the reasoning at hand is valid irrespective of the actual color assignment and solely assuming that $\Phi \neq \Psi$.

Calculi

In this chapter, we introduce the calculi that are used subsequently. These are resolution and sequent calculus.

2.1 Resolution

Resolution calculus, in the formulation as given here, is a sound and complete calculus for first-order logic with equality. Due to the simplicity of its rules, it is widely used in the area of automated deduction.

Definition 8. A *clause* is a finite set of literals. The empty clause will be denoted by \square . A *resolution refutation* of a set of clauses Γ is a derivation of \square consisting of applications of resolution rules (cf. Figure 2.1) starting from clauses in Γ . \triangle

Clauses will usually be denoted by C , D or E , literals by l , l' or λ and positions by p .

$$\begin{aligned}
 \text{Resolution:} \quad & \frac{C \vee l \quad D \vee \neg l'}{(C \vee D)\sigma} \text{ res} \quad \sigma = \text{mgu}(l, l') \\
 \text{Factorisation:} \quad & \frac{C \vee l \vee l'}{(C \vee l)\sigma} \text{ fac} \quad \sigma = \text{mgu}(l, l') \\
 \text{Paramodulation:} \quad & \frac{D \vee s = t \quad E[r]_p}{(D \vee E[t]_p)\sigma} \text{ par} \quad \sigma = \text{mgu}(s, r)
 \end{aligned}$$

Figure 2.1: The rules of resolution calculus

Theorem 9. *A clause set Γ is unsatisfiable if and only if there is resolution refutation of Γ .*

Proof. See [Rob65]. □

Definition 10 (Tree refutations). A resolution refutation is a *tree refutation* if every clause is used at most once. △

The following lemma shows that the restriction to tree refutations does not restrict the calculus given that we allow multisets as initial clause sets.

Lemma 11. *Every resolution refutation can be transformed into a tree refutation.*

Proof. Let π be a resolution refutation of a set of clauses Φ . We show that π can be transformed into a tree refutation by induction on the number of clauses that are used multiple times.

Suppose that no clause is used more than once in π . Then π is a tree refutation.

Otherwise let Ψ be the set of clauses which is used multiple times. Let $C \in \Psi$ be such that no clause $D \in \Psi$ is used in the derivation leading to C . Let χ be the derivation leading to C .

Suppose C is used m times. We create another resolution refutation π' from π which contains m copies of χ and replaces the i th use of the clause C by the final clause of the i th copy of χ , $1 \leq i \leq m$. In order to ensure that the sets of variables of the input clauses are disjoint, we rename the variables in each copy of χ and adapt π' accordingly. Hence π' is a resolution refutation of Φ where $m - 1$ clauses are used more than once. □

2.2 Resolution and Interpolation

In order to apply resolution to arbitrary first-order formulas, they have to be converted to clauses first. This usually makes use of intermediate normal forms which are defined as follows:

Definition 12. A formula is in *Negation Normal Form (NNF)* if negations only occur directly before of atoms. A formula is in *Conjunctive Normal Form (CNF)* if it is a conjunction of disjunctions of literals. △

In this context, the conjuncts of a CNF-formula are interpreted as clauses. A well-established procedure for the translation to CNF is comprised of the following steps:

1. NNF-Transformation
2. Skolemisation
3. CNF-Transformation

Step 1 can be achieved by solely pushing the negation inwards. As this transformation yields logically equivalent formulas without affecting the language, by Lemma 6, the set of interpolants remains unchanged. Step 2 and 3 on the other hand do not produce logically equivalent formulas since they introduce new symbols. In this section, we will show that they nonetheless do preserve the set of interpolants. This fact is vital for the use of resolution-based methods for interpolant computation of arbitrary formulas.

2.2.1 Interpolation and Skolemisation

Skolemisation is a procedure for replacing existential quantifiers by Skolem terms:

Definition 13. Let $V_{\exists x}$ be the set of universally bound variables whose scope includes the occurrence of $\exists x$ in a formula. The skolemisation of a formula A in NNF, denoted by $\text{sk}(A)$, is the result of replacing every occurrence of an existential quantifier $\exists x$ in A by a term $f(y_1, \dots, y_n)$ where f is a new Skolem function symbol and $V_{\exists x} = \{y_1, \dots, y_n\}$. In case $V_{\exists x}$ is empty, the occurrence of $\exists x$ is replaced by a new Skolem constant symbol c .

The skolemisation of a set of formulas Φ is defined to be $\text{sk}(\Phi) = \{\text{sk}(A) \mid A \in \Phi\}$. \triangle

Note that due to the introduction of Skolem symbols, it is not the case that $\Phi \Leftrightarrow \text{sk}(\Phi)$.

Proposition 14. *Let $\Gamma \cup \Delta$ be unsatisfiable. Then I is an interpolant for $\Gamma \cup \Delta$ if and only if it is an interpolant for $\text{sk}(\Gamma) \cup \text{sk}(\Delta)$.*

Proof. Since $\text{sk}(\cdot)$ adds fresh symbols to both Γ and Δ individually, none of them are contained in $L(\text{sk}(\Gamma)) \cap L(\text{sk}(\Delta))$. Therefore the language condition on the interpolant is satisfied in both directions.

We conclude the proof by showing that $\Phi \models A$ iff $\text{sk}(\Phi) \models A$ for $\Phi \in \{\Gamma, \Delta\}$ and $A \in \{I, \neg I\}$.

Suppose that for a model that $M \models \text{sk}(\Phi)$ and $\Phi \models A$. Note that the interpretation of the skolem symbols of $\text{sk}(\Phi)$ in M presents witnesses for the corresponding existential quantifiers in Φ . Hence $M \models \Phi$ and consequently $M \models A$.

On the other hand, suppose that $M \models \Phi$ and $\text{sk}(\Phi) \models A$. We assume that $\text{sk}(\Phi)$ only uses Skolem terms which are fresh with respect to M . Then we can

extend M to a model M' of $\text{sk}(\phi)$ by encoding the witness terms for the existential quantifiers in Φ in the Skolem terms of $\text{sk}(\Phi)$ in M' . Then $M' \models \text{sk}(\Phi)$ and thus $M' \models A$. But as $L(A) \subseteq L(M) \subseteq L(M')$, M and M' agree on the interpretation of A , hence $M \models A$. \square

2.2.2 Interpolation and structure-preserving Normal Form Transformation

In the following, we describe a common method for transforming a formula A without existential quantifiers into CNF while preserving its structure. Note that the restriction to formulas without existential quantifiers can easily be established for arbitrary formulas by means of skolemisation and therefore does not limit the applicability of this procedure.

In the following, we use the notational convention that $\{\bar{y}\} \cup \{\bar{z}\} = \{\bar{x}\}$ expressing the intuition that the free variables \bar{x} of a formula B are comprised of the not necessarily disjoint free variables \bar{y} and \bar{z} of B 's direct subformulas.

Definition 15. For every occurrence of a subformula B of a formula A without existential quantifiers, introduce a new atom $L_B(\bar{x})$, where \bar{x} are the free variables occurring in B . This atom acts as a label for the subformula. For each of them, create a defining clause D_B :

If B is atomic:

$$D_B \equiv \forall \bar{x} (\neg B \vee L_B(\bar{x})) \wedge \forall \bar{x} (B \vee \neg L_B(\bar{x}))$$

If B is of the form $\neg G$:

$$D_B \equiv \forall \bar{x} (L_B(\bar{x}) \vee L_G(\bar{x})) \wedge \forall \bar{x} (\neg L_B(\bar{x}) \vee \neg L_G(\bar{x}))$$

If B is of the form $G \wedge H$:

$$D_B \equiv \forall \bar{x} (\neg L_B(\bar{x}) \vee L_G(\bar{y})) \wedge \forall \bar{x} (\neg L_B(\bar{x}) \vee L_H(\bar{z})) \wedge \forall \bar{x} (L_B(\bar{x}) \vee \neg L_G(\bar{y}) \vee \neg L_H(\bar{z}))$$

If B is of the form $G \vee H$:

$$D_B \equiv \forall \bar{x} (L_B(\bar{x}) \vee \neg L_G(\bar{y})) \wedge \forall \bar{x} (L_B(\bar{x}) \vee \neg L_H(\bar{z})) \wedge \forall \bar{x} (\neg L_B(\bar{x}) \vee L_G(\bar{y}) \vee L_H(\bar{z}))$$

If B is of the form $G \supset H$:

$$D_B \equiv \forall \bar{x} (L_B(\bar{x}) \vee L_G(\bar{y})) \wedge \forall \bar{x} (L_B(\bar{x}) \vee \neg L_H(\bar{z})) \wedge \forall \bar{x} (\neg L_B(\bar{x}) \vee \neg L_G(\bar{y}) \vee L_H(\bar{z}))$$

If B is of the form $\forall xG$:

$$D_B \equiv \forall \bar{x} \forall x (\neg L_B(\bar{x}) \vee L_G(\bar{x}, x)) \wedge \forall \bar{x} \forall x (L_B(\bar{x}) \vee \neg L_G(\bar{x}, x))$$

Let $D_{\Sigma(A)}$ be defined as $\bigwedge_{B \in \Sigma(A)} D_B$ and $\delta(A)$ as $D_{\Sigma(A)} \wedge \forall \bar{x} L_A(\bar{x})$, where $\Sigma(A)$ denotes the set of occurrences of subformulas of A . For a set of formulas without existential quantifiers Φ , let $\delta(\Phi) = \{\delta(B) \mid B \in \Phi\}$. \triangle

Note that each of the D_B is in CNF, hence also $\delta(A)$ for any formula A without existential quantifiers. We continue by working out the logical relations of formulas and their image under A :

Lemma 16. *Let M be a model of $\delta(A)$ for a formula A without existential quantifiers. Then $M \models A$.*

Proof. We show that $M \models B \leftrightarrow L_B(\bar{x})$ for $B \in \Sigma(A)$. As $M \models \delta(A)$ directly implies that $M \models L_A$, this proves the lemma. Note that also $M \models D_{\Sigma(A)}$.

The proof is by induction on the structure of B . For the base case, let B be an atom. Then $D_B \equiv \forall \bar{x} (\neg B \vee L_B(\bar{x})) \wedge \forall \bar{x} (B \vee \neg L_B(\bar{x}))$, which due to $M \models D_B$ immediately yields $M \models B \leftrightarrow L_B(\bar{x})$.

For the induction step, we illustrate a few cases as the remaining ones are similar.

- Suppose B is of the form $\neg G$. Then:

$$D_B \equiv \forall \bar{x} (L_B(\bar{x}) \vee L_G(\bar{x})) \wedge \forall \bar{x} (\neg L_B(\bar{x}) \vee \neg L_G(\bar{x}))$$

By the induction hypothesis, $M \models G \leftrightarrow L_G(\bar{x})$. As $M \models D_B$, it follows that $M \models \neg L_G(\bar{x}) \leftrightarrow L_B(\bar{x})$, so $M \models \neg G \leftrightarrow L_B(\bar{x})$ and $M \models B \leftrightarrow L_B(\bar{x})$.

- Suppose B is of the form $G \vee H$. Then:

$$D_B \equiv \forall \bar{x} (L_B(\bar{x}) \vee \neg L_G(\bar{y})) \wedge \forall \bar{x} (L_B(\bar{x}) \vee \neg L_H(\bar{z})) \wedge \forall \bar{x} (\neg L_B(\bar{x}) \vee L_G(\bar{y}) \vee L_H(\bar{z}))$$

We can assume by the induction hypothesis that $M \models G \leftrightarrow L_G(\bar{x})$ as well as $M \models H \leftrightarrow L_H(\bar{x})$. As $M \models D_B$, we get that $M \models L_G(\bar{y}) \supset L_B(\bar{x})$, $M \models L_H(\bar{z}) \supset L_B(\bar{x})$ and $M \models L_B(\bar{x}) \supset (L_G(\bar{y}) \vee L_H(\bar{z}))$. Therefore $M \models L_B(\bar{x}) \leftrightarrow (G \vee H)$ and consequently $M \models L_B(\bar{x}) \leftrightarrow B$.

- Suppose B is of the form $\forall xG$. Then:

$$D_B \equiv \forall \bar{x} \forall x (\neg L_B(\bar{x}) \vee L_G(\bar{x}, x)) \wedge \forall \bar{x} \forall x (L_B(\bar{x}) \vee \neg L_G(\bar{x}, x))$$

By the induction hypothesis, $M \models G \leftrightarrow L_G(\bar{x}, x)$. Since $M \models D_B$ and as x does not occur in $L_B(\bar{x})$, $M \models L_B(\bar{x}) \leftrightarrow \forall xG$, which is nothing else than $M \models L_B(\bar{x}) \leftrightarrow B$. \square

Lemma 17. *Let A be a formula without existential quantifiers and M_A a model in the language $L(A)$. Extend M_A to a model M'_A in the language $L(\delta(A))$ such that for $B \in \Sigma(A)$, $M_A \models L_B(\bar{x})$ if and only if $M_A \models B$. Then $M'_A \models D_{\Sigma(A)}$.*

Proof. We proceed by induction on the structure of A . For the base case, suppose that A is an atom. Then $D_{\Sigma(A)} = D_A = \forall \bar{x}(\neg A \vee L_A(\bar{x})) \wedge \forall \bar{x}(A \vee \neg L_A(\bar{x}))$. Consider the case that $M'_A \models A$. Then by construction of M'_A , $M'_A \models L_A(\bar{x})$, hence D_A holds. In the case where $M'_A \not\models A$, we know that $M'_A \not\models L_A$, so D_A holds as well.

For the induction step, consider the following cases. The remaining cases can be argued analogously.

- A is of the form $G \supset H$. Then $D_{\Sigma(A)} = D_{\Sigma(G)} \wedge D_{\Sigma(H)} \wedge D_A$. By the induction hypothesis, we get that $M'_A \models D_{\Sigma(G)}$ as well as $M'_A \models D_{\Sigma(H)}$. It remains to show that $M'_A \models D_A$, i.e. $M'_A \models \forall \bar{x}(L_A(\bar{x}) \vee L_G(\bar{y})) \wedge \forall \bar{x}(L_A(\bar{x}) \vee \neg L_H(\bar{z})) \wedge \forall \bar{x}(\neg L_A(\bar{x}) \vee \neg L_G(\bar{y}) \vee L_H(\bar{z}))$.

Suppose that $M'_A \models A$. Then $M'_A \not\models G$ or $M'_A \models H$. By construction of M'_A , we furthermore have that $M'_A \models L_B(\bar{x})$ and $M'_A \models \neg L_G(\bar{y}) \vee L_H(\bar{z})$.

Otherwise we have that $M'_A \not\models A$, so $M'_A \models G$ and $M'_A \not\models H$. Hence $M'_A \models \neg L_A(\bar{x})$, $M'_A \models L_G(\bar{y})$ and $M'_A \not\models L_H(\bar{z})$.

- A is of the form $\forall x B$. Then $D_{\Sigma(A)} = D_{\Sigma(B)} \wedge D_A$. By the induction hypothesis, $M'_A \models D_{\Sigma(B)}$, and we conclude by showing that $M'_A \models \forall \bar{x} \forall x(\neg L_A(\bar{x}) \vee L_B(\bar{x}, x)) \wedge \forall \bar{x} \forall x(L_A(\bar{x}) \vee \neg L_B(\bar{x}, x))$:

Suppose $M'_A \models A$. Then consequently, $M'_A \models \forall x B$, so $M'_A \models L_A(\bar{x})$ and $M'_A \models L_B(\bar{x}, x)$. Otherwise $M'_A \not\models A$. In this case $M'_A \not\models \forall x B$, so $M'_A \not\models L_A(\bar{x})$ and $M'_A \not\models L_B(\bar{x}, x)$. \square

Lemma 18. *Let A be a formula and Φ a set of formulas without existential quantifiers with such that $L(A) \subseteq L(\Phi)$. Then $\Phi \models A$ if and only if $\delta(\Phi) \models A$.*

Proof. If $\Phi \models A$, then $\Phi \cup \{A\}$ is unsatisfiable and thus by the Compactness Theorem, there exists a finite $\Phi' \subseteq \Phi$ such that $\Phi' \cup \{A\}$ is unsatisfiable, or in other words $\Phi' \models A$. Extend Φ' such that $L(A) \subseteq L(\Phi')$. Let $B = \bigwedge_{C \in \Phi'} C$. We show that $B \models A$ if and only if $\delta(B) \models A$ by induction on the structure of B .

For the if-direction, assume that $\delta(B) \models A$ and let M be a model such that the $L(B)$ -reduct of M , $M|_{L(B)}$, is a model of B . Let M' extend $M|_{L(B)}$ as in Lemma 17 and hence by that lemma, $M' \models D_{\Sigma(B)}$. By the construction of M' , $M' \models L_B(\bar{x})$, therefore $M' \models \delta(B)$, so by the induction hypothesis $M' \models A$. As $L(A) \subseteq L(B)$ and $M'|_{L(B)} = M|_{L(B)}$, $M \models A$.

For the only if-direction, assume that $B \models A$ and let M be a model such that $M \models \delta(B)$. By Lemma 16, $M \models B$ and hence $M \models A$. \square

Proposition 19. *Let $\Gamma \cup \Delta$ be unsatisfiable and contain no existential quantifiers. Then I is an interpolant for $\Gamma \cup \Delta$ if and only if I is an interpolant for $\delta(\Gamma) \cup \delta(\Delta)$.*

Proof. As δ introduces fresh symbols for each Γ and Δ , they do not occur in any interpolant for Γ and Δ . This establishes the language condition in both directions.

Furthermore, Lemma 18 is applicable to interpolants I for $\Gamma \cup \Delta$ due to the language condition and demonstrates that $\Gamma \models I$ if and only if $\delta(\Gamma) \models I$ as well as $\Delta \models \neg I$ if and only if $\delta(\Gamma) \models \neg I$, which gives the result. \square

At this point, we can summarize the results which enable the use of resolution based methods for calculating interpolants:

Theorem 20. *Let $\Gamma \cup \Delta$ be unsatisfiable. Then I is an interpolant for $\Gamma \cup \Delta$ if and only if I is an interpolant for $\delta(\text{sk}(\Gamma)) \cup \delta(\text{sk}(\Delta))$.*

Proof. Immediate by Proposition 19 and Proposition 14. \square

2.3 Sequent Calculus

The famous sequent calculus was introduced in [Gen35]. Its use of sequents in lieu of plain formulas allows for a natural mapping of the logical relations expressed by the connectives to the structure of proofs.

Definition 21. For multisets of first-order formulas Γ and Δ , $\Gamma \vdash \Delta$ is called a *sequent*. In this context Γ forms the *antecedent*, whereas Δ is referred to as *succedent*.

A sequent $\Gamma \vdash \Delta$ is called *provable* if there is a sequent calculus proof of $\Gamma \vdash \Delta$. \triangle

The rules of sequent calculus are as follows:

Axioms

$$A \vdash A$$

$$\vdash t = t$$

Cut

$$\frac{\Gamma \vdash \Delta, A \quad A, \Sigma \vdash \Pi}{\Gamma, \Sigma \vdash \Delta, \Pi}$$

Structural rules

- Contraction

$$\frac{\Gamma, A, A \vdash \Delta}{\Gamma, A \vdash \Delta} \text{c} : l$$

$$\frac{\Gamma \vdash \Delta, A, A}{\Gamma \vdash \Delta, A} \text{c} : r$$

- Weakening

$$\frac{\Gamma \vdash \Delta}{\Gamma, A \vdash \Delta} \text{w} : l$$

$$\frac{\Gamma \vdash \Delta}{\Gamma \vdash \Delta, A} \text{w} : r$$

Propositional rules

- Negation

$$\frac{\Gamma \vdash \Delta, A}{\neg A, \Gamma \vdash \Delta} \neg : l$$

$$\frac{A, \Gamma \vdash \Delta}{\Gamma \vdash \Delta, \neg A} \neg : r$$

- Conjunction

$$\frac{\Gamma, A, B \vdash \Delta}{\Gamma, A \wedge B \vdash \Delta} \wedge : l$$

$$\frac{\Gamma \vdash \Delta, A \quad \Sigma \vdash \Pi, B}{\Gamma, \Sigma \vdash \Delta, \Pi, A \wedge B} \wedge : r$$

TODO: format pagebreaks nicely in final version

- Disjunction

$$\frac{\Gamma, A \vdash \Delta \quad \Sigma, B \vdash \Pi}{\Gamma, \Sigma, A \vee B \vdash \Delta, \Pi} \vee : l$$

$$\frac{\Gamma \vdash \Delta, A, B}{\Gamma \vdash \Delta, A \vee B} \vee : r$$

- Implication

$$\frac{\Gamma \vdash A, \Delta \quad \Sigma, B \vdash \Pi}{\Gamma, \Sigma, A \supset B \vdash \Delta, \Pi} \supset : l$$

$$\frac{\Gamma, A \vdash \Delta, B}{\Gamma \vdash \Delta, A \supset B} \supset : r$$

Quantifier rules

- Universal

$$\frac{\Gamma, A[x/t] \vdash \Delta}{\Gamma, \forall x A \vdash \Delta} \forall : l \qquad \frac{\Gamma \vdash \Delta, A[x/y]}{\Gamma \vdash \Delta, \forall x A} \forall : r$$

- Existential

$$\frac{\Gamma, A[x/y] \vdash \Delta}{\Gamma, \exists x A \vdash \Delta} \exists : l \qquad \frac{\Gamma \vdash \Delta, A[x/t]}{\Gamma \vdash \Delta, \exists x A} \exists : r$$

(provided no free variable of t becomes bound in $A[x/t]$ and y does not occur free in Γ, Δ or A)

Equality rules

- Left rules

$$\frac{\Gamma, A[t]_p \vdash \Delta \quad \Sigma \vdash \Pi, s = t}{\Gamma, \Sigma, A[s]_p \vdash \Delta, \Pi} = : l_1 \quad \frac{\Gamma, A[s]_p \vdash \Delta \quad \Sigma \vdash \Pi, s = t}{\Gamma, \Sigma, A[t]_p \vdash \Delta, \Pi} = : l_2$$

- Right rules

$$\frac{\Gamma \vdash \Delta, A[t]_p \quad \Sigma \vdash \Pi, s = t}{\Gamma, \Sigma \vdash \Delta, \Pi, A[s]_p} = : r_1 \quad \frac{\Gamma \vdash \Delta, A[s]_p \quad \Sigma \vdash \Pi, s = t}{\Gamma, \Sigma \vdash \Delta, \Pi, A[t]_p} = : r_2$$

(provided no free variable of s or t becomes bound in $A[t]_p$ or $A[s]_p$)

Figure 2.2: The rules of sequent calculus

For the purposes of this thesis, we usually consider the cut-free fragment of sequent calculus.

Theorem 22. *Cut-free sequent calculus is sound and complete.*

Reduction to First-Order Logic without Equality

A common theme of proofs is to avoid the tedious effort of proving the result from first principles by reducing the problem to one that is easier to solve. In this instance, we are able to give a reduction for finding interpolants in first-order logic *with* equality to first-order logic *without* equality, where it is simpler to give an appropriate algorithm. This method is due to Craig ([Cra57b]).

In order to simplify notation, we shall consider constant symbols to be function symbols of arity 0 in this section. The general layout of this approach is the following: From two sets Γ and Δ , where $\Gamma \cup \Delta$ is unsatisfiable, we compute two sets Γ' and Δ' which do not make use of equality but simulate the effects of equality in Γ and Δ via axioms. In the process of this transformation, also function symbols are replaced by predicate symbols with appropriate axioms to make sure that the behaviour of these function-representing predicates is compatible to the one of actual functions. Now an interpolant for Γ' and Δ' can be derived using an algorithm that is only capable of handling predicate symbols as all other non-logical symbols have been removed. Since the additional axioms ensure that the newly added predicate symbols mimic equality and functions respectively, we will see that the occurrences of these predicates in the interpolant can be translated back to occurrences of equality and function symbols in first-order logic with equality in the language of Γ and Δ , thereby yielding the originally desired interpolant.

3.1 Translation of formulas

As we shall see in this section, first-order formulas with equality can be transformed into first-order formulas without equality in a way that is satisfiability-preserving, which is sufficient for our purposes.

First, we define axioms in a language with fresh symbols which allows for simulation of equality and functions in first order logic without equality and function symbols:

Definition 23 (Translation of languages). For a first-order language \mathcal{L} and fresh predicate symbols E and F_f for $f \in \text{FS}(\mathcal{L})$, $T(\mathcal{L})$ denotes $(\mathcal{L} \cup \{E\} \cup \{F_f \mid f \in \text{FS}(\mathcal{L})\}) \setminus (\{=\} \cup \text{FS}(\mathcal{L}))$. \triangle

Definition 24 (Equality and function axioms). For a first-order language \mathcal{L} we define the following axioms in $T(\mathcal{L})$:

$$\begin{aligned}
F_{\text{Ax}}(\mathcal{L}) &\stackrel{\text{def}}{=} \bigcup_{f \in \text{FS}(\mathcal{L})} \forall \bar{x} \exists y (F_f(\bar{x}, y) \wedge (\forall z (F_f(\bar{x}, z) \supset E(y, z)))) \\
\text{Refl}(P) &\stackrel{\text{def}}{=} \forall x P(x, x) \\
\text{Congr}(P) &\stackrel{\text{def}}{=} \forall x_1 \forall y_1 \dots \forall x_{\text{ar}(P)} \forall y_{\text{ar}(P)} ((E(x_1, y_1) \wedge \dots \wedge E(x_{\text{ar}(P)}, y_{\text{ar}(P)})) \supset \\
&\quad (P(x_1, \dots, x_{\text{ar}(P)}) \supset P(y_1, \dots, y_{\text{ar}(P)}))) \\
E_{\text{Ax}}(\mathcal{L}) &\stackrel{\text{def}}{=} \text{Refl}(E) \cup \bigcup_{\substack{P \in \text{PS}(\mathcal{L}) \cup \{E\} \cup \\ \{F_f \mid f \in \text{FS}(\mathcal{L})\}}} \text{Congr}(P) \quad \triangle
\end{aligned}$$

$\text{Refl}(P)$ will be referred to as reflexivity axiom of P , $\text{Congr}(P)$ as congruence axiom of P . As any model of $E_{\text{Ax}}(\mathcal{L})$ requires $\text{Refl}(E)$ and $\text{Congr}(E)$, E is also symmetric and transitive in the model:

Proposition 25. *In every model of $\text{Refl}(E)$ and $\text{Congr}(E)$, E is an equivalence relation.*

Proof. Let M be a model of $\text{Refl}(E)$ and $\text{Congr}(E)$. Then M clearly is reflexive. Due to $M \models \text{Congr}(E)$, $M \models \forall x \forall y (E(x, y) \wedge E(x, x) \supset (E(x, x) \supset E(y, x)))$. As we know that E is reflexive, this simplifies to $M \models \forall x \forall y (E(x, y) \supset E(y, x))$, i.e. E is symmetric in M . We show the transitivity of E by another instance of $\text{Congr}(E)$: $M \models \forall x \forall y \forall z ((E(y, x) \wedge E(y, z)) \supset (E(y, y) \supset E(x, z)))$. As E is reflexive and symmetric, we get that $M \models \forall x \forall y \forall z ((E(x, y) \wedge E(y, z)) \supset E(x, z))$. \square

We continue by defining the translation procedure for formulas:

Definition 26 (Translation and inverse translation of formulas). Let A be a first-order formula and E and F_f for $f \in \text{FS}(A)$ be fresh predicate symbols. Then $T(A)$ is the result of applying the following algorithm to A :

1. Replace every occurrence of $s = t$ in A by $E(s, t)$
2. As long as there is an occurrence of a function symbol f in A :
 Let B be the atom in which f occurs as outermost symbol of a term. Then B is of the form $P(s_1, \dots, s_{j-1}, f(\bar{t}), s_{j+1}, \dots, s_m)$. Replace B in A by $\exists y(F_f(\bar{t}, y) \wedge P(s_1, \dots, s_{j-1}, y, s_{j+1}, \dots, s_m))$ for a fresh variable y .

Moreover, let the inverse operation $T^{-1}(B)$ for formulas B in the language $T(L(A))$ be defined as the result of applying the following algorithm to B :

1. Replace every occurrence of $E(s, t)$ in B by $s = t$.
2. For every $f \in \text{FS}(A)$, replace every occurrence of $\exists y(F_f(\bar{t}, y) \wedge P(s_1, \dots, s_{j-1}, y, s_{j+1}, \dots, s_m))$ in B by $P(s_1, \dots, s_{j-1}, f(\bar{t}), s_{j+1}, \dots, s_m)$
3. For every $f \in \text{FS}(A)$, replace every occurrence of $F_f(\bar{t}, s)$ by $f(\bar{t}) = s$.

For sets of first-order formulas Φ , let $T(\Phi) \stackrel{\text{def}}{=} \bigcup_{A \in \Phi} T(A)$ and $T^{-1}(\Phi) \stackrel{\text{def}}{=} \bigcup_{A \in \Phi} T^{-1}(A)$. \triangle

Remark. Let \mathcal{L} be a language. Step 2 and 3 of T^{-1} are both concerned with replacing occurrences of F_f by occurrences of f for $f \in \text{FS}(\mathcal{L})$, but are relevant in different contexts.

Step 2 of T^{-1} is the precise inverse of step 2 of T in the sense that for any formula A , $T^{-1}(T(A)) = A$ as we will show in Lemma 27. In this context, step 3 has no effect, as all occurrences of F_f have been introduced by $T(\cdot)$ and are consequently of exactly the form that is handled by step 2. So the algorithm is in this regard complete even without step 3.

On the other hand, if arbitrary formulas in the language $T(\mathcal{L})$ are given, they in general do not match that pattern and are only translated to \mathcal{L} in step 3. Note that T^{-1} without step 2 yields a complete algorithm, as any formula that is handled there can also be processed in step 3. In such a procedure, $T^{-1}(T(A))$ and A are in general not syntactically equal for formulas A but only logically equivalent. \triangle

Lemma 27. *Let A be a first-order formula and Φ be a set of first-order formulas. Then $T^{-1}(T(A)) = A$ and $T^{-1}(T(\Phi)) = \Phi$.*

Proof. Step 1 and 2 in the algorithms T and T^{-1} are each concerned with a different set of symbols and therefore do not interfere with each other. Moreover, the respective steps in both algorithms are the inverse of each other. For step 1, this is immediate and for step 2, consider that all occurrences of F_f for $f \in \text{FS}(A)$ in $T(A)$ have been introduced by T and are consequently of the form

$\exists y(F_f(\bar{t}, y) \wedge P(s_1, \dots, s_{j-1}, y, s_{j+1}, \dots, s_m))$, which is replaced by $P(s_1, \dots, s_{j-1}, f(\bar{t}), s_{j+1}, \dots, s_m)$ by T^{-1} . As no occurrences of F_f remain, step 3 of T^{-1} leaves the formula unchanged. \square

Definition 28 (Translation of formulas including axioms). For first-order formulas A , let $T_{Ax}(A) \stackrel{\text{def}}{=} \left(\bigwedge_{B \in F_{Ax}(L(A))} B \right) \wedge \left(\bigwedge_{B \in E_{Ax}(L(A))} B \right) \wedge T(A)$ and for sets of first-order formulas Φ , let $T_{Ax}(\Phi) \stackrel{\text{def}}{=} F_{Ax}(L(\Phi)) \cup E_{Ax}(L(\Phi)) \cup T(\Phi)$. \triangle

Note that $T_{Ax}(A)$ contains neither the equality predicate nor function symbols but additional predicate symbols instead. More formally:

Lemma 29.

1. Let Φ be a set of first-order formulas. Then $T_{Ax}(\Phi)$ is in the language $T(L(\Phi))$.
2. If Ψ is in the language $T(\mathcal{L})$, then $T^{-1}(\Psi)$ is in the language \mathcal{L} .

Proposition 30. Let Φ be a set of first-order formulas.

1. If Φ is satisfiable, then so is $T_{Ax}(\Phi)$.
2. Let \mathcal{L} be a first-order language and Φ a set of first-order formulas in the language $T(\mathcal{L})$. If $F_{Ax}(\mathcal{L}) \cup E_{Ax}(\mathcal{L}) \cup \Phi$ is satisfiable, then so is $T^{-1}(\Phi)$.

Proof. Suppose Φ is satisfiable. Let M be a model of Φ . We show that $T_{Ax}(\Phi)$ is satisfiable by extending M to the language $L(\Phi) \cup \{E\} \cup \{F_f \mid f \in FS(A)\}$ and proving that the extended model satisfies $T_{Ax}(\Phi)$.

First, let $M \models E(s, t)$ if and only if $M \models s = t$. By reflexivity of equality, it follows that $M \models \text{Refl}(E)$. As any predicate, in particular E and F_f for every $f \in FS(\Phi)$, satisfy the congruence axiom with respect to $=$, by the definition of E in M , they satisfy the congruence axiom with respect to E . Therefore M is a model of $E_{Ax}(L(\Phi))$.

Second, let $M \models F_f(\bar{x}, y)$ if and only if $M \models f(\bar{x}) = y$ for all $f \in FS(\Phi)$. Since M is a model of Φ , it maps every function symbol f to a function, which by definition returns a unique result for every combination of parameters. This however is precisely the logical requirement on F_f stated by $F_{Ax}(L(\Phi))$, hence M is a model of $F_{Ax}(L(\Phi))$.

Lastly, we show that $M \models T(A)$ for all $A \in \Phi$. By the above definition of E in M , step 1 of the algorithm in Definition 26 yields a formula that is satisfied by M as it satisfies every formula of Φ . For step 2, suppose $P(s_1, \dots, s_{j-1}, f(\bar{t}), s_{j+1}, \dots, s_m)$ does (not) hold under M . Let y be such that $M \models f(\bar{t}) = y$. By our definition of F_f under M , $M \models F_f(\bar{t}, y)$ with this unique y . Hence $\exists y(F_f(\bar{t}, y) \wedge P(s_1, \dots, s_{j-1}, y, s_{j+1}, \dots, s_m))$ does (not) hold under M .

For 2, suppose $F_{Ax}(\mathcal{L}) \cup E_{Ax}(\mathcal{L}) \cup \Phi$ is satisfiable and let M be a model of it.

First, note that as $M \models E_{Ax}(\mathcal{L})$, by Proposition 25, E^M is an equivalence relation. Let D be the domain of M . We build a model M' with the domain $D' = D/E^M$, i.e. the congruence relation of D modulo E^M . The interpretation \mathcal{I}' of M' is obtained from the interpretation \mathcal{I} of M by replacing every occurrence of a domain element d by its respective congruence class $\{x \mid (d, x) \in E^M\}$. As $M \models E_{Ax}(\mathcal{L})$, it satisfies the congruence axioms with respect to every function and predicate symbol, and therefore \mathcal{I}' is well-defined. Due to this construction, $M' \models s = t$ if and only if $M \models E(s, t)$ for all terms s and t .

how to write this in a nicer way?

Second, let $M \models f(\bar{t}) = s$ if and only if $M \models F_f(\bar{t}, s)$ for all $f \in FS(\mathcal{L})$. As by assumption M is a model of $F_{Ax}(A)$, we know that for every \bar{t} , some s with $M \models F(\bar{t}, s)$ exists and is uniquely defined. Hence f in M refers to a well-defined function.

Lastly, to show that $M \models T^{-1}(\Phi)$, consider that the interpretations of the predicates E and $=$ coincide in M . Furthermore, let B be an occurrence of $\exists y(F_f(\bar{t}, y) \wedge P(s_1, \dots, s_{j-1}, y, s_{j+1}, \dots, s_m))$ for some $f \in FS(\mathcal{L})$ in Φ . Then by the above definition of f in M , we have that B is in M equivalent to $\exists y(f(\bar{t}) = y) \wedge P(s_1, \dots, s_{j-1}, y, s_{j+1}, \dots, s_m)$, which due to f being a function is equivalent to $M \models P(s_1, \dots, s_{j-1}, f(\bar{t}), s_{j+1}, \dots, s_m)$.

Similarly, let B be an occurrence of $F_f(\bar{t}, s)$ in Φ . Then by our above definition of f in M , we have that $M \models f(\bar{t}) = s$ iff $M \models B$. \square

Corollary 31. *Let Φ be a set of first-order formulas. Then Φ is satisfiable if and only if $T_{Ax}(\Phi)$ is satisfiable.*

Proof. The left-to-right direction is directly given in Proposition 30. For the other direction, consider that by Proposition 30, $T^{-1}(T(\Phi))$ is satisfiable, which by Lemma 27 is nothing else than Φ . \square

3.2 Computation of interpolants

For the proof of the interpolation theorem by reduction we require an algorithm that operates in first-order logic without equality and function symbols, which we describe in this section.

Remark. As the idea of this reduction is to simplify the problem by amongst others not considering function symbols, resolution-based methods can not be employed in a direct manner. This is because function symbols appear naturally in them as they usually handle existential quantification by means of skolemisation, i.e. a new function symbol is introduced for every occurrence of an existential quantifier in the scope of a universal quantifier. Translating the skolemised formulas to a language without function symbols as described in Definition 26 is

of no avail since this translation introduces new existential quantifiers for every function symbol it encounters, necessitating skolemisation yet again. \triangle

Lemma 32. *Let Γ and Δ be sets of first-order formulas such that the equality symbol does not occur in them and $\Gamma \vdash \Delta$ is provable in sequent calculus. Then there exists a proof of $\Gamma \vdash \Delta$ that does not contain the equality symbol.*

Proof. Let π be a proof of $\Gamma \vdash \Delta$.

Suppose π contains an instance of the equality axiom $\vdash t = t$ for a term t . As no equality symbol is contained in the end sequent, there has to be a rule application in π which removes either $t = t$. Only instances of equality rules or cut are capable of this.

Consider the case that an equality rule removed $t = t$. As the cases for $= : l_1$, $= : l_2$, $= : r_1$ and $= : r_2$ are similar, we only consider the case of $= : l_1$. The proof π' leading up to the rule application that removes the occurrence of $t = t$ is of the form:

$$\frac{\frac{\varphi}{\Gamma, A[t]_p \vdash \Delta} \quad \frac{\psi}{\Sigma \vdash \Pi, t = t}}{\Gamma, \Sigma, A[t]_p \vdash \Delta, \Pi} = : l_1$$

We can replace π' in π by the following to obtain a proof without an occurrence of the equality symbol:

$$\frac{\frac{\varphi}{\Gamma, A[t]_p \vdash \Delta}}{\Gamma, \Sigma, A[t]_p \vdash \Delta} w : l \quad \frac{\Gamma, \Sigma, A[t]_p \vdash \Delta}{\Gamma, \Sigma, A[t]_p \vdash \Delta, \Pi} w : r$$

TODO: CUT

Suppose π contains an instance of the axiom $A \vdash A$ such that the equality symbol occurs in A . Then A is of the form $s = t$ for terms s and t . While the occurrence in the consequent might be eliminated by an equality rule application, due to the subformula property, there is no rule in cut-free sequent calculus such that the occurrence in the antecedent is removed. Hence it appears in the final sequent, which contradicts the assumption.

Suppose π contains an instance of $w : l$ such that the equality symbol occurs in the principal formula A . This case can be argued similarly as for occurrences of A as antecedent of an axiom $A \vdash A$.

Suppose π contains an instance of $w : r$ such that the equality symbol occurs in the principal formula A . Then as it does not occur in the end sequent, it is removed by either an instance of an equality rule or the cut rule. Suppose it is removed via an equality rule. We consider the case of $= : l_1$.

this is
not fin-
ished

$$\frac{\frac{\varphi}{\Gamma \vdash \Delta} \text{ w : } r}{\Gamma \vdash \Delta, s = t} \quad \vdots$$

$$\frac{\Lambda, A[t]_p \vdash \Theta \quad \Sigma \vdash \Pi, s = t}{\Lambda, \Sigma, A[s]_p \vdash \Theta, \Pi} = : l_1$$

TODO

(proof might become longer, but how does that work? don't inessential cuts remain?)

□

We now show that interpolants can be computed by means of a sequent calculus based procedure by Maehara. It is slightly stronger than the required statement as it allows for interpolants of partitions of sequents:

Definition 33 (Partition of sequents). A partition of a sequent $\Gamma \vdash \Delta$ is denoted by $\langle (\Gamma_1; \Delta_1), (\Gamma_2; \Delta_2) \rangle$, where $\Gamma_1 \uplus \Gamma_2 = \Gamma$ and $\Delta_1 \uplus \Delta_2 = \Delta$. \triangle

Lemma 34 (Maehara). *Let Γ and Δ be sets of first-order clauses without equality and function symbols such that $\Gamma \vdash \Delta$ is provable in cut-free sequent calculus. Then for any partition $\langle (\Gamma_1; \Delta_1), (\Gamma_2; \Delta_2) \rangle$ there is an interpolant I such that*

1. $\Gamma_1 \vdash \Delta_1, I$ is provable
2. $\Gamma_2, I \vdash \Delta_2$ is provable
3. $L(I) \subseteq L(\Gamma_1, \Delta_1) \cap L(\Gamma_2, \Delta_2)$

Proof. We prove this lemma by induction on the number of inferences in a cut-free proof of $\Gamma \vdash \Delta$. By Lemma 32, we can assume that no equality symbol occurs in the proof, so equality rules need not be considered.

Base case. Suppose no rules were applied. Then $C \vdash D$ is of one of the form $A \vdash A$. We give interpolants for any of the four possible partitions:

1. $\langle (A; A), (;) \rangle: I = \perp$
2. $\langle (;), (A; A) \rangle: I = \top$
3. $\langle (; A), (A;) \rangle: I = \neg A$
4. $\langle (A;), (; A) \rangle: I = A$

Structural rules. Suppose the property holds for n rule applications and the $(n + 1)$ th rule is a structural one.

- The last rule application is an instance of $c : l$. Then it is of the form:

$$\frac{\Gamma, A, A \vdash \Delta}{\Gamma, A \vdash \Delta} c : l$$

There are two possible partition schemes: of $\Gamma, A \vdash \Delta$:

1. $\chi = \langle (\Gamma_1, A; \Delta_1), (\Gamma_2; \Delta_2) \rangle$. By the induction hypothesis, we know that there is an interpolant I for the partition $\langle (\Gamma_1, A, A; \Delta_1), (\Gamma_2; \Delta_2) \rangle$ of the upper sequent. I serves as interpolant for χ as well.
2. $\chi = \langle (\Gamma_1; \Delta_1), (\Gamma_2, A; \Delta_2) \rangle$. By a similar argument, we get that there is an interpolant I for $\langle (\Gamma_1; \Delta_1), (\Gamma_2, A, A; \Delta_2) \rangle$, which again is also an interpolant for χ .

The case of $c : r$ is analogous.

- The last rule application is an instance of $w : r$. Then it is of the form:

$$\frac{\Gamma \vdash \Delta}{\Gamma \vdash \Delta, A} w : r$$

By the induction hypothesis, there exists an interpolant I for any partition $\langle (\Gamma_1; \Delta_1), (\Gamma_2; \Delta_2) \rangle$ of $\Gamma \vdash \Delta$. Clearly I remains an interpolant when adding A to either Δ_1 or Δ_2 .

The case of $w : l$ is analogous.

Propositional rules. Suppose the property holds for n rule applications and the $(n + 1)$ th rule is a propositional one.

- The last rule application is an instance of $\neg : l$. Then it is of the form:

$$\frac{\Gamma \vdash \Delta, A}{\neg A, \Gamma \vdash \Delta} \neg : l$$

There are two possible partition schemes of $\Gamma, \neg A \vdash \Delta$:

1. $\chi = \langle (\Gamma_1, \neg A; \Delta_1), (\Gamma_2; \Delta_2) \rangle$. By the induction hypothesis, there exists an interpolant I for the partition $\langle (\Gamma_1; \Delta_1, A), (\Gamma_2; \Delta_2) \rangle$ of the upper sequent. Clearly I is an interpolant for χ as well.
2. $\chi = \langle (\Gamma_1; \Delta_1), (\Gamma_2, \neg A; \Delta_2) \rangle$. A similar argument goes through.

The case of $\neg : r$ is analogous.

- The last rule application is an instance of $\supset : l$. Then it is of the form:

$$\frac{\Gamma \vdash \Delta, A \quad \Sigma, B \vdash \Pi}{\Gamma, \Sigma, A \supset B \vdash \Delta, \Pi} \supset : l$$

There are two possible partition schemes of $\Gamma, A \supset B \vdash \Delta$:

1. $\chi = \langle (\Gamma_1, \Sigma_1, A \supset B; \Delta_1, \Pi_1), (\Gamma_2, \Sigma_2; \Delta_2, \Pi_2) \rangle$. By the induction hypothesis, there is an interpolant I_1 for the partition $\langle (\Gamma_1; \Delta_1, A), (\Gamma_2; \Delta_2) \rangle$ of the left upper sequent. Hence for I_1 , we have that $\Gamma_1 \vdash \Delta_1, A, I_1$ and $I_1, \Gamma_2 \vdash \Delta_2$ are provable. Moreover, we also get by the induction hypothesis that there is an interpolant I_2 for the partition $\langle (\Sigma_1, B; \Pi_1), (\Sigma_2; \Pi_2) \rangle$ of the right upper sequent. Therefore $\Sigma_1, B \vdash \Pi_1, I_2$ and $I_2, \Sigma_2 \vdash \Pi_2$ are provable.

Using these prerequisites, we first establish that $I_1 \vee I_2$ fulfills conditions 1 and 2 of an interpolant for χ :

$$\frac{\frac{\Gamma_1 \vdash \Delta_1, A, I_1 \quad \Sigma_1, B \vdash \Pi_1, I_2}{\Gamma_1, \Sigma_1, A \supset B \vdash \Delta_1, \Pi_1, I_1, I_2} \supset : l}{\Gamma_1, \Sigma_1, A \supset B \vdash \Delta_1, \Pi_1, I_1 \vee I_2} \vee : r$$

$$\frac{I_1, \Gamma_2 \vdash \Delta_2 \quad I_2, \Sigma_2 \vdash \Pi_2}{I_1 \vee I_2, \Gamma_2, \Sigma_2 \vdash \Delta_2, \Pi_2} \vee : l$$

To show that also condition 3 is satisfied, consider that by the induction hypothesis, it holds that:

$$\begin{aligned} L(I_1) &\subseteq L(\Gamma_1, \Delta_1, A) \cap L(\Gamma_2, \Delta_2) \\ L(I_2) &\subseteq L(\Sigma_1, B, \Pi_1) \cap L(\Sigma_2, \Pi_2) \end{aligned}$$

Therefore

$$\begin{aligned} L(I_1) \cup L(I_2) &\subseteq (L(\Gamma_1, \Delta_1, A) \cap L(\Gamma_2, \Delta_2)) \cup (L(\Sigma_1, B, \Pi_1) \cap L(\Sigma_2, \Pi_2)) \\ &\Downarrow \\ L(I_1) \cup L(I_2) &\subseteq (L(\Gamma_1, \Delta_1, A) \cup L(\Sigma_1, B, \Pi_1)) \cap (L(\Gamma_2, \Delta_2) \cup L(\Sigma_2, \Pi_2)) \\ &\Updownarrow \\ L(I_1 \vee I_2) &\subseteq L(\Gamma_1, \Sigma_1, A \supset B, \Delta_1, \Pi_1) \cap L(\Gamma_2, \Sigma_2, \Delta_2, \Pi_2) \end{aligned}$$

2. $\chi = \langle (\Gamma_1, \Sigma_1; \Delta_1, \Pi_1), (\Gamma_2, \Sigma_2, A \supset B; \Delta_2, \Pi_2) \rangle$. The argument for this case is similar using $I_1 \wedge I_2$ as interpolant.

For the other binary connectives $\wedge : l$, $\wedge : r$, $\vee : l$, $\vee : r$ and $\supset : r$, similar arguments go through, where the interpolant is always either the conjunction or the disjunction of the interpolants of partitions of the preceding sequents.

Quantifier rules. Suppose the property holds for n rule applications and the $(n + 1)$ th rule is a quantifier rule.

- The last rule application is an instance of $\forall : l$. Then it is of the form:

$$\frac{\Gamma, A[x/y] \vdash \Delta}{\Gamma, \forall x A \vdash \Delta} \forall : l$$

Note that since we have excluded function symbols to occur in the final sequent (and constant symbols are treated as function symbols of arity 0),

it is only possible to find a proof where these don't occur, but they might occur in a stupid proof. add a lemma as for equality?

There are two possible partition schemes of $\Gamma, \forall x A \vdash \Delta$:

1. $\langle(\Gamma_1, \forall x A; \Delta_1), (\Gamma_2; \Delta_2)\rangle$. By the induction hypothesis, there is an interpolant I of the partition $\langle(\Gamma_1, A[x/y]; \Delta_1), (\Gamma_2; \Delta_2)\rangle$. Hence for I , $\Gamma_1, A[x/y] \vdash \Delta_1, I$ and $I, \Gamma_2 \vdash \Delta_2$ are provable. By an application of $\forall : l$ to the first sequent we get $\Gamma_1, \forall x A \vdash \Delta_1, I$, so I satisfies conditions 1 and 2 of being an interpolant for χ .

In order to show that also $L(I) \subseteq L(\Gamma_1, \forall x A, \Delta_1) \cap L(\Gamma_2, \Delta_2)$, consider that by the induction hypothesis, $L(I) \subseteq L(\Gamma_1, A[x/y], \Delta_1) \cap L(\Gamma_2, \Delta_2)$.

As free variables are not considered to be part of the language, $L(\forall x A) = L(A[x/y])$.

2. $\langle(\Gamma_1; \Delta_1), (\Gamma_2, \forall x A; \Delta_2)\rangle$. This case can be argued analogously.

In the case of $\exists : r$, a similar argument goes through.

- The last rule application is an instance of $\forall : r$. Then it is of the form:

$$\frac{\Gamma \vdash \Delta, A[x/y]}{\Gamma \vdash \Delta, \forall x A} \forall : r$$

where y does not appear in Γ, Δ or A .

There are two possible partition schemes of $\Gamma \vdash \Delta, \forall x A$:

1. $\chi = \langle(\Gamma_1; \Delta_1, \forall x A), (\Gamma_2; \Delta_2)\rangle$. By the induction hypothesis, there exists an interpolant I of the partition $\langle(\Gamma_1; \Delta_1, A[x/y]), (\Gamma_2; \Delta_2)\rangle$ of the upper sequent. Hence for I , $\Gamma_1 \vdash \Delta_1, A[x/y], I$ and $I, \Gamma_2 \vdash \Delta_2$ are provable.

As y does not occur in Γ or Δ and consequently by condition 3 does not occur in I , we may apply the $\forall : r$ rule to the former sequent to obtain $\Gamma_1 \vdash \Delta_1, \forall x A, I$. Hence I is an interpolant for χ as well.

2. $\langle(\Gamma_1; \Delta_1), (\Gamma_2; \Delta_2, \forall x A)\rangle$. This case can be argued analogously.

In the case of $\exists : l$, a similar argument goes through. \square

This allows us to state the central theorem of this section:

Theorem 35. *Let Γ and Δ be sets of closed first-order formulas without equality and function symbols such that $\Gamma \cup \Delta$ is unsatisfiable. Then there is an interpolant for Γ and Δ .*

Proof. We show that there is an interpolant for $\Gamma \models \neg\Delta$, which by Proposition 5 proves the theorem. By the completeness of cut-free sequent calculus, there is a cut-free proof of $\Gamma \vdash \neg\Delta$. By Lemma 34, there is an interpolant I for the partition $\langle(\Gamma;), (; \neg\Delta)\rangle$. I is the desired interpolant for $\Gamma \models \neg\Delta$. \square

3.3 Proof by reduction

Using the results of the previous sections, we can now give a proof of the interpolation theorem:

Theorem 4 (Reverse Interpolation). *Let Γ and Δ be sets of first-order formulas such that $\Gamma \cup \Delta$ is unsatisfiable. Then there exists a reverse interpolant for Γ and Δ .*

Proof. Since $\Gamma \cup \Delta$ is unsatisfiable, by Proposition 30, $T_{Ax}(\Gamma \cup \Delta)$ is unsatisfiable.

$$\begin{aligned}
 T_{Ax}(\Gamma \cup \Delta) &\Leftrightarrow \{F_{Ax}(L(\Gamma \cup \Delta)), E_{Ax}(L(\Gamma \cup \Delta))\} \cup T(\Gamma \cup \Delta) \\
 &\Leftrightarrow \{F_{Ax}(L(\Gamma) \cup L(\Delta)), E_{Ax}(L(\Gamma) \cup L(\Delta))\} \cup T(\Gamma) \cup T(\Delta) \\
 &\Leftrightarrow \{F_{Ax}(L(\Gamma)) \wedge F_{Ax}(L(\Delta)), E_{Ax}(L(\Gamma)) \wedge E_{Ax}(L(\Delta))\} \cup T(\Gamma) \cup T(\Delta) \\
 &\Leftrightarrow \{F_{Ax}(L(\Gamma)), E_{Ax}(L(\Gamma))\} \cup T(\Gamma) \cup \{F_{Ax}(L(\Delta)), E_{Ax}(L(\Delta))\} \cup T(\Delta) \\
 &\Leftrightarrow T_{Ax}(\Gamma) \cup T_{Ax}(\Delta)
 \end{aligned}$$

Hence $T_{Ax}(\Gamma) \cup T_{Ax}(\Delta)$ is unsatisfiable as well. By Lemma 29.1 $T_{Ax}(\Gamma)$ and $T_{Ax}(\Delta)$ contain neither function symbols nor the equality symbol. Hence by Theorem 35, there is an interpolant I such that

1. $T_{Ax}(\Gamma) \models I$
2. $T_{Ax}(\Delta) \models \neg I$
3. $L(I) \subseteq L(T_{Ax}(\Gamma)) \cap L(T_{Ax}(\Delta))$

We now show that $T^{-1}(I)$ is an interpolant for Γ and Δ .

$T_{Ax}(\Gamma) \models I$ is equivalent to $T_{Ax}(\Gamma) \cup \{\neg I\}$ being unsatisfiable. Through the unfolding of $T_{Ax}(\Gamma)$, we get that $\{F_{Ax}(L(\Gamma)), E_{Ax}(L(\Gamma))\} \cup T(\Gamma) \cup \{\neg I\}$ is

unsatisfiable. This set of formulas can now be translated back to the original language with the equality symbol and function symbols. More formally, since $L(\neg I) \subseteq L(T_{Ax}(\Gamma))$, we can apply Proposition 30.2 by considering $T(\Gamma) \cup \{\neg I\}$ as Φ to conclude that $T^{-1}(T(\Gamma) \cup \{\neg I\})$ is unsatisfiable. By pulling T^{-1} inward and an application of Lemma 27, we get that $\Gamma \cup \{T^{-1}(\neg I)\} = \Gamma \cup \{\neg T^{-1}(I)\}$ is unsatisfiable. Therefore $\Gamma \models T^{-1}(I)$.

For Δ , an analogous argument goes through and so from $T_{Ax}(\Gamma) \models \neg I$ we can deduce that $\Delta \models \neg T^{-1}(I)$.

By item 3, I is in the language $L(T_{Ax}(\Gamma)) \cap L(T_{Ax}(\Delta))$, which by Lemma 29.1 is $T(L(\Gamma)) \cap T(L(\Delta))$.

$$\begin{aligned}
T(L(\Gamma)) \cap T(L(\Delta)) &= \\
&\left(L(\Gamma) \cup \{E\} \cup \{F_f \mid f \in FS(\Gamma)\} \right) \setminus \left(\{=\} \cup FS(\Gamma) \right) \cap \\
&\left(L(\Delta) \cup \{E\} \cup \{F_f \mid f \in FS(\Delta)\} \right) \setminus \left(\{=\} \cup FS(\Delta) \right) \\
&= \left((L(\Gamma) \cap L(\Delta)) \cup \{E\} \cup \{F_f \mid f \in FS(\Gamma) \cap FS(\Delta)\} \right) \setminus \left(\{=\} \cup FS(\Gamma) \cup FS(\Delta) \right) \\
&= \left((L(\Gamma) \cap L(\Delta)) \cup \{E\} \cup \{F_f \mid f \in FS(L(\Gamma) \cap L(\Delta))\} \right) \setminus \left(\{=\} \cup FS(L(\Gamma) \cap L(\Delta)) \right) \\
&= T(L(\Gamma) \cap L(\Delta))
\end{aligned}$$

As I is in the language $T(L(\Gamma) \cap L(\Delta))$, by Lemma 29.2, $T^{-1}(I)$ is in the language $L(\Gamma) \cap L(\Delta)$. \square

Interpolant extraction from resolution proofs in two phases

In [Hua95], Huang proposes an algorithm for computing interpolants of two sets of first-order formulas Γ and Δ , where $\Gamma \cup \Delta$ is unsatisfiable, by traversing a resolution refutation of $\Gamma \cup \Delta$. We present his proof in a modified form. The central difference lies in the treatment of the interplay of substitutions and liftings. While in [Hua95], propositional deductions are employed where only trivial substitutions occur, we provide a method which allows for commuting substitutions and liftings under certain conditions.

See also **TODO:** for a comments on the original proof.

4.1 Layout of the proof

The underlying algorithm produces in the first phase propositional interpolants inductively for every clause which occurs in the resolution refutation. These interpolants are propositional in the sense that they only obey the language restriction on predicates and may contain colored terms. The propositional interpolant assigned to the last clause, the empty clause, is a propositional interpolant for the initial clause sets.

The second phase of the algorithm addresses the colored terms still contained in the propositional interpolant. These are eliminated (lifted) by replacing them with bound variables whose quantifiers are subject to a certain ordering.

4.2 Extraction of propositional interpolants

We define a procedure PI, which produces propositional interpolants from resolution refutations and is based on the “Interpolation algorithm” in [Hua95].

Definition 36 (Propositional interpolant extraction.). Let π be a resolution refutation of $\Gamma \cup \Delta$. $\text{PI}(\pi)$ is defined to be $\text{PI}(\square)$, where \square is the empty clause derived in π .

For a clause C in π , $\text{PI}(C)$ is defined as follows:

Base case. If $C \in \Gamma$, $\text{PI}(C) \stackrel{\text{def}}{=} \perp$. If otherwise $C \in \Delta$, $\text{PI}(C) \stackrel{\text{def}}{=} \top$.

Resolution. If the clause C is the result of a resolution step of $C_1 : D \vee l$ and $C_2 : E \vee \neg l'$ using a unifier σ such that $l\sigma = l'\sigma$, then $\text{PI}(C)$ is defined as follows:

1. If l is Γ -colored: $\text{PI}(C) \stackrel{\text{def}}{=} [\text{PI}(C_1) \vee \text{PI}(C_2)]\sigma$
2. If l is Δ -colored: $\text{PI}(C) \stackrel{\text{def}}{=} [\text{PI}(C_1) \wedge \text{PI}(C_2)]\sigma$
3. If l is grey: $\text{PI}(C) \stackrel{\text{def}}{=} [(l \wedge \text{PI}(C_2)) \vee (\neg l' \wedge \text{PI}(C_1))]\sigma$

Factorisation. If the clause C is the result of a factorisation of $C_1 : l \vee l' \vee D$ using a unifier σ such that $l\sigma = l'\sigma$, then $\text{PI}(C) = \text{PI}(C_1)\sigma$.

Paramodulation. Suppose the clause C is the result of a paramodulation of $C_1 : s = t \vee D$ and $C_2 : E[r]$ using a unifier σ such that $r\sigma = s\sigma$. Let $h[r]$ be the maximal colored term in which r occurs in $E[r]$. Then $\text{PI}(C)$ is defined according to the following case distinction:

1. If $h[r]$ is Δ -colored and $h[r]$ occurs more than once in $E[r] \vee \text{PI}(E[r])$:

$$\text{PI}(C) \stackrel{\text{def}}{=} [(s = t \wedge \text{PI}(C_2)) \vee (s \neq t \wedge \text{PI}(C_1))]\sigma \vee (s = t \wedge h[s] \neq h[t])\sigma$$

2. If $h[r]$ is Γ -colored and $h[r]$ occurs more than once in $E[r] \vee \text{PI}(E[r])$:

$$\text{PI}(C) \stackrel{\text{def}}{=} [(s = t \wedge \text{PI}(C_2)) \vee (s \neq t \wedge \text{PI}(C_1))]\sigma \wedge (s \neq t \vee h[s] = h[t])\sigma$$

3. If r does not occur in a colored term in $E[r]$ which occurs more than once in $E[r]$:

$$\text{PI}(C) \stackrel{\text{def}}{=} [(s = t \wedge \text{PI}(C_2)) \vee (s \neq t \wedge \text{PI}(C_1))]\sigma \quad \triangle$$

4.3 Lifting of colored symbols

As PI only fixes the propositional structure of the interpolant but still contains colored symbol, we define a procedure which replaces colored terms by lifting variables.

Definition 37 (Lifting). Let φ a formula or a term and $Z = \{\zeta_1, \dots, \zeta_n\}$ the maximal Φ -colored terms in φ .

Let furthermore $z_{\text{unfold-lift}(\zeta_1)}, \dots, z_{\text{unfold-lift}(\zeta_n)}$ be fresh variables, referred to as Φ -lifting variables or lifting variables if the coloring is clear from the context. The function unfold-lift replaces lifting variables occurring in colored terms by the term they lift in order to avoid lifting variables in the index of other lifting variables and is defined as follows:

$$\text{unfold-lift}(t) \stackrel{\text{def}}{=} \begin{cases} f(\text{unfold-lift}(t_1), \dots, \text{unfold-lift}(t_m)) & \text{if } t = f(t_1, \dots, t_m) \\ t & \text{if } t \text{ is a non-lifting variable } x \\ \text{unfold-lift}(s) & \text{if } t \text{ is a lifting variable } z_s \end{cases}$$

The lifting of φ , denoted by $\ell_\Phi^z[\varphi]$, is an abbreviation of $\ell_\Phi^z[\varphi, Z]$ which is defined as follows:

$$\ell_\Phi^z[\varphi, Z] \stackrel{\text{def}}{=} \begin{cases} \varphi & Z = \emptyset \\ \ell_\Phi^z[\varphi\{\zeta_i/z_{\zeta_i}\}, Z \setminus \{\zeta_i\}] & \zeta_i \in Z \text{ such that } \zeta_i \text{ is not sub-} \\ & \text{term of another term in } Z \end{cases}$$

To simplify the syntax, we sometimes write $\ell_\Phi[\varphi]$ or $\ell[\varphi]$ if the lifting variables or the lifting variables and the color of the terms to lift respectively is clear from the context or not of the essence. \triangle

We usually lift Δ -terms by variables with the letter x and Γ -terms with the letter y . If the lifting is not specific to a color, we use variables with the letter z .

Some elementary properties of liftings are described by the following lemmas:

Lemma 38 (Commutativity of lifting and logical operators). *Let A and B be first-order formulas and s and t be terms. Then it holds that:*

1. $\ell_\Phi^z[\neg A] \Leftrightarrow \neg \ell_\Phi^z[A]$
2. $\ell_\Phi^z[A \circ B] \Leftrightarrow (\ell_\Phi^z[A] \circ \ell_\Phi^z[B])$ for $\circ \in \{\wedge, \vee\}$
3. $\ell_\Phi^z[s = t] \Leftrightarrow (\ell_\Phi^z[s] = \ell_\Phi^z[t])$

For the proof, we also require a means of commuting substitutions and liftings. This however can not be achieved in a direct manner. The following examples illustrate that in general for a term t , it is not the case that $\ell_{\Phi}^z[t\sigma] = \ell_{\Phi}^z[t]\sigma$. However Lemma 40 defines a substitution σ' such that $\ell_{\Phi}^z[t\sigma] = \ell_{\Phi}^z[t]\sigma'$.

Example 39.A. Let $t = f(x)$ be a Φ -term and $\sigma = \{x \mapsto a\}$. Then $\ell_{\Phi}^z[t\sigma] = \ell_{\Phi}^z[f(x)\sigma] = \ell_{\Phi}^z[f(a)] = z_{f(a)}$. However $\ell_{\Phi}^z[t]\sigma = \ell_{\Phi}^z[f(x)]\sigma = z_{f(x)}\sigma = z_{f(x)}$. \triangle

Example 39.B. Let $t = x$ be a variable and $\sigma = \{x \mapsto c\}$, where c is a Φ -term. Then $\ell_{\Phi}^z[t\sigma] = \ell_{\Phi}^z[x\sigma] = \ell_{\Phi}^z[c] = z_c$. But $\ell_{\Phi}^z[t]\sigma = \ell_{\Phi}^z[x]\sigma = x\sigma = c$. \triangle

Lemma 40 (Commutativity of lifting and substitution). *Let C be a clause and σ a substitution such that no lifting variable occurs in C or σ . Define σ' with $\text{dom}(\sigma') = \text{dom}(\sigma) \cup \{z_t \mid t\sigma \neq t\}$ such that for a variable z ,*

$$x\sigma' = \begin{cases} z_{t\sigma} & \text{if } x = z_t \text{ and } t\sigma \neq t \\ \ell_{\Phi}^z[x\sigma] & \text{otherwise} \end{cases}$$

Then $\ell_{\Phi}^z[C\sigma] = \ell_{\Phi}^z[C]\sigma'$.

Proof. As substitutions and liftings only affect the terms of a clause, it suffices to show that $\ell_{\Phi}^z[t\sigma] = \ell_{\Phi}^z[t]\sigma'$ for a term t in C . More precisely, only variables of $\text{dom}(\sigma)$ and maximal Δ -terms are affected. We show that for terms t of either kind that $\ell_{\Phi}^z[t\sigma] = \ell_{\Phi}^z[t]\sigma'$ holds, which proves the lemma.

Let y be a variable in $\text{dom}(\sigma)$, which occurs in C . Then $\ell_{\Phi}^z[y]\sigma' = y\sigma' = \ell_{\Phi}^z[y\sigma]$.

Let t be a maximal Φ -term in C . Then $\ell_{\Phi}^z[t\sigma] = z_{t\sigma}$. We show that $z_{t\sigma} = \ell_{\Phi}^z[t]\sigma'$.

Suppose that $t\sigma = t$. Then $\ell_{\Phi}^z[t]\sigma' = z_t\sigma' = z_t = z_{t\sigma}$. Note that z_r must not occur in t for some term r , as $z_r\sigma = z_r$, but potentially $z_r\sigma' \neq z_r$.

Otherwise it is the case that $t\sigma \neq t$. Then $\ell_{\Phi}^z[t]\sigma' = z_t\sigma'$, and by the definition of σ' , $z_t\sigma' = z_{t\sigma}$. \square

Example 41. Let M be a model and c and d be Φ -colored constants such that $M \models c = d$. Then

?????

\triangle

Lemma 42. *Let M be a model, E a formula and s and t terms such that $M \models \ell_{\Delta}^x[s] = \ell_{\Delta}^x[t]$. Let $h[t]$ be a maximal Δ -colored term containing t at p in $E[t]_p$, if such a term exists. Then it holds that:*

- If $h[t]$ does not exist, then $M \models \ell_\Delta^x[E[s]_p] \Leftrightarrow M \models \ell_\Delta^x[E[t]_p]$.
- Otherwise $M \models \ell_\Delta^x[E[s]_p] \Leftrightarrow M \models \ell_\Delta^x[E[t]_p]$ or $M \models \ell_\Delta^x[h[s]] \neq \ell_\Delta^x[h[t]]$ holds.

Proof. Suppose that the position p in $E[s]_p$ is not contained in a Δ -colored term. Then $\ell_\Delta^x[E[t]_p]$ and $\ell_\Delta^x[E[s]_p]$ only differ at position p , where for the first, $\ell_\Delta^x[t]$ is at p , and for the latter, $\ell_\Delta^x[s]$ is at p . But in M , they are interpreted the same way, hence $M \models \ell_\Delta^x[E[s]_p] \Leftrightarrow M \models \ell_\Delta^x[E[t]_p]$.

Otherwise the position p in $E[t]_p$ is contained in the maximal Δ -colored term $h[t]$. Suppose that $M \models \ell_\Delta^x[h[s]] = \ell_\Delta^x[h[t]]$ as otherwise we would be done. But then $M \models \ell_\Delta^x[E[s]_p] \Leftrightarrow M \models \ell_\Delta^x[E[t]_p]$. \square

4.4 Symmetry of the extracted interpolants

The interpolant extraction procedure PI exhibits a convenient property which is termed *symmetry* in [DKPW10, Definition 5] and will be used to show that results concerning Γ can easily be generalised to results for Δ . The symmetry-property of PI can be stated formally as follows:

Lemma 43. *Let π be a resolution refutation of $\Gamma \cup \Delta$ and $\hat{\pi}$ be π with $\hat{\Gamma} = \Delta$ and $\hat{\Delta} = \Gamma$. Then $\text{PI}(\pi) \Leftrightarrow \neg \text{PI}(\hat{\pi})$.*

Proof. We prove this lemma by induction on π . Let $\hat{\varphi}$ denote the clause/formula/literal/term in $\hat{\pi}$ corresponding to the clause/formula/literal/term φ in π .

Base case. If $C \in \Gamma$, then $C' \in \Delta'$ and $\text{PI}(C) = \perp \Leftrightarrow \neg \top = \neg \text{PI}(C')$. The case for $C \in \Delta$ is analogous.

Resolution. If the clause C is the result of a resolution step of $C_1 : D \vee l$ and $C_2 : E \vee \neg l'$ using a unifier σ such that $l\sigma = l'\sigma$, then by induction hypothesis, we get that $\text{PI}(C_i) = \neg \text{PI}(C'_i)$ for $i \in \{1, 2\}$.

We distinguish the following cases:

1. l is Γ -colored. Then \hat{l} is Δ -colored.

$$\begin{aligned}
 \text{PI}(C) &= \text{PI}(C_1) \vee \text{PI}(C_2) \\
 &\Leftrightarrow \neg(\neg \text{PI}(C_1) \wedge \neg \text{PI}(C_2)) \\
 &= \neg(\text{PI}(\hat{C}_1) \wedge \text{PI}(\hat{C}_2)) \\
 &= \neg \text{PI}(\hat{C})
 \end{aligned}$$

2. l is Δ -colored. This case can be argued analogously.
3. l is grey. Then \hat{l} is grey. Note that $l\sigma \equiv l'\sigma$ (*).

$$\begin{aligned}
\text{PI}(C) &= [(l \wedge \text{PI}(C_2)) \vee (\neg l' \wedge \text{PI}(C_1))]\sigma \\
&= (l\sigma \wedge \text{PI}(C_2)\sigma) \vee (\neg l'\sigma \wedge \text{PI}(C_1)\sigma) \\
&\stackrel{(*)}{\Leftrightarrow} (\neg l\sigma \vee \text{PI}(C_2)\sigma) \wedge (l'\sigma \vee \text{PI}(C_1)\sigma) \\
&\Leftrightarrow \neg[(l\sigma \wedge \neg \text{PI}(C_2)\sigma) \vee (\neg l'\sigma \wedge \neg \text{PI}(C_1)\sigma)] \\
&= \neg[(\hat{l}\sigma \wedge \neg \text{PI}(C_2)\sigma) \vee (\neg \hat{l}'\sigma \wedge \neg \text{PI}(C_1)\sigma)] \\
&= \neg[(\hat{l} \wedge \neg \text{PI}(\hat{C}_2)) \vee (\neg \hat{l}' \wedge \neg \text{PI}(\hat{C}_1))]\sigma \\
&= \neg[(\hat{l} \wedge \text{PI}(\hat{C}_2)) \vee (\neg \hat{l}' \wedge \text{PI}(\hat{C}_1))]\sigma \\
&= \neg \text{PI}(\hat{C})
\end{aligned}$$

Factorisation. Suppose the clause C is the result of a factorisation of $C_1 : l \vee l' \vee D$. Then $\text{PI}(C) = \text{PI}(C_1)\sigma$ and the induction hypothesis gives the result.

Paramodulation. Suppose the clause C is the result of a paramodulation of $C_1 : s = t \vee D$ and $C_2 : E[r]$ using a unifier σ such that $r\sigma = s\sigma$. We distinguish the following cases:

1. r occurs in a maximal Δ -term $h[r]$ in $E[r]$ and $h[r]$ occurs more than once in $E[r] \vee \text{PI}(E[r])$. Then \hat{r} occurs in a maximal Γ -term $\hat{h}[r]$ in $\hat{E}[r]$ and $\hat{h}[r]$ occurs more than once in $\hat{E}[r] \vee \text{PI}(\hat{E}[r])$.

$$\begin{aligned}
\text{PI}(C) &= [(s = t \wedge \text{PI}(C_2)) \vee (s \neq t \wedge \text{PI}(C_1))]\sigma \vee (s = t \wedge h[s] \neq h[t])\sigma \\
&= [(s = t \wedge \neg \text{PI}(\hat{C}_2)) \vee (s \neq t \wedge \neg \text{PI}(\hat{C}_1))]\sigma \vee (s = t \wedge h[s] \neq h[t])\sigma \\
&\Leftrightarrow \neg[(s \neq t \vee \text{PI}(\hat{C}_2)) \wedge (s = t \vee \text{PI}(\hat{C}_1))]\sigma \wedge \neg(s \neq t \vee h[s] = h[t])\sigma \\
&\Leftrightarrow \neg[(s = t \wedge \text{PI}(\hat{C}_2)) \vee (s \neq t \wedge \text{PI}(\hat{C}_1))]\sigma \wedge \neg(s \neq t \vee h[s] = h[t])\sigma \\
&= \neg \text{PI}(\hat{C})
\end{aligned}$$

2. r occurs in a maximal Γ -term $h[r]$ in $E[r]$ and $h[r]$ occurs more than once in $E[r] \vee \text{PI}(E[r])$. This case can be argued analogously.

3. Otherwise:

$$\begin{aligned}
\text{PI}(C) &= [(s = t \wedge \text{PI}(C_2)) \vee (s \neq t \wedge \text{PI}(C_1))] \sigma \\
&= [(s = t \wedge \neg \text{PI}(\hat{C}_2)) \vee (s \neq t \wedge \neg \text{PI}(\hat{C}_1))] \sigma \\
&\Leftrightarrow \neg [(s \neq t \vee \text{PI}(\hat{C}_2)) \wedge (s = t \vee \text{PI}(\hat{C}_1))] \sigma \\
&\Leftrightarrow \neg [(s = t \wedge \text{PI}(\hat{C}_2)) \vee (s \neq t \wedge \text{PI}(\hat{C}_1))] \sigma \\
&= \neg \text{PI}(\hat{C}) \quad \square
\end{aligned}$$

Corollary 44. *Let π be a resolution refutation of $\Gamma \cup \Delta$. Then $\Delta \models \ell_{\Gamma}^x[\neg \text{PI}(C) \vee C]$ for C in π .*

Proof. Build $\hat{\pi}$ from π using $\hat{\Gamma} = \Delta$ and $\hat{\Delta} = \Gamma$ as initial clause sets. By Lemma 45, $\hat{\Gamma} \models \ell_{\hat{\Delta}}^x[\text{PI}(\hat{C}) \vee \hat{C}]$ for \hat{C} in $\hat{\pi}$. By Lemma 43, $\hat{\Gamma} \models \ell_{\hat{\Delta}}^x[\neg \text{PI}(C) \vee \hat{C}]$ for the clause C in π corresponding to \hat{C} in $\hat{\pi}$. This however is nothing else than $\Delta \models \ell_{\Gamma}^x[\neg \text{PI}(C) \vee C]$. \square

4.5 Main lemma

By lifting the propositional interpolant, we are able to already obtain a formula partially fulfilling the requirements for interpolants:

Lemma 45. *Let π be a resolution refutation of $\Gamma \cup \Delta$. Then for a clause C in π , $\Gamma \models \ell_{\Delta}^x[\text{PI}(C) \vee C]$.*

Before we proceed with the proof of this lemma, we give a corollary which demonstrates that PI extracts propositional interpolants in the sense that besides possibly containing colored terms, they are proper interpolants:

Corollary 46. *Let π be a resolution refutation of $\Gamma \cup \Delta$. Then*

1. $\Gamma \models \text{PI}(\pi)$
2. $\Delta \models \neg \text{PI}(\pi)$
3. $\text{PS}(\text{PI}(\pi)) \subseteq \text{PS}(\Gamma) \cap \text{PS}(\Delta)$.

Proof. By the definition of PI, $\text{PI}(\pi)$ denotes $\text{PI}(\square)$, where \square is the empty clause derived in PI. By Lemma 45, we get that $\Gamma \models \ell_{\Delta}^x[\text{PI}(\pi)]$. As the lifting replaces terms by variables which are then implicitly universally quantified, $\text{PI}(\pi)$ is an instance of $\ell_{\Delta}^x[\text{PI}(\pi)]$. Therefore $\Gamma \models \text{PI}(\pi)$.

By Corollary 44, $\Delta \models \neg \ell_{\Gamma}^y[\text{PI}(\pi)]$, thus by a similar argument as above, $\Delta \models \neg \text{PI}(\pi)$.

By the construction of PI, only grey predicates are added. \square

Proof of Lemma 45. We proceed by induction on the resolution refutation of the strengthening $\Gamma \models \ell_{\Delta}^x[\text{PI}(C) \vee C_{\Gamma}]$, where D_{Φ} denotes the clause created from D by removing all literals which are not contained $L(\Phi)$.

Base case. Either $C \in \Gamma$, then $\ell_{\Delta}^x[C] = C$ and $\Gamma \models C$. Otherwise $C \in \Delta$ and $\text{PI}(C) = \top$.

Resolution. Suppose the last rule application is an instance of resolution. Then it is of the following form:

$$\frac{C_1 : D \vee l \quad C_2 : E \vee \neg l'}{C : (D \vee E)\sigma} \quad l\sigma = l'\sigma$$

By the induction hypothesis, we can assume that $\Gamma \models \ell_{\Delta}^x[\text{PI}(C_1) \vee (D \vee l)_{\Gamma}]$ and $\Gamma \models \ell_{\Delta}^x[\text{PI}(C_2) \vee (E \vee \neg l')_{\Gamma}]$, which by Lemma 38 implies $(\circ) \Gamma \models \ell_{\Delta}^x[\text{PI}(C_1)] \vee \ell_{\Delta}^x[D_{\Gamma}] \vee \ell_{\Delta}^x[l_{\Gamma}]$ and $(*) \Gamma \models \ell_{\Delta}^x[\text{PI}(C_2)] \vee \ell_{\Delta}^x[E_{\Gamma}] \vee \neg \ell_{\Delta}^x[l'_{\Gamma}]$. Let σ' be defined as in Lemma 40.

We proceed by a case distinction on the color of l :

1. l is Γ -colored. Then $\text{PI}(C) = [\text{PI}(C_1) \vee \text{PI}(C_2)]\sigma$.

Since $\sigma = \text{mgu}(l, l')$, $l\sigma \equiv l'\sigma$ and therefore $\ell_{\Delta}^x[l\sigma] = \ell_{\Delta}^x[l'\sigma]$. As by Lemma 40 $\ell_{\Delta}^x[l\sigma] = \ell_{\Delta}^x[l]\sigma'$ and $\ell_{\Delta}^x[l'\sigma] = \ell_{\Delta}^x[l']\sigma'$, we get $\ell_{\Delta}^x[l]\sigma' = \ell_{\Delta}^x[l']\sigma'$. Hence by applying σ' to (\circ) and $(*)$ (note that $l_{\Gamma} = l$ and $l'_{\Gamma} = l'$ as they are Γ -colored), we can perform a resolution step on $\ell_{\Delta}^x[l]\sigma'$ and obtain $\Gamma \models \ell_{\Delta}^x[\text{PI}(C_1)]\sigma' \vee \ell_{\Delta}^x[D_{\Gamma}]\sigma' \vee \ell_{\Delta}^x[\text{PI}(C_2)]\sigma' \vee \ell_{\Delta}^x[E_{\Gamma}]\sigma'$. Now we apply Lemma 40 and Lemma 38 in the other direction and get that $\Gamma \models \ell_{\Delta}^x[\text{PI}(C_1)\sigma \vee \text{PI}(C_2)\sigma \vee (D \vee E)_{\Gamma}\sigma]$. This however is nothing else than $\Gamma \models \ell_{\Delta}^x[\text{PI}(C) \vee C_{\Gamma}]$.

2. l is Δ -colored. Then $\text{PI}(C) = [\text{PI}(C_1) \wedge \text{PI}(C_2)]\sigma$.

As l and l' are Δ -colored, we can simplify (\circ) and $(*)$ as follows and apply σ' : $\Gamma \models \ell_{\Delta}^x[\text{PI}(C_1)]\sigma' \vee \ell_{\Delta}^x[D_{\Gamma}]\sigma'$ and $\Gamma \models \ell_{\Delta}^x[\text{PI}(C_2)]\sigma' \vee \ell_{\Delta}^x[E_{\Gamma}]\sigma'$. These together imply that $\Gamma \models \left(\ell_{\Delta}^x[\text{PI}(C_1)]\sigma' \wedge \ell_{\Delta}^x[\text{PI}(C_2)]\sigma' \right) \vee \ell_{\Delta}^x[D_{\Gamma}]\sigma' \vee \ell_{\Delta}^x[E_{\Gamma}]\sigma'$. By Lemma 38 and Lemma 40, this is equivalent to $\Gamma \models \ell_{\Delta}^x[(\text{PI}(C_1) \wedge \text{PI}(C_2))\sigma \vee (D \vee E)_{\Gamma}\sigma]$, which is nothing else than $\Gamma \models \ell_{\Delta}^x[\text{PI}(C) \vee C_{\Gamma}]$.

3. l is grey. Then $\text{PI}(C) = [(l \wedge \text{PI}(C_2)) \vee (\neg l' \wedge \text{PI}(C_1))]\sigma$.

We show that $\Gamma \models \ell_\Delta^x[(l \wedge \text{PI}(C_2)) \vee (\neg l' \wedge \text{PI}(C_2)) \vee D_\Gamma \vee E_\Gamma]\sigma$, for which by Lemma 38 and Lemma 40 it suffices to show that $\Gamma \models (\ell_\Delta^x[l]\sigma' \wedge \ell_\Delta^x[\text{PI}(C_2)]\sigma') \vee (\neg \ell_\Delta^x[l']\sigma' \wedge \ell_\Delta^x[\text{PI}(C_2)]\sigma') \vee \ell_\Delta^x[D_\Gamma]\sigma' \vee \ell_\Delta^x[E_\Gamma]\sigma'$.

Suppose for a model M of Γ that $M \not\models \ell_\Delta^x[D_\Gamma]\sigma'$ and $M \not\models \ell_\Delta^x[E_\Gamma]\sigma'$ as otherwise we are done. But then by (\circ) and $(*)$, we get that $M \models \ell_\Delta^x[\text{PI}(C_1)]\sigma' \vee \ell_\Delta^x[l]\sigma'$ and $M \models \ell_\Delta^x[\text{PI}(C_2)]\sigma' \vee \neg \ell_\Delta^x[l']\sigma'$. As observed in case 1, $\ell_\Delta^x[l]\sigma' = \ell_\Delta^x[l']\sigma'$. We obtain the result by a case distinction on the truth value of $\ell_\Delta^x[l]\sigma'$.

Factorisation. Suppose the last rule application is an instance of factorisation. Then it is of the following form:

$$\frac{C_1 : l \vee l' \vee D}{C : (l \vee D)\sigma} \quad \sigma = \text{mgu}(l, l')$$

The induction hypothesis gives that $\Gamma \models \ell_\Delta^x[\text{PI}(C_1) \vee l \vee l' \vee D]$. Let σ' be defined as in Lemma 40. Then $\Gamma \models \ell_\Delta^x[\text{PI}(C_1) \vee l \vee l' \vee D]\sigma'$ and by Lemma 40 and Lemma 38, $\Gamma \models \ell_\Delta^x[\text{PI}(C_1)\sigma] \vee \ell_\Delta^x[l\sigma] \vee \ell_\Delta^x[l'\sigma] \vee \ell_\Delta^x[D\sigma]$.

As $\sigma = \text{mgu}(l, l')$, $l\sigma \equiv l'\sigma$ and hence $\ell_\Delta^x[l\sigma] = \ell_\Delta^x[l'\sigma]$. But then we can apply a factorisation step and get that $\Gamma \models \ell_\Delta^x[\text{PI}(C_1)\sigma] \vee \ell_\Delta^x[l\sigma] \vee \ell_\Delta^x[D\sigma]$ which by Lemma 40 and Lemma 38 is equivalent to $\Gamma \models \ell_\Delta^x[\text{PI}(C_1)\sigma \vee l\sigma \vee D\sigma]$. This in turn is nothing else than $\Gamma \models \ell_\Delta^x[\text{PI}(C) \vee C]$.

Paramodulation. Suppose the last rule application is an instance of paramodulation. Then it is of the following form:

$$\frac{C_1 : D \vee s = t \quad C_2 : E[r]_p}{C : (D \vee E[t]_p)\sigma} \quad \sigma = \text{mgu}(s, r)$$

By the induction hypothesis, we get that $\Gamma \models \ell_\Delta^x[\text{PI}(C_1) \vee (D \vee s = t)_\Gamma]$ and $\Gamma \models \ell_\Delta^x[\text{PI}(C_2) \vee (E[r]_p)_\Gamma]$. This is by Lemma 38 equivalent to (\circ) $\Gamma \models \ell_\Delta^x[\text{PI}(C_1)] \vee \ell_\Delta^x[D_\Gamma] \vee \ell_\Delta^x[s] = \ell_\Delta^x[t]$ and $(*)$ $\Gamma \models \ell_\Delta^x[\text{PI}(C_2)] \vee \ell_\Delta^x[(E[r]_p)_\Gamma]$ respectively.

We distinguish two cases:

1. Suppose s does not occur in a maximal Δ -term $h[s]$ in $E[s]_p$, which occurs more than once in $\text{PI}(C_2) \vee (E[r]_p)_\Gamma$.

Let M be a model of Γ . First, assume that $M \models \ell_\Delta^x[s] \neq \ell_\Delta^x[t]$. Then by (\circ) we have that $M \models \ell_\Delta^x[\text{PI}(C_1)] \vee \ell_\Delta^x[D_\Gamma]$. By applying σ' and Lemma 40, we hence can conclude from $M \models \ell_\Delta^x[s\sigma] \neq \ell_\Delta^x[t\sigma]$ that $M \models \ell_\Delta^x[\text{PI}(C_1)\sigma] \vee \ell_\Delta^x[D_\Gamma\sigma]$.

Second, assume to the contrary that $M \models \ell_\Delta^x[s] = \ell_\Delta^x[t]$. We distinguish cases:

- Suppose that s does not occur in a maximal Δ -term in $(E[s]_p)_\Gamma$. Then by Lemma 42 $M \models \ell_\Delta^x[(E[t]_p)_\Gamma] \Leftrightarrow M \models \ell_\Delta^x[(E[s]_t)_\Gamma]$. Due to $\sigma = \text{mgu}(s, r)$, $s\sigma \equiv r\sigma$. Suppose they are both not Δ -colored. Then the lifting does not affect them and $\ell_\Delta^x[(E[s]_p)_\Gamma\sigma] \equiv \ell_\Delta^x[(E[r]_p)_\Gamma\sigma]$. Otherwise the lifting will replace them with the same variable and we as well get that $\ell_\Delta^x[(E[s]_p)_\Gamma\sigma] \equiv \ell_\Delta^x[(E[r]_p)_\Gamma\sigma]$. Hence $M \models \ell_\Delta^x[(E[t]_p)_\Gamma\sigma] \Leftrightarrow M \models \ell_\Delta^x[(E[r]_p)_\Gamma\sigma]$
- Otherwise r occurs in a maximal Δ -term $h[r]$ in $(E[r]_p)_\Gamma$, but $h[r]$ does not occur elsewhere in $\text{PI}(C_2) \vee (E[r]_p)_\Gamma$. Then the lifting variable $x_{h[r]}$ occurs only once in $(*)$. Hence Γ does not pose any restriction on $x_{h[r]}$ and we can substitute it by $x_{h[t]}$. So in M we have that $M \models \ell_\Delta^x[(E[t]_p)_\Gamma] \Leftrightarrow M \models \ell_\Delta^x[(E[r]_p)_\Gamma]$.

Hence in any of the cases, by $(*)$ and by applying σ' and Lemma 40 it follows from $M \models \ell_\Delta^x[s\sigma] = \ell_\Delta^x[t\sigma]$ that $M \models \ell_\Delta^x[\text{PI}(C_2)\sigma] \vee \ell_\Delta^x[(E[t]_p)_\Gamma\sigma]$.

In conclusion, we can derive that $\Gamma \models (\ell_\Delta^x[s\sigma] \neq \ell_\Delta^x[t\sigma] \wedge (\ell_\Delta^x[\text{PI}(C_1)\sigma] \vee \ell_\Delta^x[D_\Gamma\sigma])) \vee (\ell_\Delta^x[s\sigma] = \ell_\Delta^x[t\sigma] \wedge (\ell_\Delta^x[\text{PI}(C_2)\sigma] \vee \ell_\Delta^x[(E[t]_p)_\Gamma\sigma]))$. This however implies that $\Gamma \models (\ell_\Delta^x[s\sigma] \neq \ell_\Delta^x[t\sigma] \wedge (\ell_\Delta^x[\text{PI}(C_1)\sigma])) \vee (\ell_\Delta^x[s\sigma] = \ell_\Delta^x[t\sigma] \wedge (\ell_\Delta^x[\text{PI}(C_2)\sigma])) \vee (\ell_\Delta^x[D_\Gamma\sigma] \vee \ell_\Delta^x[(E[t]_p)_\Gamma\sigma])$, which by Lemma 38 is nothing else than $\Gamma \models \ell_\Delta^x[\text{PI}(C) \vee C]$.

2. Otherwise s occurs in a maximal Δ -term $h[s]_q$ in $E[s]_p$, which occurs more than once in $E[s]_p$.

Then a similar line of argument as in case 1 can be employed, with the difference that the application of Lemma 42 in the case of $M \models \ell_\Delta^x[s] = \ell_\Delta^x[t]$ yields the additional possibility that $M \models \ell_\Delta^x[h[s]] \neq \ell_\Delta^x[h[t]]$. Hence we arrive at:

$$\begin{aligned} \Gamma \models & \left(\ell_\Delta^x[s]\sigma' = \ell_\Delta^x[t]\sigma' \wedge \ell_\Delta^x[\text{PI}(C_2)]\sigma' \right) \vee \left(\ell_\Delta^x[s]\sigma' \neq \ell_\Delta^x[t]\sigma' \wedge \right. \\ & \left. \ell_\Delta^x[\text{PI}(C_1)]\sigma' \right) \vee \left(\ell_\Delta^x[s]\sigma' = \ell_\Delta^x[t]\sigma' \wedge (\ell_\Delta^x[h[s]]\sigma' \neq \ell_\Delta^x[h[t]]\sigma') \right) \vee \\ & \left(\ell_\Delta^x[D_\Gamma]\sigma' \vee \ell_\Delta^x[(E[t]_p)_\Gamma]\sigma' \right) \end{aligned}$$

This however is by Lemma 40 and Lemma 38 equivalent to $\Gamma \models \ell_\Delta^x[\text{PI}(C) \vee C]$. \square

4.6 Quantifying over lifting variables

As we have already seen in Corollary 46 that $\text{PI}(\pi)$ forms a propositional interpolant, it remains to lift all colored terms and quantify over the resulting lifting variables in a viable order.

Lemma 47. *For a formula or term φ , $\ell_\Gamma^y[\ell_\Delta^x[\varphi]] = \ell_\Delta^x[\ell_\Gamma^y[\varphi]]$.*

Proof. Let φ be a mixed-colored term as otherwise we are done. Suppose without loss of generality that it is a Γ -term which contains a maximal Δ -term t at position p . Then $\ell_\Delta^x[\ell_\Gamma^y[\varphi]] = \ell_\Delta^x[y_\varphi] = y_\varphi$.

On the other hand $\ell_\Gamma^y[\ell_\Delta^x[\varphi]] = \ell_\Gamma^y[\psi]$ such that ψ is equal to φ besides having x_t at position p . But $\ell_\Gamma^y[\psi] = y_{\text{unfold-lift}(\psi)} = y_\varphi$. \square

Theorem 48. *Let π be a resolution refutation of $\Gamma \cup \Delta$ and z_1, \dots, z_n be the variables which replace the colored terms in $\ell_\Gamma^y[\ell_\Delta^x[\text{PI}(\pi)]]$ ordered by their length. Then $Q_1 z_1 \dots Q_n z_n \ell_\Gamma^y[\ell_\Delta^x[\text{PI}(\pi)]]$, where Q_i is \forall (\exists) if z_i replaces a Δ (Γ)-term, is an interpolant.*

Proof. By Lemma 45, $\Gamma \models \forall x_1 \dots \forall x_m \ell_\Delta^x[\text{PI}(\pi)]$ where m is the number of maximal Δ -colored terms in $\text{PI}(\pi)$.

A term in $\ell_\Delta^x[\text{PI}(\pi)]$ is either x_i , $1 \leq i \leq m$, a grey term or a Γ -terms. Let t be a maximal Γ -term in $\ell_\Delta^x[\text{PI}(\pi)]$ and x_{j_1}, \dots, x_{j_k} the variables replacing Δ -terms in t . Note that the Δ -terms, which are replaced by x_{j_1}, \dots, x_{j_k} respectively, are each of strictly smaller size than t as they are strict subterms of t .

In $\ell_\Gamma^y[\ell_\Delta^x[\text{PI}(\pi)]]$, t is replaced by some z_j , which is existentially quantified. Hence t is a witness for z_j as due to the quantifier ordering, the existential quantification of z_j is in the scope of the quantifiers of x_{j_1}, \dots, x_{j_k} respectively. Therefore $\Gamma \models Q_1 z_1 \dots Q_n z_n \ell_\Gamma^y[\ell_\Delta^x[\text{PI}(\pi)]]$.

By Corollary 44 $\Delta \models \forall y_1 \dots \forall y_k \neg \ell_\Gamma^y[\text{PI}(\pi)]$, where k is the number of Γ -colored terms in $\text{PI}(\pi)$. By a similar line of argumentation as above, we can replace the maximal Δ -terms by existentially quantified variables and arrive at $\Delta \models \overline{Q}_1 z_1 \dots \overline{Q}_n z_n \neg \ell_\Delta^x[\ell_\Gamma^y[\text{PI}(\pi)]]$ where $\overline{Q}_i = \exists$ (\forall) if $Q_i = \forall$ (\exists). Therefore also $\Delta \models \neg Q_1 z_1 \dots Q_n z_n \ell_\Delta^x[\ell_\Gamma^y[\text{PI}(\pi)]]$. By Lemma 47, $\Delta \models \neg Q_1 z_1 \dots Q_n z_n \ell_\Gamma^y[\ell_\Delta^x[\text{PI}(\pi)]]$.

As it is now easy to see that $Q_1 z_1 \dots Q_n z_n \ell_\Gamma^y[\ell_\Delta^x[\text{PI}(\pi)]]$ contains no colored symbol, it is an interpolant. \square

WT: Huang's proof

A.1 Propositional interpolants

Let $\Gamma \cup \Delta$ be unsatisfiable. Let π be a proof of the empty clause from $\Gamma \cup \Delta$. Then PI is a function that returns a interpolant with respect to the current clause.

Definition 49 (Propositional interpolant). Let π be a resolution refutation of $\Gamma \cup \Delta$. A formula A is a *propositional interpolant* if

1. $\Gamma \models A$
2. $\Delta \models \neg A$
3. $\text{PS}(A) \subseteq (\text{PS}(\Gamma) \cap \text{PS}(\Delta)) \cup \{\top, \perp\}$.

For a clause C in π , a formula A_C is a *propositional interpolant relative to C* if

1. $\Gamma \models A_C \vee C$
2. $\Delta \models \neg A_C \vee C$
3. $\text{PS}(A_C) \subseteq (\text{PS}(\Gamma) \cap \text{PS}(\Delta)) \cup \{\top, \perp\}$.

The propositional interpolant of the empty clause derived in π is denoted by $\text{PI}(\pi)$. \triangle

The third condition of a propositional interpolant will sometimes be referred to as *language restriction*. It is easy to see that the propositional interpolant relative to the empty clause of a resolution refutation is a propositional interpolant.

We proceed by defining a procedure PI which extracts propositional interpolants from a resolution refutation.

Definition 50 (Propositional interpolant extraction.). Let π be a resolution refutation of $\Gamma \cup \Delta$. $\text{PI}(\pi)$ is defined to be $\text{PI}(\square)$, where \square is the empty clause derived in π .

For a clause C in π , $\text{PI}(C)$ is defined as follows:

Base case. If $C \in \Gamma$, $\text{PI}(C) = \perp$. If otherwise $C \in \Delta$, $\text{PI}(C) = \top$.

Resolution. If the clause C is the result of a resolution step of $C_1 : D \vee l$ and $C_2 : E \vee \neg l'$ using a unifier σ such that $l\sigma = l'\sigma$, then $\text{PI}(C)$ is defined as follows:

1. If l is Γ -colored: $\text{PI}(C) = [\text{PI}(C_1) \vee \text{PI}(C_2)]\sigma$
2. If l is Δ -colored: $\text{PI}(C) = [\text{PI}(C_1) \wedge \text{PI}(C_2)]\sigma$
3. If l is grey: $\text{PI}(C) = [(l \wedge \text{PI}(C_2)) \vee (\neg l' \wedge \text{PI}(C_1))]\sigma$

Factorisation. If the clause C is the result of a factorisation of $C_1 : l \vee l' \vee D$ using a unifier σ such that $l\sigma = l'\sigma$, then $\text{PI}(C) = \text{PI}(C_1)\sigma$.

Paramodulation. Suppose the clause C is the result of a paramodulation of $C_1 : s = t \vee D$ and $C_2 : E[r]$ using a unifier σ such that $r\sigma = s\sigma$. Let $h[r]$ be the maximal colored term in which r occurs in $E[r]$. Then $\text{PI}(C)$ is defined according to the following case distinction:

1. If $h[r]$ is Δ -colored and $h[r]$ occurs more than once in $E[r] \vee \text{PI}(E[r])$:

$$\text{PI}(C) = [(s = t \wedge \text{PI}(C_2)) \vee (s \neq t \wedge \text{PI}(C_1))]\sigma \vee (s = t \wedge h[s] \neq h[t])\sigma$$

2. If $h[r]$ is Γ -colored and $h[r]$ occurs more than once in $E[r] \vee \text{PI}(E[r])$:

$$\text{PI}(C) = [(s = t \wedge \text{PI}(C_2)) \vee (s \neq t \wedge \text{PI}(C_1))]\sigma \wedge (s \neq t \vee h[s] = h[t])\sigma$$

3. Otherwise:

$$\text{PI}(C) = [(s = t \wedge \text{PI}(C_2)) \vee (s \neq t \wedge \text{PI}(C_1))]\sigma \quad \triangle$$

Proposition 51. Let C be a clause of a resolution refutation of $\Gamma \cup \Delta$. Then $\text{PI}(C)$ is a propositional interpolant with respect to C .

Proof. Proof by induction on the number of rule applications including the following strengthenings: $\Gamma \models \text{PI}(C) \vee C_\Gamma$ and $\Delta \models \neg \text{PI}(C) \vee C_\Delta$, where C_Φ denotes the clause D with only the literals which are contained in $L(\Phi)$. They clearly imply conditions 1 and 2 of definition 49.

Base case. Suppose no rules were applied. We distinguish two possible cases:

1. $C \in \Gamma$. Then $\text{PI}(C) = \perp$. Clearly $\Gamma \models \perp \vee C_\Gamma$ as $C_\Gamma = C \in \Gamma$, $\Delta \models \neg \perp \vee C_\Delta$ and \perp satisfies the restriction on the language.
2. $C \in \Delta$. Then $\text{PI}(C) = \top$. Clearly $\Gamma \models \top \vee C_\Gamma$, $\Delta \models \neg \top \vee C_\Delta$ as $C_\Delta = C \in \Delta$ and \top satisfies the restriction on the language.

Suppose the property holds for n rule applications. We show that it holds for $n + 1$ applications by considering the last one:

Resolution. Suppose the last rule application is an instance of resolution. Then it is of the form:

$$\frac{C_1 : D \vee l \quad C_2 : E \vee \neg l'}{C : (D \vee E)\sigma} \quad l\sigma = l'\sigma$$

By the induction hypothesis, we can assume that:

$$\begin{aligned} \Gamma &\models \text{PI}(C_1) \vee (D \vee l)_\Gamma \\ \Delta &\models \neg \text{PI}(C_1) \vee (D \vee l)_\Delta \\ \Gamma &\models \text{PI}(C_2) \vee (E \vee \neg l')_\Gamma \\ \Delta &\models \neg \text{PI}(C_2) \vee (E \vee \neg l')_\Delta \end{aligned}$$

We consider the respective cases from definition 50:

1. l is Γ -colored. Then $\text{PI}(C) = [\text{PI}(C_1) \vee \text{PI}(C_2)]\sigma$.
As $\text{PS}(l) \in L(\Gamma)$, $\Gamma \models (\text{PI}(C_1) \vee D_\Gamma \vee l)\sigma$ as well as $\Gamma \models (\text{PI}(C_2) \vee E_\Gamma \vee \neg l')\sigma$. By a resolution step, we get $\Gamma \models (\text{PI}(C_1) \vee \text{PI}(C_2))\sigma \vee ((D \vee E)\sigma)_\Gamma$.
Furthermore, as $\text{PS}(l) \notin L(\Delta)$, $\Delta \models (\neg \text{PI}(C_1) \vee D_\Delta)\sigma$ as well as $\Delta \models (\neg \text{PI}(C_2) \vee E_\Delta)\sigma$. Hence it certainly holds that $\Delta \models (\neg \text{PI}(C_1) \vee \neg \text{PI}(C_2))\sigma \vee (D \vee E)\sigma_\Delta$.
The language restriction clearly remains satisfied as no non-logical symbols are added.
2. l is Δ -colored. Then $\text{PI}(C) = [\text{PI}(C_1) \wedge \text{PI}(C_2)]\sigma$.
As $\text{PS}(l) \notin L(\Gamma)$, $\Gamma \models (\text{PI}(C_1) \vee D_\Gamma)\sigma$ as well as $\Gamma \models (\text{PI}(C_2) \vee E_\Gamma)\sigma$. Suppose that in a model M of Γ , $M \not\models D_\Gamma$ and $M \not\models E_\Gamma$. Then $M \models \text{PI}(C_1) \wedge \text{PI}(C_2)$. Hence $\Gamma \models (\text{PI}(C_1) \wedge \text{PI}(C_2))\sigma \vee ((D \vee E)\sigma)_\Gamma$.
Furthermore due to $\text{PS}(l) \in L(\Delta)$, $\Delta \models (\neg \text{PI}(C_1) \vee D_\Delta \vee l)\sigma$ as well as $\Delta \models (\neg \text{PI}(C_2) \vee E_\Delta \vee \neg l')\sigma$. By a resolution step, we get $\Delta \models (\neg \text{PI}(C_1) \vee \neg \text{PI}(C_2))\sigma \vee (D_\Delta \vee E_\Delta)\sigma$ and hence $\Delta \models \neg(\text{PI}(C_1) \wedge \text{PI}(C_2))\sigma \vee (D_\Delta \vee E_\Delta)\sigma$.

The language restriction again remains intact.

3. l is grey. Then $\text{PI}(C) = [(l \wedge \text{PI}(C_2)) \vee (\neg l' \wedge \text{PI}(C_1))]\sigma$

First, we have to show that $\Gamma \models [(l \wedge \text{PI}(C_2)) \vee (l' \wedge \text{PI}(C_1))]\sigma \vee ((D \vee E)\sigma)_\Gamma$. Suppose that in a model M of Γ , $M \not\models D_\Gamma$ and $\Gamma \not\models E$. Otherwise we are done. The induction assumption hence simplifies to $M \models \text{PI}(C_1) \vee l$ and $M \models \text{PI}(C_2) \vee \neg l'$ respectively. As $l\sigma = l'\sigma$, by a case distinction argument on the truth value of $l\sigma$, we get that either $M \models (l \wedge \text{PI}(C_2))\sigma$ or $M \models (\neg l' \wedge \text{PI}(C_1))\sigma$.

Second, we show that $\Delta \models ((l \vee \neg \text{PI}(C_1)) \wedge (\neg l' \vee \neg \text{PI}(C_2)))\sigma \vee ((D \vee E)\sigma)_\Delta$. Suppose again that in a model M of Δ , $M \not\models D_\Delta$ and $\Gamma \not\models E_\Delta$. Then the required statement follows from the induction hypothesis.

The language condition remains satisfied as only the common literal l is added to the interpolant.

Factorisation. Suppose the last rule application is an instance of factorisation. Then it is of the form:

$$\frac{C_1 : l \vee l' \vee D}{C : (l \vee D)\sigma} \quad \sigma = \text{mgu}(l, l')$$

Then the propositional interpolant $\text{PI}(C)$ is defined as $\text{PI}(C_1)$. By the induction hypothesis, we have:

$$\Gamma \models \text{PI}(C_1) \vee (l \vee l' \vee D)_\Gamma$$

$$\Delta \models \text{PI}(C_1) \vee (l \vee l' \vee D)_\Delta$$

It is easy to see that then also:

$$\Gamma \models (\text{PI}(C_1) \vee (l \vee D))_\Gamma \sigma$$

$$\Delta \models (\text{PI}(C_1) \vee (l \vee D))_\Delta \sigma$$

The restriction on the language trivially remains intact.

Paramodulation. Suppose the last rule application is an instance of paramodulation. Then it is of the form:

$$\frac{C_1 : D \vee s = t \quad C_2 : E[s]_p}{C : D \vee E[t]_p} \quad \sigma = \text{mgu}(s, r)$$

By the induction hypothesis, we have:

$$\Gamma \models \text{PI}(C_1) \vee (D \vee s = t)_\Gamma$$

$$\Delta \models \neg \text{PI}(C_1) \vee (D \vee s = t)_\Delta$$

$$\Gamma \models \text{PI}(C_2) \vee (E[r])_\Gamma$$

$$\Delta \models \neg \text{PI}(C_2) \vee (E[r])_\Delta$$

First, we show that $\text{PI}(C)$ as constructed in case 3 of the definition is a propositional interpolant in any of these cases:

$$\text{PI}(C) = (s = t \wedge \text{PI}(C_2)) \vee (s \neq t \wedge \text{PI}(C_1))$$

Suppose that in a model M of Γ , $M \not\models D\sigma$ and $M \not\models E[t]_p\sigma$. Otherwise we are done. Furthermore, assume that $M \models (s = t)\sigma$. Then $M \not\models E[r]_p\sigma$, but then necessarily $M \models \text{PI}(C_2)\sigma$.

On the other hand, suppose $M \models (s \neq t)\sigma$. As also $M \not\models D\sigma$, $M \models \text{PI}(C_1)\sigma$. Consequently, $M \models [(s = t \wedge \text{PI}(C_2)) \vee (s \neq t \wedge \text{PI}(C_1))]\sigma \vee [(D \vee E)_\Gamma]\sigma$

By an analogous argument, we get $\Delta \models [(s = t \wedge \neg \text{PI}(C_2)) \vee (s \neq t \wedge \neg \text{PI}(C_1))]\sigma \vee [(D \vee E)_\Delta]\sigma$, which implies $\Delta \models [(s \neq t \vee \neg \text{PI}(C_2)) \wedge (s = t \vee \neg \text{PI}(C_1))]\sigma \vee ((D \vee E)_\Delta)\sigma$

The language restriction again remains satisfied as the only predicate, that is added to the interpolant, is $=$.

This concludes the argumentation for case 3.

The interpolant for case 1 differs only by an additional formula added via a disjunction and hence condition 1 of definition 49 holds by the above reasoning. As the adjoined formula is a contradiction, its negation is valid which in combination with the above reasoning establishes condition 2. Since no new predicated are added, the language condition remains intact.

The situation in case 2 is somewhat symmetric: As a tautology is added to the interpolant with respect to case 1, condition 1 is satisfied by the above reasoning. For condition 2, consider that the negated interpolant for case 1 implies the negated interpolant for this case. The language condition again remains intact. \square

A.2 Propositional refutations

Before we are able to specify a procedure to transform the propositional interpolant generated by PI into a proper interpolant without any colored terms, we need to make some observations about tree refutations.

In a tree refutation where the input clauses have a disjoint sets of variables, every variable has a unique ancestor which traces back to an input clause and hence appears only along a certain path. This insight allows us to push substitutions of the variables upwards along this path and arrive at the following definition and lemma:

Definition 52. A resolution refutation is a *propositional refutation* if no non-trivial substitutions are employed. \triangle

Lemma 53. *Let Φ be unsatisfiable. Then there is a propositional refutation of Φ which starts from instances of Φ .*

Proof. Let π be a resolution refutation of Φ . By Lemma 11, we can assume without loss of generality that π is a tree refutation where the sets of variables of the input clauses are disjoint. Furthermore, we can assume that only most general unifiers are employed in π .

Then any unifier in π is either trivial on x or there is one unique unifier σ in π with $x\sigma = t$ where x does not occur in t . Hence along the path through the deduction where x occurs, it remains unchanged. Therefore we can create a new resolution refutation π' from π where x is replaced by t . Clearly π' is rooted in instances of Φ .

By application of this procedure to all variable occurring in π , we obtain a desired resolution refutation. \square

Even though propositional refutations have nice properties for theoretical analysis, their use in practise is not desired as its construction involves a considerable blowup of the refutation. But its use is still justified in this instance as we can show for arbitrary refutations π that the algorithm stated in 50 gives closely related results for both π and its corresponding propositional refutation.

Lemma 54. *Let π be a resolution refutation of Φ and π' a propositional refutation corresponding to π . Then for every clause C in π and its corresponding clause C' in π' , $\text{PI}(C)\sigma = \text{PI}(C')$, where σ is the composition of the unifications of π which are applied to the variables occurring in C .*

Proof. For the construction of the propositional skeleton of $\text{PI}(\cdot)$ only the coloring of the clauses is relevant and since this is the same in both π and π' , it coincides for $\text{PI}(C)$ and $\text{PI}(C')$.

Hence $\text{PI}(C)$ and $\text{PI}(C')$ differ only in their term structure. To be more specific, in $\text{PI}(C')$, the composition of substitutions that are applied in π have already been applied to the initial clauses of π' . Note that substitution commutes with the rules of resolution. Therefore the only difference between $\text{PI}(C)$ and $\text{PI}(C')$ is that at certain term positions, there are variables in $\text{PI}(C)$ where in $\text{PI}(C')$ by some substitution a different term is located. But these substitutions are certainly applied by σ , hence $\text{PI}(C)\sigma = \text{PI}(C')$. \square

this part occurs in other proof

A.3 Lifting of colored symbols

This establishes the theoretical framework which is required to define and show the correctness of the procedure to construct a proper interpolant from the propositional interpolant. The idea of this procedure will be to replace colored terms still occurring in the propositional interpolant with variables and quantifying them appropriately. This replacement is referred to as lifting:

Definition 55 (Lifting). Let Γ and Δ be sets of first-order formulas, ϕ a formula or a term, $Z = \{\zeta_1, \dots, \zeta_n\}$ the maximal Φ -terms for $\Phi \in \{\Gamma, \Delta\}$ in ϕ and z_1, \dots, z_n fresh variables, referred to as *lifting variables*.

Then $\ell_\Phi^z[\phi]$ denotes $\ell_\Phi^z[\phi, Z]$ which is defined as follows:

$$\ell_\Phi^z[\phi, Z] \stackrel{\text{def}}{=} \begin{cases} \phi & Z = \emptyset \\ \ell_\Phi^z[\phi\{\zeta_i/z_i\}, Z \setminus \{\zeta_i\}] & \zeta_i \in Z \text{ such that } \zeta_i \text{ is not subterm of another term in } Z \end{cases}$$

We also denote ϕ lifted of both Γ - and Δ -terms by $\ell[\phi]$ if the variables replacing the colored terms are clear from the context or are not crucial. \triangle

We usually lift Δ -terms by indexed variables with letter x and Γ -terms with letter y . If the lifting is not specific to a color, we use indexed variables with letter z .

Some elementary properties of liftings are described by the following lemmas:

Lemma 56. *Let A and B be first-order formulas and s and t be terms. Then it holds that:*

1. $\ell_\Phi^z[\neg A] \Leftrightarrow \neg \ell_\Phi^z[A]$
2. $\ell_\Phi^z[A \circ B] \Leftrightarrow (\ell_\Phi^z[A] \circ \ell_\Phi^z[B])$ for $\circ \in \{\wedge, \vee\}$
3. $\ell_\Phi^z[s = t] \Leftrightarrow (\ell_\Phi^z[s] = \ell_\Phi^z[t])$

Lemma 57. $\ell_\Gamma^y[\ell_\Delta^x[C]]$ and $\ell_\Delta^{x'}[\ell_\Gamma^{y'}[C]]$ differ only in the naming of the variables replacing maximal colored terms.

not quite right with unfold-lift, see proof without P_P

Proof. Suppose a term t in C is affected by a lifting. We only need to consider maximal colored terms as grey terms are not affected by the liftings. Without loss of generality let t be a maximal Δ -colored term.

Let Φ be the positions of maximal occurrences of t . Then in the left hand side, exactly all terms at positions Φ are replaced by x_i for some i .

In the right hand side, all terms at positions Φ are replaced by $\ell_\Gamma^{y'}[t]$ first. However after this step, all these terms are equal to $\ell_\Gamma^{y'}[t]$, and as all distinct maximal Γ -terms are replaced by distinct variables, no other maximal colored term is equal to $\ell_\Gamma^{y'}[t]$. Hence exactly the terms at positions Φ are replaced by the same variable x'_j for some j . \square

First, we consider the lifting of the Δ -terms:

Lemma 58. *Let π be a resolution refutation of $\Gamma \cup \Delta$. Then $\Gamma \models \ell_\Delta^x[\text{PI}(C) \vee C]$ for C in π .*

Proof. We proof this result by induction on the number of rule applications in the propositional refutation corresponding to π . Similar to the proof of 51, we show the strengthening: $\Gamma \models \ell_\Delta^x[\text{PI}(C) \vee C_\Gamma]$ for C in π .

Base case. If no rules have been applied, C is an instance of a clause of either Γ or Δ . In the former case, all Δ -terms of C were added by unification, hence by replacing them with variables, we obtain a clause C' which still is an instance of C and consequently is implied by Γ . In the latter case, $\text{PI}(C) = \top$.

Resolution. Suppose the last rule application is an instance of resolution. Then it is of the form:

$$\frac{C_1 : D \vee l \quad C_2 : E \vee \neg l}{C : D \vee E}$$

By the induction hypothesis,

$$\Gamma \models \ell_\Delta^x[\text{PI}(C_1) \vee (D \vee l)_\Gamma] \text{ and}$$

$$\Gamma \models \ell_\Delta^x[\text{PI}(C_2) \vee (E \vee \neg l)_\Gamma]$$

which by Lemma 56 is equivalent to

$$\Gamma \models \ell_\Delta^x[\text{PI}(C_1)] \vee \ell_\Delta^x[D_\Gamma] \vee \ell_\Delta^x[l_\Gamma]^{(\circ)} \text{ and}$$

$$\Gamma \models \ell_\Delta^x[\text{PI}(C_2)] \vee \ell_\Delta^x[E_\Gamma] \vee \neg \ell_\Delta^x[l_\Gamma]^{(*)}.$$

1. Suppose l is Γ -colored. Then $\text{PI}(C) = \text{PI}(C_1) \vee \text{PI}(C_2)$. By using resolution of $(*)$ and (\circ) on $\ell_\Delta^x[l_\Gamma]$, we get that

$$\Gamma \models \ell_\Delta^x[\text{PI}(C_1)] \vee \ell_\Delta^x[\text{PI}(C_2)] \vee \ell_\Delta^x[D_\Gamma] \vee \ell_\Delta^x[E_\Gamma].$$

Several applications of Lemma 56 give $\Gamma \models \ell_\Delta^x[\text{PI}(C_1) \vee \text{PI}(C_2) \vee (D \vee E)_\Gamma]$.

2. Suppose l is Δ -colored. Then $\text{PI}(C) = \text{PI}(C_1) \wedge \text{PI}(C_2)$.
 As l and $\neg l$ are not contained in $L(\Gamma)$, we get that
 $\Gamma \models \ell_\Delta^x[\text{PI}(C_1)] \vee \ell_\Delta^x[D_\Gamma]$ and
 $\Gamma \models \ell_\Delta^x[\text{PI}(C_2)] \vee \ell_\Delta^x[E_\Gamma]$.
 So if in a model M of Γ we have that $M \not\models \ell_\Delta^x[D_\Gamma]$ and $M \not\models \ell_\Delta^x[E_\Gamma]$,
 it follows that $M \models \ell_\Delta^x[\text{PI}(C_1)]$ and $M \models \ell_\Delta^x[\text{PI}(C_2)]$. Hence by
 Lemma 56 $M \models \ell_\Delta^x[\text{PI}(C_1) \wedge \text{PI}(C_2)] \vee \ell_\Delta^x[(D \vee E)_\Gamma]$.
3. Suppose l is grey. Then $\text{PI}(C) = (l \wedge \text{PI}(C_2)) \vee (\neg l \wedge \text{PI}(C_1))$.
 We show that $\Gamma \models \ell_\Delta^x[(l \wedge \text{PI}(C_2)) \vee (\neg l \wedge \text{PI}(C_1)) \vee (D \vee E)_\Gamma]$.
 Suppose that for a model M of Γ that $M \not\models \ell_\Delta^x[D_\Gamma]$ and $M \not\models \ell_\Delta^x[E_\Gamma]$.
 Then by (\circ) and $(*)$, we get that
 $M \models \ell_\Delta^x[\text{PI}(C_1)] \vee \ell_\Delta^x[l_\Gamma]$ as well as
 $M \models \ell_\Delta^x[\text{PI}(C_2)] \vee \neg \ell_\Delta^x[l_\Gamma]$.
 So $M \models \ell_\Delta^x[l_\Gamma]$ implies that $M \models \ell_\Delta^x[\text{PI}(C_2)]$ and $M \models \neg \ell_\Delta^x[l_\Gamma]$ implies
 that $M \models \ell_\Delta^x[\text{PI}(C_1)]$ and
 Therefore $M \models (\ell_\Delta^x[l] \wedge \ell_\Delta^x[\text{PI}(C_2)]) \vee (\neg \ell_\Delta^x[l] \wedge \ell_\Delta^x[\text{PI}(C_1)]) \vee (\ell_\Delta^x[D_\Gamma] \vee \ell_\Delta^x[E_\Gamma])$,
 and several applications of Lemma 56 give $M \models \ell_\Delta^x[(l \wedge \text{PI}(C_2)) \vee (\neg l \wedge \text{PI}(C_1)) \vee (D_\Gamma \vee E_\Gamma)]$.

Factorisation. Suppose the last rule application is an instance of factorisation.
 Then it is of the form:

$$\frac{C_1 : l \vee l \vee D}{C : l \vee D}$$

The propositional interpolant directly carried over from C_1 , i.e. $\text{PI}(C) = \text{PI}(C_1)$.

By the induction hypothesis, we get that $\Gamma \models \ell_\Delta^x[\text{PI}(C_1) \vee (l \vee l \vee D)_\Gamma]$. By Lemma 56,

$$\Gamma \models \ell_\Delta^x[\text{PI}(C_1)] \vee (\ell_\Delta^x[l_\Gamma] \vee \ell_\Delta^x[l_\Gamma] \vee \ell_\Delta^x[D_\Gamma]),$$

which clearly is equivalent to

$$\Gamma \models \ell_\Delta^x[\text{PI}(C_1)] \vee (\ell_\Delta^x[l_\Gamma] \vee \ell_\Delta^x[D_\Gamma]),$$

so by again applying Lemma 56, we arrive at

$$\Gamma \models \ell_\Delta^x[\text{PI}(C_1) \vee (l \vee D)_\Gamma].$$

Paramodulation. Suppose the last rule application is an instance of paramodulation. Then it is of the form:

$$\frac{C_1 : D \vee s = t \quad C_2 : E[s]_p}{C : D \vee E[t]_p}$$

By the induction hypothesis, we have that

$\Gamma \models \ell_\Delta^x[\text{PI}(C_1) \vee (D \vee s = t)]_\Gamma$ and

$\Gamma \models \ell_\Delta^x[\text{PI}(C_2) \vee (E[s]_p)]_\Gamma$.

By Lemma 56, we get that

$\Gamma \models \ell_\Delta^x[\text{PI}(C_1)] \vee \ell_\Delta^x[D_\Gamma] \vee \ell_\Delta^x[s] = \ell_\Delta^x[t]$ and

$\Gamma \models \ell_\Delta^x[\text{PI}(C_2)] \vee \ell_\Delta^x[(E[s]_p)_\Gamma]$.

We distinguish two cases:

1. Suppose s does not occur in a maximal Δ -term $h[s]$ in $E[s]_p$ which occurs more than once in $\text{PI}(E(s)) \vee E[s]_p$.

We show that $\Gamma \models \ell_\Delta^x[(s = t \wedge \text{PI}(C_2)) \vee (s \neq t \wedge \text{PI}(C_1)) \vee (D \vee E[t]_p)]_\Gamma$, which subsumes the cases 2 and 3 of Definition 50. By Lemma 56, this is equivalent to

$$\Gamma \models (\ell_\Delta^x[s] = \ell_\Delta^x[t] \wedge \ell_\Delta^x[\text{PI}(C_2)]) \vee (\ell_\Delta^x[s] \neq \ell_\Delta^x[t] \wedge \ell_\Delta^x[\text{PI}(C_1)]) \vee (\ell_\Delta^x[D_\Gamma] \vee \ell_\Delta^x[(E[t]_p)_\Gamma])$$

Suppose that in a model M of Γ , $M \not\models \ell_\Delta^x[D_\Gamma]$ and $M \not\models \ell_\Delta^x[(E[t]_p)_\Gamma]$. We show that then, depending on whether $\ell_\Delta^x[s] = \ell_\Delta^x[t]$ holds in M , one of the first two disjuncts holds in M .

Then in case $M \models \ell_\Delta^x[s] = \ell_\Delta^x[t]$ we also get $M \not\models \ell_\Delta^x[(E[s]_p)_\Gamma]$ and consequently by the induction hypothesis $M \models \ell_\Delta^x[\text{PI}(C_2)]$.

However in case $M \models \ell_\Delta^x[s] \neq \ell_\Delta^x[t]$ we get by the induction hypothesis that $M \models \ell_\Delta^x[\text{PI}(C_1)]$.

2. Otherwise s occurs in a maximal Δ -term $h[s]$ in $E[s]_p$ which occurs more than once in $\text{PI}(E(s)) \vee E[s]_p$. This reflects case 1 of Definition 50.

Then models are possible in which $s = t$ and therefore $\ell_\Delta^x[s] = \ell_\Delta^x[t]$ holds, while at the same time $\ell_\Delta^x[h[s]] \neq \ell_\Delta^x[h[t]]$ does not as $h[s]$ and $h[t]$ are replaced by distinct variables due to being different Δ -terms.

Therefore we amend the proof of case 1 as follows:

In case $M \models \ell_\Delta^x[s] = \ell_\Delta^x[t]$ (otherwise proceed as in case 1), one of the following cases holds:

- $M \models \ell_\Delta^x[h[s]] = \ell_\Delta^x[h[t]]$. From this, it follows that as in the proof of case 1, $M \not\models \ell_\Delta^x[(E[s]_p)_\Gamma]$ and consequently $M \models \ell_\Delta^x[\text{PI}(C_2)]$ again by the induction hypothesis.

- $M \models \ell_{\Delta}^x[h[s]] \neq \ell_{\Delta}^x[h[t]]$. However as here $\text{PI}(C)$ contains the with respect to case 1 additional disjunct $s = t \wedge h[s] \neq h[t]$, $M \models \ell_{\Delta}^x[\text{PI}(C)]$ due to $M \models \ell_{\Delta}^x[s] = \ell_{\Delta}^x[t] \wedge \ell_{\Delta}^x[h[s]] \neq \ell_{\Delta}^x[h[t]]$ \square

The interpolant extraction procedure PI exhibits a convenient property which is termed *symmetry* in [DKPW10, Definition 5] and can be stated formally as follows:

Lemma 59. *Let π be a resolution refutation of $\Gamma \cup \Delta$ and $\hat{\pi}$ be π with $\hat{\Gamma} = \Delta$ and $\hat{\Delta} = \Gamma$. Then $\text{PI}(\pi) \Leftrightarrow \neg \text{PI}(\hat{\pi})$.*

Proof. We prove this lemma by induction on π . Let $\hat{\varphi}$ denote the clause/formula/literal/term in $\hat{\pi}$ corresponding to the clause/formula/literal/term φ in π .

Base case. If $C \in \Gamma$, then $C' \in \Delta'$ and $\text{PI}(C) = \perp \Leftrightarrow \neg \top = \neg \text{PI}(C')$. The case for $C \in \Delta$ is analogous.

Resolution. If the clause C is the result of a resolution step of $C_1 : D \vee l$ and $C_2 : E \vee \neg l'$ using a unifier σ such that $l\sigma = l'\sigma$, then by induction hypothesis, we get that $\text{PI}(C_i) = \neg \text{PI}(C'_i)$ for $i \in \{1, 2\}$.

We distinguish the following cases:

1. l is Γ -colored. Then \hat{l} is Δ -colored.

$$\begin{aligned} \text{PI}(C) &= \text{PI}(C_1) \vee \text{PI}(C_2) \\ &\Leftrightarrow \neg(\neg \text{PI}(C_1) \wedge \neg \text{PI}(C_2)) \\ &= \neg(\text{PI}(\hat{C}_1) \wedge \text{PI}(\hat{C}_2)) \\ &= \neg \text{PI}(\hat{C}) \end{aligned}$$

2. l is Δ -colored. This case can be argued analogously.

3. l is grey. Then \hat{l} is grey.

$$\begin{aligned} \text{PI}(C) &= [(l \wedge \text{PI}(C_2)) \vee (\neg l' \wedge \text{PI}(C_1))]\sigma \\ &= (l\sigma \wedge \text{PI}(C_2)\sigma) \vee (\neg l'\sigma \wedge \text{PI}(C_1)\sigma) \\ &\Leftrightarrow (\neg l\sigma \vee \text{PI}(C_2)\sigma) \wedge (l'\sigma \vee \text{PI}(C_1)\sigma) \\ &\Leftrightarrow \neg[(l\sigma \wedge \neg \text{PI}(C_2)\sigma) \vee (\neg l'\sigma \wedge \neg \text{PI}(C_1)\sigma)] \\ &= \neg[(\hat{l}\sigma \wedge \neg \text{PI}(\hat{C}_2)\sigma) \vee (\neg \hat{l}'\sigma \wedge \neg \text{PI}(\hat{C}_1)\sigma)] \\ &= \neg[(\hat{l} \wedge \neg \text{PI}(\hat{C}_2)) \vee (\neg \hat{l}' \wedge \neg \text{PI}(\hat{C}_1))]\sigma \\ &= \neg[(\hat{l} \wedge \text{PI}(\hat{C}_2)) \vee (\neg \hat{l}' \wedge \text{PI}(\hat{C}_1))]\sigma \\ &= \neg \text{PI}(\hat{C}) \end{aligned}$$

Factorisation. Suppose the clause C is the result of a factorisation of $C_1 : l \vee l' \vee D$. Then $\text{PI}(C) = \text{PI}(C_1)\sigma$ and the induction hypothesis gives the result.

Paramodulation. Suppose the clause C is the result of a paramodulation of $C_1 : s = t \vee D$ and $C_2 : E[r]$ using a unifier σ such that $r\sigma = s\sigma$. We distinguish the following cases:

1. r occurs in a maximal Δ -term $h[r]$ in $E[r]$ and $h[r]$ occurs more than once in $E[r] \vee \text{PI}(E[r])$. Then \hat{r} occurs in a maximal Γ -term $\hat{h}[r]$ in $\hat{E}[r]$ and $\hat{h}[r]$ occurs more than once in $\hat{E}[r] \vee \text{PI}(\hat{E}[r])$.

$$\begin{aligned} \text{PI}(C) &= [(s = t \wedge \text{PI}(C_2)) \vee (s \neq t \wedge \text{PI}(C_1))]\sigma \vee (s = t \wedge h[s] \neq h[t])\sigma \\ &= [(s = t \wedge \neg \text{PI}(\hat{C}_2)) \vee (s \neq t \wedge \neg \text{PI}(\hat{C}_1))]\sigma \vee (s = t \wedge h[s] \neq h[t])\sigma \\ &\Leftrightarrow \neg[(s \neq t \vee \text{PI}(\hat{C}_2)) \wedge (s = t \vee \text{PI}(\hat{C}_1))]\sigma \wedge \neg(s \neq t \vee h[s] = h[t])\sigma \\ &\Leftrightarrow \neg[(s = t \wedge \text{PI}(\hat{C}_2)) \vee (s \neq t \wedge \text{PI}(\hat{C}_1))]\sigma \wedge \neg(s \neq t \vee h[s] = h[t])\sigma \\ &= \neg \text{PI}(\hat{C}) \end{aligned}$$

2. r occurs in a maximal Γ -term $h(r)$ in $E[r]$ and $h(r)$ occurs more than once in $E[r] \vee \text{PI}(E[r])$. This case can be argued analogously.
3. Otherwise:

$$\begin{aligned} \text{PI}(C) &= [(s = t \wedge \text{PI}(C_2)) \vee (s \neq t \wedge \text{PI}(C_1))]\sigma \\ &= [(s = t \wedge \neg \text{PI}(\hat{C}_2)) \vee (s \neq t \wedge \neg \text{PI}(\hat{C}_1))]\sigma \\ &\Leftrightarrow \neg[(s \neq t \vee \text{PI}(\hat{C}_2)) \wedge (s = t \vee \text{PI}(\hat{C}_1))]\sigma \\ &\Leftrightarrow \neg[(s = t \wedge \text{PI}(\hat{C}_2)) \vee (s \neq t \wedge \text{PI}(\hat{C}_1))]\sigma \\ &= \neg \text{PI}(\hat{C}) \end{aligned} \quad \square$$

This lemma can be leveraged to show a counterpart of Lemma 58 for Δ :

Corollary 60. *Let π be a resolution refutation of $\Gamma \cup \Delta$. Then $\Delta \models \ell_{\Gamma}^x[\neg \text{PI}(C) \vee C]$ for C in π .*

Proof. Build $\hat{\pi}$ from π using $\hat{\Gamma} = \Delta$ and $\hat{\Delta} = \Gamma$ as initial clause set partition.

By Lemma 58, $\hat{\Gamma} \models \ell_{\hat{\Delta}}^x[\text{PI}(\hat{C}) \vee \hat{C}]$ for \hat{C} in $\hat{\pi}$.

By Lemma 59, $\hat{\Gamma} \models \ell_{\hat{\Delta}}^x[\neg \text{PI}(C) \vee \hat{C}]$ for the clause C in π corresponding to \hat{C} in $\hat{\pi}$. This however is nothing else than $\Delta \models \ell_{\Gamma}^x[\neg \text{PI}(C) \vee C]$. \square

Theorem 61. *Let π be a resolution refutation of $\Gamma \cup \Delta$ and z_1, \dots, z_n be the variables which replace the colored terms in $\ell_\Gamma^y[\ell_\Delta^x[\text{PI}(\pi)]]$ ordered by their length. Then $Q_1 z_1 \dots Q_n z_n \ell_\Gamma^y[\ell_\Delta^x[\text{PI}(\pi)]]$, where Q_i is \forall (\exists) if z_i replaces a Δ (Γ)-term, is an interpolant.*

Proof. By Lemma 58, $\Gamma \models \forall x_1 \dots \forall x_m \ell_\Delta^x[\text{PI}(\pi)]$ where m is the number of maximal Δ -colored terms in $\text{PI}(\pi)$.

A term in $\ell_\Delta^x[\text{PI}(\pi)]$ is either x_i , $1 \leq i \leq m$, a grey term or a Γ -terms. Let t be a maximal Γ -term in $\ell_\Delta^x[\text{PI}(\pi)]$ and x_{j_1}, \dots, x_{j_k} the variables replacing Δ -terms in t . Note that the Δ -terms, which are replaced by x_{j_1}, \dots, x_{j_k} respectively, are each of strictly smaller size than t as they are strict subterms of t .

In $\ell_\Gamma^y[\ell_\Delta^x[\text{PI}(\pi)]]$, t is replaced by some z_j , which is existentially quantified. Hence t is a witness for z_j as due to the quantifier ordering, the existential quantification of z_j is in the scope of the quantifiers of x_{j_1}, \dots, x_{j_k} respectively. Therefore $\Gamma \models Q_1 z_1 \dots Q_n z_n \ell_\Gamma^y[\ell_\Delta^x[\text{PI}(\pi)]]$.

By Corollary 60 $\Delta \models \forall y_1 \dots \forall y_m \neg \ell_\Gamma^y[\text{PI}(\pi)]$, where m is the number of Γ -colored terms in $\text{PI}(\pi)$. By a similar line of argumentation as above, we can replace the maximal Δ -terms by existentially quantified variables and arrive at $\Delta \models \overline{Q}_1 z_1 \dots \overline{Q}_n z_n \neg \ell_\Delta^x[\ell_\Gamma^y[\text{PI}(\pi)]]$ where $\overline{Q}_i = \exists$ (\forall) if $Q_i = \forall$ (\exists). Therefore also $\Delta \models \neg Q_1 z_1 \dots Q_n z_n \ell_\Delta^x[\ell_\Gamma^y[\text{PI}(\pi)]]$. By Lemma 57 and as all variables which replace colored terms are bound, $\Delta \models \neg Q_1 z_1 \dots Q_n z_n \ell_\Gamma^y[\ell_\Delta^x[\text{PI}(\pi)]]$.

As it is now easy to see that $Q_1 z_1 \dots Q_n z_n \ell_\Gamma^y[\ell_\Delta^x[\text{PI}(\pi)]]$ contains no colored symbol, it is an interpolant. \square

Bibliography

- [BJ13] Maria Paola Bonacina and Moa Johansson. On Interpolation in Automated Theorem Proving. Technical Report 86/2012, Dipartimento di Informatica, Università degli Studi di Verona, 2013. Submitted to journal August 2013.
- [BL11] Matthias Baaz and Alexander Leitsch. *Methods of Cut-Elimination*. Trends in Logic. Springer, 2011.
- [CK90] C.C. Chang and H.J. Keisler. *Model Theory*. Studies in Logic and the Foundations of Mathematics. Elsevier Science, 1990.
- [Cra57a] William Craig. Linear Reasoning. A New Form of the Herbrand-Gentzen Theorem. *The Journal of Symbolic Logic*, 22(3):250–268, September 1957.
- [Cra57b] William Craig. Three Uses of the Herbrand-Gentzen Theorem in Relating Model Theory and Proof Theory. *The Journal of Symbolic Logic*, 22(3):269–285, September 1957.
- [DKPW10] Vijay D’Silva, Daniel Kroening, Mitra Purandare, and Georg Weissenbacher. Interpolant Strength. In *Proceedings of the International Conference on Verification, Model Checking, and Abstract Interpretation (VMCAI)*, volume 5944 of *Lecture Notes in Computer Science*, pages 129–145. Springer, January 2010.
- [Gen35] Gerhard Gentzen. Untersuchungen über das logische Schließen II. *Mathematische Zeitschrift*, 39, 1935.

-
- [Hua95] Guoxiang Huang. Constructing Craig Interpolation Formulas. In *Proceedings of the First Annual International Conference on Computing and Combinatorics*, COCOON '95, pages 181–190, London, UK, UK, 1995. Springer-Verlag.
- [Kle67] Stephen Cole Kleene. *Mathematical logic*. Wiley, New York, NY, 1967.
- [Kra97] Jan Krajíček. Interpolation Theorems, Lower Bounds for Proof Systems, and Independence Results for Bounded Arithmetic. *Journal of Symbolic Logic*, pages 457–486, 1997.
- [Lyn59] Roger C. Lyndon. An Interpolation Theorem in the Predicate Calculus. *Pacific Journal of Mathematics*, 9(1):129–142, 1959.
- [McM03] Kenneth L. McMillan. Interpolation and SAT-Based Model Checking. In Jr. Hunt, Warren A. and Fabio Somenzi, editors, *Computer Aided Verification*, volume 2725 of *Lecture Notes in Computer Science*, pages 1–13. Springer Berlin Heidelberg, 2003.
- [Pud97] Pavel Pudlák. Lower Bounds for Resolution and Cutting Plane Proofs and Monotone Computations. *J. Symb. Log.*, 62(3):981–998, 1997.
- [Rob65] J. A. Robinson. A machine-oriented logic based on the resolution principle. *J. ACM*, 12(1):23–41, January 1965.
- [Sho67] Joseph R. Shoenfield. *Mathematical logic*. Addison-Wesley series in logic. Addison-Wesley Pub. Co., 1967.
- [Sla70] James R. Slagle. Interpolation theorems for resolution in lower predicate calculus. *J. ACM*, 17(3):535–542, July 1970.
- [Tak87] Gaisi Takeuti. *Proof Theory*. Studies in logic and the foundations of mathematics. North-Holland, 1987.
- [Wei10] Georg Weissenbacher. *Program Analysis with Interpolants*. PhD thesis, 2010.